

CERTAIN IMPLICATIONS FOR HIGH ENERGY TOTAL  
CROSS SECTIONS FROM THE  
MULTIPERIPHERAL MODEL

Peter D. Ting

RECEIVED  
LAWRENCE  
RADIATION LABORATORY

December 31, 1968

JAN 30, 1969

LIBRARY  
DOCUMENTS

TWO-WEEK LOAN COPY

This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545

LAWRENCE RADIATION LABORATORY  
UNIVERSITY of CALIFORNIA BERKELEY

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Phys. Rev.

UCRL-18670  
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory  
Berkeley, California

AEC Contract No. W-7405-eng-48

CERTAIN IMPLICATIONS FOR HIGH ENERGY TOTAL CROSS SECTIONS  
FROM THE MULTIPERIPHERAL MODEL

Peter D. Ting

December 31, 1968

CERTAIN IMPLICATIONS FOR HIGH ENERGY TOTAL CROSS SECTIONS  
FROM THE MULTIPERIPHERAL MODEL\*

Peter D. Ting

Lawrence Radiation Laboratory  
University of California  
Berkeley, California

December 31, 1968

ABSTRACT

The multiperipheral model is used to relate the experimental fact,  $\sigma_{N\bar{N}}^{\text{tot}} > \sigma_{NN}^{\text{tot}}$ , to the possibility that the  $N\bar{N}$  channel can annihilate into purely meson final states by exchange of  $N$  or  $\bar{N}$  trajectories, whereas the  $NN$  channel cannot do so. Our argument shows that from the multi-Regge bootstrap point of view, the combination of baryon and antibaryon trajectories will generate a pair of exchange-degenerate trajectories  $P'$  and  $\omega$ . The same line of reasoning can easily be generalized to include strange particles, as well as the  $(K^-N, K^+N)$  and  $(\pi^-p, \pi^+p)$  systems.

## I. INTRODUCTION

It is a familiar empirical fact that, if a larger number of two-particle channels communicate with channel  $a\bar{b}$  than with channel  $ab$ , then the total cross section  $\sigma_{a\bar{b}}$  tends to be larger than  $\sigma_{ab}$  at nonasymptotic energies. For example, the  $p\bar{p}$  channel can communicate with  $p\bar{p}$ ,  $\pi^0\pi^0$ ,  $\pi^+\pi^-$ ,  $K^0\bar{K}^0$ ,  $K^+K^-$ ,  $\Lambda\bar{\Lambda}$ ,  $\Sigma^0\bar{\Sigma}^0$ , etc., whereas the  $pp$  channel can only communicate with  $pp$ . It is well known that  $\sigma_{p\bar{p}} > \sigma_{pp}$ . A similar correlation exists for the paired channels,  $(K^-p, K^+p)$ ,  $(\pi^-p, \pi^+p)$ . In this note we propose to relate these phenomena to the generalized multiperipheral model recently developed by Chew, Goldberger, and Low<sup>1</sup> (CGL). The type of reasoning to be applied is illustrated by the following argument: If more two-particle channels communicate with  $a\bar{b}$  than with  $ab$ , we will get more unitarity box diagrams (two-particle unitarity contributions) in  $a\bar{b}$  than  $ab$ . The multiperipheral model iterates the box diagrams to produce a series of multiparticle contributions to the cross section (i.e., to the forward-elastic unitarity integral). Since each such contribution is positive definite, it follows that the channel with the larger number of box diagrams has the larger total cross section, if we can demonstrate an equal magnitude for those box diagrams shared by the two channels.

In Section II the multiperipheral model of CGL has been generalized to exchange several trajectories. In Section III the multiperipheral model is applied to  $NN$  and  $N\bar{N}$  channels. Some Reggeon bootstrap arguments are given in Section IV. In the last section some general remarks are made to include the strangeness and the isospin in the model.

## II. THE MULTIPERIPHERAL MODEL

Let us briefly review the multiperipheral model. A function called  $B_a$  was introduced in CGL by

$$\begin{aligned} A_{ab}(p_a, p_b) &\equiv A_{ab}(s, 0) \\ &= \int B_a(p_a, p_b; Q') (G^b(Q'))^2 \delta^+[(Q' - p_0)^2 - \mu'^2] d^4Q' , \end{aligned} \quad (1)$$

where  $A_{ab}(p_a, p_b)$  is the absorptive part of the  $ab \rightarrow ab$  elastic amplitude at forward direction. The function  $B_a$  satisfies the integral equation

$$\begin{aligned} B_a(p_a, p_b; Q') &= I_a(p_a, p_b; Q') + \int B_a(p_a, Q'; Q) K(Q, Q', p_b) \delta^+[(Q - Q')^2 - \mu^2] d^4Q , \end{aligned} \quad (2)$$

where  $I_a$  corresponds to two-particle unitarity, namely

$$I_a(p_a, p_b; Q') = (G^a(Q'))^2 |f(p_a, Q', p_b)|^2 , \quad (3)$$

while the kernel  $K$  is given by

$$K(Q, Q', p_b) = \beta^2(Q, Q') |f(Q, Q', p_b)|^2 . \quad (4)$$

In a multi-Regge model  $G$  and  $\beta$  are the external and internal vertex couplings respectively, and  $f$  is the factor  $\left(\frac{s_i}{s_0}\right)^\alpha$ . The kernel

can take other forms in other multiperipheral models. Equation (1) and Eq. (2) can be written symbolically as

$$A_{ab} = \int B_a G^b \quad \text{and} \quad B_a = I_a + \int B_a K .$$

The CGL equations assumed only one kind of trajectory to be exchanged. If several Reggeons  $\alpha_1, \alpha_2, \dots, \alpha_m$  are exchanged, these equations should be written

$$A_{ab} = \sum_x \sum_{i=1}^m \int B_a^{\alpha_i} (G_x^{\alpha_i b})^2 , \quad (5)$$

$$B_a^{\alpha_i} = I_a^{\alpha_i} + \sum_y \sum_{j=1}^m \int B_a^{\alpha_j} K_y^{\alpha_j \alpha_i} , \quad (6)$$

with

$$I_a^{\alpha_i} = \sum_z (G_z^{\bar{a}\alpha_i})^2 |f^{\alpha_i}|^2 \quad (\text{see Fig. 1b}) , \quad (7)$$

$$K_y^{\alpha_j \alpha_i} = \left( \beta_y^{\alpha_j \alpha_i} \right)^2 |f^{\alpha_i}|^2 , \quad (8)$$

where

$G_x^{\alpha_i b}$  is the external coupling of particles b and x to the Reggeon  $\alpha_i$ , in the sense of Fig. 1a, and similarly for  $G_z^{\bar{a}\alpha_i}$ ;

$\beta_y^{\alpha_j \alpha_i}$  is the internal coupling of the two Reggeons  $\alpha_i$  and  $\alpha_j$  to the particle  $y$ , in the sense of Fig. 1b; and

$\sum_x$  means to sum over all stable particles  $x$  which conserve baryon number, strangeness, and isospin<sup>2</sup> at the end vertex coupling  $b$ , and similarly for the symbols  $\sum_y$  and  $\sum_z$ .



### III. MULTIPERIPHERAL MODEL IN NN AND $N\bar{N}$ CHANNELS

For the sake of simplicity and clarity, we begin with the special case of NN and  $N\bar{N}$  elastic scattering and include as exchanged trajectories only the nucleon N (or  $\Delta$ ), the antinucleon  $\bar{N}$  (or  $\bar{\Delta}$ ), and an averaged effective nonstrange meson M (including both signatures).<sup>3</sup> The justification for using only one kind of meson is that the important nonstrange mesons like P',  $\rho$ ,  $\omega$ , and  $A_2$  have roughly the same intercept and slope at  $t = 0$  on the Chew-Frautschi plot. The Pomeron P is different but there are arguments<sup>4,5</sup> against the importance of Pomeron repetition in the multiperipheral chain. Of course in the lowest-order unitarity box diagrams, we have to consider the Pomeron as an important trajectory. Hence, in the following discussion the Pomeron appears only in unitarity box diagrams. To make this feature explicit, we rewrite Eqs. (5) and (6) as

$$A_{NN}^i = A_{NN}^{iP} + \sum_{x, i=M, N, \bar{N}} \int B_N^i (G_x^{iN})^2, \quad (9a)$$

$$A_{N\bar{N}}^i = A_{N\bar{N}}^{iP} + \sum_{x, i=M, N, \bar{N}} \int B_N^i (G_x^{i\bar{N}})^2, \quad (9b)$$

$$B_N^i = I_N^i + \sum_{y, j=M, N, \bar{N}} \int B_N^j K_y^{ji}, \quad i = M, N, \bar{N}, \quad (10)$$

where

$$A_{ab}^P = \int \left[ |T_{ab}^P|^2 + (T_{ab}^P)^*(T_{ab}^M) + (T_{ab}^M)^*(T_{ab}^P) \right] \quad (11)$$

and

$$T_{ab}^\alpha = \sum_x I_a^\alpha (G_x^{\alpha b})^2 = \sum_{z,x} (G_z^{\alpha a})^2 |f^\alpha|^2 (G_x^{\alpha b})^2, \quad (12)$$

where  $T_{ab}^\alpha$  is the part of the elastic amplitude where only the  $\alpha$  trajectory is exchanged, and P denotes "Pomeron" and M our average effective meson.

Now we propose to argue that  $A_{NN}^P \approx A_{N\bar{N}}^P$ . From the experimental data, the total elastic cross sections<sup>6</sup> for NN and  $N\bar{N}$  are roughly equal<sup>7</sup> for  $p_L > 2$  GeV/c, i.e.,  $\sigma_{NN}^{el} \approx \sigma_{N\bar{N}}^{el}$ , but by definition

$$\sigma_{NN}^{el} \equiv A_{NN}^P + \int |T_{NN}^M|^2,$$

$$\sigma_{N\bar{N}}^{el} \equiv A_{N\bar{N}}^P + \int |T_{N\bar{N}}^M|^2.$$

Thus our first task is to show that  $\int |T_{NN}^M|^2 = \int |T_{N\bar{N}}^M|^2$ .

Rewrite

$$T_{ab}^M = T_{ab}^{M(+)} + T_{ab}^{M(-)},$$

where

$M(+)$  denote the effective meson trajectory with signature +,

$M(-)$  denote the effective meson trajectory with signature -.

Now  $M(+)$  and  $M(-)$  have the same intercept and slope at  $t = 0$ , so the usual Regge signature factor  $1 \pm e^{-i\pi\alpha}$  leads to a 90-degree phase difference between  $T_{ab}^{M(+)}$  and  $T_{ab}^{M(-)}$ . Thus  $|T_{ab}^{M(\pm)}|^2 = |T_{ab}^{M(+)}|^2 + |T_{ab}^{M(-)}|^2$ . Then, since the CPT theorem implies  $|T_{ab}^{M(\pm)}|^2 = |T_{ab}^{M(\pm)}|^2$ , it follows that  $A_{NN}^P = A_{\overline{NN}}^P$ .

Next if we subtract Eq. (9a) from (9b), we get

$$A_{\overline{NN}} - A_{NN} = \sum_{i=M,N,\overline{N}} \int B_N^i \left[ \sum_x (G_x^{i\overline{N}})^2 - \sum_{x'} (G_{x'}^{iN})^2 \right]. \quad (13)$$

From the dearth of stable  $B = 2$ ,  $s = 0$  baryons produced in high energy collisions and conversely the abundance of nonstrange meson ( $B = 0$ ,  $s = 0$ ) production, we conclude that  $G_x^{\overline{NN}} \approx 0$ ,  $K_y^{\overline{NN}} \approx 0$ , and  $I_N^{\overline{N}} \approx 0$ . By the CPT theorem we also know  $(G_x^{\overline{NN}})^2 = (G_x^{\overline{NN}})^2$ ,  $(G_x^{\overline{NN}})^2 = (G_x^{\overline{NN}})^2$ ,  $\alpha_N = \alpha_{\overline{N}}$ ,  $K_x^{\overline{NN}} = K_y^{\overline{NN}}$ , and  $K_y^{\overline{NN}} = K_x^{\overline{NN}}$ .

Equation (10) and Eq. (13) can then be reduced to

$$A_{\overline{NN}} - A_{NN} \approx \int B_N^M (G_N^{M\overline{N}})^2 - (G_N^{MN})^2 + \sum_x \int (B_N^{\overline{N}} - B_N^N) (G_x^{\overline{NN}})^2, \quad (14)$$

$$B_N^{\overline{N}} - B_N^N \approx I_N^{\overline{N}} + \int B_N^M (K_N^{M\overline{N}} - K_N^{MN}) + \sum_y \int (B_N^{\overline{N}} - B_N^N) K_y^{\overline{NN}}. \quad (15)$$

Because the symbol  $M$  in our model contains both positive and negative signature mesons, in general the CPT theorem will not give

$(G_N^{MN})^2 = (G_N^{\overline{MN}})^2$ ,  $K_N^{MN} = K_N^{\overline{MN}}$ , and  $K_N^{\overline{NM}} = K_N^{\overline{NM}}$ . However, from the rough equality  $\sigma_{NN}^{el} \approx \sigma_{\overline{NN}}^{el}$ , exchange degeneracy among mesons, and other experimental results,<sup>8,9,10</sup> we believe it would be a reasonable 0th order approximation to take  $\int_{B_N^M} (G_N^{MN})^2 \approx \int_{B_N^M} (G_N^{\overline{MN}})^2$  and  $\int_{B_N^M} K_N^{MN} \approx \int_{B_N^M} K_N^{\overline{MN}}$  as our input. Then Eqs. (14) and (15) will become

$$A_{\overline{NN}} - A_{NN} = \sum_x \int (B_N^{\overline{N}} - B_N^N) (G_x^{\overline{NN}})^2, \quad (16)$$

$$B_N^{\overline{N}} - B_N^N = L_N^{\overline{N}} + \sum_y \int (B_N^{\overline{N}} - B_N^N) K_y^{\overline{NN}}. \quad (17)$$

Since all quantities in Eq. (16) and Eq. (17) are evaluated at forward direction both  $L_N^{\overline{N}}$  and  $K_y^{\overline{NN}}$  are positive definite; Eq. (17) can be solved by iteration, so both  $(B_N^{\overline{N}} - B_N^N)$  and  $(A_{\overline{NN}} - A_{NN})$  must be positive. From the optical theorem we have thus shown that  $\sigma_{\overline{NN}}^- > \sigma_{NN}^-$  at nonasymptotic energies.

The physical meaning behind Eqs. (16) and (17) is very simple. Within our model the difference  $\sigma_{\overline{NN}}^- - \sigma_{NN}^-$  arises because the  $\overline{NN}$  channel can annihilate into pure meson final states<sup>11</sup> by exchange of  $N$  (or  $\overline{N}$ ) trajectories while the  $NN$  channel cannot.

## IV. REGGEON BOOTSTRAP

In order to calculate  $A_{\overline{NN}} - A_{NN}$  explicitly we must assume explicit forms for the kernel  $K_y^{\overline{NN}}$  as well as for  $I_N^{\overline{N}}$  and  $G^{\overline{NN}}$ , a step we do not propose to take here. However, we wish to discuss this problem from the bootstrap point of view. In the usual Regge theory, if we neglect isospin dependence in  $\overline{NN}$  and  $NN$ , Regge fitting says

$$A_{\overline{NN}} \approx \beta_P \left( \frac{s}{s_0} \right)^{\alpha_P(0)} + \beta_{P'} \left( \frac{s}{s_0} \right)^{\alpha_{P'}(0)} + \beta_\omega \left( \frac{s}{s_0} \right)^{\alpha_\omega(0)}, \quad (18)$$

$$A_{NN} \approx \beta_P \left( \frac{s}{s_0} \right)^{\alpha_P(0)} + \beta_{P'} \left( \frac{s}{s_0} \right)^{\alpha_{P'}(0)} - \beta_\omega \left( \frac{s}{s_0} \right)^{\alpha_\omega(0)}, \quad (19)$$

at  $t = 0$ , where  $s_0$  is a constant. Then

$$A_{\overline{NN}} - A_{NN} \approx 2\beta_\omega \left( \frac{s}{s_0} \right)^{\alpha_\omega(0)}. \quad (20)$$

Comparing Eq. (20) with Eqs. (16) and (17), we get a result of interest to bootstrap arguments, since it can be proved that equations of the latter form lead to Regge poles. According to Eqs. (16) and (17) our model forms a combination of baryon and antibaryon trajectories. The meson-meson contribution, we suggest, is small.

Furthermore, we notice that in high energy production process, mainly mesons are produced;  $(\overline{NN})$  pairs are rare.<sup>12</sup> This implies

$\beta_N^{\bar{N}M}, \beta_N^{NM} \ll \beta_x^{MM}$ , so  $K_N^{JM}, K_N^{NM} \ll K_{y'}^{MM}$ , and

$K_N^{MN}, K_N^{M\bar{N}} \ll K_y^{\bar{N}\bar{N}}$ . If we accept this experimental fact, the meson

and nucleon channels are decoupled. That is, from Eqs. (9a), (9b),

and (10), we get

$$A_{\bar{N}\bar{N}} + A_{NN} \approx 2A_{NN}^P + 2 \int B_N^M (G_N^{\bar{M}\bar{N}})^2 + \sum_x \int (B_N^{\bar{N}} + B_N^N) (G_x^{\bar{N}\bar{N}})^2, \quad (21)$$

$$\begin{aligned} B_N^M &\approx I_N^M + \sum_{y'} \int B_N^M K_{y'}^{MM} + \int (B_N^{\bar{N}} K_N^{\bar{N}M} + B_N^N K_N^{NM}) \\ &\approx I_N^M + \sum_{y'} \int B_N^M K_{y'}^{MM}, \end{aligned} \quad (22)$$

$$\begin{aligned} (B_N^{\bar{N}} + B_N^N) &\approx I_N^{\bar{N}} + \int B_N^M (K_N^{MN} + K_N^{M\bar{N}}) + \sum_y \int (B_N^{\bar{N}} + B_N^N) K_y^{\bar{N}\bar{N}} \\ &\approx I_N^{\bar{N}} + \sum_y \int (B_N^{\bar{N}} + B_N^N) K_y^{\bar{N}\bar{N}}. \end{aligned} \quad (23)$$

By adding Eqs. (18) and (19), we get

$$A_{\bar{N}\bar{N}} + A_{NN} \approx 2\beta_P \left(\frac{s}{s_0}\right)^{\alpha_P(0)} + 2\beta_{P'} \left(\frac{s}{s_0}\right)^{\alpha_{P'}(0)}. \quad (24)$$

Because the kernels of Eqs. (23) and (17) are identical, it follows that besides the  $\omega$  trajectory, the combination of  $N$  and  $\bar{N}$  trajectories will generate another meson trajectory  $\alpha_P$ , which has the same intercept and slope as  $\omega$ . The  $\alpha_P$  contributes only to  $(A_{\bar{N}N} + A_{N\bar{N}})$ , not to  $(A_{\bar{N}N} - A_{N\bar{N}})$ , whereas for  $\omega$  the converse is true; i.e.,  $\alpha_P$  has positive signature,  $\alpha_\omega$  has negative signature. Thus our model gives an explanation of the exchange degeneracy between  $\alpha_P$  and  $\alpha_\omega$ .

The effective meson trajectory  $M$  may be expected to generate an even signature Regge pole, which Chew and Pignotti<sup>4</sup> (CP) have identified with Pomeron. For more discussion of bootstrap questions, see CGL<sup>1</sup> and CP.<sup>4</sup>

### V. SUMMARY AND DISCUSSION

The above line of reasoning is easily generalized to include strange particles. The key points in our previous argument, leading to partial approximate diagonalization of the multiperipheral kernel, were the low rates of production of  $B = 2$  particles and  $B\bar{B}$  pairs. These points remain true in general. Furthermore, the low rate of production of mesons with  $B = 0$ ,  $|s| = 2$ ;  $B = 0$ ,  $s = 0$ ,  $I = 2$ , together with the fact that the nucleon and kaon have  $I = 1/2$ , allows a partial diagonalization with respect to  $A_{KN}^- + A_{KN}$  and with respect to  $A_{\pi N}^- + A_{\pi N}$ , by arguments which completely parallel the preceding. We then can conclude that  $\sigma_{K^-p} > \sigma_{K^+p}$ ,  $\sigma_{\pi^-p} > \sigma_{\pi^+p}$ . We do not attempt here to exhaust all applications of our general argument. The principle should be clear from the above examples.



ACKNOWLEDGMENTS

It is a pleasure to thank Professor Geoffrey Chew for suggesting this problem and for his unfailing advice and encouragement. I would also like to thank Dr. A. Pignotti, <sup>Dr.</sup> G. Fox, and Mr. C. K. Chen for helpful discussions.

## FOOTNOTES AND REFERENCES

- \* This work was supported in part by the U.S. Atomic Energy Commission.
1. G. Chew, M. Goldberger, and F. Low, An Integral Equation for Scattering Amplitudes, Princeton University preprint, 1968.
  2. We treat all particles as spinless. This is equivalent to summing and averaging over spin orientations.
  3. For simplicity we neglect isospin in the calculation, because the isospin dependence of  $NN$  and  $N\bar{N}$  elastic amplitude is very weak, but we should consider isospin in calculating  $A_{\pi^+p}$  and  $A_{\pi^-p}$ .
  4. G. Chew and A. Pignotti, A Multiperipheral Bootstrap Model, Lawrence Radiation Laboratory Report UCRL-18275 (to be published in Phys. Rev.).
  5. Chew and Pignotti, (Ref. 4) have argued that the internal coupling of the Pomeron is small compared with internal couplings which involve  $P'$ ,  $\rho$ ,  $\omega$ , and  $A_2$ . Some physicists even think that, because the Pomeron trajectory is associated with diffractive phenomena, multiple Pomeron exchange is meaningless.
  6. Yoshio Sumi and Toshihiro Yoshida, Suppl. of Progr. Theoret. Phys. (Kyoto) 41 and 42, 53 (1967).
  7. The combination of  $\sigma_{N\bar{N}} > \sigma_{NN}$  with  $\sigma_{N\bar{N}}^{el} \approx \sigma_{NN}^{el}$  implies the crossover in the elastic scattering angular distribution. We wish to thank Dr. G. Fox for pointing out this fact.
  8. Atsuko Iwaki, Suppl. of Progr. Theoret. Phys. (Kyoto) 41 and 42, 316 (1967).

9. G. Alexander, O. Benary, and U. Maor, Nucl. Phys. B5, 1 (1968).  
 10. From Ref. (8) and Ref. (9), we get

$$\sum_{m=1}^{\infty} \sigma_{pp \rightarrow NN+m\pi} \approx \sum_{m=1}^{\infty} \sigma_{pp \rightarrow NN+m\pi} \quad \text{for } 2 \text{ GeV/c} < p_L \quad ,$$

$$\sigma_{pp \rightarrow pp\pi^+\pi^-} \approx \sigma_{pp \rightarrow pp\pi^+\pi^-} \quad \text{for } 2 \text{ GeV/c} < p_L \quad ,$$

$$\sigma_{pp \rightarrow NN+3\pi} \approx \sigma_{pp \rightarrow NN+3\pi} \quad \text{for } 2 \text{ GeV/c} < p_L \quad ,$$

$$\sum_{m=2}^{\infty} \sigma_{pp \rightarrow m\pi} \approx (\sigma_{pp} - \sigma_{pp}) \quad 3 \text{ GeV/c} < p_L < 7 \text{ GeV/c} \quad ,$$


$$\sum_{m=2}^{\infty} \sigma_{pp \rightarrow m\pi} \gg \sum_{m=1}^{\infty} (\sigma_{pp \rightarrow NN+m\pi} - \sigma_{pp \rightarrow NN+m\pi})$$

$$3 \text{ GeV/c} < p_L < 7 \text{ GeV/c} \quad .$$

11. Because we have not considered  $K^*$  and  $Y^*$  (or  $\Lambda$ ,  $\Sigma$ ) exchange in this calculation, we neglect those final states which involve strange particles. See the concluding paragraph of this paper.  
 12. V. Barashenkov, M. Valtsev, I. Patera, and V. Toneev, Fortschr. Phys. 14, 357 (1966).

FIGURE CAPTIONS

- Fig. 1.(a) The external couplings  $G_x^{\alpha_i b}$  and  $G_z^{\bar{a}\alpha_i}$ .
- (b) The internal coupling  $\beta_y^{\alpha_j \alpha_i}$ , and the two-particle unitarity contribution  $I_a^{\alpha_i}$ .



XBL691-1679

Fig. 1

LEGAL NOTICE

*This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:*

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or*
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.*

*As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.*

TECHNICAL INFORMATION DIVISION  
LAWRENCE RADIATION LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720