

UCSF

UC San Francisco Previously Published Works

Title

Hospitalist utilization and hospital performance on 6 publicly reported patient outcomes

Permalink

<https://escholarship.org/uc/item/21w5c60j>

Journal

Journal of Hospital Medicine, 7(6)

ISSN

1553-5592

Authors

Goodrich, Kate
Krumholz, Harlan M
Conway, Patrick H
et al.

Publication Date

2012-07-01

DOI

10.1002/jhm.1943

Peer reviewed



Published in final edited form as:

J Hosp Med. 2012 ; 7(6): 482–488. doi:10.1002/jhm.1943.

Hospitalist Utilization and Hospital Performance on Six Publicly Reported Patient Outcomes

Kate Goodrich, MD, MHS^{1,2}, Harlan Krumholz, MD, SM^{3,4}, Patrick Conway, MD, MSc^{1,5}, Peter Lindenauer, MD, MSc^{6,7}, and Andrew D. Auerbach, MD, MPH⁸

¹Office of Clinical Standards and Quality, Centers for Medicare and Medicaid Services, Baltimore, MD

²Division of Hospital Medicine, Department of Medicine, George Washington University School of Medicine, Washington, DC

³Section of Health Policy and Administration, Yale School of Public Health, Center for Outcomes Research and Evaluation, Yale-New Haven Hospital, New Haven, CT

⁴Section of Cardiovascular Medicine and the Robert Wood Johnson Clinical Scholars Program, Department of Internal Medicine, Yale University School of Medicine, New Haven, CT

⁵Division of Hospital Medicine, Cincinnati Children's Hospital Medical Center, Cincinnati, OH

⁶Center for Quality of Care Research, Baystate Medical Center, Springfield, MA

⁷Department of Medicine, Tufts University School of Medicine, Springfield, MA

⁸Division of Hospital Medicine, Department of Medicine, University of California, San Francisco, San Francisco, CA

Abstract

BACKGROUND—The increase in hospitalist-provided inpatient care may be accompanied by an expectation of improvement on patient outcomes. To date the association between utilization of hospitalists and the publicly reported patient outcomes is unknown.

OBJECTIVE—Assess the relationship between hospitalist utilization and performance on six publicly reported patient outcomes.

DESIGN—Cross-sectional study

PARTICIPANTS—Representatives of 598 hospitals in the United States with direct knowledge of inpatient service models.

INTERVENTION—Survey of hospital personnel with knowledge of hospitalist use and hospitalist programs

MEASUREMENTS—Six publicly reported quality outcome measures across 3 medical conditions: acute myocardial infarction (AMI); congestive heart failure (HF); and pneumonia. Using multivariable regression models, we assessed the relationship between presence of hospitalists and performance on each outcome measures; we further assessed the relationship between the percentage of patients admitted by hospitalists and each outcome measure.

Disclosures

Dr. Krumholz is the chair of the Cardiac Scientific Advisory Board for United Health and has a research grant with Medtronic through Yale University; Dr. Auerbach has a grant through the National Heart, Lung and Blood Institute (NHLBI). The authors have no other disclosures to report.

RESULTS—Of 598 respondents, 429 (72%) reported the use of hospitalist services. In the comparison of hospitals with and without hospitalists, there was no statistically significant difference on any of the mortality or readmissions measures with the exception of the risk-stratified readmission rate for heart failure. For hospitals that used hospitalists, there was no significant change in any of the outcome measures with increasing percentage of patients admitted by hospitalists.

CONCLUSIONS—The presence of hospitalists is not an independent predictor of performance on publicly reported mortality and readmissions measures for AMI, HF or pneumonia. It is likely that broader system or organizational interventions are required to improve performance on patient outcomes.

Keywords

acute myocardial infarction; cross-sectional studies; heart failure; hospitalist; hospital medicine; outcome measures; pneumonia; quality of care

Introduction

The past several years have seen a dramatic increase in the percentage of patients cared for by hospitalists, yet an emerging body of literature examining the association between care given by hospitalists and performance on a number of process measures has shown mixed results. Hospitalists do not appear to provide higher quality of care for pneumonia [1,2] while results in heart failure are mixed [3-5]. Each of these studies was conducted at a single site, and examined patient-level effects. More recently, Vasilevskis et al assessed the association between the intensity of hospitalist use (measured as the percent of patients admitted by hospitalists) and performance on process measures. In a cohort of 208 California hospitals they found a significant improvement in performance on process measures in patients with acute myocardial infarction, heart failure and pneumonia with increasing percentages of patients admitted by hospitalists[6].

To date, no study has examined the association between the use of hospitalists and the publicly reported 30-day mortality and readmission measures. Specifically, the Centers for Medicare and Medicaid Services (CMS) has developed and now publicly reports risk-standardized 30-day mortality (RSMR) and readmission rates (RSRR) for Medicare patients hospitalized for three common and costly conditions – acute myocardial infarction (AMI), heart failure (HF), and pneumonia[7]. Performance on these hospital-based quality measures varies widely, and vary by hospital volume, ownership status, teaching status, and nurse staffing levels[8-13]. However, even accounting for these characteristics leaves much of the variation in outcomes unexplained. We hypothesized that the presence of hospitalists within a hospital would be associated with higher performance on 30-day mortality and 30-day readmission measures for AMI, HF, and pneumonia. We further hypothesized that for hospitals using hospitalists, there would be a positive correlation between increasing percentage of patients admitted by hospitalists and performance on outcome measures. To test these hypotheses, we conducted a national survey of hospitalist leaders, linking data from survey responses to data on publicly reported outcome measures for AMI, HF, and pneumonia.

Materials and Methods

Study Sites

Of the 4289 hospitals in operation in 2008, 1945 had 25 or more AMI discharges. We identified hospitals using American Hospital Association (AHA) data, calling hospitals up to

6 times each until we reached our target sample size of 600. Using this methodology, we contacted 1558 hospitals of a possible 1920 with AHA data; of the 1558 called, 598 provided survey results.

Survey Data

Our survey was adapted from the survey developed by Vasilevskis et al [6]. The entire survey can be found in the Appendix. Our key questions were: 1) Does your hospital have at least one hospitalist program or group? 2) Approximately what percent of all medical patients in your hospital are admitted by hospitalists? The latter question was intended as an approximation of the intensity of hospitalist use, and has been used in prior studies [6,14]. A more direct measure was not feasible given the complexity of obtaining admission data for such a large and diverse set of hospitals. Respondents were also asked about hospitalist care of AMI, HF and pneumonia patients. Given the low likelihood of precise estimation of hospitalist participation in care for specific conditions, the response choices were divided into percentage quartiles: 0 to 25, 26-50, 51-75, and 76 to 100. Finally, participants were asked a number of questions regarding hospitalist organizational and clinical characteristics.

Survey Process

We obtained data regarding presence or absence of hospitalists and characteristics of the hospitalist services via phone- and fax-administered survey (see Appendix). Telephone and faxed surveys were administered between February 2010 and January 2011. Hospital telephone numbers were obtained from the 2008 AHA survey database and from a review of each hospital's website. Up to six attempts were made to obtain a completed survey from non-respondents unless participation was specifically refused. Potential respondents were contacted in the following order: hospital medicine department leaders, hospital medicine clinical managers, vice president for medical affairs, chief medical officers and other hospital executives with knowledge of the hospital medicine services. All respondents agreed with a question asking whether they had direct working knowledge of their hospital medicine services; contacts who said they did not have working knowledge of their hospital medicine services were asked to refer our surveyor to the appropriate person at their site. Absence of a hospitalist program was confirmed by contacting the Medical Staff Office.

Hospital Organizational and Patient-Mix Characteristics

Hospital-level organizational characteristics (e.g. bed size, teaching status) and patient-mix characteristics (e.g. Medicare and Medicaid inpatient days) were obtained from the 2008 AHA survey database.

Outcomes Performance Measures

The 30-day risk-standardized mortality and readmission rates (RSMR and RSRR) for 2008 for AMI, HF and pneumonia were calculated for all admissions for people age 65 and over with traditional fee-for-service Medicare. Beneficiaries had to be enrolled for 12 months prior to their hospitalization for any of the 3 conditions, and had to have complete claims data available for that 12 month period [7]. These six outcomes measures were constructed using hierarchical generalized linear models[15-20]. Using the RSMR for AMI as an example, for each hospital the measure is estimated by dividing the predicted number of deaths within 30 days of admission for AMI by the expected number of deaths within 30 days of admission for AMI. This ratio is then divided by the national unadjusted 30-day mortality rate for AMI, which is obtained using data on deaths from the Medicare beneficiary denominator file. Each measure is adjusted for patient characteristics such as age, gender and co-morbidities. All six measures are endorsed by the National Quality Forum (NQF) and are reported publicly by CMS on the Hospital Compare website.

Statistical Analysis

Comparison of hospital- and patient-level characteristics between hospitals with and without hospitalists was performed using Chi-square tests and Student's t-tests.

The primary outcome variables are the RSMRs and RSRRs for AMI, HF and pneumonia. Multivariable linear regression models were used to assess the relationship between hospitals with at least one hospitalist group and each dependent variable. Models were adjusted for variables previously reported to be associated with quality of care. Hospital-level characteristics included core-based statistical area, teaching status, number of beds, region, safety-net status, nursing staff ratio (# registered nurse FTEs/# hospital FTEs), and presence or absence of cardiac catheterization and coronary bypass capability. Patient-level characteristics included Medicare and Medicaid inpatient days as a percentage of total inpatient days and percent of admissions by race (black vs. non-black). The presence of hospitalists was correlated with each of the hospital and patient-level characteristics. Further analyses of the subset of hospitals that use hospitalists included construction of multivariable linear regression models to assess the relationship between the percentage of patients admitted by hospitalists and the dependent variables. Models were adjusted for the same patient- and hospital-level characteristics.

The institutional review boards at Yale University and University of California, San Francisco approved the study. All analyses were performed using Statistical Analysis Software (SAS) version 9.1 (SAS Institute, Inc., Cary, NC).

Results

Characteristics of Participating Hospitals

Telephone, fax and email surveys were attempted with 1558 hospitals; we received 598 completed surveys for a response rate of 40%. There was no difference between responders and non-responders on any of the six outcome variables, the number of Medicare or Medicaid inpatient days and the percent of admissions by race. Responders and non-responders were also similar in size, ownership, safety net and teaching status, nursing staff ratio, presence of cardiac catheterization and coronary bypass capability and core-based statistical area. They differed only on region of the country, where hospitals in the West North Central and Pacific regions of the country had larger overall proportions of respondents. All hospitals provided information about the presence or absence of hospitalist programs. The majority of respondents were hospitalist clinical or administrative managers (n=220) followed by hospitalist leaders (n=106), other executives (n=58), vice-presidents for medical affairs (n=39) and chief medical officers (n=15). Each respondent indicated a working knowledge of their site's hospitalist utilization and practice characteristics. Absence of hospitalist utilization was confirmed by contact with the Medical Staff Office.

Comparisons of Sites with Hospitalists and Those without Hospitalists

Hospitals with and without hospitalists differed by a number of organizational characteristics (Table 1). Sites with hospitalists were more likely to be larger, non-profit teaching hospitals, located in metropolitan regions, and have cardiac surgical services. There was no difference in the hospitals' safety-net status or RN staffing ratio. Hospitals with hospitalists admitted lower percentages of black patients.

Characteristics of Hospitalist Programs and Responsibilities

Of the 429 sites reporting use of hospitalists, the median percentage of patients admitted by hospitalists was 60% with an inter-quartile range (IQR) of 35% to 80%. The median number of full time equivalent hospitalists per hospital was 8 with an IQR of 5 to 14. The IQR

reflects the middle 50% of the distribution of responses, and is not affected by outliers or extreme values. Additional characteristics of hospitalist programs can be found in Table 2. The estimated percent of patients with AMI, HF and pneumonia cared for by hospitalists varied considerably, with fewer patients with AMI and more patients with pneumonia under hospitalist care. Overall a majority of hospitalist groups provided the following services: care of critical care patients, emergency department admission screening, observation unit coverage, coverage for cardiac arrests and rapid response teams, quality improvement or utilization review activities, development of hospital practice guidelines and participation in implementation of major hospital system projects (such as implementation of an electronic health record system).

Relationship between Hospitalist Utilization and Outcomes

Tables 3 and 4 show the comparisons between hospitals with and without hospitalists on each of the six outcome measures. In the bivariate analysis (Table 3), there was no statistically significant difference between groups on any of the outcomes measures with the exception of the risk-stratified readmission rate for heart failure. Sites with hospitalists had a lower RSRR for HF than sites without hospitalists (24.7% vs. 25.4%, $p < 0.0001$). These results were similar in the multivariable models as seen in Table 4, in which the beta estimate (slope) was not significantly different for hospitals utilizing hospitalists compared to those that did not on all measures except the RSRR for HF. For the subset of hospitals that used hospitalists, there was no statistically significant change in any of the six outcome measures with increasing percentage of patients admitted by hospitalists. Table 5 demonstrates that for each RSMR and RSRR, the slope did not consistently increase or decrease with incrementally higher percentages of patients admitted by hospitalists and the confidence intervals for all estimates crossed zero.

Discussion

In this national survey of hospitals, we did not find a significant association between the use of hospitalists and hospitals' performance on 30-day mortality or readmissions measures for AMI, HF or pneumonia. While there was a statistically lower 30-day risk-standardized readmission rate measure for the heart failure measure among hospitals that use hospitalists, the effect size was small. The survey response rate of 40% is comparable to other surveys of physicians and other health care personnel; however, there were no significant differences between responders and non-responders so the potential for response bias, while present, is small.

Contrary to the findings of a recent study [21], we did not find a higher readmission rate for any of the 3 conditions in hospitals with hospitalist programs. One advantage of our study is the use of more robust risk-adjustment methods. Our study used NQF-endorsed risk-standardized measures of readmission which capture readmissions to any hospital for common, high priority conditions where the impact of care coordination and discontinuity of care are paramount. The models use administrative claims data, but have been validated by medical record data. Another advantage is that our study focused on a time period when hospital readmissions were a standard quality benchmark and increasing priority for hospitals, hospitalists, and community-based care delivery systems. While our study is not able to discern whether patients had primary care physicians or the reason for admission to a hospitalist's care, our data do suggest that hospitalists continue to care for a large percentage of hospitalized patients. Moreover, increasing the proportion of patients being admitted to hospitalists did not affect the risk for readmission, providing perhaps reassuring evidence (or lack of proof) for a direct association between use of hospitalist systems and higher risk for readmission.

While hospitals with hospitalists clearly did not have better mortality or readmission rates, an alternate viewpoint might hold that, despite concerns that hospitalists negatively impact care continuity, our data do not demonstrate an association between readmission rates and use of hospitalist services. It is possible that hospitals that have hospitalists may have more ability to invest in hospital-based systems of care [22], an association which may incorporate any 'hospitalist effect', but our results were robust even after testing whether adjustment for hospital factors (such as profit status, size) affected our results.

It is also possible that secular trends in hospitals or hospitalist systems affected our results. A handful of single-site studies carried out soon after the hospitalist model's earliest descriptions found a reduction in mortality and readmission rates with the implementation of a hospitalist program [23-25]. Alternatively, it may be that there has been a dilution of the effect of hospitalists as often occurs when any new innovation is spread from early adopter sites to routine practice. Consistent with other multicenter studies from recent eras [21, 26] our paper's findings do not demonstrate an association between hospitalists and improved outcomes. Unlike other multicenter studies, we had access to disease-specific risk adjustment methodologies which may partially account for referral biases related to patient-specific measures of acute or chronic illness severity.

Changes in the 'hospitalist effect' over time have a number of explanations, some of which are relevant to our study. Recent evidence suggests that complex organizational characteristics such as organizational values and goals may contribute to performance on 30-day mortality for AMI rather than specific processes and protocols [27]; intense focus on AMI as a quality improvement target is emblematic of a number of national initiatives that may have affected our results. Interestingly, hospitalist systems have changed over time as well. Early in the hospitalist 'movement' hospitalist systems were implemented largely at the behest of hospitals trying to reduce costs. In recent years however, hospitalist systems are at least as frequently being implemented because outpatient-based physicians or surgeons request hospitalists; hospitalists have been focused on care of 'uncovered' patients since the model's earliest description. In addition, some hospitals invest in hospitalist programs based on perceived ability of hospitalists to improve quality and achieve better patient outcomes in an era of payment increasingly being linked to quality of care metrics.

Our study has several limitations, six of which are noted here. First, while the hospitalist model has been widely embraced in the adult medicine field, in the absence of board certification there is no gold standard definition of a hospitalist. It is therefore possible that some respondents may have represented groups that were identified incorrectly as hospitalists. Second the data for the primary independent variable of interest was based upon self-report and therefore subject to recall bias and potential misclassification of results. Respondents were not aware of our hypothesis, so the bias should not have been in one particular direction. Third, the data for the outcome variables are from 2008. They may therefore not reflect organizational enhancements related to use of hospitalists that are in process and take years to yield downstream improvements on performance metrics. In addition, of the 429 hospitals that have hospitalist programs, 46 programs were initiated after 2008. While national performance on the 6 outcomes variables has been relatively static over time [7], any significant change in hospital performance on these metrics since 2008 could suggest an over- or under-estimation of the effect of hospitalist programs on patient outcomes. Fourth, we were not able to adjust for additional hospital or health system level characteristics that may be associated with hospitalist use or patient outcomes. Fifth, our regression models had significant collinearity in that the presence of hospitalists was correlated with each of the covariates. However, this finding would indicate that our estimates may be overly conservative and could have contributed to our non-significant findings. Finally, outcomes for two of the three clinical conditions measured are ones for

which hospitalists may less frequently provide care: acute myocardial infarction and heart failure. An outcome measure more relevant for hospitalists may be all-condition, all-cause 30-day mortality and readmission.

This work adds to the growing body of literature examining the impact of hospitalists on quality of care. To our knowledge, it is the first study to assess the association between hospitalist use and performance on outcomes metrics at a national level. While our findings suggest that use of hospitalists alone may not lead to improved performance on outcomes measures, a parallel body of research is emerging implicating broader system and organizational factors as key to high performance on outcome measures. It is likely that multiple factors contribute to performance on outcome measures, including type and mix of hospital personnel, patient care processes and workflow, and system level attributes. Comparative effectiveness and implementation research that assess the contextual factors and interventions that lead to successful system improvement and better performance is increasingly needed. It is unlikely that a single factor such as hospitalist use will significantly impact 30-day mortality or readmission and therefore multifactorial interventions are likely required. In addition, hospitalist use is a complex intervention as the structure, processes, training, experience, role in the hospital system, and other factors (including quality of hospitalists or the hospitalist program) vary across programs. Rather than focusing on the volume of care delivered by hospitalists, hospitals will likely need to support hospital medicine programs that have the time and expertise to devote to improving the quality and value of care delivered across the hospital system. This study highlights that interventions leading to improvement on core outcomes measures are more complex than simply having a hospital medicine program.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors wish to acknowledge Judy Maselli, MPH, Division of General Internal Medicine, Department of Medicine, University of California, San Francisco, for her assistance with statistical analyses and preparation of tables.

Funding Sources

Supported by the Robert Wood Johnson Clinical Scholars Program (K.G.). A.D.A.'s work on this project was supported by a California Healthcare Foundation grant (15763). H.M.K.'s work on this project was supported by a grant from the National Heart, Lung, and Blood Institute (NHLBI), study# 1U01HL105270-02.

References

1. Rifkin WD, Burger A, Holmboe ES, Sturdevant B. Comparison of hospitalists and nonhospitalists regarding core measures of pneumonia care. *Am J of Manag Care*. 2007; 13:129–32. [PubMed: 17335356]
2. Rifkin WD, Conner D, Silver A, Eichorn A. Comparison of Processes and Outcomes of Pneumonia Care Between Hospitalists and Community-Based Primary Care Physicians. *Mayo Clin Proc*. 2002; 77(10):1053–1058. [PubMed: 12374249]
3. Lindenauer PK, Chehabbedine R, Pekow P, Fitzgerald J, Benjamin EM. Quality of Care for Patients Hospitalized With Heart Failure: Assessing the Impact of Hospitalists. *Arch Intern Med*. 2002; 162(11):1251–1256. [PubMed: 12038943]
4. Vasilevskis EE, Meltzer D, Schnipper J, Kaboli P, Wetterneck T, Gonzales D, Arora V, Zhang J, Auerbach AD. Quality of Care for Decompensated Heart Failure: Comparable Performance

- between Academic Hospitalists and Non-hospitalists. *J Gen Intern Med.* 2008; 23(9):1399–1406. [PubMed: 18592321]
5. Roytman MM, Thomas SM, Jiang CS. Comparison of practice patterns of hospitalists and community physicians in the care of patients with congestive heart failure. *J Hosp Med.* 2008; 3(1): 35–41. [PubMed: 18257099]
 6. Vasilevskis EE, Knebel RJ, Dudley RA, Wachter RM, Auerbach AD. Cross-Sectional Analysis of Hospitalist Prevalence and Quality of Care in California. *JHosp Med.* 2010; 5(4):200–207. [PubMed: 20394024]
 7. Hospital Compare. Department of Health and Human Services; Available at www.hospitalcompare.hhs.gov
 8. Ayanian JZ, Weissman JS. Teaching Hospitals and Quality of Care: A Review of the Literature. *Milbank Quarterly.* 2002; 80(3):569–593. [PubMed: 12233250]
 9. Devereaux PJ, Choi PT, Lacchetti C, Weaver B, Schunemann HJ, Haines T, Lavis JN, Grant BJ, Haslam DR, Bhandari M, Sullivan J, Cook DJ, Walter SD, Meade M, Khan H, Bhatnagar N, Guyatt GH. A systematic review and meta-analysis of studies comparing mortality rates of private for-profit and private not-for-profit hospitals. *CMAJ.* 2002; 166(11):1399–1406. [PubMed: 12054406]
 10. Fine JM, Fine MJ, Galusha D, Patrillo M, Meehan TP. Patient and Hospital Characteristics Associated With Recommended Processes of Care for Elderly Patients Hospitalized With Pneumonia: Results From the Medicare Quality Indicator System Pneumonia Module. *Arch Intern Med.* 2002; 162(7):827–833. [PubMed: 11926859]
 11. Jha AK, Li Z, Orav EJ, Epstein AM. Care in U.S. Hospitals -- The Hospital Quality Alliance Program. *N Engl J Med.* 2005; 353(3):265–274. [PubMed: 16034012]
 12. Keeler EB, Rubenstein LV, Khan KL, Draper D, Harrison ER, McGinty MJ, Rogers WH, Brook RH. Hospital Characteristics and Quality of Care. *JAMA.* 1992; 268(13):1709–1714. [PubMed: 1527880]
 13. Needleman J, Buerhaus P, Mattke S, Stewart M, Zelevinsky K. Nurse-Staffing Levels and the Quality of Care in Hospitals. *N Engl J Med.* 2002; 346(22):1715–1722. [PubMed: 12037152]
 14. Pham HH, Devers KJ, Kuo S, Berenson R. Health care market trends and the evolution of hospitalist use and roles. *J Gen Intern Med.* 2005; 20:101–107. [PubMed: 15836541]
 15. Krumholz HM, Wang Y, Mattera JA, Han LF, Ingber MJ, Roman S, Normand SL. An administrative claims model suitable for profiling hospital performance based on 30-day mortality rates among patients with an acute myocardial infarction. *Circulation.* 2006; 113:1683–1692. [PubMed: 16549637]
 16. Krumholz HM, Lin Z, Drye EE, Desai MM, Han LF, Rapp MT, Mattera JA, Normand SL. An administrative claims measure suitable for profiling hospital performance based on 30-day all-cause readmission rates among patients with acute myocardial infarction. *Circulation.* 2011; 4:243–252. [PubMed: 21406673]
 17. Keenan PS, Normand SL, Lin Z, Drye EE, Bhat KR, Ross JS, Schuur JD, Stauffer BD, Bernheim SM, Epstein AJ, Wang Y, Herrin J, Chen J, Federer JJ, Mattera JA, Krumholz HM. An administrative claims measure suitable for profiling hospital performance on the basis of 30-day all-cause readmission rates among patients with heart failure. *Circ Cardiovasc Qual Outcomes.* 2008; 1:29–37. [PubMed: 20031785]
 18. Krumholz HM, Wang Y, Mattera JA, Han LF, Ingber MJ, Roman S, Normand SL. An administrative claims model suitable for profiling hospital performance based on 30-day mortality rates among patients with heart failure. *Circulation.* 2006; 113:1693–1701. [PubMed: 16549636]
 19. Bratzler DW, Normand SL, Wang Y, O'Donnell WJ, Metersky M, Han LF, Rapp MT, Krumholz HM. An administrative claims model for profiling hospital 30-day mortality rates for pneumonia patients. *PLoS ONE.* 2011; 6(4):e17401. [PubMed: 21532758]
 20. Lindenauer PK, Normand SL, Drye EE, Lin Z, Goodrich K, Desai MM, Bratzler DW, O'Donnell WJ, Metersky ML, Krumholz HM. Development, validation and results of a measure of 30-day readmission following hospitalization for pneumonia. *J Hosp Medicine.* 2011; 6:142–150.
 21. Kuo YF, Goodwin JS. Association of hospitalist care with medical utilization after discharge: evidence of cost shift from a cohort study. *Ann Intern Med.* 2011; 155:152–159. [PubMed: 21810708]

22. Vasilevskis EE, Knebel RJ, Wachter RM, Auerbach AD. California hospital leaders' views of hospitalists: meeting needs of the present and future. *J Hosp Medicine*. 2009; 4:528–534.
23. Meltzer D, Manning WG, Morrison J, Shah MN, Jin L, Guth T, Levinson W. Effects of physician experience on costs and outcomes on an academic general medicine service: results of a trial of hospitalists. *Ann Intern Med*. 2002; 137:866–874. [PubMed: 12458986]
24. Auerbach AD, Wachter RM, Katz P, Showstack J, Baron RB, Goldman L. Implementation of a voluntary hospitalist service at a community teaching hospital: improved clinical efficiency and patients outcomes. *Ann Intern Med*. 2002; 137:859–865. [PubMed: 12458985]
25. Palacio C, Alexandraki I, House J, Mooradian A. A comparative study of unscheduled hospital readmissions in a resident-staffed teaching service and a hospitalist-based service. *South Med J*. 2009; 102:145–149. [PubMed: 19139694]
26. Lindenauer P, Rothberg M, Pekow P, Kenwood C, Benjamin E, Auerbach A. Outcomes of care by hospitalists, general internists, and family physicians. *N Engl J Med*. 2007; 357:2589–2600. [PubMed: 18094379]
27. Curry LA, Spatz E, Cherlin E, Thompson JW, Berg D, Ting HH, Decker C, Krumholz HM, Bradley EH. What distinguishes top-performing hospitals in acute myocardial infarction mortality rates? *Ann Intern Med*. 2011; 154:384–390. [PubMed: 21403074]

Table 1

Hospital Characteristics

	Hospitalist Program N=429 N (%)	No Hospitalist Program n=169 N (%)	P value
Core-Based Statistical Area			<0.0001
Division	94 (21.9%)	53 (31.4%)	
Metro	275 (64.1%)	72 (42.6%)	
Micro	52 (12.1%)	38 (22.5%)	
Rural	8 (1.9%)	6 (3.6%)	
Owner			0.0003
Public	47 (11.0%)	20 (11.8%)	
Non-profit	333 (77.6%)	108 (63.9%)	
Private	49 (11.4%)	41 (24.3%)	
Teaching Status			<0.0001
COTH	54 (12.6%)	7 (4.1%)	
Teaching	110 (25.6%)	26 (15.4%)	
Other	265 (61.8%)	136 (80.5%)	
Cardiac Type			0.0003
CABG	286 (66.7%)	86 (50.9%)	
CATH	79 (18.4%)	36 (21.3%)	
Other	64 (14.9%)	47 (27.8%)	
Region			0.007
New England	35 (8.2%)	3 (1.8%)	
Middle Atlantic	60 (14.0%)	29 (17.2%)	
South Atlantic	78 (18.2%)	23 (13.6%)	
E. N. Central	60 (14.0%)	35 (20.7%)	
E. S. Central	31 (7.2%)	10 (5.9%)	
W. N. Central	38 (8.9%)	23 (13.6%)	
W. S. Central	41 (9.6%)	21 (12.4%)	
Mountain	22 (5.1%)	3 (1.8%)	
Pacific	64 (14.9%)	22 (13.0%)	
Safety-Net			0.53
Yes	72 (16.8%)	32 (18.9%)	
No	357 (83.2%)	137 (81.1%)	
	Mean (SD)	Mean (SD)	P value
RN Staffing Ratio (n=455)	27.3 (17.0)	26.1 (7.6)	0.28
Total Beds	315.0 (216.6)	214.8 (136.0)	<0.0001
% Medicare Inpatient Days	47.2 (42)	49.7 (41)	0.19
% Medicaid Inpatient Days	18.5 (28)	21.4 (46)	0.16
% Black	7.6 (9.6)	10.6 (17.4)	0.03

Table 2

Hospitalist Program and Responsibility Characteristics

	N (%)
Date Program Established	
1987-1994	9 (2.2%)
1995-2002	130 (32.1%)
2003-2011	266 (65.7%)
Missing date	24
Number of Hospitalist FTEs	8 (5, 14)
Median (IQR)	
Percent of Medical Patients Admitted by Hospitalists	60% (35, 80)
Median (IQR)	
Number of Hospitalists Groups	
1	333 (77.6%)
2	54 (12.6%)
>=3	36 (8.4%)
Don't know	6 (1.4%)
Employment of Hospitalists (not mutually exclusive)	
Hospital system	98 (22.8%)
Hospital	185 (43.1%)
Local physician practice group	62 (14.5%)
Hospitalist physician practice group (local)	83 (19.3%)
Hospitalist physician practice group (national/regional)	36 (8.4%)
Other/unknown	36 (8.4%)
Any 24-hour in-House Coverage by Hospitalists	
Yes	329 (76.7%)
No	98 (22.8%)
3	1 (0.2%)
Unknown	1 (0.2%)
Number of Hospitalist International Medical Graduates	3 (1, 6)

	N (%)
Median (IQR)	
Number of Hospitalists that are <1 Year out of Residency	1 (0, 2)
Median (IQR)	
Percent of Patients with AMI Cared for by Hospitalists	
0-25%	148 (34.5%)
26-50%	67 (15.6%)
51-75%	50 (11.7%)
76-100%	54 (12.6%)
Don't know	110 (25.6%)
Percent of Patients with Heart Failure Cared for by Hospitalists	
0-25%	79 (18.4%)
26-50%	78 (18.2%)
51-75%	75 (17.5%)
76-100%	84 (19.6%)
Don't know	113 (26.3%)
Percent of Patients with Pneumonia Cared for by Hospitalists	
0-25%	47 (11.0%)
26-50%	61 (14.3%)
51-75%	74 (17.3%)
76-100%	141 (32.9%)
Don't know	105 (24.5%)
Hospitalist Provision of Services	
Care of Critical Care Patients	
Hospitalists provide service	346 (80.7%)
Hospitalists do not provide service	80 (18.7%)
Don't know	3 (0.7%)
Emergency Department Admission Screening	
Hospitalists provide service	281 (65.5%)

	N (%)
Hospitalists do not provide service	143 (33.3%)
Don't know	5 (1.2%)
Observation Unit Coverage	
Hospitalists provide service	359 (83.7%)
Hospitalists do not provide service	64 (14.9%)
Don't know	6 (1.4%)
Emergency Department Coverage	
Hospitalists provide service	145 (33.8%)
Hospitalists do not provide service	280 (65.3%)
Don't know	4 (0.9%)
Coverage for Cardiac Arrests	
Hospitalists provide service	283 (66.0%)
Hospitalists do not provide service	135 (31.5%)
Don't know	11 (2.6%)
Rapid Response Team Coverage	
Hospitalists provide service	240 (55.9%)
Hospitalists do not provide service	168 (39.2%)
Don't know	21 (4.9%)
Quality Improvement or Utilization Review	
Hospitalists provide service	376 (87.7%)
Hospitalists do not provide service	37 (8.6%)
Don't know	16 (3.7%)
Hospital Practice Guideline Development	
Hospitalists provide service	339 (79.0%)
Hospitalists do not provide service	55 (12.8%)
Don't know	35 (8.2%)
Implementation of Major Hospital System Projects	
Hospitalists provide service	309 (72.0%)

	N (%)
Hospitalists do not provide service	96 (22.4%)
Don't know	24 (5.6%)

\$watermark-text

\$watermark-text

\$watermark-text

Table 3

Bivariate Analysis of Hospitalist Utilization and Outcomes

Outcome Measure	Hospitalist Program N=429 Mean % (SD)	No Hospitalist Program N=169 Mean (SD)	P value
MI RSMR	16.0 (1.6)	16.1 (1.5)	0.56
MI RSRR	19.9 (.88)	20.0 (0.86)	0.16
HF RSMR	11.3 (1.4)	11.3 (1.4)	0.77
HF RSRR	24.7 (1.6)	25.4 (1.8)	<0.0001
Pneumonia RSMR	11.7 (1.7)	12.0 (1.7)	0.08
Pneumonia RSRR	18.2 (1.2)	18.3 (1.1)	0.28

Table 4

Multivariable Analysis of Hospitalist Utilization and Outcomes

	Adjusted beta estimate (95% CI)
MI RSMR	
Hospitalist	0.001 (-0.002, 0.004)
MI RSRR	
Hospitalist	-0.001 (-0.002, 0.001)
HF RSMR	
Hospitalist	0.0004 (-0.002, 0.003)
HF RSRR	
Hospitalist	-0.006 (-0.009, -0.003)
Pneumonia RSMR	
Hospitalist	-0.002 (-0.005, 0.001)
Pneumonia RSRR	
Hospitalist	0.00001 (-0.002, 0.002)

Table 5

Percent of Patients Admitted by Hospitalists and Outcomes

MI RSMR	Adjusted Beta Estimate (95% CI)
Percent admit	
0-30%	-0.003 (-0.007, 0.002)
32-48%	0.001 (-0.005, 0.006)
50-66%	Ref
70-80%	0.004 (-0.001, 0.009)
>=85%	-0.004 (-0.009, 0.001)
MI RSRR	
Percent admit	
0-30%	0.001 (-0.002, 0.004)
32-48%	-0.001 (-0.004, 0.004)
50-66%	Ref
70-80%	0.001 (-0.002, 0.004)
>=85%	0.001 (-0.002, 0.004)
HF RSMR	
Percent admit	
0-30%	-0.001 (-0.005, 0.003)
32-48%	-0.002 (-0.007, 0.003)
50-66%	Ref
70-80%	-0.002 (-0.006, 0.002)
>=85%	0.001 (-0.004, 0.005)
HF RSRR	
Percent admit	
0-30%	0.002 (-0.004, 0.007)
32-48%	0.0003 (-0.005, 0.006)
50-66%	Ref
70-80%	-0.001 (-0.005, 0.004)
>=85%	-0.002 (-0.007, 0.003)
Pneumonia RSMR	
Percent admit	
0-30%	0.001 (-0.004, 0.006)
32-48%	-0.00001 (-0.006, 0.006)
50-66%	Ref
70-80%	0.001 (-0.004, 0.006)
>=85%	-0.001 (-0.006, 0.005)
Pneumonia RSRR	
Percent admit	
0-30%	-0.0002 (-0.004, 0.003)

MI RSMR	Adjusted Beta Estimate (95% CI)
32-48%	0.004 (-0.0003, 0.008)
50-66%	Ref
70-80%	0.001 (-0.003, 0.004)
>=85%	0.002 (-0.002, 0.006)

\$watermark-text

\$watermark-text

\$watermark-text