UCLA

UCLA Previously Published Works

Title

Dihydromyricetin prevents fetal alcohol exposure-induced behavioral and physiological deficits: the roles of GABAA receptors in adolescence.

Permalink

<https://escholarship.org/uc/item/22h9n3h6>

Journal

Neurochemical research, 39(6)

ISSN

0364-3190

Authors

Liang, Jing
Shen, Yi
Shao, Xuesi M
et al.

Publication Date

2014-06-01

DOI

10.1007/s11064-014-1291-5

Peer reviewed

Dihydromyricetin Prevents Fetal Alcohol Exposure-Induced Behavioral and Physiological Deficits: The Roles of GABA_A Receptors in Adolescence

Jing Liang · Yi Shen · Xuesi M. Shao · Michael B. Scott · Eddie Ly ·
Stephanie Wong · Albert Nguyen · Kevin Tan · Bill Kwon · Richard W. Olsen ·
Igor Spigelman

Received: 27 December 2013 / Revised: 18 March 2014 / Accepted: 20 March 2014 / Published online: 28 March 2014
© Springer Science+Business Media New York 2014

Abstract Fetal alcohol exposure (FAE) can lead to a variety of behavioral and physiological disturbances later in life. Understanding how alcohol (ethanol, EtOH) affects fetal brain development is essential to guide the development of better therapeutics for FAE. One of EtOH's many pharmacological targets is the γ -aminobutyric acid type A receptor (GABA_AR), which plays a prominent role in early brain development. Acute EtOH potentiates inhibitory currents carried by certain GABA_AR subtypes, whereas chronic EtOH leads to persistent alterations in GABA_AR subunit composition, localization and function. We recently introduced a flavonoid compound, dihydromyricetin (DHM), which selectively antagonizes EtOH's intoxicating effects in vivo and in vitro at enhancing GABA_AR function as a candidate for alcohol abuse pharmacotherapy. Here, we studied the effect of FAE on

physiology, behavior and GABA_AR function of early adolescent rats and tested the utility of DHM as a preventative treatment for FAE-induced disturbances. Gavage administration of EtOH (1.5, 2.5, or 5.0 g/kg) to rat dams on day 5, 8, 10, 12, and 15 of pregnancy dose-dependently reduced female/male offspring ratios (largely through decreased numbers of female offspring) and offspring body weights. FAE (2.5 g/kg) rats tested on postnatal days (P) 25–32 also exhibited increased anxiety and reduced pentylenetetrazol (PTZ)-induced seizure threshold. Patch-clamp recordings from dentate gyrus granule cells (DGCs) in hippocampal slices from FAE (2.5 g/kg) rats at P25–35 revealed reduced sensitivity of GABAergic miniature inhibitory postsynaptic currents (mIPSCs) and tonic current (I_{tonic}) to potentiation by zolpidem (0.3 μM). Interestingly, potentiation of mIPSCs by gaboxadol increased, while potentiation of I_{tonic} decreased in DGCs from FAE rats. Co-administration of EtOH (1.5 or 2.5 g/kg) with DHM (1.0 mg/kg) in pregnant dams prevented all of the behavioral, physiological, and pharmacological alterations observed in FAE offspring. DHM administration alone in pregnant rats had no adverse effect on litter size, progeny weight, anxiety level, PTZ seizure threshold, or DGC GABA_AR function. Our results indicate that FAE induces long-lasting alterations in physiology, behavior, and hippocampal GABA_AR function and that these deficits are prevented by DHM co-treatment of EtOH-exposed dams. The absence of adverse side effects and the ability of DHM to prevent FAE consequences suggest that DHM is an attractive candidate for development as a treatment for prevention of fetal alcohol spectrum disorders.

J. Liang (✉) · Y. Shen · M. B. Scott · E. Ly · S. Wong ·
A. Nguyen · K. Tan · B. Kwon · R. W. Olsen
Department of Molecular and Medical Pharmacology, David
Geffen School of Medicine, University of California,
Los Angeles, CA 90095, USA
e-mail: jliang@ucla.edu

Present Address:

Y. Shen
Department of Neurobiology, Zhejiang University School of
Medicine, Hangzhou 310058, Zhejiang, People's Republic of
China

X. M. Shao
Department of Neurobiology, David Geffen School of Medicine,
University of California, Los Angeles, CA, USA

I. Spigelman
Division of Oral Biology and Medicine, School of Dentistry,
University of California, Los Angeles, CA, USA

Keywords Fetal alcohol spectrum disorders · Pregnancy ·
Brain development · Anxiety · Seizure · Tolerance ·
Ethanol · Zolpidem · Gaboxadol · Dihydromyricetin

Introduction

Fetal alcohol exposure (FAE) leads to a variety of behavioral disturbances later in life collectively referred to as fetal alcohol spectrum disorders (FASD) [1, 2]. The severity of the disturbances depends on pregnant mothers' alcohol (ethanol, EtOH) drinking patterns and doses. For example, approximately 5–20 % of pregnant women engage in moderate drinking (≥ 5 drinks/week) [3]. Many women keep drinking until they realize they are pregnant (4–6 weeks), and some of them quit or decrease their alcohol use only by mid-pregnancy [4–6], therefore many fetuses are exposed to EtOH during early stages of pregnancy. EtOH readily crosses the placenta; consequently, peak fetal blood EtOH levels are similar to the mother [7]. Although EtOH clearance is increased in pregnancy [8, 9], EtOH elimination capacity of the fetus is low, particularly in the early stages of pregnancy [10], and EtOH remains trapped in the amniotic fluid leading to reabsorption by the fetus, thereby prolonging exposure time [9, 11].

Neuro-pathophysiological mechanisms underlying EtOH's deleterious effects on brain development are poorly understood. One of EtOH's several major pharmacological targets is GABA_ARs [12–14]. GABA_ARs are heteropentameric ligand-gated ion channels formed from a family of 19 related subunits ($\alpha 1$ –6, $\beta 1$ –3, $\gamma 1$ –3, δ , ϵ , θ , π , and $\rho 1$ –3), some of which are more sensitive to acute potentiation by EtOH than others [15–17]. In the mature brain, chronic intermittent EtOH (CIE) administration leads to persistent alterations in GABA_AR subunit composition, localization, and function, reviewed in [13, 18]. GABA_AR plasticity is an important underlying mechanism of alcohol withdrawal symptoms, such as anxiety, hyperexcitability and tolerance to alcohol, as well as alcohol dependence [19–21]. By contrast, FAE effects on GABA_ARs are not well understood, in part because GABA_AR subunit composition and function is highly developmentally regulated, with many subunit combinations expressed only postnatally [22, 23]. Better understanding of FAE-induced changes in brain development is essential to guide the development of better prevention and treatment strategies.

While numbers of women drinking alcohol during pregnancy appear to be decreasing in the US, as 12.2 % were reported in 1991–2005 [24] compared to 7.6 % in 2006–2010 [3], FASD collectively still represent the largest preventable set of birth defects [1, 2]. Furthermore, available medications for alcohol use disorders (acamprostate, naltrexone, and disulfiram) are not approved by the US Food and Drug Administration for use during pregnancy due to their multiple side effects, and no agents have yet progressed to the point of consideration for human trials [25].

We recently introduced a flavonoid compound, dihydromyricetin (DHM), as a candidate for alcohol abuse pharmacotherapy. We demonstrated that DHM is a weak positive modulator acting on GABA_ARs, and that in vivo, DHM ameliorates acute alcohol intoxication and reduces alcohol withdrawal symptoms, while in vitro, DHM counteracts EtOH effects on GABA_ARs and antagonizes EtOH exposure/withdrawal-induced GABA_AR plasticity [26]. In the present study, we test the hypothesis that FAE-induced neurobehavioral deficits involve alteration of GABA_AR function and that these deficits can be prevented by DHM administration.

Materials and Methods

Animals

All animal experiments followed The Institutional Animal Care and Use Committee approved protocols. The timed first pregnancy (~ 24 h accuracy) Sprague–Dawley young rats (~ 10 –12 weeks old) were purchased and shipped from the supplier (Charles River Laboratories, Hollister CA) to the University of California, Los Angeles. Rats were housed in the vivarium under a 12 h light/dark cycle and had free access to food and water. Groups of pregnant dams received the following treatments: (1) EtOH (1.5 g/kg, gavage); (2) EtOH (2.5 g/kg); and (3) EtOH (5 g/kg); (4) EtOH (1.5 g/kg) + DHM (1 mg/kg); (5) EtOH (2.5 g/kg) + DHM (1 mg/kg); (6) EtOH (5.0 g/kg) + DHM (1 mg/kg); (7) DHM (1 mg/kg); (8) vehicle (water, 20 ml/kg). Dams were weighed and treated as above on the days 5, 8, 10, 12, and 15 of pregnancy but were otherwise undisturbed except for cage cleaning twice per week. Close to the parturition day, investigators took turn to continuously observe dams for signs of parturition, counted the number of newborn rats, and verified that no newborns were eaten. Offspring were counted on their birth date. On postnatal day 21 (P21), the offspring were weighed, separated by gender, and weaned. There were no offspring deaths between birth and weaning day in any of the experimental groups. Given the difficulty in determining the gender of offspring of dams treated with EtOH (5.0 g/kg), they were returned to their dams and weaned on P25 and their gender determined then. Subsequently, offspring were group-housed by gender. Behavioral and electrophysiological experiments were performed on P25–35 offspring.

Drugs

EtOH (Pharmco Products, Brookfield, CT) was administered by oral intubation (gavage) to dams at 1.5, 2.5 or 5 g/

kg of body weight as a 25 % (w/v) solution in drinking water. The control group received water (20 ml/kg of body weight). DHM (dihydromyricetin, (2R,3R)-3,5,7-trihydroxy-2-(3,4,5-trihydroxyphenyl)-2,3-dihydrochromen-4-one. HPLC purified ≥ 98 %, Blue California, USA. Lot No. 2731-052708) was administered at 1 mg/kg in water by gavage.

Behavioral Assays

Offspring from treated dams were studied in behavioral assays on P25–32. The elevated plus maze assay (EPM) was performed in a quiet and dark room with only a low power red light. Rats were placed on the central area of the maze, and their behavior video recorded for 5 min. The number of entries into open arms, closed arms, or center platform and time spent in each of these areas were scored during off-line analysis. Data were reported as % of number of entries in arms, % of time spent in different arms, and number of total entries. For the pentylenetetrazol (PTZ) seizure assay, rats received an injection of PTZ (42 mg/kg, i.p.). The time to onset and the duration of tonic-clonic seizures were determined. For the loss of righting reflex (LORR) assay, rats were injected with EtOH (3 g/kg, i.p.) to induce LORR. After LORR onset, rats were placed on their backs in home cage and a timer started. The LORR period ends when the animal was able to flip over three times in 30 s.

Plasma EtOH Concentration Assay

Blood samples were collected from the tail vein of pregnant rats at different time points (30, 60, 120, and 180 min) after EtOH or EtOH + DHM gavage treatment as follows. The rat was briefly (<5 min) placed into a restraint tube; the tail vein at the tip of the tail was pricked with a sharp blade. Approximately 0.2 ml venous blood was placed into a capillary blood collection tube containing lithium heparin (Ram Scientific). Blood samples were centrifuged at 2,500 rpm for 20 min. The supernatant was collected and stored at -80°C until assay. The EtOH content of each sample was measured in duplicate along with EtOH standards using the alcohol oxidase reaction procedure (GM7 Micro-Stat, Analox Instruments) [27].

Brain Slice Preparation and Recordings

Coronal slices (400 μm thick) of rat brain at the level of dorsal hippocampus were obtained using standard techniques [27]. Briefly, P25–35 rats were decapitated under isoflurane anesthesia, brains were quickly removed,

trimmed with a razor blade and glued to the base of a cutting chamber (Leica VT1200S) filled with cold ($\sim 4^{\circ}\text{C}$) artificial cerebrospinal fluid (ACSF) composed of 125 mM NaCl, 2.5 mM KCl, 2 mM CaCl_2 , 2 mM MgCl_2 , 26 mM NaHCO_3 , and 10 mM D-glucose (Sigma-Aldrich Co, MO). The ACSF was continuously bubbled with a 95/5 % mixture of O_2/CO_2 to ensure adequate oxygenation of slices and a pH of 7.4. Patch electrode filling solutions contained: 135 mM CsCl, 2 mM MgCl_2 , 1 mM CaCl_2 , 11 mM ethylene glycol-bis (β -aminoethyl ether)-N,N,N',N'-tetraacetic acid (EGTA), 10 mM HEPES, 2 mM K_2ATP , 0.2 mM Na_2GTP ; pH adjusted to 7.25 with CsOH. Whole-cell patch clamp recordings were obtained from dentate granule cells (DGCs) during perfusion with oxygenated ACSF at $34 \pm 0.5^{\circ}\text{C}$. DGCs were voltage-clamped at -70 mV (CsCl) with the aid of a Multiclamp 700B amplifier and a Digidata 1440 A/D converter (Molecular Devices, Sunnyvale, CA). GABA_AR-mediated currents were separated pharmacologically by application of TTX (0.5 μM), CNQX (10 μM), APV (40 μM), and CGP54626 (1 μM) in the ACSF. Picrotoxin, zolpidem and gaboxadol (4,5,6,7 tetrahydroisoxazolo [5,4-c] pyridin-3-ol, THIP) (all from Sigma-Aldrich) were applied after appropriate dilution in the ACSF. The recordings were low-pass filtered (Clampfit software, Molecular Devices, Sunnyvale, CA) at 2 kHz and analyzed with the aid of the Mini Analysis Program (Synaptosoft Inc., Fort Lee, NJ). The mIPSCs were detected with threshold criteria of: 5 pA, amplitude and 20 fC, charge transfer. Frequency of mIPSCs was determined from all automatically detected events in a given recording period. The tonic current magnitudes were obtained from the mean baseline current of a given recording period. For kinetic analysis of mIPSCs, single events with a stable baseline, sharp rising phase and exponential decay were chosen. Double and multiple peak mIPSCs were excluded. The mIPSC kinetics were obtained from analysis of the averaged chosen single events (>120 events/100 s recording period) aligned with half rise-time in each cell.

Statistics

The investigators performing the behavioral tests, electrophysiological recordings and analysis were blind to treatments (vehicle, EtOH, DHM, or EtOH + DHM) received by the dams. All summary values are presented as mean $\pm$ SEM. Group differences were evaluated by one-way or two-way analysis of variance (ANOVA) or repeated measures (RM) ANOVA, followed by a post hoc multiple comparison based on Holm-Sidak method when ANOVA showed statistical significance. $p < 0.05$ was considered statistically significant.

Results

Fetal Alcohol Exposure Induces Deficits in Growth and Gender-Specific Birth Rates

We first assessed the dose-dependent effects of fetal EtOH exposure on the numbers, gender, and body weights (BW) of offspring from dams treated with vehicle (water) or EtOH (1.5, 2.5, or 5.0 g/kg) on day 5, 8, 10, 12, and 15 of pregnancy (Fig. 1a).

Pregnancy lasted 21–22 days. There was a trend to decreased body weight of dams treated with various EtOH

Fig. 1 Fetal EtOH exposure dose-dependently induces growth deficits and decreases numbers of female offspring. **a** Pregnant dam treatment groups and dam numbers. **b** Average male and female offspring numbers per dam. A two-way repeated measures (RM) ANOVA with repeated measures on one factor, “gender”, was performed. The interaction between gender and treatment was highly significant ($p < 0.01$). Therefore, post hoc multiple comparisons with Holm-Sidak method was performed. **c** Average male and female offspring body weights at weaning (P21). A two-way ANOVA followed by a post hoc multiple comparison with Holm-Sidak method was performed. For **b** and **c**: * $p < 0.05$ treatment versus the vehicle group of the corresponding gender. [†] $p < 0.05$, female versus male for each specific treatment

doses which reached statistical significance for dams treated with 1.5 or 2.5 g/kg of EtOH by day 15 of pregnancy (Table 1). Dams built nests prior to parturition. Stretching and hindleg extension were usually signs of impending birth. Parturition occurred at night with 8–15 pups being born over a course of 1–4 h. Pups were born either head or tail first (breech). The dams ate the placenta. Offspring were counted on their birth date whereas their body weights and gender were determined on weaning day (P21) in order to minimize interference with dams and their newborns. Compared to offspring of vehicle-treated dams (14.5 ± 0.24), the number of offspring from each of the three groups of EtOH-treated dams (1.5 g/kg: 12.6 ± 0.34 ; 2.5 g/kg: 12.3 ± 0.29 ; 5 g/kg: 9.3 ± 0.88) was significantly decreased ($p < 0.001$, one-way ANOVA). All offspring survived to their weaning day. The number of male offspring from the 2.5 and 5.0 g/kg EtOH dams slightly increased, while the female offspring numbers significantly decreased in all EtOH groups (Fig. 1b). Surprisingly, two of three dams treated with 5 g/kg EtOH delivered male offspring only. The results indicate that FAE not only reduces litter size, but also affects offspring gender ratios (male vs., female: vehicle: 1.02; EtOH (1.5 g/kg): 2:1; EtOH (2.5 g/kg): 3:1; EtOH (5 g/kg): 13:1). Both male and female offspring of EtOH-treated dams exhibited lower body weights compared to the appropriate gender offspring of vehicle-treated dams (Fig. 1c). The decreases in offspring body weight were a function of increasing EtOH concentrations; the greatest decreases were observed in offspring of dams treated with 5 g/kg of EtOH (Fig. 1c).

DHM Co-administration Prevents FAE-Induced Birth and Growth Deficits

In previous studies, we showed that dihydromyricetin (DHM) could counteract the effects of EtOH intoxication in adult rats [26]. Here we tested whether DHM could prevent the detrimental consequences of fetal alcohol exposure on offspring. We co-administered EtOH at 1.5 or 2.5 g/kg with DHM 1 mg/kg and DHM alone to dams on day 5, 8, 10, 12, and 15 of pregnancy (Fig. 2a). The offspring body weight from the DHM alone or the EtOH (1.5 and 2.5 g/kg) plus DHM groups did not differ from that of the vehicle group (Fig. 2b). Furthermore, the number of offspring of dams treated with DHM alone or with EtOH + DHM, as well as their male: female ratios did not differ from those of the vehicle group (Fig. 2c). These results indicate that DHM administered alone has no detrimental effects on offspring number, body weights, or gender ratios and that when co-administered with EtOH, DHM prevents FAE-induced birth and growth deficits.

Table 1 Dam body weights on drug administration days of pregnancy (E)

Group	Dam body weight (g)					
	E5	E8	E10	E12	E15	n
Vehicle	235.3 ± 2.5	241.0 ± 3.1	249.5 ± 3.4	260.0 ± 3.5	271.0 ± 4.1	9
EtOH (1.5 g/kg)	234.6 ± 2.6	240.8 ± 3.5	248.5 ± 3.0	259.0 ± 3.5	267.5 ± 3.6	9
EtOH (2.5 g/kg)	235.5 ± 3.1	240.0 ± 3.2	245.5 ± 3.0	252.0 ± 2.9	261.0 ± 3.5*	7
EtOH (5.0 g/kg)	240.0 ± 2.5	246.1 ± 2.2	253.3 ± 3.5	257.5 ± 3.5	262.4 ± 3.7*	3
EtOH (1.5 g/kg) + DHM (1 mg/kg)	235.3 ± 2.8	240.9 ± 3.2	249.7 ± 3.3	260.2 ± 3.1	270.9 ± 3.6	8
EtOH (2.5 g/kg) + DHM (1 mg/kg)	233.7 ± 2.5	240.2 ± 3.0	247.5 ± 3.5	256.2 ± 2.5	267.8 ± 3.1	7
DHM (1 mg/kg)	240.0 ± 1.5	246.2 ± 2.5	255.2 ± 3.1	261.5 ± 3.5	272.4 ± 3.8	6

* $p < 0.05$ versus Vehicle group (two-way RM ANOVA) followed by a post hoc multiple comparison with Holm-Sidak method

Fig. 2 DHM co-administration with EtOH in dams prevents EtOH-induced decreases in offspring numbers and weights. **a** Pregnant dam treatment groups and dam numbers. **b** Average male and female offspring numbers per dam. The same two-way RM ANOVA was performed as described in Fig. 1b legend. No significant differences were found. **c** Average male and female offspring body weights. The same two-way ANOVA was performed as described in Fig. 1c legend. No significant differences were found

DHM Co-administration Decreases Blood EtOH Levels in Pregnant Dams

Previous studies demonstrated that DHM co-administration decreased plasma EtOH concentrations in adult male rats

[26]. Since the reduction in plasma EtOH concentrations could contribute to the mechanisms of DHM's prevention of FAE-induced birth and growth deficits, we tested if DHM would similarly affect plasma EtOH levels of pregnant dams. We took venous blood samples at 3, 15, 60, 120, and 180 min from dams treated with EtOH alone and EtOH plus DHM on their last treatment day (15th day of pregnancy) and processed them for plasma [EtOH] measurements. After EtOH administration (2.5 and 5 g/kg, gavage), plasma [EtOH] increased slowly to peak at 60 min, followed by a very slow gradual decline (Fig. 3). The peak plasma [EtOH] of ~160 mg/dl after the 2.5 g/kg dose is similar to that achieved in humans after a moderate level of social drinking. Co-administration of EtOH (2.5 g/kg) with DHM (1 mg/kg), resulted in significant decreases of plasma [EtOH] at the 120, and 180 min time points compared to the EtOH alone group. These results suggest that DHM co-administration decreases blood EtOH levels in pregnant dams and that this can contribute to the protective effects of DHM against FAE-induced abnormalities.

DHM Co-administration Prevents FAE-Induced Behavioral Alterations

GABA_AR functional alterations are well-known to affect anxiety behaviors [28–30], seizure susceptibility [31–33], and sensitivity to anesthetic/soporific effects of acute EtOH administration in the adult brain [13, 18]. To determine the effect of FAE on these parameters later in life we first assessed offspring anxiety levels in the elevated plus maze (EPM). For each experimental group there were 2–4 males and 1–3 females chosen from the same litter. Both male and female offspring of the EtOH (2.5 g/kg) group tested at P25–32 spent significantly less time in the open arms and more time in closed arms, an indication of increased anxiety levels, compared to the offspring from the vehicle-treated group (Fig. 4a). By contrast, the offspring of the DHM only and the EtOH + DHM-treated dams spent similar time in the open and closed arms as the offspring of the vehicle-treated group (Fig. 4a).

Fig. 3 Co-administration of DHM with EtOH decreases plasma [EtOH]. Dams were treated with EtOH (2.5 or 5 g/kg, gavage) or EtOH (2.5 g/kg) with DHM (1 mg/kg). Blood samples were collected at 3, 15, 60, 120 and 180 min after EtOH administration. Two-way RM ANOVA revealed highly significant ($p < 0.01$) interaction of time by treatment. Therefore, post hoc multiple comparisons with Holm-Sidak method followed. The value at every time point of every treatment group versus the corresponding pre-treatment (time 0) value was significant (for simplicity, we did not put symbols to show this). * $p < 0.05$ EtOH (2.5 g/kg) + DHM group versus EtOH (2.5 g/kg) group ($n = 3$ –4 dams per group) at the time point

The sensitivity to the acute anesthetic/soporific effects of EtOH (3 g/kg, i.p.) was addressed using the loss of righting reflex (LORR) assay in male and female offspring previously studied in the EPM. There was a significant difference in mean LORR duration between male and female FAE offspring (Fig. 4b), consistent with the known gender differences in sensitivity to EtOH [34, 35]. Compared with the LORR duration in the vehicle group, FAE offspring of both genders tended to have a reduced duration of LORR (Fig. 4b. Not statistically significant; $p = 0.25$). By contrast, the offspring of the DHM only and the EtOH + DHM-treated dams did not show this trend (Fig. 4b).

Hyperexcitability was assessed with the PTZ-induced seizure assay in P27–34 offspring 2 days after the LORR assays. Overall, female rats tended to have longer time to onset (Fig. 4c) and shorter duration (Fig. 4d) of PTZ-induced seizures (not statistically significant), consistent with known gender differences in PTZ sensitivity [36]. We also observed a trend of shorter seizure onset time after PTZ injection in the EtOH group offspring compared to the offspring from the vehicle-treated groups of both genders which did not reach statistical significance (Fig. 4c). By contrast, the offspring of the DHM alone and the EtOH + DHM groups showed seizure onset times similar to the offspring of the vehicle-treated group (Fig. 4c). PTZ-induced seizure duration was significantly longer in the offspring of EtOH-treated dams compared to that of the vehicle group offspring (Fig. 4d). By contrast, seizure duration in the offspring of DHM only and the

EtOH + DHM-treated dams was similar to the offspring of the vehicle-treated group (Fig. 4d). These results suggest that FAE induces long-lasting increases in postnatal anxiety, tolerance to acute sedative/anesthetic effects of EtOH, and increases seizure susceptibility. All of these FAE-induced changes were prevented by DHM co-administration in dams.

DHM Prevents FAE-Induced Changes in Postsynaptic GABA_AR Function

The observed behavioral and pharmacological manifestations of FAE were suggestive of altered GABAergic inhibition. The plasticity of GABA_AR in hippocampal neurons induced by acute and chronic EtOH is likely to be representative of the same sort of EtOH-sensitive GABA_AR in many brain regions including those cells responsible for the behavioral observations. Therefore, we assessed the kinetics and pharmacological sensitivity of GABA_AR-mediated miniature postsynaptic inhibitory currents (mIPSCs) and tonic inhibitory current (I_{tonic}) in dentate gyrus granule cells (DGCs) in hippocampal slices from offspring (P25–35) of dams exposed to vehicle, EtOH (2.5 g/kg), DHM (1 mg/kg), or EtOH + DHM treatments. All recordings were performed in hippocampal slices from offspring that were not previously subjected to behavioral and pharmacological tests. First we compared the basal magnitude of the picrotoxin-sensitive I_{tonic} and mIPSC kinetics for differences between genders and different treatment groups (Table 2). In the vehicle-treated groups, the only difference between genders was a smaller mIPSC charge transfer in the female progeny. In the FAE groups, there was a trend to decreased magnitude of I_{tonic} in both male and female FAE progeny, but it did not reach statistical significance. The frequency of mIPSCs was not altered in offspring from EtOH-treated dams or other treatment groups (Table 2). Similarly, the rise-time, amplitude, and decay time constant (τ_1) were not affected by EtOH, DHM, or EtOH + DHM treatments. However, the decay τ_2 of mIPSCs from FAE rats of both genders was significantly increased, suggestive of changes in postsynaptic GABAergic function. The total charge transfer of mIPSCs was also significantly larger in FAE rats. All of the FAE-induced changes in GABA_AR currents were prevented by DHM co-administration in dams, while DHM treatment alone did not alter any of the recorded parameters (Table 2).

DHM Prevents FAE-Induced Decreases in Zolpidem Potentiation of Hippocampal GABA_ARs

Postsynaptic function and pharmacological sensitivity of GABA_ARs is largely dictated by their subunit composition

Fig. 4 Co-administration of DHM with EtOH in pregnant dams prevents FAE-induced behavioral and physiological abnormalities in their offspring. **a** Anxiety in male and female offspring was measured by elevated plus maze ($n = 8$ offspring per group). Both male and female offspring of EtOH (2.5 g/kg)-treated pregnant dams spent less time in the open arms and more time in the closed arms compared to the vehicle group. The EtOH + DHM group spent a similar amount of time in each arm as the vehicle group. Two-way ANOVA for open and separately for closed arm entry durations followed by post hoc multiple comparisons based on Holm-Sidak method were performed. * $p < 0.05$ treatment versus vehicle group. **b** Sensitivity to anesthetic effect of EtOH (3 g/kg, i.p.) was measured by LORR in rats from **a**. There was a significant difference in mean LORR duration between male and female FAE offspring (overall gender difference $p = 0.047$). The EtOH groups of both genders showed a trend to

shorter duration of acute EtOH-induced LORR which did not reach statistical significance (overall group effects: $p = 0.25$). The DHM alone and EtOH + DHM groups did not show this trend. **c** At 2 days after LORR testing, the offspring of EtOH-treated dams showed a trend to shorter time to onset of PTZ-induced seizures which did not reach statistical significance (overall group effects: $p = 0.627$). The DHM and EtOH + DHM groups showed similar seizure onset time as the vehicle groups. **d** The duration of PTZ-induced seizures was significantly longer in the offspring of EtOH-treated dams. The DHM alone and EtOH + DHM groups showed similar duration of seizures as the vehicle group. For **b–d**, two-way ANOVA followed by post hoc multiple comparisons based on Holm-Sidak were performed. * $p < 0.05$ treatment versus the vehicle control group. † $p < 0.05$ females versus males

Table 2 Tonic current magnitude and kinetic properties of mIPSCs from offspring DGC recordings

Gender	Group	I_{tonic} (pA)	Frequency (Hz)	Rise-time (ms)	Amplitude (pA)	Decay τ_1 (ms)	Decay τ_2 (ms)	Area (fC)	Cells/progeny
Male	Vehicle	27.8 ± 1.5	13.3 ± 0.1	0.98 ± 0.05	33.6 ± 3.5	6.7 ± 0.5	27.7 ± 4.0	465.9 ± 21.3	11/3
	EtOH (2.5 g/kg)	$22.5 \pm 1.3^*$	13.0 ± 0.2	1.02 ± 0.03	33.8 ± 2.2	7.3 ± 0.4	$48.3 \pm 6.1^*$	$549.2 \pm 19.6^*$	18/4
	EtOH (2.5 g/kg) + DHM (1 mg/kg)	26.0 ± 1.3	13.3 ± 0.1	0.97 ± 0.05	33.9 ± 3.1	6.7 ± 0.3	27.2 ± 3.6	469.4 ± 18.3	19/4
	DHM (1 mg/kg)	28.1 ± 1.7	13.3 ± 0.1	0.99 ± 0.05	33.7 ± 2.6	6.7 ± 0.3	30.6 ± 6.3	465.18 ± 25.0	10/3
Female	Vehicle	26.7 ± 1.5	13.4 ± 0.1	0.98 ± 0.11	32.0 ± 0.5	6.6 ± 0.4	21.3 ± 3.2	$398.0 \pm 21.3^\dagger$	8/2
	EtOH (2.5 g/kg)	$21.1 \pm 1.4^*$	13.0 ± 0.1	1.01 ± 0.03	34.3 ± 3.4	6.5 ± 0.3	$41.4 \pm 7.4^*$	$503.1 \pm 20.4^*$	15/3
	EtOH (2.5 g/kg) + DHM (1 mg/kg)	26.2 ± 1.5	13.3 ± 0.1	1.01 ± 0.03	32.8 ± 2.2	6.6 ± 0.3	24.3 ± 2.0	438.9 ± 21.3	25/5
	DHM (1 mg/kg)	27.5 ± 1.7	13.3 ± 0.1	0.99 ± 0.03	33.2 ± 2.3	6.4 ± 0.4	26.3 ± 3.3	448.4 ± 25.0	9/2

* $p < 0.05$, treatment group versus vehicle group

† $p < 0.05$, female versus male with the same treatment (two-way ANOVA followed by a post hoc multiple comparison with Holm-Sidak method)

Fig. 5 Co-administration of DHM with EtOH in pregnant dams prevents FAE-induced decreases in potentiation of synaptic and extrasynaptic GABA_ARs by zolpidem. **a–d** Recordings (left traces) of pharmacologically-isolated GABA_AR currents in dentate granule cells from male offspring of differentially-treated dams. Gray dashed lines show the picrotoxin (PTX, 50 μ M)-sensitive tonic current (I_{tonic}). Superimposed mIPSCs (right traces) are averaged from the 100 s recording periods before (Ctrl) and during zolpidem (ZP, 0.3 μ M) application. **e** The recording protocol. **f** Summary of group

differences in I_{tonic} sensitivity to ZP for male (left graph) and female (right graph) progeny. **g** Summary of group differences in mIPSC sensitivity to ZP for male (left graph) and female (right graph) progeny. $\dagger p < 0.05$, versus Ctrl; $* p < 0.05$, versus offspring of vehicle-treated dams ($n = 4$ –6 cells from 2 to 3 male offspring per group; $n = 4$ cells from 1 to 2 female offspring per group), two-way RM ANOVA followed by a post hoc multiple comparison with Holm-Sidak method

[37] and subject to some fine tuning by phosphorylation/dephosphorylation states, e.g., [38]. Therefore, we tested GABA_AR-currents for sensitivity to zolpidem (ZP), a benzodiazepine site positive modulator of GABA_AR function with selectivity for the $\alpha 1$ -containing GABA_ARs, intermediate potency at $\alpha 2/3$ -containing GABA_ARs, and inactivity at $\alpha 4/5/6$ -containing GABA_ARs [39]. In DGCs from vehicle-treated dams bath application of ZP potentiated both I_{tonic} magnitude (Fig. 5a, f) and mIPSC charge transfer (Fig. 5a, g). The ZP responsiveness of both I_{tonic} and mIPSCs was significantly reduced in recordings from DGCs of FAE offspring compared to offspring of vehicle-treated dams (Fig. 5b, g, f). These decreases in ZP responsiveness were observed in recordings from DGCs of both male and female offspring (Fig. 5f, g). By contrast, ZP responsiveness of I_{tonic} and mIPSCs from DHM alone and EtOH + DHM groups was indistinguishable from that of the vehicle group offspring (Fig. 5a, c, d, g, f). These results indicate that FAE alters postsynaptic GABA_AR function (Table 2) and ZP sensitivity (Fig. 5) and that these alterations are prevented by co-administration of DHM in dams. These results further suggest that the FAE-induced decreases in ZP sensitivity are likely due to decreased function and/or expression of $\alpha 1$ subunit-containing synaptic and/or extrasynaptic GABA_ARs.

DHM Prevents FAE-Induced Changes in Responsiveness of Synaptic and Extrasynaptic GABA_ARs to THIP

Previous studies have demonstrated that chronic EtOH exposure in adult rodents leads not only to decreases in $\alpha 1$ and δ subunit-containing GABA_ARs, but also to increased expression and altered localization of $\alpha 4$ -containing GABA_ARs [13, 21]. To determine if similar alterations in expression and localization of $\alpha 4$ -containing GABA_ARs occur after FAE we tested GABA_AR-currents for sensitivity to gaboxadol (4,5,6,7 tetrahydroisoxazolo [5,4-*c*] pyridin-3-ol, THIP), which is a selective GABA_AR agonist with a preference for $\alpha 4/\delta$ -containing extrasynaptic GABA_ARs [40–45]. In DGCs from offspring of vehicle-treated dams, bath application of THIP (0.1 μ M) potentiated both I_{tonic} magnitude (Fig. 6a, f) and mIPSC charge transfer (Fig. 6a, g). The potentiation of I_{tonic} by THIP was significantly reduced in DGCs of FAE offspring compared to offspring of vehicle-treated dams (Fig. 5b, f, g). By contrast, THIP potentiation of mIPSCs was increased in offspring of EtOH-treated dams. These alterations in THIP responsiveness of synaptic and extrasynaptic GABA_AR currents were observed in DGCs of both male and female offspring (Fig. 6f, g). In recordings from the offspring of

Fig. 6 Co-administration of DHM with EtOH in pregnant dams prevents FAE-induced changes in sensitivity of synaptic and extrasynaptic GABA_ARs to THIP. **a–d** Recordings (left traces) of pharmacologically-isolated GABA_AR currents in dentate granule cells from male offspring of differentially-treated dams. Gray dashed lines show the picrotoxin (PTX, 50 μM)-sensitive tonic current (I_{tonic}). Superimposed mIPSCs (right traces) are averaged from the 100 s recording periods before (Ctrl) and during THIP (0.3 μM)

the DHM alone and EtOH + DHM groups, THIP responsiveness of I_{tonic} and mIPSCs was indistinguishable from that of the vehicle group offspring (Fig. 6a, c, d, f, g). These results indicate that the FAE-induced alterations in THIP sensitivity are prevented by co-administration of DHM. These results also suggest that FAE-induced differential changes in THIP sensitivity of I_{tonic} and mIPSCs are likely due to alterations in expression and/or localization of α4/δ-containing GABA_ARs at extrasynaptic and synaptic locations.

Discussion

FAE and DHM Effects on Birth and Growth Deficits

We show that intermittent EtOH (1.5, 2.5, or 5.0 g/kg) administration on day 5, 8, 10, 12, and 15 of pregnancy in rats leads to dose-dependent decreases in litter sizes and weights at weaning. This is consistent with numerous reports of FAE-induced decreases in rat offspring numbers and birth weights [46–51]. In humans, FAE is also correlated with low birth weight, growth, and morphological abnormalities [4], and in higher rates of spontaneous abortions [52, 53]. Various potential mechanisms have been proposed to account for the birth and growth deficits

application. **e** The recording protocol. **f** Summary of group differences in I_{tonic} sensitivity to THIP. **g** Summary of group differences in mIPSC area sensitivity to THIP. †*p* < 0.05 versus Ctrl; **p* < 0.05 versus offspring of vehicle-treated dams (*n* = 4 cells from two male offspring per group; *n* = 4 cells from two female offspring per group), two-way RM ANOVA followed by a post hoc multiple comparison with Holm-Sidak method

of FAE rats including, but not limited to, interference with chromosome segregation [54], epigenetic reprogramming [55–57], deficient choline metabolism [58], blunting of cellular immunity [59], oxidative stress [60], as well as decreased production and release of various growth factors and hormones [61–64]. We also observed that the FAE-induced decreases in litter sizes were primarily due to decreased numbers of female offspring. Since gender is determined at conception and embryos were first exposed to EtOH only 5 days later, the lower female/male ratios may have resulted from preferential decreases in the in utero viability of female offspring. While we have not found references to FAE-induced changes in gender ratios in the literature, previous studies have demonstrated an in utero “feminization” of FAE male rodents and “masculinization” of female rodents, attributable to EtOH-induced derangements of in utero androgenesis [65, 66]. Studies have also shown that maternal exposure to EtOH during the critical period of rat organogenesis preferentially affects body weights and increases malformations in offspring dependent on their in utero contiguity to siblings of the opposite sex [67]. Thus, the observed decreases in male offspring body weights could have been exaggerated due to reduced in utero influence of viable female siblings.

We showed that the FAE-induced birth and growth deficits were completely prevented by co-administration of

DHM with EtOH to pregnant rats. We also showed that DHM (1 mg/kg) co-administration with EtOH (2.5 g/kg) decreases plasma [EtOH] in dams, consistent with similar findings in adult male rats [26]. Some flavonoids do affect ethanol metabolism, probably peripherally. It has been reported that *Hovenia*, the herb from which we originally extracted DHM, promotes alcohol clearance, and enhances the activity of alcohol metabolic enzymes, alcohol dehydrogenase and aldehyde dehydrogenase [68, 69], thereby decreasing blood EtOH concentrations. We think that the reversal of birth and growth deficits by DHM is more likely pharmacodynamic than pharmacokinetic. Previous studies have demonstrated that administering EtOH at doses as low as 1 g/kg to pregnant rat dams produces behavioral and physiological abnormalities in their offspring, including higher neonatal mortality and lower birth weights [48]. In our study, DHM reduced plasma [EtOH] to levels that might be achieved after administration of a lower (1.5–2 g/kg) EtOH dose [70–72]. However, we also showed that birth and growth deficits were still evident in the offspring of the 1.5 g/kg EtOH alone group without DHM (Fig. 1), whereas such deficits were prevented after DHM co-treatment of the 2.5 g/kg EtOH group (Fig. 2). This suggests that the complete prevention of FAE-induced decreases in litter sizes and weights cannot be attributed solely to DHM-induced decreases in plasma [EtOH]. One plausible mechanism is the ability of DHM to block EtOH actions at GABA_ARs [26].

GABA and GABA_ARs are essential regulators of early brain development and disruption of GABAergic signaling is suspected in numerous neurodevelopmental disorders [73–78]. We observed increased anxiety in the EPM and faster onset, as well as longer duration of PTZ-induced seizures after FAE in P25–34 rats (Fig. 4), which corresponds to the beginning (~P28) of the adolescent period in rats [79]. To our knowledge, this is the first demonstration of heightened anxiety in FAE rats during early adolescence. Previous studies have demonstrated increased anxiety and depressive behaviors in adult FAE rats [60, 80–82], however see [83]. Children diagnosed with FASD exhibit high anxiety [84, 85], as well as a remarkably high prevalence of epilepsy [86]. While other mechanisms may also be involved, decrements in GABAergic neurotransmission have been associated with increased anxiety [29, 30, 81] and reduced seizure thresholds [33, 87].

Cognitive impairment is also prominent in children diagnosed with FASD [1, 2]. Numerous studies have demonstrated impairment in various learning and memory tasks in adolescent and adult FAE rats, including hippocampal-mediated spatial reference memory [88–93]. Both DGCs and hilar GABAergic interneurons of the dentate gyrus were demonstrated to be integral neural substrates for spatial reference and spatial working memory [94–97].

Interestingly, THIP-sensitive GABA_ARs containing the δ subunit promote memory and neurogenesis in the dentate gyrus [98]. We observed a trend to decreased magnitude of I_{tonic} and significant increases in charge transfer (and decay τ_2) of mIPSCs in DGCs of both male and female FAE progeny (Table 2). We also observed altered pharmacological responsiveness of DGC GABA_ARs from FAE rats to zolpidem and THIP (Figs. 5, 6). It is therefore likely that the FAE-induced alterations in hippocampal GABA_ARs contribute to the deficits in hippocampal learning and memory in FAE rats.

Prior studies also have shown that the responsiveness of GABA-stimulated $^{36}\text{Cl}^-$ flux to benzodiazepine site ligands and to neurosteroids is altered in a brain region-specific manner in FAE rats [99]. Studies have also demonstrated FAE-induced decreases in allopregnanolone potentiation of [^3H]flunitrazepam binding to hippocampal GABA_ARs in guinea pigs [100], age-dependent changes in $\alpha 5$ GABA_AR subunit expression in mice [101], and increased δ GABA_AR subunit expression and sensitivity to THIP in cerebellar granule neurons of rats after early postnatal (P2–12) EtOH vapor exposure [102]. Neurosteroids, such as $3\alpha, 5\alpha$ -3-hydroxypregnan-20-one ($3\alpha, 5\alpha$ -THP or allopregnanolone) and $3\alpha, 5\alpha$ -tetrahydrodeoxycorticosterone (THDOC), are potent endogenous modulators of δ subunit-containing GABA_ARs [103]. Acute EtOH dynamically regulates local $3\alpha, 5\alpha$ -THP levels in several subcortical regions of adult rats, independent of the adrenal glands [104, 105]. Several lines of evidence also suggest that $3\alpha, 5\alpha$ -THP and THDOC contribute to EtOH's pharmacological effects in rats [106]. The levels of these and other neurosteroids change dramatically during embryonic development, parturition, and postnatal development [107–109]. Other studies showed a direct link between neurosteroid concentrations and alterations in GABA_AR expression and function [110–114]. Importantly, FAE has been shown to affect neurosteroid synthesis [115] and to decrease both cellular and behavioral responsiveness to neurosteroids [116, 117]. Collectively these studies demonstrate that fetal and early postnatal EtOH exposure leads to brain region-selective alterations in subunit composition and function of GABA_ARs and that while alterations in neurosteroidogenesis likely play a role, the precise molecular and cellular mechanisms by which FAE exposure alters GABA_AR function remain to be determined.

The decreases in zolpidem sensitivity of adult CNS neurons have been related to chronic EtOH-induced impairment of $\alpha 1$ -containing GABA_ARs, usually as a result of decreases in membrane surface (functional) expression of $\alpha 1$ -containing GABA_ARs [13, 18]. By contrast, the EtOH-induced switch in the relative responsiveness of synaptic and tonic GABA_AR currents to THIP has been previously related to decreases in membrane surface

expression of extrasynaptic $\alpha 4\beta\delta$ GABA_ARs and compensatory increases in $\alpha 4\beta\gamma 2$ GABA_ARs at synaptic and extrasynaptic locations [27, 42, 118, 119]. The transcriptionally-mediated increases in $\alpha 4$ were shown to be due to EtOH-induced activation of heat shock factor 1 and its binding to a short downstream DNA sequence, the alcohol response element on the *Gabra4* gene [120]. Protein kinase C γ was later shown to be selectively involved in the incorporation of such EtOH-up-regulated $\alpha 4$ -GABA_ARs into the plasma membrane [121]. Analogous decreases in $\alpha 1$ - and increases in $\alpha 4$ -containing GABA_ARs have been observed in DGCs in a rat model of temporal lobe epilepsy [122, 123], later shown to involve inducible early growth response factor 3 (*Egr3*) up-regulation of the *Gabra4* minimal promoter [124]. Consistent with the similarity of the GABA_AR alterations, the heightened anxiety, depression, and spatial learning deficits in epilepsy models [125–128] are also observed in rodent FAE models [60, 64, 80, 129, 130]. Collectively, these studies provide a strong link between altered GABA_AR function and the cognitive and behavioral abnormalities associated with FAE.

As with neurobehavioral deficits, the FAE-induced changes in GABA_AR responsiveness to zolpidem and THIP were also prevented by DHM co-administration with EtOH in pregnant rats. Importantly, DHM administration alone in pregnant rats had no adverse effect on litter size, progeny weight, performance in the EPM, EtOH-induced LORR, PTZ-induced seizure threshold, or DGC GABA_AR function. This is consistent with the previous observations that on its own, DHM is not anxiolytic, or sedative, or pro-convulsant at the dose (1 mg/kg) that blocks EtOH effects on GABA_ARs in adult male rats [26]. This contrasts DHM with Ro15-4513, which was shown to block EtOH's enhancement of GABA_ARs in vitro and in vivo [131, 132] and shown to also possess both pro-convulsant and anxiogenic effects due to its ability to block most GABA_AR subtypes [133]. DHM was also shown to greatly reduce EtOH consumption when high voluntary EtOH consumption is already established in adult male rats [26], suggesting that similar reductions in voluntary EtOH consumption could be achieved in females. The absence of adverse side effects and the ability of DHM to completely prevent FAE consequences suggest that DHM is an attractive candidate for development as a treatment for prevention of FASD.

References

- Riley EP, Infante MA, Warren KR (2011) Fetal alcohol spectrum disorders: an overview. *Neuropsychol Rev* 21:73–80
- May PA, Gossage JP, Kalberg WO, Robinson LK, Buckley D, Manning M, Hoyme HE (2009) Prevalence and epidemiologic characteristics of FASD from various research methods with an emphasis on recent in-school studies. *Dev Disabil Res Rev* 15:176–192
- Centers for Disease Control and Prevention (CDC) (2012) Alcohol use and binge drinking among women of childbearing age—United States, 2006–2010. *Morb Mortal Wkly Rep* 61:534–538
- Day NL, Jasperse D, Richardson G, Robles N, Sambamoorthi U, Taylor P, Scher M, Stoffer D, Cornelius M (1989) Prenatal exposure to alcohol: effect on infant growth and morphologic characteristics. *Pediatrics* 84:536–541
- Floyd RL, Weber MK, Denny C, O'Connor MJ (2009) Prevention of fetal alcohol spectrum disorders. *Dev Disabil Res Rev* 15:193–199
- Floyd RL, O'Connor MJ, Sokol RJ, Bertrand J, Cordero JF (2005) Recognition and prevention of fetal alcohol syndrome. *Obstet Gynecol* 106:1059–1064
- Thomas JD, Riley EP (1998) Fetal alcohol syndrome: does alcohol withdrawal play a role? *Alcohol Health Res World* 22:47–53
- Shankar K, Ronis MJ, Badger TM (2007) Effects of pregnancy and nutritional status on alcohol metabolism. *Alcohol Res Health* 30:55–59
- Nava-Ocampo AA, Velazquez-Armenta Y, Brien JF, Koren G (2004) Elimination kinetics of ethanol in pregnant women. *Reprod Toxicol* 18:613–617
- Pikkarainen PH (1971) Metabolism of ethanol and acetaldehyde in perfused human fetal liver. *Life Sci* 10:1359–1364
- Brien JF, Loomis CW, Tranmer J, McGrath M (1983) Disposition of ethanol in human maternal venous blood and amniotic fluid. *Am J Obstet Gynecol* 146:181–186
- Koob GF (2004) A role for GABA mechanisms in the motivational effects of alcohol. *Biochem Pharmacol* 68:1515–1525
- Kumar S, Porcu P, Werner DF, Matthews DB, Diaz-Granados JL, Helfand RS, Morrow AL (2009) The role of GABA_A receptors in the acute and chronic effects of ethanol: a decade of progress. *Psychopharmacology* 205:529–564
- Ticku MK (1990) Alcohol and GABA-benzodiazepine receptor function. *Ann Med* 22:241–246
- Macdonald RL, Olsen RW (1994) GABA_A receptor channels. *Annu Rev Neurosci* 17:569–602
- Olsen RW, Hancher HJ, Meera P, Wallner M (2007) GABA_A receptor subtypes: the “one glass of wine” receptors. *Alcohol* 41:201–209
- Olsen RW, Sieghart W (2008) International union of pharmacology. LXX. Subtypes of gamma-aminobutyric acid_A receptors: classification on the basis of subunit composition, pharmacology, and function. Update. *Pharmacol Rev* 60:243–260
- Spigelman I, Olsen RW, Liang J, Suryanarayanan A, Lindemeyer AK, Meyer EM, Shen Y, Bagnara R, Marty VN (2013) Molecular and functional changes in receptors: GABA and chronic alcohol consumption. In: *Biological research on addiction: comprehensive addictive behaviors and disorders*. Elsevier Inc., Academic Press, New York, 219–230
- Enoch MA (2008) The role of GABA_A receptors in the development of alcoholism. *Pharmacol Biochem Behav* 90:95–104
- Frey GD (1990) γ -Aminobutyric acid changes in alcohol withdrawal. In: Porter RJ, Mattson RH, Cramer JA, Diamond I (eds) *Alcohol and seizures: basic mechanisms and clinical concepts*. FA Davis Company, Philadelphia, pp 87–101
- Olsen RW, Spigelman I (2012) GABA_A receptor plasticity in alcohol withdrawal. In: Noebels JL, Avoli M, Rogawski MA, Olsen RW, Delgado-Escueta AV (eds) *Jasper's basic mechanisms of the epilepsies*. Oxford University Press, New York, pp 562–573

22. Fritschy J-M, Paysan J, Enna A, Mohler H (1994) Switch in the expression of rat GABA_A-receptor subtypes during postnatal development: an immunohistochemical study. *J Neurosci* 14:5302–5324
23. Laurie DJ, Wisden W, Seeburg PH (1992) The distribution of thirteen GABA_A receptor subunit mRNAs in the rat brain. III. Embryonic and postnatal development. *J Neurosci* 12:4151–4172
24. Centers for Disease Control and Prevention (CDC) (2009) Alcohol use among pregnant and nonpregnant women of childbearing age—United States, 1991–2005. *Morb Mortal Wkly Rep* 58:529–532
25. National Institute on Alcohol Abuse and Alcoholism (2006) 5 Year strategic plan FY07–11. National Institute on Alcohol Abuse and Alcoholism
26. Shen Y, Lindemeyer AK, Gonzalez C, Shao XM, Spigelman I, Olsen RW, Liang J (2012) Dihydromyricetin as a novel anti-alcohol intoxication medication. *J Neurosci* 32:390–401
27. Liang J, Suryanarayanan A, Abriam A, Snyder B, Olsen RW, Spigelman I (2007) Mechanisms of reversible GABA_A receptor plasticity after ethanol intoxication. *J Neurosci* 27:12367–12377
28. Gulinello M, Smith SS (2003) Anxiogenic effects of neurosteroid exposure: sex differences and altered GABA_A receptor pharmacology in adult rats. *J Pharmacol Exp Ther* 305:541–548
29. Quirk GJ, Gehlert DR (2003) Inhibition of the amygdala: key to pathological states? *Ann N Y Acad Sci* 985:263–272
30. Sanders SK, Shekhar A (1995) Regulation of anxiety by GABA_A receptors in the rat amygdala. *Pharmacol Biochem Behav* 52:701–706
31. Olsen RW, DeLorey TM, Gordey M, Kang MH (1999) GABA receptor function and epilepsy. *Adv Neurol* 79:499–510
32. Coulter DA (2001) Epilepsy-associated plasticity in gamma-aminobutyric acid receptor expression, function, and inhibitory synaptic properties. *Int Rev Neurobiol* 45:237–252
33. Sperk G, Furtinger S, Schwarzer C, Pirker S (2004) GABA and its receptors in epilepsy. *Adv Exp Med Biol* 548:92–103
34. Webb B, Burnett PW, Walker DW (2002) Sex differences in ethanol-induced hypnosis and hypothermia in young Long-Evans rats. *Alcohol Clin Exp Res* 26:695–704
35. Silveri MM, Spear LP (1998) Decreased sensitivity to the hypnotic effects of ethanol early in ontogeny. *Alcohol Clin Exp Res* 22:670–676
36. Kokka N, Sapp DW, Witte U, Olsen RW (1992) Sex differences in sensitivity to pentylenetetrazol but not in GABA_A receptor binding. *Pharmacol Biochem Behav* 43:441–447
37. Olsen RW, Sieghart W (2008) International union of pharmacology. LXX. Subtypes of gamma-aminobutyric acid_A receptors: classification on the basis of subunit composition, pharmacology, and function. Update. *Pharmacol Rev* 60:243–260
38. Houston CM, He Q, Smart TG (2009) CaMKII phosphorylation of the GABA_A receptor: receptor subtype- and synapse-specific modulation. *J Physiol* 587:2115–2125
39. Olsen RW, Sieghart W (2008) International union of pharmacology. LXX. Subtypes of γ -aminobutyric acid_A receptors: classification on the basis of subunit composition, pharmacology, and function. Update. *Pharmacol Rev* 60:243–260
40. Adkins CE, Pillai GV, Kerby J, Bonnert TP, Haldon C, McKernan RM, Gonzalez JE, Oades K, Whiting PJ, Simpson PB (2001) $\alpha 4\beta 3\delta$ GABA_A receptors characterized by fluorescence resonance energy transfer-derived measurements of membrane potential. *J Biol Chem* 276:38934–38939
41. Krogsgaard-Larsen P, Frolund B, Liljefors T, Ebert B (2004) GABA_A agonists and partial agonists: THIP (Gaboxadol) as a non-opioid analgesic and a novel type of hypnotic. *Biochem Pharmacol* 68:1573–1580
42. Liang J, Cagetti E, Olsen RW, Spigelman I (2004) Altered pharmacology of synaptic and extrasynaptic GABA_A receptors on hippocampal CA1 pyramidal neurons is consistent with subunit changes in a model of alcohol withdrawal and dependence. *J Pharmacol Exp Ther* 310:1234–1245
43. Boehm SL, Homanics GE, Blednov YA, Harris RA (2006) δ -Subunit containing GABA_A receptor knockout mice are less sensitive to the actions of 4,5,6,7-tetrahydroisoxazolo-[5,4-c]pyridin-3-ol. *Eur J Pharmacol* 541:158–162
44. Chandra D, Jia F, Liang J, Peng Z, Suryanarayanan A, Werner DF, Spigelman I, Houser CR, Olsen RW, Harrison NL, Homanics GE (2006) GABA_A receptor $\alpha 4$ subunits mediate extrasynaptic inhibition in thalamus and dentate gyrus and the action of gaboxadol. *Proc Natl Acad Sci USA* 103:15230–15235
45. Meera P, Wallner M, Otis TS (2011) Molecular basis for the high THIP/gaboxadol sensitivity of extrasynaptic GABA_A receptors. *J Neurophysiol* 106:2057–2064
46. Testar X, Lopez D, Llobera M, Herrera E (1986) Ethanol administration in the drinking fluid to pregnant rats as a model for the fetal alcohol syndrome. *Pharmacol Biochem Behav* 24:625–630
47. Sherwin BT, Jacobson S, Troxell SL, Rogers AE, Pelham RW (1979) A rat model (using a semipurified diet) of the fetal alcohol syndrome. *Curr Alcohol* 7:15–30
48. Vaglenova J, Petkov VV (1998) Fetal alcohol effects in rats exposed pre- and postnatally to a low dose of ethanol. *Alcohol Clin Exp Res* 22:697–703
49. Buts JP, Sokal EM, Van HF (1992) Prenatal exposure to ethanol in rats: effects on postnatal maturation of the small intestine and liver. *Pediatr Res* 32:574–579
50. Henderson GI, Schenker S (1977) The effect of maternal alcohol consumption on the viability and visceral development of the newborn rat. *Res Commun Chem Pathol Pharmacol* 16:15–32
51. Lopez-Tejero D, Ferrer I, Llobera M, Herrera E (1986) Effects of prenatal ethanol exposure on physical growth, sensory reflex maturation and brain development in the rat. *Neuropathol Appl Neurobiol* 12:251–260
52. Windham GC, Von BJ, Fenster L, Schaefer C, Swan SH (1997) Moderate maternal alcohol consumption and risk of spontaneous abortion. *Epidemiology* 8:509–514
53. Kline J, Shrout P, Stein Z, Susser M, Warburton D (1980) Drinking during pregnancy and spontaneous abortion. *Lancet* 2:176–180
54. Kaufman MH, Bain IM (1984) Influence of ethanol on chromosome segregation during the first and second meiotic divisions in the mouse egg. *J Exp Zool* 230:315–320
55. Balaraman S, Tingling JD, Tsai PC, Miranda RC (2013) Dysregulation of microRNA expression and function contributes to the etiology of fetal alcohol spectrum disorders. *Alcohol Res* 35:18–24
56. Laufer BI, Mantha K, Kleiber ML, Diehl EJ, Addison SM, Singh SM (2013) Long-lasting alterations to DNA methylation and ncRNAs could underlie the effects of fetal alcohol exposure in mice. *Dis Model Mech* 6:977–992
57. Haycock PC, Ramsay M (2009) Exposure of mouse embryos to ethanol during preimplantation development: effect on DNA methylation in the h19 imprinting control region. *Biol Reprod* 81:618–627
58. Zeisel SH (2011) What choline metabolism can tell us about the underlying mechanisms of fetal alcohol spectrum disorders. *Mol Neurobiol* 44:185–191
59. Chiappelli F, Taylor AN (1995) The fetal alcohol syndrome and fetal alcohol effects on immune competence. *Alcohol Alcohol* 30:259–262
60. Brocardo PS, Boehme F, Patten A, Cox A, Gil-Mohapel J, Christie BR (2012) Anxiety- and depression-like behaviors are

- accompanied by an increase in oxidative stress in a rat model of fetal alcohol spectrum disorders: protective effects of voluntary physical exercise. *Neuropharmacology* 62:1607–1618
61. Conway S, Ling SY, Leidy JW Jr, Blaine K, Holtzman T (1997) Effect of fetal ethanol exposure on the in vitro release of growth hormone, somatostatin and growth hormone-releasing factor induced by clonidine and growth hormone feedback in male and female rats. *Alcohol Clin Exp Res* 21:826–839
 62. Davis MI (2008) Ethanol-BDNF interactions: still more questions than answers. *Pharmacol Ther* 118:36–57
 63. Moore DB, Madorsky I, Paiva M, Barrow HM (2004) Ethanol exposure alters neurotrophin receptor expression in the rat central nervous system: effects of prenatal exposure. *J Neurobiol* 60:101–113
 64. Wilcoxon JS, Kuo AG, Disterhoft JF, Redei EE (2005) Behavioral deficits associated with fetal alcohol exposure are reversed by prenatal thyroid hormone treatment: a role for maternal thyroid hormone deficiency in FAE. *Mol Psychiatry* 10:961–971
 65. McGivern RF, Clancy AN, Hill MA, Noble EP (1984) Prenatal alcohol exposure alters adult expression of sexually dimorphic behavior in the rat. *Science* 224:896–898
 66. Udani M, Parker S, Gavalier J, Van Thiel DH (1985) Effects of in utero exposure to alcohol upon male rats. *Alcohol Clin Exp Res* 9:355–359
 67. Mankes RF, Glick SD (1986) Preferential alcoholic embryopathy among contiguous siblings of Long-Evans rats. *Alcohol Clin Exp Res* 10:388–392
 68. Chen SH, Zhong GS, Li AL, Li SH, Wu LK (2006) Influence of *Hovenia dulcis* on alcohol concentration in blood and activity of alcohol dehydrogenase (ADH) of animals after drinking. *Zhongguo Zhong. Yao Za Zhi* 31:1094–1096
 69. Kim MH, Chung YT, Lee JH, Park YS, Shin MK, Kim HS, Kim DH, Lee HY (2000) Hepatic detoxification activity and reduction of serum alcohol concentration of *Hovenia dulcis* Thunb from Korea and China. *Korean J Med Crop Sci* 8:225–233
 70. Badger TM, Hildestrand M, Shankar K, McGuinn WD, Ronis MJ (2005) The effects of pregnancy on ethanol clearance. *Life Sci* 77:2111–2126
 71. Bielawski DM, Abel EL (2002) The effect of administering ethanol as single vs. divided doses on blood alcohol levels in the rat. *Neurotoxicol Teratol* 24:559–562
 72. Kelly SJ, Bonthius DJ, West JR (1987) Developmental changes in alcohol pharmacokinetics in rats. *Alcohol Clin Exp Res* 11:281–286
 73. Kilb W, Kirischuk S, Luhmann HJ (2013) Role of tonic GABAergic currents during pre- and early postnatal rodent development. *Front. Neural Circuits* 7:1–13
 74. Simeone TA, Donevan SD, Rho JM (2003) Molecular biology and ontogeny of gamma-aminobutyric acid (GABA) receptors in the mammalian central nervous system. *J Child Neurol* 18:39–48
 75. Rudolph U, Mohler H (2014) GABA receptor subtypes: therapeutic potential in Down syndrome, affective disorders, schizophrenia, and autism. *Annu Rev Pharmacol Toxicol* 54:1–25
 76. Salazar P, Velasco-Velazquez MA, Velasco I (2008) GABA effects during neuronal differentiation of stem cells. *Neurochem Res* 33:1546–1557
 77. Represa A, Ben-Ari Y (2005) Trophic actions of GABA on neuronal development. *Trends Neurosci* 28:278–283
 78. Paysan J, Fritschy JM (1998) GABA_A-receptor subtypes in developing brain. Actors or spectators? *Perspect Dev Neurobiol* 5:179–192
 79. Spear LP (2000) The adolescent brain and age-related behavioral manifestations. *Neurosci Biobehav Rev* 24:417–463
 80. Osborn JA, Kim CK, Steiger J, Weinberg J (1998) Prenatal ethanol exposure differentially alters behavior in males and females on the elevated plus maze. *Alcohol Clin Exp Res* 22:685–696
 81. Zhou R, Wang S, Zhu X (2010) Prenatal ethanol exposure attenuates GABAergic inhibition in basolateral amygdala leading to neuronal hyperexcitability and anxiety-like behavior of adult rat offspring. *Neuroscience* 170:749–757
 82. Cullen CL, Burne TH, Lavidis NA, Moritz KM (2013) Low dose prenatal ethanol exposure induces anxiety-like behaviour and alters dendritic morphology in the basolateral amygdala of rat offspring. *PLoS ONE* 8:1–12
 83. Carneiro LM, Diogenes JP, Vasconcelos SM, Aragao GF, Noronha EC, Gomes PB, Viana GS (2005) Behavioral and neurochemical effects on rat offspring after prenatal exposure to ethanol. *Neurotoxicol Teratol* 27:585–592
 84. Hellemans KG, Sliwowska JH, Verma P, Weinberg J (2010) Prenatal alcohol exposure: fetal programming and later life vulnerability to stress, depression and anxiety disorders. *Neurosci Biobehav Rev* 34:791–807
 85. Steinhausen HC, Willms J, Metzke CW, Spohr HL (2003) Behavioural phenotype in foetal alcohol syndrome and foetal alcohol effects. *Dev Med Child Neurol* 45:179–182
 86. Bell SH, Stade B, Reynolds JN, Rasmussen C, Andrew G, Hwang PA, Carlen PL (2010) The remarkably high prevalence of epilepsy and seizure history in fetal alcohol spectrum disorders. *Alcohol Clin Exp Res* 34:1084–1089
 87. Carlen PL, Davies MF, Rougier-Naquet I, Reynolds JN, Spigelman I (1990) Sedative drug withdrawal seizures: cellular electrophysiological mechanisms. In: Avoli M, Gloor P, Kostopoulos G, Naquet R (eds) *Generalized epilepsy neurobiological approaches*. Birkhauser, Boston, pp 460–470
 88. Kim CK, Kalynchuk LE, Kornecook TJ, Mumby DG, Dadgar NA, Pineda JP, Weinberg J (1997) Object-recognition and spatial learning and memory in rats prenatally exposed to ethanol. *Behav Neurosci* 111:985–995
 89. Gabriel KI, Johnston S, Weinberg J (2002) Prenatal ethanol exposure and spatial navigation: effects of postnatal handling and aging. *Dev Psychobiol* 40:345–357
 90. Thomas JD, La Fiette MH, Quinn VR, Riley EP (2000) Neonatal choline supplementation ameliorates the effects of prenatal alcohol exposure on a discrimination learning task in rats. *Neurotoxicol Teratol* 22:703–711
 91. Matthews DB, Simson PE (1998) Prenatal exposure to ethanol disrupts spatial memory: effect of the training-testing delay period. *Physiol Behav* 64:63–67
 92. Zimmerberg B, Weston HE (2002) Postnatal stress of early weaning exacerbates behavioral outcome in prenatal alcohol-exposed juvenile rats. *Pharmacol Biochem Behav* 73:45–52
 93. Neese S, La GL, Trujillo E, Romero D (2004) The effects of ethanol and silymarin treatment during gestation on spatial working memory. *BMC Complement. Altern Med* 4:4
 94. Andrews-Zwilling Y, Gillespie AK, Kravitz AV, Nelson AB, Devidze N, Lo I, Yoon SY, Bien-Ly N, Ring K, Zwilling D, Potter GB, Rubenstein JL, Kreitzer AC, Huang Y (2012) Hilar GABAergic interneuron activity controls spatial learning and memory retrieval. *PLoS ONE* 7:e40555
 95. Jessberger S, Clark RE, Broadbent NJ, Clemenson GD Jr, Consiglio A, Lie DC, Squire LR, Gage FH (2009) Dentate gyrus-specific knockdown of adult neurogenesis impairs spatial and object recognition memory in adult rats. *Learn Mem* 16:147–154
 96. Kee N, Teixeira CM, Wang AH, Frankland PW (2007) Preferential incorporation of adult-generated granule cells into spatial memory networks in the dentate gyrus. *Nat Neurosci* 10:355–362
 97. Jeltsch H, Bertrand F, Lazarus C, Cassel JC (2001) Cognitive performances and locomotor activity following dentate granule

- cell damage in rats: role of lesion extent and type of memory tested. *Neurobiol Learn Mem* 76:81–105
98. Whissell PD, Rosenzweig S, Lecker I, Wang DS, Wojtowicz JM, Orser BA (2013) γ -Aminobutyric acid type A receptors that contain the δ subunit promote memory and neurogenesis in the dentate gyrus. *Ann Neurol* 74:611–621
 99. Allan AM, Wu H, Paxton LL, Savage DD (1998) Prenatal ethanol exposure alters the modulation of the gamma-aminobutyric acidA1 receptor-gated chloride ion channel in adult rat offspring. *J Pharmacol Exp Ther* 284:250–257
 100. Iqbal U, Dringenberg HC, Brien JF, Reynolds JN (2004) Chronic prenatal ethanol exposure alters hippocampal GABA_A receptors and impairs spatial learning in the guinea pig. *Behav Brain Res* 150:117–125
 101. Toso L, Roberson R, Woodard J, Abebe D, Spong CY (2006) Prenatal alcohol exposure alters GABA_A $\alpha 5$ expression: a mechanism of alcohol-induced learning dysfunction. *Am J Obstet Gynecol* 195:522–527
 102. Diaz MR, Vollmer CC, Zamudio-Bulcock PA, Vollmer W, Blomquist S, Morton RA, Everett JC, Zurek AA, Yu J, Orser BA, Valenzuela CF (2014) Repeated intermittent alcohol exposure during the third trimester-equivalent increases expression of the GABA_A receptor δ subunit in cerebellar granule neurons and delays motor development in rats. *Neuropharmacology* 79:262–274
 103. Wang M (2011) Neurosteroids and GABA-A receptor function. *front. Endocrinology (Lausanne)* 2:44
 104. Cook JB, Dumitru AM, O'Buckley TK, Morrow AL (2014) Ethanol administration produces divergent changes in GABAergic neuroactive steroid immunohistochemistry in the rat brain. *Alcohol Clin Exp Res* 38:90–99
 105. Cook JB, Nelli SM, Neighbors MR, Morrow DH, O'Buckley TK, Maldonado-Devincini AM, Morrow AL (2014) Ethanol alters local cellular levels of (3 α ,5 α)-3-hydroxypregnan-20-one (3 α ,5 α -THP) independent of the adrenals in subcortical brain regions. *Neuropsychopharmacology*. doi:10.1038/npp.2014.46
 106. Morrow AL (2007) Recent developments in the significance and therapeutic relevance of neuroactive steroids—Introduction to the special issue. *Pharmacol Ther* 116:1–6
 107. Concas A, Follesa P, Barbaccia ML, Purdy RH, Biggio G (1999) Physiological modulation of GABA_A receptor plasticity by progesterone metabolites. *Eur J Pharmacol* 375:225–235
 108. Grobin AC, Morrow AL (2001) 3 α -hydroxy-5 α -pregnan-20-one levels and GABA_A receptor-mediated ³⁶Cl[−] flux across development in rat cerebral cortex. *Brain Res Dev Brain Res* 131:31–39
 109. Kellogg CK, Frye CA (1999) Endogenous levels of 5 alpha-reduced progestins and androgens in fetal vs. adult rat brains. *Brain Res Dev Brain Res* 115:17–24
 110. Biggio F, Gorini G, Caria S, Murru L, Mostallino MC, Sanna E, Follesa P (2006) Plastic neuronal changes in GABA_A receptor gene expression induced by progesterone metabolites: in vitro molecular and functional studies. *Pharmacol Biochem Behav* 84:545–554
 111. Follesa P, Biggio F, Caria S, Gorini G, Biggio G (2004) Modulation of GABA_A receptor gene expression by allopregnanolone and ethanol. *Eur J Pharmacol* 500:413–425
 112. Yu R, Follesa P, Ticku MK (1996) Down-regulation of the GABA receptor subunits mRNA levels in mammalian cultured cortical neurons following chronic neurosteroid treatment. *Brain Res Mol Brain Res* 41:163–168
 113. Yu R, Hay M, Ticku MK (1996) Chronic neurosteroid treatment attenuates single cell GABA_A response and its potentiation by modulators in cortical neurons. *Brain Res* 706:160–162
 114. Modol L, Casas C, Navarro X, Llido A, Vallee M, Pallares M, Darbra S (2014) Neonatal finasteride administration alters hippocampal $\alpha 4$ and δ GABA_AR subunits expression and behavioural responses to progesterone in adult rats. *Int J Neuropsychopharmacol* 17:259–273
 115. Caldeira JC, Wu Y, Mameli M, Purdy RH, Li PK, Akwa Y, Savage DD, Engen JR, Valenzuela CF (2004) Fetal alcohol exposure alters neurosteroid levels in the developing rat brain. *J Neurochem* 90:1530–1539
 116. Zimmerberg B, McDonald BC (1996) Prenatal alcohol exposure influences the effects of neuroactive steroids on separation-induced ultrasonic vocalizations in rat pups. *Pharmacol Biochem Behav* 55:541–547
 117. Costa ET, Olivera DS, Meyer DA, Ferreira VM, Soto EE, Frausto S, Savage DD, Browning MD, Valenzuela CF (2000) Fetal alcohol exposure alters neurosteroid modulation of hippocampal N-methyl-D-aspartate receptors. *J Biol Chem* 275:38268–38274
 118. Liang J, Spigelman I, Olsen RW (2009) Tolerance to sedative/hypnotic actions of GABAergic drugs correlates with tolerance to potentiation of extrasynaptic tonic currents of alcohol-dependent rats. *J Neurophysiol* 102:224–233
 119. Liang J, Zhang N, Cagett E, Houser CR, Olsen RW, Spigelman I (2006) Chronic intermittent ethanol-induced switch of ethanol actions from extrasynaptic to synaptic hippocampal GABA_A receptors. *J Neurosci* 26:1749–1758
 120. Pignataro L, Miller AN, Ma L, Midha S, Protiva P, Herrera DG, Harrison NL (2007) Alcohol regulates gene expression in neurons via activation of heat shock factor 1. *J Neurosci* 27:12957–12966
 121. Werner DF, Kumar S, Criswell HE, Suryanarayanan A, Alex FJ, Comerford CE, Morrow AL (2011) PKC γ is required for ethanol-induced increases in GABA_A receptor $\alpha 4$ subunit expression in cultured cerebral cortical neurons. *J Neurochem* 116:554–563
 122. Brooks-Kayal AR, Shumate MD, Jin H, Rikhter TY, Coulter DA (1998) Selective changes in single cell GABA_A receptor subunit expression and function in temporal lobe epilepsy. *Nat Med* 4:1166–1172
 123. Raol YH, Lund IV, Bandyopadhyay S, Zhang G, Roberts DS, Wolfe JH, Russek SJ, Brooks-Kayal AR (2006) Enhancing GABA_A receptor $\alpha 1$ subunit levels in hippocampal dentate gyrus inhibits epilepsy development in an animal model of temporal lobe epilepsy. *J Neurosci* 26:11342–11346
 124. Roberts DS, Raol YH, Bandyopadhyay S, Lund IV, Budreck EC, Passini MA, Wolfe JH, Brooks-Kayal AR, Russek SJ (2005) Egr3 stimulation of GABRA4 promoter activity as a mechanism for seizure-induced up-regulation of GABA_A receptor $\alpha 4$ subunit expression. *Proc Natl Acad Sci USA* 102:11894–11899
 125. Castelano AS, Cassane GS, Scorza FA, Cysneiros RM (2013) Altered anxiety-related and abnormal social behaviors in rats exposed to early life seizures. *Front Behav Neurosci* 7:1–8
 126. Duarte FS, Duzzioni M, Hoeller AA, Silva NM, Ern AL, Piermartiri TC, Tasca CI, Gavioli EC, Lemos T, Carobrez AP, De Lima TC (2013) Anxiogenic-like profile of Wistar adult rats based on the pilocarpine model: an animal model for trait anxiety? *Psychopharmacology* 227:209–219
 127. Sayin U, Sutula TP, Stafstrom CE (2004) Seizures in the developing brain cause adverse long-term effects on spatial learning and anxiety. *Epilepsia* 45:1539–1548
 128. Pineda E, Shin D, Sankar R, Mazarati AM (2010) Comorbidity between epilepsy and depression: experimental evidence for the involvement of serotonergic, glucocorticoid, and neuroinflammatory mechanisms. *Epilepsia* 51(Suppl 3):110–114
 129. Kleiber ML, Laufer BI, Wright E, Diehl EJ, Singh SM (2012) Long-term alterations to the brain transcriptome in a maternal voluntary consumption model of fetal alcohol spectrum disorders. *Brain Res* 1458:18–33

130. Hellemans KG, Verma P, Yoon E, Yu W, Weinberg J (2008) Prenatal alcohol exposure increases vulnerability to stress and anxiety-like disorders in adulthood. *Ann N Y Acad Sci* 1144:154–175
131. Suzdak PD, Glowa JR, Crawley JN, Schwartz RD, Skolnick P, Paul SM (1986) A selective imidazobenzodiazepine antagonist of ethanol in the rat. *Science* 234:1243–1247
132. Wallner M, Hancher HJ, Olsen RW (2006) Low-dose alcohol actions on $\alpha 4\beta 3\delta$ GABA_A receptors are reversed by the behavioral alcohol antagonist Ro15-4513. *Proc Natl Acad Sci USA* 103:8540–8545
133. Paul SM (2006) Alcohol-sensitive GABA receptors and alcohol antagonists. *Proc Natl Acad Sci USA* 103:8307–8308