

UCSF

UC San Francisco Previously Published Works

Title

Dental x-rays and risk of meningioma: Response to Drs. Calnon, Jorgensen, and White

Permalink

<https://escholarship.org/uc/item/22n4k1jr>

Journal

Cancer, 119(2)

ISSN

1097-0142

Authors

Claus, Elizabeth B

Wiemels, Joseph

Wrensch, Margaret

Publication Date

2013-01-15

DOI

10.1002/cncr.27708

Peer reviewed



Published in final edited form as:

Cancer. 2013 January 15; 119(2): 465–466. doi:10.1002/cnrc.27708.

Dental X-rays and Risk of Meningioma: Response to Drs. Jorgenson and White

Elizabeth B. Claus, MD, PhD [Professor and Director of Medical Research],

Department of Epidemiology and Public Health, Yale University School of Public Health New Haven, Connecticut Attending Neurosurgeon Brigham and Women's Hospital Boston, Massachusetts

Joseph L. Wiemels, PhD. [Associate Professor], and

Department of Epidemiology and Biostatistics University of California San Francisco San Francisco, CA

Margaret Wrensch, PhD. [Professor and Lewis Chair in Brain Tumor Research]

Department of Neurological Surgery University of California San Francisco San Francisco, CA

Abstract

Report of past dental x-rays has been reported to be associated with the diagnosis of intracranial meningioma. The authors respond to commentary regarding their recent case/control study of the topic.

Letter

We thank Drs. Jorgensen and White for re-iterating two important considerations presented in the discussion section of our manuscript¹ that relate to interpretation of our results, i.e. recall bias and the fact that exposures reported by our study subjects are from the past when exposure to ionizing radiation associated with dental x-rays was higher than in the current era. Recall bias is an important issue in every case/control study when validation of reported exposures is not available. In our study, presuming an average of 4-5 dentists seen per study subject,² validation of dental records would require contact with a minimum of 12,000 dentists, a task not technically or financially possible. As cohort datasets containing information on both dental x-rays and meningioma are, to our knowledge, limited at present, the case/control design (with its associated limitations) remains one important tool to gain information regarding associations between this exposure and a diagnosis of intra-cranial meningioma. Validation of dental x-ray reports has been previously addressed by Dr. White and his colleagues² and suggests that while under- and over-reporting of this exposure does occur, recall of this exposure is unbiased with a similar measure of agreement for cases and controls. The authors thus concluded that interview data alone may be used for case/control comparisons of dental x-ray exposure.²

Address reprint requests to Dr. Elizabeth B. Claus at the Department of Epidemiology and Public Health, Yale University School of Medicine, 60 College St., P.O. Box 208034, New Haven, CT 06520-8034 or at elizabeth.claus@yale.edu..

Financial Disclosures: None

The association between therapeutic radiation to the head (whether for benign or malignant conditions) and meningioma risk is consistently reported and our study confirms these findings.¹ The suggestion that oral-facial region pain associated with meningioma may lead to increased use of imaging is interesting; in our data only one study subject to date reported such a presenting symptom.

It is important for readers to note that the amount of radiation exposure associated with dental x-rays has markedly decreased over the past several decades. As Dr. White and his colleagues state, current exposure levels for any given dental x-ray are likely to be low. As receipt of medical imaging (including dental x-rays) remains one of the few modifiable exposures to ionizing radiation and the effect of interaction of cumulative exposure with patient genotype is unclear it is prudent for patients and health-care providers to converse regarding use of such technology. We concur with the American Dental Association guidelines and recent reaffirmation that dental x-rays, an important component of good dental care, should be ordered for patients only when necessary for diagnosis and treatment.³

Acknowledgments

Funding Sources: This work was supported by NIH R01 grants CA109468, CA109461, CA109745, CA108473, and CA109475 as well as by the Brain Science Foundation and the Meningioma Mommas.

Reference

1. Claus EB, Calvocoressi L, Bondy M, Schildkraut J, Wiemels J, Wrensch M. Dental X-rays and Risk of Meningioma. *Cancer*. 2012 (In press).
2. Preston-Martin, s; Bernstein, L.; Maldonado, AA.; Henderson, BE.; White, SC. A dental xray validation study. Comparison of information from patient interviews and dental charts. *Am J Epidemiol*. 1985; 121(3):430–439. [PubMed: 4014133]
3. [accessed 5/1/2012] <http://www.ada.org/>