UC Irvine

UC Irvine Previously Published Works

Title

Isotopic signatures (^{14}C , ^{13}C , ^{15}N) as tracers of sources and cycling of soluble and particulate organic matter in the Santa Monica Basin, California

Permalink

<https://escholarship.org/uc/item/22p776wq>

Journal

Progress in Oceanography, 30(1-4)

ISSN

0079-6611

Authors

Williams, PM
Robertson, KJ
Soutar, A
et al.

Publication Date

1992

DOI

10.1016/0079-6611(92)90015-R

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, availalbe at
<https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Isotopic signatures (^{14}C , ^{13}C , ^{15}N) as tracers of sources and cycling of soluble and particulate organic matter in the Santa Monica Basin, California

P.M. WILLIAMS¹, K.J. ROBERTSON¹, A. SOUTAR¹, S.M. GRIFFIN² and E.R.M. DRUFFEL²

¹*Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California 92093, USA*

²*Radiocarbon Laboratory, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts 02543, USA*

Abstract – Measurements of $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ are reported for dissolved (plus colloidal), suspended and sinking particulate, and total sedimentary organic matter in the Santa Monica Basin (mid-basin and shelf sites) on CaBS cruises 1, 3, 4, 7, 8 and 10. These isotopic signatures were indicative of the following processes occurring within the basin regime: (1) terrestrial inputs of organic matter to the sinking and suspended particulate organic matter were of the order of 10% or less, and as high as 25% for the sedimentary organic matter; (2) $\Delta^{14}\text{C}$ values of the UV-oxidizable dissolved organic matter below 5m were similar to those measured in open ocean waters, while the $\Delta^{14}\text{C}$ values in the suspended, and, to a lesser degree, in the sinking organic matter decreased markedly with depth. This latter decrease was primarily attributed to episodic resuspension of shelf and slope sedimentary organic matter, and secondarily to natural and/or anthropogenic petroleum inputs; and (3) the isotopic signatures of the UV-oxidizable dissolved organic matter, coupled with total dissolved carbon and amino acid and carbohydrate concentrations were strikingly similar in the deep basin and at an open-ocean site, suggesting a common history for the subsurface (>300m) and deep water at both sites.

In addition, total mass and organic carbon and nitrogen fluxes from five particle trap deployments are described in detail. Mass fluxes increased with depth, especially on the shelf, suggesting that particle input from the basin slopes may reach the mid-basin site.

We conclude that there is minimal perturbation of all organic phases in the basin from terrestrial sources, and that the properties of the UV-oxidizable dissolved organic matter are not greatly influenced by particles of local origin.

CONTENTS

1.	Introduction	254
2.	Background	255
3.	Methodology	261
3.1	Sampling	261
3.2	Analyses	263
3.3	Suspended filterable POM [POC(susp)] and PON [PON(susp)]	263
3.4	Sinking POM [POC(sink)] and PON [PON(sink)]	263
4.	Results	265
4.1	Dissolved organic and inorganic carbon (DOC, DIC)	266
4.2	Suspended particulate organic matter: POC(susp) and PON(susp)	272
4.3	Sinking particulate organic matter [POC(sink) and PON(sink)], and total mass and organic carbon and nitrogen fluxes	274
4.4	Sedimentary organic matter (SOC)	276
5.	Discussion	277
5.1	Terrestrial (sewage, storm run-off, industrial organics contribution to the SMB organic phases)	277
5.2	Origin of $\Delta^{14}\text{C}$ gradients in mid-basin and shelf organic phases (water column)	280
5.3	Origin of the sedimentary organic matter	282
5.4	Phase inter-relationships deduced from isotopic signatures	283
5.5	C/N ratios in the sinking organic matter	283
5.6	Comparison of isotopic signatures and other properties between the Santa Monica Basin and the North Central Pacific gyre	284
6.	Summary	286
7.	Acknowledgements	287
8.	References	287

1. INTRODUCTION

The principal goal of the California Basin Study (CaBS) is to delineate the sources, transformation, dispersal and ultimate fate of particulate organic particles derived from marine and terrestrial sources. An understanding of particle dynamics is essential in predicting the impact of energy-derived, possibly deleterious, substances on the southern California borderland regime. Inherent in understanding particle dynamics are concurrent studies on all organic phases in the basin system. This includes sinking and suspended particulate, dissolved (plus colloidal), and sedimentary organic matter. Especially important are transformations between these phases.

This report focuses on the use of multiple isotopic signatures (^{14}C , ^{13}C , ^{15}N) to trace the origin and cycling of organic phases in the Santa Monica-San Pedro basin system.

Radiocarbon (^{14}C) is derived from two sources: natural cosmogenic production in the stratosphere, and atmospheric bomb production during the late 1950s and early 1960s. The high levels of ^{14}C in surface dissolved inorganic carbon (DIC) subsequent to 1960 (NYDAL, LOVESETH and SKOGSETH, 1980) result in organic matter produced in surface waters during photosynthesis also containing high levels of ^{14}C ($\Delta^{14}\text{C}^a = +50$ to $+150\text{‰}$). Thus, surface-derived organic matter photosynthetically-fixed since 1957, and found in sub-surface waters and in the sediments, contains bomb ^{14}C . Lower levels of ^{14}C ($\Delta^{14}\text{C} < -75\text{‰}$) indicate pre-bomb levels (BERGER, TAYLOR and LIBBY, 1966; ROBINSON and THOMPSON, 1981), which in turn, suggests a minimal

turnover of carbon in the past 35 years. Thus, ^{14}C is useful as both a long and short-term tracer because of the natural and bomb-produced components, respectively. The high ^{14}C activity of recently synthesized (<35y) terrestrial organic matter ($\Delta^{14}\text{C} \approx +200\text{‰}$) can also be reflected in marine organic phases impacted from terrestrial (and/or atmospheric) sources.

Carbon-13 ($\delta^{13}\text{C}$)^a in conjunction with nitrogen-15 ($\delta^{15}\text{N}$)^a signatures reflect both the origin and fractionation of organic matter as it flows through ecosystems and undergoes phase transformations. Measurements of both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ on the same organic components are obligatory in sorting out organic matter cycles because of the wide spread in the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures for the various terrestrial and marine end-members and within specific trophic levels. Generally speaking, ^{13}C isotopic compositions have proven most valuable for delineating dietary sources and less so in following isotopic fractionation along food-chains, while ^{15}N reflects trophic levels and specific processes such as nitrogen fixation and denitrification and particle transformations (see, for example, FRY and SHERR, 1984; FRY, 1988; ALTABET, 1988).

2. BACKGROUND

Table 1 summarizes previously-published $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values measured in the Southern California Bight regime for various organic pools and end-members which are relevant for interpreting the measurements presented here for the Santa Monica Basin. Some $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ data are omitted for the sedimentary column below the upper 0-1cm horizon.

It is evident that $\Delta^{14}\text{C}$ values reported to date are sparse for all the organic phases. Few DIC ^{14}C activities were measured in California coastal waters from 1959 to 1987, and no values are available for sewage or river (storm water) soluble or particulate organic components. Thus it is difficult to rigorously assign $\Delta^{14}\text{C}$ values to these important end-members. The $\Delta^{14}\text{C}$ measurements of EMERY and BRAY (1962) for sedimentary organic carbon (SOC) are primarily from subsurface samples where surficial $\Delta^{14}\text{C}$ values represent extrapolations. Nonetheless, these numbers are valuable in that they represent pre-bomb activities.

There are no reported measurements of $\delta^{15}\text{N}$ in dissolved organic nitrogen (DON) - off southern California or anywhere else owing to the difficulty of performing such measurements - and only a few results for suspended particulate organic matter (POM) at a location seaward of the Southern California Bight (SAINO and HATTORI, 1987). There are, however, some $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements which can be used to assign limits to the relevant end-member values necessary for estimating marine versus terrestrial contributions to the Santa Monica mid-basin organic phases.

^a $\Delta^{14}\text{C}$ (per mil deviation from the ^{14}C activity of 19th century wood) = $\delta^{14}\text{C}\text{‰} - 2[\delta^{13}\text{C}\text{‰} + 25.0] [1 + \delta^{14}\text{C}\text{‰} \cdot 10^{-3}]$; where the Age (BP) = $8033 \ln [1 + \Delta^{14}\text{C} \cdot 10^{-3}]^{-1}$, and $\delta^{13}\text{C}\text{‰} = [({}^{13}\text{C}/{}^{12}\text{C})_{\text{sample}} / ({}^{13}\text{C}/{}^{12}\text{C})_{\text{PDB}}] / ({}^{13}\text{C}/{}^{12}\text{C})_{\text{PDB}} \times 10^3$. All $\Delta^{14}\text{C}$ values are corrected for: $\delta^{13}\text{C}$ of the sample, age to 1950AD (STUIVER and POLACH, 1977), and relevant blanks for quartz filters and/or combustion tubes. $\delta^{15}\text{N}\text{‰} = [({}^{15}\text{N}/{}^{14}\text{N})_{\text{SAMPLE}} - ({}^{15}\text{N}/{}^{14}\text{N})_{\text{Air}}] / ({}^{15}\text{N}/{}^{14}\text{N})_{\text{Air}} \times 10^3$.

TABLE 1. Pertinent data from previous studies and publications on $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures of organic and inorganic pools in the S. California Bight regime and adjacent areas if applicable

C Pool	Collection date	Location	Depth (m)	$\Delta^{14}\text{C}$ (‰)	Remarks	References
DIC	1959	32°52'N, 117°44'W	Surface	-88 ± 10	SIO pier off Baja, California	1
	1959	30°04'N, 118°02'W	10	-13 ± 8		2
			75	-32 ± 8		
			120	-38 ± 8		
			390	-100 ± 8		
			600	-169 ± 8		
			2000	-239 ± 8		
Atmospheric CO ₂	1959	32°52'N, 117°44'W	Surface	-90 ± 10	SIO pier	3
	1965	"	"	-17 ± 15	"	"
	1969	"	"	49 ± 8	"	"
	1974	28°30'N, 121°29'W	5	138 ± 15	Geosecs: Pacific	4
	1980	31°9.8'N, 117°12'W	2	195 ± 4	South of San Diego	5
				109 ± 8		
	1986-87	N and S hemispheres	Earth's surface	ca 200		6
DOC	1986	SIO pier	~5	-168 ± 18	Replicates	7
	"	"	"	-166 ± 14		
	1988	1km off Del Mar, CA	Surface film	-124 ± 5	Upper 250mm	"
	"	"	10cm	-189 ± 5	5-15cm subsurface	"
Phytoplankton	1958	Off La Jolla, CA	Surface	-35 ± 10	<i>Gonyaulax polyedra</i>	1
Zooplankton	1967	3 miles off SIO pier	Surface	127 ± 10	70% Copepods	8
	"	"	"	140 ± 12	"	"
Surface sediment (total organic carbon)	1956	Santa Monica Basin	1cm	-147 ± 19*	Grab sampler, bottom=825m	9
	"	"	15cm	-362 ± 23*	Open barrel corer, bottom=907m	"
	"	San Pedro Basin	15cm	-258 ± 19*	Piston corer, bottom=900m	"
	"	"	15cm	-195 ± 21*	Open barrel corer, bottom=900m	"
	"	Santa Catalina Basin	1cm	-230 ± 37*	Grab sampler, bottom=1134m	"
	"	San Nicolas Basin	2cm	-273 ± 14*	Piston corer, bottom=1774m	"

*EMERY and BRAY ¹⁴C ages converted to $\Delta^{14}\text{C}$ values using Libby $t^{1/2}=5568\text{y(BP)}$.

C and N Pools	Collection date	Location	Depth (m)	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{C}$ (‰)	Remarks	References
NO_3^-	1976	32°35.5'N, 119°38.5'W	0-200 200-700 1000-2000	-	11.4 to 9.6 9.6 to 8.0 7.0 to 6.8	Open ocean station	10
	1978	San Pedro Basin	0-350 350-600 600-850	0	7.8 to 11.7 11.7 to 9.2 9.2 to 8.3		11
	1974-76 (sediment pore waters)	Santa Barbara Basin	Surface to 890cm	-	8.2 to 12.4 (mean=10.2)	38 samples	12
DON DOC	1976-78	Santa Barbara Basin	Denitrifying zones	-	7.5 to 16.3 (mean=9)		10
	1967	No measurements to date					
		San Diego Trough 32°35'N, 118°35'W	5 98 280 435 750 970	-22.5 -22.5 -22.4 -22.8 -22.7 -22.9	-	bottom-1300m	13
POC(susp), PON(susp)	1968	SIO pier	Surface	-22.0, -22.1	-		"
	1968	SIO pier	~5	-20.07	-		7
	"	"	"	-22.79	-		"
	1988	1km off Del Mar, CA	Surface film	-21.44	-	Upper 250mm	"
	1968	"	5-15cm	-21.32	-	Film subsurface	"
POC(sink), PON(sink)	1968	30°47'N, 120°04'W (no measurements in the Basins)	10-2940	-22.0 to -24.3	-	200 miles SW of San Diego bottom=4000m bottom=>4000m	13
	1983	31°40'N, 124°22'W	80 300 1000-3500	-	8 14 14 to 17		14
	1983	Santa Monica - San Pedro Basin system	100 300 500	-	8.1 7.1 8.0	Deployment 1(2.5d) "	15
(Kerogen)	1977	Santa Monica Basin	100 300 500	-	7.9 7.8 8.0	Deployment 2(2.5d) "	"
	"	San Pedro Basin	830-860 820-850	-21.6 -22.0	-	Values also given for Santa Barbara and San Nicolas Basins	16
	"	Santa Monica Basin San Pedro Basin	830-860 820-850	-22.6 -22.3	13.5		"
(Humic acids)	"				9.4		"

C and N Pools	Collection date	Location	Depth (m)	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)	Remarks	References
Floculents from outfalls	1972-73	White's Point effluent	-	-23.7 ± 0.5 (n=11)	-		17
	1976	"	-	-	3.0, 2.0		18
	"	"	-	-	2.4, 2.5		"
	1976	San Pedro shelf	7, 13, 20	-	3.0 to 7.4 (n=16)	Sampled with slurp guns	19
Suspended particles, Los Angeles River storm waters	1979	White's Point effluent	-	-23.86 ± 0.27 (n=11)	-		20
	1978-79	CHCl ₃ extractables		-26.07 ± 0.15 (n=11)	-		21
		Humic acids		-24.02 ± 0.53 (n=9)	5.06 ± 1.69 (n=9)		"
		Insolubles		-26.37 ± 0.71 (n=7)	12.80 ± 2.07 (n=7)		"
		Total suspended solids		-24.86 ± 0.26 (n=11)			"
	1956	Santa Catalina Basin	3-13	-20.8	-		22
	"	San Pedro Basin	5-25	-21.0	-		"
Surface sediment (total organic carbon)	"	San Nicolas Basin	0-5	-20.6	-		"
	"	Santa Barbara Basin	0-19	-20.4	-		"
	1972	San Pedro shelf	Surface grab sample	-	1.8 to 10.6 (n=31, mean=5.6)		18
	1975	Santa Barbara Basin	2	-	7.5		12
	1981	San Pedro shelf	0-2	-23.6	6.0		23
Surface sediment (humic acids)	<1972	Santa Monica Basin	0-10	-27.4	-		24
	"	San Pedro Basin	"	-22.7	-		"
(Humic acids) (Kerogen)	"	Santa Barbara Basin	"	-22.5	-		"
	1977	San Pedro Basin	0-5	-22.4	10.2		25
	"	"	"	-22.4	9.6		"

C and N Pools	Collection date	Location	Depth (m)	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)	Remarks	References
Zooplankton	1967	3 miles off SIO pier	Surface	-19.7	-	1 meter net	13
	"	"	"	-18.6	-	"	"
	1975	30°-35°N, 118°-120°W	-	-	6.5 to 12.1 (n=7, mean=8.9)	"	12
	1975	S California Bight copepods	0-50	-21.9 ± 0.65 (n=16)	8.1 to 11.0 (n=16, mean=9.8)	Preserved samples	26
	1975	S California Bight chaetognaths	"	-22.1 ± 0.64 (n=12)	9.5 to 12.7 (n=12, mean=11.2)	"	"
Zooplankton (500-1000mm and 1000-2000mm sizes)	1981	White's Point outfall area (unsorted samples)	-	-19.7 to -21.1 (n=9, mean=-20)	8.5 to 9.9 (n=9, mean=9.2)	Fresh-frozen samples	"
	1987	Santa Monica Basin	25	-	9.9, 9.9	Net tows	27
	"	33°46'N, 118°48'W	50	-	9.4, 11.0	"	"
	"	"	100	-	11, 10.7, 11	"	"
	"	"	25	-	8.0, 8.1	Fecal pellets from above zooplankton	"
Zooplankton fecal pellets (500-1000mm and 1000-2000mm sizes)	"	"	50	-	8.6, 9.2	"	"
	"	"	100	-	8.5, 8.6, 8.8	"	"
	1989	Santa Barbara Basin	-	-21.1 to -14.3 (n=6, mean=-17.1±2.8)	-	"	28
Dover sole	1979	White's Point outfall area	<61	-17.7 to -18.2 (n=6, mean=-18)	8.7 to 11.4 (n=5, mean=10.1)	"	29
	"	Control, 130km NW of White's Point	<58	-16.3 to -17.0 (n=5, mean=-16.6)	11.6 to 16.7 (n=5, mean=14.8)	"	"
	1979	White's Point outfall area	<61	-16.7 to -17.6 (n=3, mean=-17.2)	8.5 to 11.8 (n=3, mean=9.6)	"	"
Ridgeback prawn	"	Control, 130km NW of White's Point	<58	-16.4 to -16.5 (n=3, mean=-16.5)	13.2 to 15.3 (n=3, mean=14.2)	"	"

REFERENCES FOR TABLE 1:

1. HUBBS, BIEN and SUESS (1965)
2. BIEN, RAKESTRAW and SUESS (1965)
3. LINICK (1975)
4. ÖSTLUND and STUIVER (1980)
5. E. DRUFFEL (unpublished results)
6. MANNING *et al* (1990)
7. P.M. WILLIAMS (unpublished results)
8. WILLIAMS, MCGOWAN and STUIVER (1970)
9. EMERY and BRAY (1962)
10. LIU and KAPLAN (1989)
11. LIU and KAPLAN (1982)
12. SWEENEY and KAPLAN (1980a)
13. WILLIAMS and GORDON (1970)
14. SAINO and HATTORI (1987)
15. NELSON *et al* (1987)
16. CRISP *et al* (1979)
17. MYERS (1974)
18. SWEENEY *et al* (1980)
19. SWEENEY and KAPLAN (1980b)
20. EGANHOUSE (unpublished results, 1988, Personal communication)
21. EGANHOUSE (1982)
22. EMERY (1960)
23. EGANHOUSE and KAPLAN (1988)
24. NISSENBAUM and KAPLAN (1972)
25. VENKATESAN, BRENNER, RUTH, BONILLA and KAPLAN (1980)
26. MULLIN, RAU and EPPLEY (1984)
27. ALTABET and SMALL (1990)
28. SCHIMMELMANN and TEGNER (unpublished results)
29. RAU, SWEENEY, KAPLAN, MEARNS and YOUNG (1981)

3. METHODOLOGY

3.1 Sampling

Table 2 lists the cruises, dates and samples taken. Note that all mid-basin and shelf samples are from the same two locations for all cruises. The box core (BC-89, CaBS-1) was taken ~10km southwest of the main mid-basin station.

Dissolved organic carbon (DOC) and DIC were sampled with a new, 12 l Go-Flo Niskin bottle on CaBS-7 (trace ^{14}C contaminations of DOC samples from CaBS-4 may have resulted from a "used" 30 l Niskin). Aliquots for DOC were passed directly from the Niskin through ignited GF/C glass fiber filters (nominal pore diameter - 1.0 μm) into 1 gallon brown glass bottles and frozen at -20°C. Unfiltered, 500ml samples for DIC were taken initially from the same Niskin bottle by overflowing three successive volumes in a 500ml brown bottle. These samples were preserved with 0.5ml of saturated HgCl_2 solution and stored at 3°C.

Suspended particulate organic matter was collected using an *in situ* filtration system (WILLIAMS, CARLUCCI and OLSON, 1980) based on an earlier version of LAIRD, JONES and JENTSCH (1967). The current system employs two Pelagic Electronics, sealed *in situ* pumps powered by two marine high-ampereage, 12-volt lead-acid batteries to pull seawater through four 142mm diameter Whatman, quartz fiber filters (nominal pore diameter 0.8 μm) pre-ignited at 800°C) held in PVC holders. Flow rates are initially 10-15 l m^{-1} and total filtration volumes are measured using individual ARAD® flowmeters. The pumps are turned on and off at depth with a mechanical timer, and at deployment are filled with distilled water which is displaced by ambient seawater after dissolution of a soluble wafer (Lifesaver). The intake water is prefiltered through 505 μm pore diameter nitex netting. Contamination is minimized by coating the supporting framework with heat-cured epoxy resin and using stainless steel hardware. The amounts of seawater filtered per single 142mm filter unit varied from 650 l at 100m to 1440 l at depth for 3 to 6h of pumping, respectively. After deployment, each filter pad was folded and frozen at -20°C in cleaned glass jars.

Sinking particles were collected using paired, fiberglass "Soutar" cones, each with an effective mouth opening of 0.062 m^2 (see BRULAND, FRANKS, LANDING and SOUTAR, 1981, for basic trap design). The mouth of each trap was baffled with a cellular grid (1cm x 1cm x 4cm deep) to restrict internal water flow and constrain particle capture to the trap mouth. The lucite cod-end collectors (2 l) were charged with 30g of solid HgCl_2 in a perforated plastic bottle prior to deployment, and filled with surface seawater at launching. On retrieval, the cod-ends were capped and stored at 3-5°C prior to splitting and processing the sample (usually 1 to 14 days later, depending on cruise duration).

The sediment core (8cm diameter by 28cm length) was subsampled from a Soutar box core (SOUTAR, 1979), capped, and frozen at -20°C (collected by R. JAHNKE).

All of the above shipboard sampling procedures were done on the R/V *New Horizon*, where ^{14}C productivity studies were routinely carried out. Thus, it was necessary to rigorously clean the samplers with dilute HCl, methanol and distilled water and to use plastic gloves, etc., when appropriate. No ^{14}C contamination was evident in the $\Delta^{14}\text{C}$ results documented in this manuscript.

TABLE 2. Sampling information. POC(sink) and PON(sink) = particulate organic carbon and nitrogen in sediment trap particulates; POC(susp) and PON(susp) = particulate organic carbon and nitrogen in filterable, suspended particles; DOC = soluble plus colloidal organic carbon passing 1 μ m pore-diameter glass fiber filters; DIC = dissolved inorganic carbon; and SOC and SON = total organic carbon and nitrogen in dried, acidified sediment samples. Note that bottom depths are ± 2 m and lat./long. are within 0.2'

Cruise Samples	Depths (m)	Deployment Dates	Bottom Depth (m)
CaBS-1			
SOC, SON (Core BC-89) 33°41.5'N, 118°56.3'W	0-20cm in 1.25 cm increments	10/14/85	908
CaBS-3			
POC, PON(sink) 33°51'N, 118°38'W	Shelf 100,350	02/09/86-05/19/86 (99d)	395
33°45'N, 118°53'W	Mid-basin 100,500,700,850	02/08/86-05/17/86 (98d)	910
CaBS-4			
POC, PON(sink) 33°51'N, 118°38'W	Shelf 100,350	05/10/86-10/22/86 (155d)	395
33°45'N, 118°53'W	Mid-basin 100,500,700,850	05/18/86-10/21/86 (156d)	910
POC, PON(susp) 33°51'N, 118°38'W	Shelf 100	10/23/86	395
33°45'N, 118°54'W	Mid-basin 83,513,698,850	10/21/86-10/22/86	905
DOC 33°51'N, 118°38'W	Shelf 100	10/24/86	395
33°45.3'N, 118°54.5'W	Mid-basin 5,100,500,700,850	10/24/86	906
CaBS-7			
POC, PON(sink) 33°51.1'N, 118°38.1'W	Shelf 200,350	04/08/87-10/11/87 (186d)	395
33°45.2'N, 118°53.5'W	Mid-basin 200,500,700,850	04/09/87-10/09/87 (183d)	911
DOC			
33°51'N, 118°38'W	Shelf 100	10/12/87	395
33°45'N, 118°53'W	Mid-basin 5,100,500,700,850	10/12/87	910
DIC 33°45'N, 118°53'W	Mid-basin 5,500,850	10/12/87	910
CaBS-8			
POC, PON(sink) 33°44.2'N, 118°50.1'W	Mid-basin 850	10/10/87-02/07/88 (120d)	910
CaBS-10			
POC, PON(sink) 33°51'N, 118°38'W	Shelf 200,350	02/11/88-10/01/88 (233d)	395
33°45'N, 118°53'W	Mid-basin 20,500,700,850	02/10/88-09/27/88 (230d)	911

3.2 Analyses

DOC was oxidized to CO_2 by high-energy UV irradiation following procedures given in WILLIAMS and GORDON (1970) as modified in DRUFFEL, WILLIAMS, ROBERTSON, GRIFFIN, JULL and DONAHUE (1989). Five litre, thawed seawater samples are acidified to $\text{pH} = 2.2\text{--}2.5$ with H_3PO_4 , sparged free of DIC with clean N_2 , saturated with O_2 , and oxidized for 6h with a 1200W Conrad-Hanovia UV lamp at 70°C . The resultant CO_2 is purified and its ^{14}C content determined at the NSF-TAMS facility at University of Arizona after conversion to graphite. $\delta^{13}\text{C}$ ($\pm 0.1\text{‰}$) values for the same DOC were measured on a V.G. Micromass 602E isotope ratio mass spectrometer at the Woods Hole Oceanographic Institution (WHOI).

DIC was collected as CO_2 by acidifying 150ml of the preserved samples with 3ml of 5% H_3PO_4 (final $\text{pH} \approx 2.3\text{--}2.4$) in a CO_2 -free closed system. The CO_2 was collected and purified as with the DOC- CO_2 . $\Delta^{14}\text{C}$ and $\delta^{13}\text{C}$ measurements were done at the University of Arizona and WHOI, respectively.

3.3 Suspended, filterable POM [POC (susp) and PON (susp)]

At WHOI the thawed quartz fiber filters were acidified with 5ml of 1% H_3PO_4 and dried *in vacuo* overnight to remove inorganic carbon and water. The filters were then oxidized for 1h with CuO (wire form) and Ag foil in double quartz tubes at 850°C following the procedure of MINAGAWA, WINTER and KAPLAN (1984). The resultant CO_2 derived from the POC (susp) was converted to graphite (JULL, DONAHUE, HATHEWAY, LINICK and TOOLIN, 1986) at the University of Arizona and the $\Delta^{14}\text{C}$ and $\delta^{13}\text{C}$ values measured at the University of Arizona and WHOI, respectively. $\Delta^{14}\text{C}$ values for CO_2 obtained from all closed tube combustions were corrected for blank CO_2 from the quartz combustion tubes and quartz filters. Blanks composed 0.3 to 5.7% of the sample CO_2 volumes. $\Delta^{14}\text{C}$ values of blank CO_2 samples averaged -480‰ . $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for replicate filters were also measured by D. Winter at the Institute of Geophysics and Planetary Physics, University of California, Los Angeles (UCLA). These latter values are those reported in Tables 3, 4 and 5, where the precision is $\pm 0.1\text{‰}$ for both the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements.

3.4 Sinking POM [POC (sink)] and PON [PON(sink)]

Approximately 1200-1500ml of the supernatant liquid in the lucite cod-ends was syphoned off, and the remaining 500-800ml mixed and subsampled for isotope analyses, mass and C and N flux measurements (SIO), and trace organic compounds (M.I. VENKATESAN, UCLA). Aliquots for $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\delta^{15}\text{N}$ analyses were acidified to a stable $\text{pH} = 1.5\text{--}3$ to remove carbonates and filtered through 42mm diameter GF/C glass fibre filters using a $\sim 15\text{psi}$ vacuum. The collected solids were briefly rinsed with double-distilled water, dried, and pulverized in a mortar. Portions of these homogenized solids were: (1) oxidized to CO_2 at WHOI as for the POC (susp), and the CO_2 - ^{14}C content measured at the University of Arizona; and (2) oxidized at UCLA and their $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures measured. Additional aliquots of the original cod-end samples were analyzed in duplicate for their total mass (dry weight) by drying at 60°C to constant weight (NELSON, BEERS, EPPLEY, JACKSON, MCCARTHY and SOUTAR, 1987) and in duplicate for their CHN content (SHARP, 1974) using either a Hewlett-Packard Model 185B or a Perkin-Elmer Model 2400 CHN analyzer.

The sediment core was extruded from its lucite liner after slightly thawing on the outside, rinsed briefly with double-distilled water, and cut into 1.25cm sections with a hacksaw. The outer edges were trimmed, and the centre portions acidified to $\text{pH} = 2.6\text{--}3.0$ by titration with successive

portions of 50% H_3PO_4 . These slurries were dried to constant weight at 60°C , pulverized, and aliquots analyzed for $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\delta^{15}\text{N}$ [as for POC(sink)].

All of the above operations concerned with isotopic measurements were carried out at either the Mt Soledad Radiocarbon Laboratory (SIO) or the WHOI Radiocarbon Laboratory. Both of these facilities are free of ^{14}C contamination.

TABLE 3. $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\Delta^{14}\text{C}$ and total organic carbon data: POC(susp) and PON(susp), DOC, and DIC. $\Delta^{14}\text{C}$ values are corrected for $\delta^{13}\text{C}$ and age to 1950 AD, according to STUIVER and POLACH (1977). S = shelf, D = mid-basin

Cruise		Depth (m)	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)	$\Delta^{14}\text{C}$ (‰)	Age (y, BP)	Σ Organic Carbon ($\mu\text{mol l}^{-1}$)
POC(susp), PON(susp)							
CaBS-4							
(10/86)	S	100	-23.61	7.74	102 ± 7	post-bomb	0.57
	D	80	-25.27	6.78	110 ± 8	"	0.50
		510	-22.22	7.60	27 ± 6	"	0.21
		700	-22.56	9.64	-46 ± 6	"	0.25
		850	-23.08	9.88	9 ± 6	"	0.36
DOC							
CaBS-4							
(10/86)	S	100	-21.83				53
	D	5	-22.19	(Contaminated with traces of ^{14}C from Niskin bottle)			76
		100	-21.20				52
		500	-21.14				42
		700	-20.86				42
		850	-21.39				42
CaBS-7							
(10/87)	S	100	-21.96		-323 ± 5	3135 ± 40	53
	D	5	-21.94		-197 ± 9	1760 ± 70	76
		100	-21.74		-305 ± 8	2920 ± 60	55
		500	—*		—*		46
		700	-21.92		-475 ± 5	5180 ± 40	45
		850	-21.61		-462 ± 5	4980 ± 40	44
*sample lost							
DIC							
CaBS-7							
(10/87)	D	5	1.9		94 ± 6	post-bomb	
		500	-0.4		-139 ± 5	1200 ± 40	
		850	-0.3		-169 ± 6	1485 ± 50	

TABLE 4. $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and $\Delta^{14}\text{C}$ data. POC(sink) and PON(sink). $\Delta^{14}\text{C}$ values are corrected for $\delta^{13}\text{C}$, preparation blanks, and age to 1950 AD. S = shelf, D = mid-basin

Cruise		Depth (m)	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)	$\Delta^{14}\text{C}$ (‰)
CaBS-3 (05/86)	S	100	-21.15	8.76	66 ± 7
		350	-21.84	3.55	-18 ± 6
	D	100	-21.73	7.80	78 ± 6
		500	-21.84	6.58	26 ± 6
		700	-20.59	7.80	25 ± 6
850		-21.12	5.50	52 ± 6	
CaBS-4 (10/86)	S	100	-22.22	8.95	68 ± 7
		350	-22.01	7.40	-24 ± 10
	D	100	-22.18	8.46	94 ± 6
		500	-22.27	8.49	117 ± 6
		700	-22.28	8.15	47 ± 7
850		-22.43	7.62	33 ± 6	
CaBS-7 (10/87)	S	200	-21.31	9.99	39 ± 6
		350	-21.29	8.06	2 ± 5
	D	200	-21.41	10.07	101 ± 6
		500	-21.78	10.20	63 ± 5
		700	-21.93	7.27	52 ± 5
		850	-21.82	8.06	39 ± 6
CaBS-8 (02/88)	D	850	-22.06	5.97	2 ± 6
CaBS-10 (10/88)	S	200	-21.63	8.37	
		350	-21.48	7.86	
	D	200	-21.52	9.16	
		500	-21.70	8.41	
		700	-21.57	7.52	
		850	-21.76	8.50	

4. RESULTS

Tables 3, 4 and 5 contain the original $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements for the dissolved, sinking, suspended, and sedimentary organic phases in the Santa Monica Basin. These same isotopic signatures are plotted in Figs 1, 2 and 3. Mass and organic C and N fluxes are contained in Tables 6, 7 and 8, and depicted in Fig. 4.

TABLE 5. $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\Delta^{14}\text{C}$ data and chronologies. Sedimentary (SOC, SON), carbonate-free organic matter subsampled from a Soutar box core, No. BC-89, Cruise CaBS-1, 10/14/85. $\Delta^{14}\text{C}$ values are corrected as in Table 4

Core depth ($\pm 0.05\text{cm}$)	$\delta^{13}\text{C}$ ($‰$)	$\delta^{15}\text{N}$ ($‰$)	$\Delta^{14}\text{C}$ ($‰$)	Age (y, BP)	Sedimentation age ¹ (y, AD)
0.00 - 1.25	-22.15	6.37	-10 $\pm$ 6	80 $\pm$ 50	1974
1.25 - 2.50	-22.62	7.18	-21 $\pm$ 6	170 $\pm$ 50	1960
2.50 - 3.75	-21.95	7.33			1939
3.75 - 5.00	-22.03	7.84			1911
5.00 - 6.25	-22.26	7.32			1879
6.25 - 7.50	-22.39	6.09	-154 $\pm$ 5	1345 $\pm$ 40	1847
7.50 - 8.75	-21.82	7.34			1818
8.75 - 10.00	-21.98	6.56			1783
12.50 - 13.75	-21.79	7.11	-307 $\pm$ 5	2945 $\pm$ 40	1637
18.75 - 20.00	-22.11	6.64	-416 $\pm$ 4	4320 $\pm$ 30	1343

¹Calculated from cumulative mass (mg cm^{-2})/mass flux ($\text{mg cm}^{-2}\text{y}^{-1}$) using mass flux = $14.1\text{mg cm}^{-2}\text{y}^{-1}$ for box core no. BC-89 (see HUH *et al.*, 1990). Cumulative mass values for 1.25cm sections interpolated graphically from 1.0cm sections measured by HUH, ZAHNLE and SMALL (1986). Ages are mean values for 1.25cm sections ± 3 years.

4.1 Dissolved organic and inorganic carbon (DOC, DIC)

The ^{14}C activities in DOC (DO^{14}C) in surface waters ($<5\text{m}$) were -167, -197 and -189 $‰$ in 1986, 1987 and 1988, respectively (Tables 1 and 3). These DO^{14}C values are 16-46 $‰$ more negative than at 3m in the North Central Pacific ($\Delta^{14}\text{C} = -151‰$) in 1985 (WILLIAMS and DRUFFEL, 1987) and reflect the lower $\Delta^{14}\text{C}$ values for the corresponding DIC in the same samples. The ^{14}C values in DIC (DI^{14}C) at 3-5m in surface waters south of San Diego and in the Santa Monica Basin (Tables 1 and 3) were 109 and 94 $‰$ in 1980 and 1987, respectively, compared to $140 \pm 10‰$ at 3m in the North Central Pacific in 1985 and 1987 (WILLIAMS and DRUFFEL, 1987; DRUFFEL and WILLIAMS, 1990). These lower DI^{14}C values in the coastal areas probably result from both sporadic upwelling and the shoaling of 100m isopycnal surfaces, resulting in low ^{14}C activity, subsurface DIC reading the surface. This low ^{14}C activity is then reflected in the carbon fixed by photosynthesis which is the source of suspended and sinking particulate organic matter and, eventually, DOC. The bomb- ^{14}C signal is apparent even in the limited measurements of DI^{14}C from surface waters off Southern California (Tables 1 and 3), where $\Delta^{14}\text{C}$ increased from $\sim -90‰$ in 1959 to $\sim +140‰$ in 1969 and then decreased to $\sim -90‰$ in 1987. The corresponding average $\Delta^{14}\text{C}$ values for atmospheric CO_2 during these same years were approximately 230, 550 and 200 $‰$, respectively, reflecting the bomb inputs in the late 1950s and early 1960s (LEVIN, KROMER, SCHOCH-FISCHER, BRUNS, MUNNICH, BERDAN, VOGEL and MUNNICH, 1985; MANNING, LOWE, MELHUISH, SPARKS, WALLACE, BRENNINKMEIJER and MCGILL, 1990). If we assume that photosynthetically-derived terrestrial inputs to the Southern California borderland have the same ^{14}C activity as atmospheric CO_2 , then the $\Delta^{14}\text{C}$ values for marine and terrestrial end-members are ~ 90 and $\sim 200‰$, respectively. At the other extreme, petroleum hydrocarbons (seeps, spills, chemicals) will probably contain little or no ^{14}C , resulting in $\Delta^{14}\text{C}$ values approaching or equal to -1000 $‰$.

FIG.1. Radiocarbon activities ($\Delta^{14}\text{C}$, per mil deviation from the activity of 19th century wood) for UV-oxidizable dissolved organic carbon (DOC; ϕ = mid-basin, $\otimes$ = shelf), suspended particulate organic carbon (POC (susp); $\circ$ = mid-basin, $\bullet$ = shelf), sinking particulate organic carbon (POC (sink); Δ , ∇ , $\diamond$, $\times$ = mid-basin, $\blacktriangle$, $\blacktriangledown$, $\blacklozenge$ = shelf), dissolved inorganic carbon (ϕ = mid-basin), and total sedimentary organic carbon ($\blacksquare$ = mid-basin).

FIG.2. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures for DOC ($\otimes$), POC (susp) ($\bullet$) and POC (sink) ($\circ$), Cruise CaBS-4; and total SOC (I), Cruise CaBS-1.

FIG.3. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures for POC (sink): o = CaBS-3, ∇ = CaBS-4, $\bullet$ = CaBS-7, + = CaBS-8, $\otimes$ = CaBS-10. The heavy dashed lines (---) represent mean values.

TABLE 6. Mass fluxes. Soutar-paired (cones A and B) sediment traps. All values are means of duplicates as $\text{mg m}^{-2}\text{d}^{-1}$ dry weight

Cruise	Deployment depth (m)							
	Shelf (395m)			Mid-basin (905-910m)				
	100	200	350	100	200	500	700	850
CaBS-3 (05/86)								
A	1040		2530	531		603	867	822
B	<u>944</u>		<u>2830</u>	<u>603</u>		<u>654</u>	<u>633</u>	<u>821</u>
Mean	992		2680	567		629	750	822
CaBS-4 (10/86)								
A	485		2410	231		467	613	701
B	<u>646</u>		<u>2360</u>	<u>253</u>		<u>359</u>	<u>468</u>	<u>457</u>
Mean	566		2385	242		413	541	579
CaBS-7 (10/87)								
A		1130	2590		354	564	750	555
B		<u>1050</u>	<u>2420</u>		<u>432</u>	<u>574</u>	<u>618</u>	<u>527</u>
Mean		1090	2505		393	569	684	541
CaBS-8 (02/88)								
A								687
B								<u>586</u>
Mean								637
CaBS-10 (10/88)								
A		1230	2680		497	495	519	545
B		<u>1120</u>	<u>2920</u>		<u>385</u>	<u>433</u>	<u>456</u>	<u>555</u>
Mean		1175	2800		441	464	488	550

TABLE 7. Organic carbon and nitrogen fluxes, Soutar-paired (cones A and B) sediment traps. All values represent duplicate measurements in $\text{mg m}^{-2}\text{d}^{-1}$ dry weight. AF = acid-fumed

	CaBS-3 (05/86)		CaBS-4 (10/86)		CaBS-7 (10/87)		CaBS-10 (10/88)	
	C	N	C	N	C	N	C	N
Shelf:								
100m								
A	83.5	11.7	51.5	7.1				
B	83.0	13.0						
A(AF)	119	19.6	52.2	8.6				
B(AF)	95.0	15.8	79.6	13.3				
200m								
A					97.3	26.3	126	16.8
B					84.1	20.9	111	16.4
A(AF)					85.1	34.8	112	19.5
B(AF)					47.9	15.1	108	17.3
350m								
A	165	21.4	89.5	12.7	160	54.5		
B	166	18.7	150	20.2	139	31.2	190	24.0
A(AF)	146	24.4	125	19.4	124	63.6		
B(AF)	139	18.2	111	17.4	80.0	29.5	160	22.6
Mid-Basin:								
100m								
A	73.4	11.0	33.8	5.5				
B	77.8	13.8	51.1	8.3				
A(AF)	72.3	12.4	44.9	8.0				
B(AF)	64.3	10.8	44.3	8.0				
200m								
A					68.9	16.2	110	17.9
B					59.8	15.8	74.1	11.8
A(AF)					70.2	32.1	126	22.5
B(AF)					31.7	10.3	51.1	10.2
500m								
A	49.0	5.5	49.3	5.5	59.6	8.4	69.2	8.7
B	49.1	5.5	25.0	3.1	48.7	7.3	68.0	8.9
A(AF)	45.4	5.9	40.0	5.0	40.8	6.8	48.7	7.0
B(AF)	52.6	6.9	42.2	6.1	40.7	8.7	40.7	6.5
700m								
A	61.4	6.4	46.0	5.0	58.6	5.1	67.5	7.3
B	50.5	5.4	40.1	4.1	53.3	6.8	54.6	5.7
A(AF)	62.8	7.5	36.5	4.8	37.9	5.7	44.0	6.0
B(AF)	41.8	5.0	48.2	6.6	36.5	5.9	33.4	4.7
850m								
A	61.8	6.2	37.0	4.1	44.4	4.1	64.5	6.8
B	59.3	5.9	50.2	4.9	34.3	3.9	62.6	7.1
A(AF)	59.5	7.3	40.2	5.0	30.9	4.3	45.6	6.1
B(AF)	50.0	5.6	38.4	5.1	29.9	4.2	46.9	6.5

TABLE 8. Organic carbon and nitrogen fluxes. Acid-fumed, mean values in $\text{mg m}^{-2}\text{d}^{-1}$ from Table 7 plus C/N ratios (atoms)

	CaBS-3 (05/86)			CaBS-4 (10/86)			CaBS-7 (10/87)			CaBS-10 (10/88)		
	C	N	C/N	C	N	C/N	C	N	C/N	C	N	C/N
Shelf												
100m	107	17.7	7.0	65.9	11.0	7.0						
200m							66.5	25.0	3.1	110	18.4	7.0
350m	<u>143</u>	<u>21.3</u>	<u>7.8</u>	<u>118</u>	<u>18.4</u>	<u>7.5</u>	<u>102</u>	<u>46.6</u>	<u>2.6</u>	<u>160*</u>	<u>22.6*</u>	<u>8.3*</u>
Mean	125	19.5	7.4	92.0	14.7	7.3	84.3	35.8	2.8	135	20.5	7.7
Mid-basin												
100m	68.3	11.6	6.9	44.6	8.0	6.5						
200m							51.0	21.2	2.8	88.6	16.4	6.3
500m	49.0	6.4	8.9	41.1	5.6	8.6	40.8	7.8	6.1	44.7	6.8	7.7
700m	52.3	6.3	9.8	42.4	5.7	8.7	37.2	5.8	7.5	38.7	5.4	8.4
850m	<u>54.8</u>	<u>6.5</u>	<u>9.9</u>	<u>39.3</u>	<u>5.1</u>	<u>9.1</u>	<u>30.4</u>	<u>4.3</u>	<u>8.3</u>	<u>46.3</u>	<u>6.3</u>	<u>8.6</u>
Mean	56.1	7.7	8.9	41.8	6.1	8.2	39.8	9.8	6.2	54.6	8.7	7.8

*single measurement

FIG.4. Mass and CN fluxes, all in $\text{mgm}^{-2}\text{d}^{-1}$, for mid-basin and shelf sites: o = CaBS-3, ● = CaBS-4, ⊙ = CaBS-7, ▲ = CaBS-10. Dashed lines (---) are N fluxes, and the heavy dashed lines (---) represent mean values.

The differences in $\Delta^{14}\text{C}$ values (Table 3) between DO^{14}C and DI^{14}C (DOC-DIC) are -291‰ at 5m and -293‰ at 850m, differences which are of the same order as those found in the NCP at these depths (WILLIAMS and DRUFFEL, 1987). This suggests that this fraction of the DOC (oxidizable by UV radiation) is recycling throughout the oceanic water column including the contiguous, semi-isolated coastal basins. However, the ^{14}C activity of the DOC (or some fraction thereof) oxidizable only by high temperature combustion (HTC) (SUGIMURA and SUZUKI, 1988) must be significantly higher than that of the UV-oxidizable fraction to account for the observed covariance between DOC_{HTC} concentration with apparent oxygen utilization (AOU) in some oceanic regimes. There is some indirect evidence for this "extra" DOC having an elevated ^{14}C content (DRUFFEL, WILLIAMS and SUZUKI, 1989), but there are no hard data. The significance of this paradox of the dissolved/colloidal organic matter (see WILLIAMS and DRUFFEL, 1988) to organic carbon cycling in coastal basins in general, and the Santa Monica Basin in particular, is in assessing the role of DOC in balancing particle fluxes, *in situ* biochemical oxidation, and sedimentary organic matter burial against carbon fixed by photosynthesis and/or entering the system from non-marine sources (JACKSON, AZAM, CARLUCCI, EPPLEY, FINNEY, GORSLINE, HICKEY, HUH, JAHNKE, KAPLAN, LANDRY, SMALL, VENKATESAN, WILLIAMS and WONG, 1989).

Perhaps the most important impact of the "old" recycled DOC is via its possible association with both suspended and sinking POC which may result in low $\Delta^{14}\text{C}$ values for these particulate phases at depth (Fig. 1) and hence, ambiguity in reading the bomb- ^{14}C signal or radiocarbon clock of these phases *vis-à-vis* their origin and eventual burial in the sediments.

$\delta^{13}\text{C}$ values for UV-oxidizable DOC (DOC-UV) in the basins (Table 3) vary little with depth or location and only slightly between cruises - the mean values for all depths for CaBS-4 and CaBS-7 samples were $-21.4 \pm .5\text{‰}$ ($n=6$) and $-21.8 \pm 2\text{‰}$ ($n=5$) respectively. The average $\delta^{13}\text{C}$ value for the North Central Pacific (3-900m, $n=17$) in 1985 and 1987 was $-21.1 \pm .3\text{‰}$ ($n=22$) (WILLIAMS and DRUFFEL, 1987, unpublished data); and for the SIO pier and San Diego Trough area (Table 1) in 1967-1968 was -22.5‰ (WILLIAMS and GORDON, 1970; results which could be 1‰ more negative because of earlier procedural problems). The DOC-UV concentrations for CaBS-4 and CaBS-7 (Table 3) were nearly identical at all mid-basin and shelf depths, and were $\sim 4\mu\text{mol}$ higher at depth than the North Central Pacific DOC-UV values. Thus, the concept of a common origin for the UV-oxidizable fraction of DOC present in both the open ocean and coastal basin is strengthened by $\delta^{13}\text{C}$ values and total DOC-UV content - in addition to the $\Delta^{14}\text{C}$ signatures.

4.2 Suspended particulate organic matter: POC(susp) and PON(susp)

Suspended POM, whose carbon concentration is of the order of 1% of the DOC-UV (Table 3), has estimated residence times in the upper mixed layer (photic zone) of 20-39 days (SMALL, LANDRY, EPPLEY, AZAM and CARLUCCI, 1989) and 3 to >100 days (EPPLEY, RINGER and BETZER, 1983) based on new production and sinking fluxes, alternatively 4 to 7 days based on thorium isotope measurements (HUH and BEASLEY, 1987). In the deeper basin waters, these POC (susp) residence times may be of the order of 5 to 10 years, as has been reported for thorium isotope measurements in the open ocean (BACON and ANDERSON, 1982). Particles collected in the SMB by *in situ* filtration were nominally $>1\mu\text{m}$ in size but would include finer particles scavenged on larger ones (McCAVE, 1984) and trapped on clogged filters. Thus, the origin of suspended POM is not necessarily localized to the basin system, but is a combination of: (1) POM advected into the basin by surface currents and deep water exchange across the sills; (2) POM derived from the basin slopes through resuspension and subsequent incorporation into various levels within the water column; and (3) POM resulting from fractionation of sinking POM into finer, suspended

particles. Transmissometer measurements within the SMB water column (NELSON *et al.*, 1987; HUH, SMALL, NIEMNIL, FINNEY, HICKEY, KACHEL, GORSLINE and WILLIAMS, 1990; B. HICKEY, personal communication) indicate that there are no permanent nepheloid layers beneath the euphotic zone but do not preclude sporadic injections of suspended particles at depth from the basin shelf or slopes. Surface and deep current measurements and tracer experiments give residence times for water parcels (with their associated suspended POM) ranging from 3-20d in the upper 200m, up to several years for sporadic renewal of deep basin waters (JACKSON *et al.*, 1989). These renewal times are short compared to estimated suspended particle residence times in the surface mixed layer, suggesting caution in relating chemical and isotopic properties of POC (susp) and PON (susp) to locally-sinking POM or other indigenous sources of suspended organic particles. Clearly, advected particles may dominate those of local origin, especially in deeper waters.

$\Delta^{14}\text{C}$ values for POC (susp) decrease significantly with depth (Table 3, Fig. 1) with a minimum value of -46‰ at 700m. Neither the $\delta^{13}\text{C}$ signature nor the POC (susp) concentration is strikingly different at 700m from those same parameters measured in the 510 and 850m samples, suggesting that this particular sample was not significantly "contaminated" with oil or other hydrocarbons. Without replicate samples, this low $\Delta^{14}\text{C}$ value cannot be attributable to the filtration of a spurious cloud of "old" suspended particles; nor can it arise from resuspended surficial sedimentary organic particles whose $\Delta^{14}\text{C}$ values were -10 and -21‰ in the upper 2.5cm of bulk sediment (Table 5). Two facts are evident from the POC (susp) ^{14}C activities: (1) Reduced carbon, recently fixed photosynthetically at the surface, whose D^{14}C value should be similar to the surface water D^{14}C value (94‰), does not constitute all of the suspended POC. Low ^{14}C -containing organic particles and/or DOC must contribute to the POC (susp) $\Delta^{14}\text{C}$ values below 100m assuming that we are not dealing with spurious oil contamination; and (2) the PO^{14}C (susp) activities are lower than those for the sinking POC collected on the same cruise at the same depths (Fig. 1). This might seem to support longer turnover times for the POC (susp) than for the POC (sink), but this is only true if inclusion of "old" sedimentary particles and/or DOC are excluded, and if the POC (susp) has a common origin with the POC (sink).

$\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for POC (susp) do not reflect the isotopic signatures in either POC (sink) or DOC (Figs 2 and 3). $\delta^{13}\text{C}$ values for POC (susp) are more negative by $\sim 1\text{‰}$ than for DOC at comparable depths (Table 3), as was earlier observed by WILLIAMS and GORDON (1970) (Table 1). JEFFREY, PFLAUM, BROOKS and SACKETT (1983), who measured $\delta^{13}\text{C}$ values for POC (susp) in numerous shallow profiles ($<400\text{m}$) from various oceans, suggested that biological reworking of POC (susp) in the pycnocline caused fractionation of POC- $\delta^{13}\text{C}$ towards lighter isotopic values so that $\delta^{13}\text{C}$ values increased with depth. The opposite was found in the SMB where the two shallow samples (mid-basin and shelf) were lighter in ^{13}C than samples at 500m and below by 3.0‰ and 1.39‰ , respectively. This depletion in $\delta^{13}\text{C}$ suggests a terrestrial contribution is possible; but, if so, none of the shallow POC (sink) $\delta^{13}\text{C}$ values (Table 4) is isotopically lighter than those measured in the deeper samples. Flocculents from the outfalls and suspended particles from Los Angeles River storm waters (Table 1) have $\delta^{13}\text{C}$ values which vary from -23.7 to -24.9‰ , and $\delta^{15}\text{N}$ values from 2.4 to 3.0‰ , but there is no compelling evidence from $\delta^{15}\text{N}$ values for PON (susp) supporting an appreciable contribution of outfall-derived suspensoids (this general question of terrestrial contributions to all organic phases in the Santa Monica Basin system is amplified in the Discussion section). The approximately 2‰ increase in $\delta^{15}\text{N}$ at 700 and 850m relative to the 80, 100 and 510m samples (Table 3, Fig.,2) follows previous measurements of lighter $\delta^{15}\text{N}$ values in PON (susp) in the euphotic zone than at depth (ALTABET, 1988).

4.3 Sinking particulate organic matter [POC(sink) and PON(sink)], and total mass and organic carbon and nitrogen fluxes

There are two caveats concerning the Soutar particle trap deployments. The first is that the shallower traps (100 and 200m) collected significantly fewer organic particles than smaller-diameter, multicylinder "Vertex" traps (WELSCHMEYER and LORENZEN, 1985; SMALL *et al.*, 1989) deployed at these same shallower depths (see LANDRY, PETERSON and ANDREWS, 1992). This apparent "under-collection" by the shallow Soutar cones is probably related to motion-induced orientation problems in the upper 200m. We consider that the deeper traps (500-700 and 850m) did collect representative flux samples based on the following observations. JAHNKE (1990) has shown that organic carbon burial rates correspond to sinking carbon fluxes at 850m adjusted for sediment oxidant consumption (see also JACKSON *et al.*, 1988). Also ratios of organic carbon fluxes to total mass fluxes (from Tables 6 and 8) decrease from 0.13-0.20 at 100 and 200m to 0.055-0.084 at 850m, primarily because of a decrease in organic matter trapped at depth coupled with an increase in total mass flux with depth (Fig. 4). These ratios do not preclude under-trapping of sinking organic matter in the upper 200m with Soutar traps, but do show an approximately two-fold decrease in the amount of organic C trapped relative to total mass at 850m compared to 100 or 200m.

The second caveat is that the 100, 200 and 350m traps contained variable but often significant numbers of swimmers, primarily copepod nauplii, all of which could not be separated from the sinking particles. Such swimmers will have $\Delta^{14}\text{C}$ values close to that of surface DIC, and $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values representative of marine end-members, so the isotopic signatures for sinking organic matter will be biased in these shallow samples. The swimmers will also lead to the total organic C and N fluxes for passively sinking particles being over-estimated. Conclusions based solely on fluxes and total organic matter collected in the shallow traps are most subject to these errors.

There are no $\Delta^{14}\text{C}$ values reported previously for POC (sink) in coastal basins. In the North Central Pacific, PO^{14}C (sink) at 4120 and 5120m were 85 ± 7 and $114 \pm 7\text{‰}$, respectively (DRUFFEL and WILLIAMS, 1990). These values are higher than those measured at depth in the SMB, possibly reflecting the higher surface DI^{14}C content in the North Central Pacific.

$\Delta^{14}\text{C}$ values for POC (sink) from five CaBS cruises (Table 4 and Fig. 1) generally decrease to 700m at the mid-basin station with no further reduction in ^{14}C at 850m. All of the $\Delta^{14}\text{C}$ values are greater than -75‰ , the maximum pre-bomb value for surface DIC prior to 1950 (AD); or, conversely, all of the POC (sink) samples contain bomb- ^{14}C - as would be expected. The interesting point is that this diminution of ^{14}C activity with depth requires that some fraction of the POC (sink) contains organic carbon which is not surface-derived and/or derived from surface waters >35 years ago. At 850m (Fig. 1) the PO^{14}C (susp) and PO^{14}C (sink) values converge to a mean $\Delta^{14}\text{C} = 27 \pm 21\text{‰}$ (range = $2-52\text{‰}$, $n=4$) from a mean value of $96 \pm 14\text{‰}$ (range = $78-110\text{‰}$, $n=4$) at 80-200m, the same value as DI^{14}C at 5m. The general correspondence between the ^{14}C contents of sinking and suspended POC suggests a common source and similar histories for these phases, although it does not rule out their having disparate sources with closely related isotopic signatures.

One clue to the source of "older" sinking POC is the lower $\Delta^{14}\text{C}$ values for all of the particle trap POM samples from the shelf. $\Delta^{14}\text{C}$ decreased from a mean of $67 \pm 7\text{‰}$ at 100m to $39 \pm 6\text{‰}$ at 200m and $-13 \pm 14\text{‰}$ at 350m, ~50m above the seafloor. Perhaps, fortuitously, the differences between the surface and deep samples at mid-basin (69‰) and on the shelf (80‰) are not significantly different. This might be interpreted as a "common source", e.g. resuspended, low- ^{14}C activity, shelf sedimentary organic matter (SOM), mixing with recently-fixed organic carbon, where the amount of common-source POM decreases with distance from its origin on the

shelf. Since persistent nepheloid layers have not been observed at depth in mid-basin waters, these low $\Delta^{14}\text{C}$ values for POC (sink) on the shelf are more likely from resuspended ^{14}C -poor SOM, while another process (e.g. adsorption of "old" DOC) may also occur with the mid-basin samples.

$\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for sinking POM (Table 4, Figs 2 and 3) vary significantly between cruises, between depths, and between the mid-basin and shelf stations. The $\delta^{15}\text{N}$ values are particularly variable at mid-basin where $\delta^{15}\text{N}$ at 850m (cruise CaBS-3) was 5.50‰, and that at 500m (cruise CaBS-7) was 10.20‰. Only the CaBS-4 deployment showed smooth $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ profiles (Figs 2 and 3). If all measurements at each depth are averaged, vertical changes in $\delta^{15}\text{N}$ are much less pronounced (see Fig. 3 and JACKSON *et al.*, 1989), and the average $\delta^{15}\text{N}$ values at 100 and 500m (8.87 and 8.42‰) are similar to those measured at the same two depths (8.0 and 8.0‰, respectively) for samples from two, 2.5 day, free-floating conical trap deployments in 1983 (NELSON *et al.*, 1987).

It is apparent from Fig. 3 that the vertical profiles of $\delta^{13}\text{C}$ (with the exception of 700 and 850m on CaBS-3) are about 1‰ heavier in ^{13}C on cruises CaBS-3, 7 and 10 than on CaBS-4, both at mid-basin and on the shelf. No similar differences occur with the $\delta^{15}\text{N}$ values. $\delta^{15}\text{N}$ on cruise CaBS-4 is not systematically lower as would be expected if the lighter $\delta^{13}\text{C}$ values on CaBS-4 originated from terrestrial inputs. It is difficult to attribute variations in $\delta^{13}\text{C}$ values, be they dissolved, suspended or sinking organic phases to terrestrial sources, since $\delta^{13}\text{C}$ values for the terrestrial end-members (-23.7 to -26.4‰, Table 1) are only about 3‰ or less depleted in ^{13}C relative to the dissolved and particulate organic phases in question. Variations in the $\delta^{13}\text{C}$ signatures for the sinking organic matter are most likely a function of differing $\delta^{13}\text{C}$ values for the various sources compromising the sinking POM, and do not arise from trophic level fractionation processes. These sources include (see Table 1): phyto- and zooplankters ($\delta^{13}\text{C} \equiv -19$ to -22 ‰); kelp ($\delta^{13}\text{C} \equiv -14$ to -21 ‰); sole and prawns ($\delta^{13}\text{C} \equiv -16$ to -18 ‰), resuspended sedimentary organic matter ($\delta^{13}\text{C} \equiv -20$ to -23 ‰), and sewage or storm water particles ($\delta^{13}\text{C} = -23.7$ to -26.4 ‰). The uniform, CaBS-4 $\delta^{13}\text{C}$ signatures for sinking POM may have been a result of predominantly one source of particles (e.g. copepod-derived organic matter) being trapped during this deployment.

Some variation in the $\delta^{13}\text{C}$ signatures could have arisen from changes in CO_2 (aq) concentrations (RAU, TAKAHASHI and DES MARAIS, 1989) which would alter the $\delta^{13}\text{C}$ values of the organic phases. However, the small interannual variations in CO_2 (aq) or temperature in this region would result only in small changes in the $\delta^{13}\text{C}$ signatures.

How then, are the corresponding $\delta^{15}\text{N}$ signatures interpreted? The large, and seemingly unrelated excursions in $\delta^{15}\text{N}$ between deep and shallow trap samples and between deployments must be a function of isotopic fractionation during phase changes, and reutilization and recycling in the euphotic zone coupled with a sporadic input of sewage particulates. Such processes as: (1) ^{15}N enrichment in higher trophic levels of the order of 3.5‰ per trophic level (FRY, 1988 and references therein); (2) ^{15}N depletion in fecal pellets (~1-2‰) relative to the parent zooplankton body tissue coupled with an approximate 2‰ fecal pellet enrichment over the apparent zooplankton food source (ALTABET and SMALL, 1990); (3) variable $\delta^{15}\text{N}$ values in photosynthetically fixed NO_3^- and NH_4^+ reflecting variations in NO_3^- and NH_4^+ during denitrification and recycling of nitrogen in the euphotic zone (LIU and KAPLAN, 1989; CHECKLEY and ENTZEROTH, 1985; CHECKLEY and MILLER, 1989); and (4) ^{15}N fractionation between sinking and suspended particles (ALTABET, 1988). Any or all of these processes could be operating within the complex nitrogen cycle in the SMB system, making it difficult, if not impossible, to relate any specific $\delta^{15}\text{N}$ profile of PON (susp) to any one (or more) direct sources or processes. What is evident is that on cruise CaBS-3, ^{15}N -depleted particles (presumably sewage-derived) dominated the trapped particles at 350m on the shelf ($\delta^{15}\text{N} = 3.55$) and even appeared to have entered the 850m trap at

the mid-basin station. Such “sewage” contributions were not apparent in the other mid-basin PON (susp) profiles (Fig. 3), but could have enhanced the regular depletion of ^{13}N in the 350m traps relative to the 100 and 200m samples on the shelf. This depletion could also arise from resuspension of sedimentary organic particles whose $\delta^{13}\text{C}$ values (e.g. 6.4 at 0–1.25cm, Table 5 for core BC-89) were more similar to the PON (sink) $\delta^{15}\text{N}$ values at 350m on the shelf (and at 850m at mid-basin) than to, for example, zooplankton tissue (Table 1). The significant increases in mass and C and N fluxes on the shelf, and in mass fluxes at mid-basin (Fig. 4), support the hypothesis for such a sedimentary contribution from the shelf/slope regions; so do the lower ^{14}C activities in the 350 and 700 and 850m trapped particles and in the surface sedimentary organic matter.

4.4 Sedimentary organic matter (SOC)

$\Delta^{14}\text{C}$, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for the upper 20cm of box core No. BC-89 are listed in Table 5 and depicted in Figs 1 and 2.

The $\Delta^{14}\text{C}$ values measured in the surficial sediments of the SMB (-10 ± 6 and $-21 \pm 6\text{‰}$, 0 to 2.5cm) clearly show the presence of bomb ^{14}C . As expected, a $\Delta^{14}\text{C}$ result obtained for a grab sample collected during the pre-bomb period of 1956 in the SMB ($-147 \pm 19\text{‰}$, 1280 $\pm$ 150y BP; EMERY and BRAY, 1962) was much lower than our 1985 values. There is, however, reasonable agreement between the 1956 $\Delta^{14}\text{C}$ values at 1cm and 15cm ($-147 \pm 19\text{‰}$ and $-258 \pm 19\text{‰}$, respectively, EMERY and BRAY, 1962) and our 1985 $\Delta^{14}\text{C}$ values at 7cm and 13cm ($-154 \pm 5\text{‰}$ and $-307 \pm 5\text{‰}$). Our $\Delta^{14}\text{C}$ value at 20cm appears to be anomalously low ($-416 \pm 4\text{‰}$) when compared with any of EMERY and BRAY’s results for the upper 90cm in three cores from the Santa Monica Basin and the adjoining San Pedro Basin. This may be evidence of “old”, redeposited carbon that EMERY and BRAY (1962) postulated to be in surficial SOC from the basins off Southern California.

For the Santa Monica and San Pedro basins, EMERY and BRAY (1962) calculated six sedimentation rates ranging from 38 to 145cm 10^{-3}y^{-1} (mean = 66cm 10^{-3}y^{-1}) based on the difference in ^{14}C activities between a surface and deep ($>300\text{cm}$) $\Delta^{14}\text{C}$ measurement. These sedimentation rates are similar to the 31cm 10^{-3}y^{-1} calculated for core BC-89 from ^{210}Pb measurements in the top 20cm (Table 5, 20cm $\times$ $10^3/642\text{y}$). The ^{14}C sedimentation rate for Core BC-89 is much less ($\sim 4\text{cm } 10^{-3}\text{y}^{-1}$) based on near-surface (6.2 to 20cm) $\Delta^{14}\text{C}$ results. This latter rate may be unique to core BC-89, as large variability in $\Delta^{14}\text{C}$ results were also observed by EMERY and BRAY (1962) in near-surface SOC.

$\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for SOC (Table 5, Fig. 2) do not show extreme excursions within the sedimentary column. $\delta^{13}\text{C}$ (-21.79 to -22.62‰ , mean = -22.11‰) and $\delta^{15}\text{N}$ (6.09 to 8.74 ‰ , mean = 6.98 ‰) values fall within the envelope of data points on a regression plot of $\delta^{13}\text{C}$ vs. $\delta^{15}\text{N}$ for sedimentary organic matter from continental shelf sediments of the northeast Pacific (PETERS, SWEENEY and KAPLAN, 1978), but are not themselves significantly correlated (Fig. 5F). There are no striking correlations of $\delta^{13}\text{C}$ or $\delta^{15}\text{N}$ with sedimentation events which have occurred during the past 500 years; e.g. the decline in sedimentation rates in core BC-89 which occurred 200 years ago (HUH, ZAHNLE and SMALL, 1990) at the time of an extensive turbidite intrusion into the basin (MALOUTA, GORSLINE and THORNTON, 1981), or with episodic, high, storm-derived run-off at 10–30y intervals (MALOUTA, GORSLINE and THORNTON, 1981). Nor are there any tight correlations of $\delta^{13}\text{C}$ or $\delta^{15}\text{N}$ with past productivity maxima (SMITH and EPPLEY, 1982) or total SOC content for the 1922–1979 period. Note that total SOC concentration profiles at mid-basin and shelf sites show a maxima in SOC content in 1965–1970, when inputs of wastewater particulates from the White’s Point outfall reached a maximum (FINNEY and HUH, 1989; EGANHOUSE and

KAPLAN, 1988). There is also no reflection in the isotopic ratios for core BC-89 of the SOC and SON concentration maxima and $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ minima observed at 6-10cm in a core taken at 60m bathymetric depth on the San Pedro shelf in 1981 (EGANHOUSE and KAPLAN, 1988). These SOC and SON maxima and isotopic minima on the shelf are a direct result of the maximum discharge of terrestrially-derived sewage particles, which occurred in the late 1960s. Thus, the terrestrial C and N isotopic signatures have been poorly preserved in the mid-basin, sedimentary column (at least in 1.25cm sections), implying that significant biochemical oxidation-solubilization of sewage particulates occurs en route to their eventual sequestration in the sediments.

With respect to the isotopic signatures for the mid-basin suspended, sinking and dissolved organic phases (Figs 1, 2 and 3), the surficial (0-1.25cm) sedimentary organic matter most closely resembles: (1) the 850m sinking POC from Cruise CaBS-8 in ^{14}C content ($\Delta^{14}\text{C}_{\text{SOC}} = -10 \pm 6\text{‰}$, $\Delta^{14}\text{C}_{\text{POC(sink)}} = 2 \pm 6\text{‰}$); and (2), the 850m suspended POC from Cruise CaBS-4 ($\Delta^{14}\text{C} = 9 \pm 6\text{‰}$). The SOC $\delta^{13}\text{C}$ value (-22.15‰) is slightly lighter than the mean ($n=5$) value for the sinking POC at 850m (-21.78‰) or the DOC value at 850m (-21.50‰ , $n=2$), but slightly heavier than the suspended POC value at 850m from Cruise CaBS-4 (-23.08‰). $\delta^{15}\text{N}$ for the 0-1.25cm SON (6.37‰) is significantly lighter than the mean ($n=5$) $\delta^{15}\text{N}$ (7.42‰) for the mid-basin sinking particles; however, this mean value is misleading because the two extreme $\delta^{15}\text{N}$ values (Table 4) 5.50 and 5.97 are indicative of contamination from terrestrial (sewage) inputs. Thus the source of total sedimentary organic carbon can not be attributed either to any one of the indigenous organic phases in the water column or to the wastewater outfalls. This result was unexpected but makes it difficult to link unequivocally the terrestrial vs. marine sources to the SOM.

5. DISCUSSION

5.1 Terrestrial (sewage, storm run-off, industrial organics) contribution to the SMB organic phases

The average $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures for White's Point sewage flocculents are -24 and 2.5‰ , respectively (Table 1). $\delta^{13}\text{C}$ in Los Angeles river storm water suspended solids is -25‰ (Table 1). No values for $\delta^{15}\text{N}$ in river solids, or for $\Delta^{14}\text{C}$ in sewage or runoff organics are available. $\Delta^{14}\text{C}$ values in natural oil seeps, anthropogenic petroleum products and spills, and petroleum-derived industrial chemicals will be -1000‰ , and in modern organic matter (grass, leaves, paper, etc.) will be taken as 200‰ , equivalent to that of contemporary atmospheric CO_2 (Table 1).

A rough estimate of the contribution of sewage-derived organic flocculents to the SMB organic phases can be calculated using the two-source mixing equation:

$$X_s [\delta^{13}\text{C}_s] + X_m [\delta^{13}\text{C}_m] = [\delta^{13}\text{C}_{\text{SMB}}]$$

$$X_s [\delta^{15}\text{N}_s] + X_m [\delta^{15}\text{N}_m] = [\delta^{15}\text{N}_{\text{SMB}}] \quad (1)$$

where X_s and X_m are the fractions of sewage and marine end-members in the SMB organic phases having the appropriate measured isotopic signatures ($\delta^{13}\text{C}_{\text{SMB}}$ and $\delta^{15}\text{N}_{\text{SMB}}$). Table 9 lists the percentages of sewage-derived organics in the SMB organic phases and includes end-members and organic phase isotopic signatures. The latter values are means of both mid-basin and shelf measurements.

FIG.5. Plots of $\Delta^{14}\text{C}$ vs. $\delta^{13}\text{C}$ or $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ vs. $\delta^{15}\text{N}$ for the various organic phases (all values). A and B: POC (sink), $\blacktriangle$ = mid-basin, $\blacktriangle$ = shelf; POC (susp), $\bullet$ = mid-basin, $\times$ = shelf. C: DOC, $\circ$ = mid-basin, $\otimes$ = shelf. D: All POC (sink) and POC (susp) values, $\bullet$ = mid-basin, $+$ = shelf. E: SOC, $\blacksquare$ = $\Delta^{14}\text{C}$ vs. $\delta^{13}\text{C}$, $\square$ = $\Delta^{14}\text{C}$ vs. $\delta^{15}\text{N}$. F: SOC, $\times$ = all values, 0 to 20cm.

TABLE 9. Percentage contribution of sewage organics to the basin organic phases calculated from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values

Mean $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for end members (‰)			
Pool	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	Reference
NO_3^-	—	~8.5	Table 1 (11)
White's Point flocculents	-24	2.5	Table 1 (17,18)
DOC	-21.6 (n=11)	—	Table 3
POC(susp)	-23.3 (n=5)	—	"
PON(susp)	—	8.3 (n=5)	"
POC(sink)	-21.7 (n=25)	—	Table 4
PON(sink)	—	8.1 (n=24) ¹	"
SOC(upper 3.75cm)	-22.2 (n=3)	—	Table 5
SON(upper 3.75cm)	—	7.0 (n=3)	"

Taking the sewage end member signatures as $\delta^{13}\text{C}_s = -24\text{‰}$ and $\delta^{15}\text{N}_s = 2.5\text{‰}$ and the marine end member signatures as $\delta^{13}\text{C}_m = -21.5\text{‰}$ and $\delta^{15}\text{N}_m = 8.5\text{‰}$, the percentage contributions of sewage organic matter, X_s , to the basin organic phases are:

Pool	From $\delta^{13}\text{C}$	From $\delta^{15}\text{N}$
DOC	4	—
POM(susp)	(72)	3
POM(sink)	8	7
SOM	28	25

¹Does not include $\delta^{15}\text{N} = 3.55\text{‰}$ from 350m, shelf station, Cruise CaBS-3.

The choice of end-member isotopic signatures is largely subjective, and does not include non-sewage terrigenous organic matter. $\delta^{15}\text{N}_s$ and $\delta^{13}\text{C}_s$ values are from SWEENEY and KAPLAN (1980b, Table 1). For the marine end-members, we have chosen 8.5‰ for $\delta^{15}\text{N}_m$, the mean value for NO_3^- in upper 100m of the San Pedro Basin (LIU and KAPLAN, 1989), and -21.5‰ for $\delta^{13}\text{C}_m$, based on DOC and POC(sink) values at 100 or 200m.

The percentage contribution of sewage flocculents, X_s , to the SMB organic phases (Table 9) is 3-8% for the sinking and suspended POM (excluding the 72% value for POC (susp) based on $\delta^{13}\text{C}$) and 25-28% for the surficial SOM. The wider spread in the $\delta^{13}\text{C}$ end-members makes the percentages calculated using $\delta^{15}\text{N}$ values more credible. The 25% calculated for surficial SOM corresponds to the value found by EGANHOUSE and KAPLAN (1988) for SOM at 23-34cm depth in a core from the San Pedro shelf at 60m, where the corresponding percentages for the 0 to 23cm sections varied from ~40% at 0-2cm to 95% at 2-4cm. Thus, the contribution of sewage flocculents to the water column organic phases in the mid-basin and shelf waters is probably less than 10%, while the surficial, mid-basin SOM could contain ~25% terrestrial components, assuming the terrestrial end-member assignments are valid.

There are also indications from $\delta^{15}\text{N}$ values for the POC (sink) of episodic inputs of terrestrial POM at 350m on the shelf ($\delta^{15}\text{N} = 3.55\text{‰}$, CaBS-3), 850m mid-basin ($\delta^{15}\text{N} = 5.50\text{‰}$, CaBS-3), and 850m, mid-basin ($\delta^{15}\text{N} = 5.97\text{‰}$, CaBS-10). Using the $\delta^{15}\text{N}_s$ and $\delta^{15}\text{N}_m$ end-member values from Table 9, the estimated contribution of sewage to the POM particle trap contents at 350 and at 850m for CaBS-3 and 850m for CaBS-10 are 83%, 50% and 42%, respectively.

However such high sewage inputs are not reflected in the $\delta^{13}\text{C}$ signatures of the POC (sink), which suggests that there is reworking and fractionation of the sewage-derived POC without corresponding changes in the $\delta^{15}\text{N}$ values (see MYERS, 1974, and SWEENEY, KALIL and KAPLAN, 1980).

Independent of the isotopic evidence, the doubling in the mean mass flux between depths of 100 and 200m to 350m on the shelf, coupled with a 40% increase in the corresponding CN flux (Fig.4), implies that there is export of POM, including terrestrial components, across the shelf. As well as the increased mass flux on the shelf, there is a 40% increase in total mass flux between depths of 100 to 850m at mid-basin (Fig.4), although its origins could be from the seaward slopes of the basin as well as the coastal shelf and slope. These inputs augment the POM derived directly from marine organisms.

With respect to the $\Delta^{14}\text{C}$ signatures, sewage inputs cannot be assessed without knowing the $\Delta^{14}\text{C}$ value for sewage flocculents and solubles. Sewage is composed mainly of faeces, cellulose (toilet paper) and variable quantities of chloroform-soluble hydrocarbons (R. EGANHOUSE, personal communication). Its $\Delta^{14}\text{C}$ signature will be a mix of recent ($\Delta^{14}\text{C} = 200\text{‰}$) and ancient ($\Delta^{14}\text{C} = -1000\text{‰}$) components. The hydrocarbon fraction, however, could contribute to lowering $\Delta^{14}\text{C}$ values in the water column and sedimentary organics (see below).

5.2 Origin of $\Delta^{14}\text{C}$ gradients in mid-basin and shelf organic phases (water column)

The most obvious carbon source resulting in the diminution of ^{14}C activities in the DOC, POC (susp) and POC (sink) is petroleum derived from natural seeps, sewage discharges, oil spills and atmospheric inputs.

As stated earlier, the DOC $\Delta^{14}\text{C}$ profile can be primarily attributed to the influx of open-ocean water into the basin, rather than to soluble hydrocarbons, whose concentrations (in the boiling point range of n-tetradecane to n-dotriacontane) in 1-3m surface waters and in samples collected 10m off the bottom have been reported to be 0.03 to 80 $\mu\text{g l}^{-1}$ (~ 0.001 to 3 $\mu\text{mol C l}^{-1}$) (RISEBROUGH and BURLINGAME, 1978). Assuming a mean total DOC concentration of 44 $\mu\text{mol l}^{-1}$ at 500 to 850m (Table 3), and a surface (5m) $\Delta^{14}\text{C}$ value of -197‰ for the source DOC, then it would require an additional 10 μmol (23% of the total UV oxidizable DOC!) of hydrocarbon-C per liter ($\Delta^{14}\text{C} = -1000\text{‰}$) to attain the measured value of $\sim -470\text{‰}$ (Table 3) for the deep water DOC. However, the resultant $\delta^{13}\text{C}$ value of this DOC would then be -23.3‰ rather than the mean measured value of -21.75 (Table 3), assuming a $\delta^{13}\text{C} = -28\text{‰}$ for petroleum products.

With respect to the sinking particles, CRISP, BRENNER, VENKATESAN, RUTH and KAPLAN (1979) reported a total hydrocarbon flux of 0.071 g ($\sim 0.006\text{mol C}$) $\text{m}^{-2}\text{y}^{-1}$ for the Santa Monica Basin at $\sim 850\text{m}$. The mean C flux at 850m (Table 7) (is $\sim 15\text{g m}^{-2}\text{y}^{-1}$ and 45 $\text{g m}^{-2}\text{y}^{-1}$ at 350m on the shelf, which gives hydrocarbon contents of $\sim 0.24\%$ (mid-basin) and 0.08% (shelf) assuming the same flux of hydrocarbons at both sites. If the $\Delta^{14}\text{C}$ value for the POC sinking from the surface was equal to surface DIC (94‰), then the $\Delta^{14}\text{C}$ for POC (sink) at 850m (or at 350m on the shelf) would be lowered only to 91.3‰ (and to 93.1‰ on the shelf) with admixture of 0.24% (or 0.08% on the shelf) petroleum hydrocarbons ($\Delta^{14}\text{C} = -1000\text{‰}$). The measured means of $\Delta^{14}\text{C}$ at 850m and 350m, however, are 41 and -13‰, respectively (Table 4).

A similar calculation for POC (susp) can be made assuming (1) the $\Delta^{14}\text{C}$ source for POC (susp) has a value of 94‰ with a mean C concentration of 0.27 $\mu\text{mol l}^{-1}$ at 500-850m (Table 3), and (2) the mean concentration of hydrocarbon -C in suspended particles is 0.005 $\mu\text{mol l}^{-1}$, based on data from RISEBROUGH and BURLINGAME (1978) and DeLAPPE, RISEBROUGH and WALKER (1983). These assumptions give a POC (susp) ^{14}C activity of 74‰ versus the mean measured $\Delta^{14}\text{C}$ value

of $\sim -3\text{‰}$ at 510 to 850m (mid-basin).

These crude estimates of the contribution of petroleum to the organic phases in mid-basin and shelf waters while not precluding its presence in these phases, do suggest that the contribution is minimal ($<1\%$) based on the data available for hydrocarbon contents of DOC, POC (susp) and POC (sink) in the basin.

If natural oil seeps and/or anthropogenic petroleum inputs are not responsible for the diminution of ^{14}C activities with depth, then the most plausible explanation (as mentioned earlier) is that there are contributions of low ^{14}C activity, surficial sedimentary organic particles which either enter the mid-basin water column sporadically from the basin slopes, or, in the case of the shelf waters, are directly injected into the overlying waters via resuspension by slumping and internal waves. Downslope resuspension of subsurface sediment (5-10cm) injected into the near-bottom water by the activities of benthic animal activity may also contribute significant amounts of low ^{14}C activity SOC. The observed increases in mass and CN fluxes with depth (Tables 6 and 7, Fig.4) are evidence for these processes, most notably at the shelf stations; although the frequency with which these inputs occur, especially the introduction of slope particulates into the deep basin waters, is unknown. The sporadic presence of nepheloid layers in the deep basin waters (B. HICKEY, personal communication) lends credence to the interdigitation mechanism.

Assuming the $\Delta^{14}\text{C}$ values of the POC (susp) derived from the slope surficial sediments to be -15‰ (the same as the mean value measured for the mid-basin 0-2.5cm SOC, Table 5), and the original source $\Delta^{14}\text{C}$ of POC (susp) to be 94‰ (same as 5m DIC), then the resultant percentage contribution of SOC to the POC (susp) at 510 and 850m is estimated to be 61 and 78%, respectively.

The corresponding percentage contributions of resuspended sedimentary POC to the POC (sink) at 500, 700 and 850m in the basin will be 27, 58 and 50% respectively, where the mean measured $\Delta^{14}\text{C}$ values at these depths are 65, 31 and 39‰ respectively (Table 4), and the two source $\Delta^{14}\text{C}$ values are again taken as 95‰ [surface POC (sink)] and -15‰ (sedimentary POC). For the 350m POC (sink) at the shelf stations, the SOC contribution would be 98% of the total POC (sink) to give its resultant mean $\Delta^{14}\text{C}$ value of -13‰ .

Obviously, the $\Delta^{14}\text{C}$ value of the SOC derived from the basin slopes and/or shelf may be higher or lower than the assumed value of -15‰ , depending upon how thick a surficial layer of sedimentary POC was injected into the deep basin or resuspended on the shelf. The $\Delta^{14}\text{C}$ values may also be a function of sedimentary grain size. Coarse particles are rich in fresh plant debris while fine-grained particles contain more fossil organic matter (THOMPSON and EGLINTON, 1978). Thus one might expect that $\Delta^{14}\text{C}$ values would vary at any given sediment depth depending on grain size. This mechanism does not account for the ^{14}C activity ($\Delta^{14}\text{C} = -46\text{‰}$) of the suspended POC at 700m (Table 3). The only explanation for this anomalously low result is contamination with "dead" (petroleum) carbon. An addition of 6.3% "dead" carbon ($\Delta^{14}\text{C} = -1000$) is needed to lower the $\Delta^{14}\text{C}$ of POC (susp) at 700m (assumed to be 20‰) to -46‰ . This invokes a spurious contamination event, probably sampling related, as opposed to the unlikely ubiquitous contribution of petroleum to the POC (susp) at 510m and 850m discussed above.

We feel that the estimated 27, 58 and 50% contributions of SOC to the POC (sink) are high, but not unreasonably so, considering the probability of sporadic SOC inputs occurring during the 98-233d particle trap deployments. However, the estimates of 61 and 78% SOC contributions to the POC (susp) are based solely on a single profile with 6h *in situ* particle filtrations at 510, 700 and 850m. This collection may have occurred fortuitously during a period of significant SOC input, so may be unrepresentative of the normal background POC (susp). If a significant fraction of the POC (susp) was derived from near-bottom waters from the slope being advected into the

basin, then the resultant isotopic signatures would be strongly influenced by the isotopic characteristics of this advected POC (susp).

There is also the intriguing possibility of adsorption of low ^{14}C activity DOC-UV onto suspended (or sinking) particles which, if it occurred during the 5 to 10 year residence time of the POC (susp) in the water column, could contribute to the gradient in $\Delta^{14}\text{C}$. It would require 12, 25 and 15% of the POC (susp) to be adsorbed DOC at 510, 700 and 850m respectively, to give the measured POC (susp) $\Delta^{14}\text{C}$ values (27, -46 and 9‰) at these depths (Table 3), assuming the $\Delta^{14}\text{C}$ of the surface-derived POC (susp) to be 94‰ (same as DIC at 5m). This calculation also assumes a $\Delta^{14}\text{C}$ of -450‰ for the DOC-UV at 500m based on the North Central Pacific value at 700m. Again, the SMB 700m POC (susp) $\Delta^{14}\text{C}$ value of -46‰ is suspect (it's $\Delta^{14}\text{C}$ value would be 12‰ assuming a 15% DOC contribution). Although a 12 to 15% contribution of DOC to the POC (susp) intuitively seems too high, it only amounts to 0.025 to $0.054\mu\text{mol}$ ($0.3\text{--}0.65\mu\text{g}$) C l^{-1} of the total 0.21 to $0.36\mu\text{mol l}^{-1}$ of POC (susp), or 0.06 to 0.13% of the total DOC-UV at those depths.

5.3 Origin of the sedimentary organic matter

The contribution of sinking POC reaching the ocean floor at mid-basin is $43\text{mg m}^{-2}\text{d}^{-1}$ (Table 8, mean value, $n=4$) which may be augmented by unknown, but presumably small, quantities of POC (susp) and DOC. JAHNKE (1990) has calculated that 43% of the total organic carbon reaching the seafloor ($18.5\text{mg C m}^{-2}\text{d}^{-1}$) is permanently buried. He concludes that carbon burial rates for the Southern California Bight overall are primarily controlled by input rates and not by variations in *in situ* redox potentials and other preservation variables. This sinking POC includes: (1) terrestrially-derived POC; (2) detritus and associated macro- and micro-organisms derived from the euphotic zone and added during transit to the seafloor; and (3) recycled SOC derived from the basin slopes via episodic injections into the deep basin water.

The contribution of terrestrial organic matter from all sources to the mid-basin sediments, calculated from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures, is about 25% (Table 9), based solely on the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ end-member signatures of sewage organics and the surficial SOM. However, the mass emission of sewage carbon ($\sim 5 \times 10^7\text{kg y}^{-1}$) into the Southern California Bight calculated from sedimentary coprostanol contents (VENKATESAN and KAPLAN, 1990) has been estimated to be $\sim 10^3$ times less than carbon fixed photosynthetically in the Bight ($\sim 1.4 \times 10^{10}\text{kg y}^{-1}$). This sewage carbon amounts to 1.6% of the mid-basin surficial SOC, which coupled with a 4% organic carbon contribution from terrestrial plants (based on lignin-derived phenol measurements), gives a total of $\sim 6\%$ terrestrially-derived organic carbon in the mid-basin sediments, excluding humic substances (VENKATESAN and KAPLAN, 1992). If isotopic fractionation, during *in situ* oxic mineralization of SOM, discriminates against both ^{13}C and ^{15}N , it could result in isotopic end-member values 0.5 to 1‰ heavier than their initial signatures as given in Table 9, so the contribution of terrestrial organic matter would then be even lower than those values calculated in Table 9. For example, if the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures in the 0-3.75cm core section were altered to -21.7 and 8.0‰ from -22.2 and 7.0‰ by mineralization, then the terrestrial contribution would be reduced from 28% to 8% and from 25% to 8%, respectively, based on these heavier $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values. If we assume isotopic fractionation occurs, the estimated upper limit to the amount of terrestrially-derived organic carbon in the surficial SOC of 10% is closer to the sewage and land plant inputs calculated by VENKATESAN and KAPLAN (1992).

Assuming that our previous estimate of recycled SOC in the POC (sink) at 850m (50% of the total) is valid, then 40% of the POC (sink) is unrecycled organic carbon of local origin. How much of each of these three fractions is mineralized and how much is buried is largely conjectural. However, if 57% ($24.5\text{mg C m}^{-2}\text{d}^{-1}$) of the organic matter reaching the ocean floor is mineralized

(JAHNKE, 1990), then there must be nearly complete oxidation of the sinking particles of local origin, plus oxidation of a significant fraction (~15%) of the "already-oxidized", slope-derived SOC and also of a small amount (~2%) of the terrestrial component of the POC (sink). Neither isotopic signatures nor carbon contents of the POC (sink) and SOC (0-2.5cm) can be used to quantify these fractions, largely because we know neither the true isotopic signatures nor the C and N concentrations of the terrestrial and slope inputs. If, for example, the slope-derived surficial SOC had a $\Delta^{14}\text{C}$ value less than that assumed (-15‰) because of contributions of "older" SOC from deeper sediment layers, then the calculated fraction of slope-derived SOC in the POC (sink) would be lower, and the proportion of recycled SOC in the basin sediment (0-2.5cm) reduced. It is clear, however, that the $\Delta^{14}\text{C}$ value for the SOC (0-2.5cm) at mid-basin (-15‰) does not arise from bioturbation in these distinctly varved sediments, and hence there has to be admixture of pre-bomb organic carbon with the sinking POC ($\Delta^{14}\text{C}$ at 850m = 32‰). In the simplest case, 57% of pre-bomb carbon ($\Delta^{14}\text{C}$ = -75‰) plus 43% modern POC (sink) ($\Delta^{14}\text{C}$ = 32‰) would give an SOC $\Delta^{14}\text{C}$ value of -29‰ for the 0-2.5cm accumulated since 1960. This is of the same order as the observed average $\Delta^{14}\text{C}$ value for this depth horizon (-15‰).

This still leaves the question: What is responsible for these "old" SO^{14}C ages in a rapidly accumulating sediment that is not appreciably mixed by bioturbation or by ventilation with near-bottom water from burrowing organisms? Radiocarbon decay at 20cm (^{210}Pb age since AD 1950 = 607 years) is equivalent to a decrease of about 73‰ in $\Delta^{14}\text{C}$, and an additional decrease of 75‰ (or 625 yrs) is accounted for in pre-bomb SOC owing to the reservoir age of the surface ocean at this site (BERGER, TAYLOR and LIBBY, 1966; ROBINSON and THOMPSON, 1981). Thus, we can account for a $\Delta^{14}\text{C}$ value for SOC at 20cm of -150‰; instead, our measured value is -416‰. EMERY and BRAY (1962) suggested that unexpectedly "old" ages for surficial SOC (mean ^{14}C age of 2500 yrs BP, $\Delta^{14}\text{C}$ = -270‰), may result in part from redeposition and mixing of ancient with modern organic carbon. An alternate explanation is to invoke the incorporation of DOC into the POC that sediments onto the sediment/water interface and becomes buried. The large gradient with depth observed in POC $\Delta^{14}\text{C}$ results suggests that "old" carbon in some form is being incorporated into POC in the water column.

5.4 Phase inter-relationships deduced from isotopic signatures

Figure 5 portrays regressions between $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ for the organic phases at the mid-basin and shelf stations. These number-6-shot patterns, in general, show common, but not distinctive, sources for the organic matter in all phases. The narrow spread in $\delta^{13}\text{C}$ values precludes any meaningful and distinctive regressions of $\delta^{13}\text{C}$ vs. $\Delta^{14}\text{C}$ for any of the organic phases, and while there is a larger spread in $\delta^{15}\text{N}$ values, no distinctive patterns in $\delta^{15}\text{N}$ vs $\Delta^{14}\text{C}$ regressions emerge which would uniquely characterize the origin(s) of any particular organic phases. There is a tight coupling of $\delta^{13}\text{C}$ vs. $\delta^{15}\text{N}$ only for the SOC. Thus, these isotopic signatures basically reflect the multiplicity of source organic matter to all phases (at least in the water column), and are not specific enough for interpreting phase inter-relationships.

5.5 C/N ratios in the sinking organic matter

C/N ratios (by atoms) at mid-basin vary from 6.1 to 9.9 (excepting at 200m, CaBS-7), and from 7.0 to 8.3 (excepting at 200 and 350m, CaBS-7) at the shelf (Table 8, Fig.4). The three exceptionally low ratios (2.6, 2.8, 3.1) cannot be attributed to analytical error, and must have resulted from both reduced carbon flux and increased nitrogen flux; all are lower than C/N ratios in phyto- and zooplankters (C/N = 5-6) and in pure protein (C/N = 3.5; GNAIGER and BITTERLICH,

1984). No concurrent analyses were done for nitrogen-rich molecules such as ammonia, urea or methyl amines, which, by sorption on highly proteinaceous particles, could have accounted for these anomalously low C/N values. The only unique isotopic signature which appears in these shallow samples from CaBS-7 is a $\Delta^{14}\text{C}$ value of 2‰ at 350m, in comparison to values of -18 and -24‰ for CaBS-3 and 4. The $\delta^{15}\text{N}$ values were relatively high for these three samples; especially at 350m, where $\delta^{15}\text{N}$ was 8.06‰ compared to the CaBS-3, 4 and 10 values of 3.55, 7.40 and 7.86‰ respectively. Thus the most likely source of these highly nitrogenous particles is from the marine organic pool, unlike the terrestrially-impacted sinking particles at 350m for CaBS-3 and 4.

5.6 Comparison of isotopic signatures and other properties between the Santa Monica Basin and the North Central Pacific gyre

One of the more striking observations to result from this study was the close correspondence of $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$, total UV-oxidizable DOC, total POC (susp), soluble free and combined amino acids, and free plus combined carbohydrates between the mid-basin station in the Santa Monica Basin and at a station at 31°N , 159°W in the oligotrophic North Central Pacific (NCP) gyre. Some of these similarities are shown for $\Delta^{14}\text{C}$ values (Fig.6) and for dissolved free amino acids (DFAA), total dissolved, hydrolyzable amino acids (THAA) and dissolved free plus hydrolyzable carbohydrates (TCHO) (Fig.7).

The amino acid analyses followed the procedure of LINDROTH and MOPPER (1979), and the carbohydrate determinations were from BURNEY and SIEBURTH (1977). These latter values are reported as glucose equivalents. DFAA, THAA and TCHO for the Santa Monica Basin are from WILLIAMS (1986), and from DRUFFEL *et al* (1989) for the NCP. All DOC-UV and POC (susp) results are from the total CO_2 collected during the UV-oxidation of DOC and the closed-tube combustion of POC (susp). $\Delta^{14}\text{C}$ values at the NCP site for DOC-UV, DIC, POC (susp) and POC (sink) are from DRUFFEL and WILLIAMS (1990).

The decrease in DOC-UV and DIC $\Delta^{14}\text{C}$ values in the upper 900m at both sites follows a similar gradient, such that the SMB values are approximately the same as those 200m deeper in the NCP. This is believed to be the result of the shoaling of isopycnals of open-ocean water entering the basin, and secondarily to local upwelling. Total DOC-UV concentrations in the SMB (Table 3) for cruises CaBS-4 and 7 were 42 to $46\mu\text{mol l}^{-1}$ from 500 to 850m; compared to 52, 47 and $42\mu\text{mol l}^{-1}$ at 450, 600 and 900m in the NCP, respectively. Mean total DOC-UV values at the two sites for the upper 100m waters were $62\mu\text{mol l}^{-1}$ (NCP, $n=3$) and $65\mu\text{mol l}^{-1}$ (SMB, $n=4$). There are also some recent, preliminary values for the DOC_{HTC} at mid-basin. These results are from D. HANSELL, UCSC (personal communication), for samples collected on CaBS-11, and analyzed for DOC_{HTC} using a "clone" of Suzuki's 3% $\text{Pt}/\text{Al}_2\text{O}_3$ -catalyzed oxidation at 680°C (SUGIMURA and SUZUKI, 1988). The SMB, DOC_{HTC} values decreased from $130\mu\text{mol l}^{-1}$ at the surface to $100\mu\text{mol l}^{-1}$ at 100m and to $85\mu\text{mol l}^{-1}$ at 850m. The results for the NCP were $210\mu\text{mol l}^{-1}$ at the surface, decreasing to $108\mu\text{mol l}^{-1}$ at 482m and to 94 and $90\mu\text{mol l}^{-1}$ at 757 and 900m, respectively (DRUFFEL, WILLIAMS and SUZUKI, 1989). Thus, the DOC-UV and DOC_{HTC} concentrations in the 500 to 900m waters in both the SMB and NCP are similar, suggesting that these soluble organic fractions have a common origin, or at least a common biochemical history, while the "extra" DOC_{HTC} in the upper waters may have origins unique to various regions, judging from its elevated concentration in central (DRUFFEL, WILLIAMS and SUZUKI, 1989) and western (SUGIMURA and SUZUKI, 1988) Pacific surface waters. Lastly, the $\delta^{13}\text{C}$ signatures for the DOC-UV fall within the ranges of -20.86 to -22.19‰ (SMB, Table 3) and -20.4 to -21.6‰ (NCP, DRUFFEL *et al*, 1989) for the upper 900m.

FIG.6. $\Delta^{14}\text{C}$ values for the Santa Monica Basin (SMB, mid-basin and shelf sites) and the north central Pacific site (NCP, $31^\circ\text{N } 159^\circ\text{W}$). DOC, ϕ = NCP, $\diamond$ = SMB (mid-basin, CaBS-7), $\blacklozenge$ = SMB (shelf, CaBS-7); DIC, $\bullet$ = NCP, $\times$ = SMB (mid-basin, CaBS-7); POC (susp), $\circ$ = NCP, $\bullet$ = SMB (mid-basin, plus one point at 100m at the shelf site, CaBS-4); POC (sink), ∇ = NCP, $\blacktriangledown$ = SMB (mid-basin, CaBS-7), $\blacktriangledown$ = SMB (shelf, CaBS-7).

With respect to the more labile DFAA, THAA and TCHO values (Fig.7), however, the close correspondence of their vertical profiles between the SMB and the NCP is more difficult to explain. The DFAA concentrations are low at both sites, where the increase below the sill (740m) in the SMB is a result of denitrification and possible benthic contributions. It appears that higher productivity in the coastal waters is not reflected in elevated concentrations of either THAA or TCHO in the surface or deep basin waters; and, in fact, TCHO in the upper 100m of the SMB is lower than in the NCP.

The $\Delta^{14}\text{C}$ values for POC (susp) in the SMB and NCP both decrease with depth, but have significantly lower ^{14}C activities in the basin. Aside from contributions of resuspended POC discussed above, some fraction of the POC (susp) in the basin may have had a lower initial $\Delta^{14}\text{C}$ value because of the lower surface DIC- ^{14}C activity (94‰ in the basin at 5m versus 140‰ at 3m in the NCP). This assumes that the source of the POC (susp) was from photosynthetically-fixed

FIG. 7. Profiles of total dissolved free amino acids (DFAA) total dissolved hydrolyzable amino acids (THAA) and total dissolved hydrolyzable carbohydrates (TCHO, as glucose equivalents) in the SMB and NCP: ● = DFAA (NCP), ○ = DFAA (SMB), ■ = THAA (NCP), □ = THAA (SMB), ▲ = TCHO (NCP), Δ = TCHO (SMB).

DIC whose value was indeed 94‰. Total concentrations of POC (susp) in the basin deep-waters (500–850m) (Table 3) were 0.21 to 0.36 $\mu\text{mol l}^{-1}$ compared to values of 0.24 and 0.15 $\mu\text{mol l}^{-1}$ at 450 and 900m in the NCP. The corresponding $\delta^{13}\text{C}$ signatures were -22.22 to -23.08‰ (SMB) and -19.35 and -21.70 at 450 and 900m in the NCP. Unlike the dissolved organic matter, these above comparisons for POC (susp) do not indicate a common origin of POC (susp) for the SMB and NCP regimes.

6. SUMMARY

Isotopic signatures ($\Delta^{14}\text{C}$, $\delta^{13}\text{C}$, $\delta^{15}\text{N}$) of the soluble and particulate organic phases measured in this study did not prove, nor were they anticipated to be, ideal proxies for unequivocally defining the sources and cycling mechanisms of organic matter within the Santa Monica basin system.

However, these isotopic signatures, coupled with organic matter concentrations and fluxes, did provide clues to the sources of certain organic components and, to some extent, set quantitative limits to these sources. For example: (1) terrestrial inputs of organic matter from both natural and anthropogenic sources were estimated to be <5% for dissolved organic matter, <10% for suspended and sinking particulate organic matter, and <25% for sedimentary organic matter; (2) the isotopic signatures and concentrations of UV-oxidizable DOC were strikingly similar in the deep basin and at an open-ocean site in the North Central Pacific, suggesting a common history for a major fraction of the DOC in the subsurface waters at both sites; and (3) episodic resuspension

of "old" (^{14}C -depleted) shelf and/or slope sedimentary organic matter appeared to be reflected in the low ^{14}C activities of the POC (susp) and POC (sink) in the basin subsurface waters.

There were no significant correlations for the $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures between the suspended and sinking particulate organic matter, sedimentary organic matter, and soluble organic matter which would have been useful in delineating phase transformations among these organic pools. The multiplicity of fractionation processes within and between organic phases, plus the wide ranges in end-member isotopic signatures, makes it difficult, if not impossible, to rely upon isotopic measurements alone for future studies of this nature; studies which must couple isotopic measurements with chemical analysis of process-oriented tracers (e.g. organic biomarkers).

7. ACKNOWLEDGEMENTS

We thank the crews of the R/V *New Horizon* and R/V *Robert G. Sproul* for their long-term shipboard and logistical support throughout the CaBS field program. Expertise in box core and particle trap deployments was provided by R. Jahnke and J. Singleton. E. Renger, SIO, measured the CN fluxes, and assisted with particle-trap sample-splitting. We also appreciate the logistical support of D. Long for all cruises.

We are particularly indebted to L. Toolin, A.J.T. Jull, D. Donahue and T. Linick, University of Arizona, for high-quality ^{14}C -AMS analyses, and to D. Winter, UCLA for similar high-quality $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements.

Financial support was primarily provided by Grant DE-FG05-85-ER60339 from the Ecological Science Division of the U.S. Department of Energy, and to a lesser extent by grants OCE84-17102 and OCE88-40531 (P.M. Williams) and OCE84-16632 and OCE87-16590 (E.R.M. Druffel) from the Ocean Sciences division of the U.S. National Science Foundation. This is Woods Hole Oceanographic Institution contribution No. 7848

8. REFERENCES

- ALTABET, M.A. (1988) variations in nitrogen isotopic composition between sinking and suspended particles: implications for nitrogen cycling and particle transformation in the open ocean. *Deep-Sea Research*, **35**, 535-554.
- ALTABET, M.A. and L.F. SMALL (1990) Nitrogen isotopic ratios in fecal pellets produced by marine zooplankton. *Geochimica et Cosmochimica Acta*, **54**, 155-163.
- BACON, M. and R.F. ANDERSON (1982) Distribution of thorium isotopes between dissolved and particulate forms in the deep sea. *Journal of Geophysical Research*, **87**, 2045-2056.
- BERGER, R., R.E. TAYLOR and W.F. LIBBY (1966) Radiocarbon content of marine shells from the California and Mexican west coast. *Science*, **153**, 864-866.
- BIEN, G.S., N.W. RAKESTRAW and H.E. SUESS (1965) Radiocarbon in the pacific and Indian oceans and its relation to deep water movements. *Limnology and Oceanography*, **10** (suppl.), R25-R38.
- BRULAND, K.W., R.P. FRANKS, W.M. LANDING and A. SOUTAR (1981) A Southern California inner basin sediment trap calibration. *Earth Planetary Science Letters*, **53**, 400-408.
- BURNEY, C.M. and J. McN. SIEBURTH (1977) Dissolved carbohydrates in seawater. II. A spectrophotometric procedure for total carbohydrate analysis and polysaccharide estimation. *Marine Chemistry*, **5**, 15-28.
- CHECKLEY, D.M.Jr., and L.C. ENTZEROOTH (1985) Elemental and isotopic fractionation of carbon and nitrogen by marine planktonic copepods and implications to the marine nitrogen cycle. *Journal of Plankton Research*, **7**, 553-568.
- CHECKLEY, D.M.Jr. and C.A. MILLER (1989) Nitrogen isotope fractionation by oceanic zooplankton. *Deep-Sea Research*, **36**, 1449-1456.
- CRISP, P.T., S. BRENNER, M.I. VENKATESAN, E. RUTH and I.R. KAPLAN (1979) Organic chemical characterization of sediment-trap particulates from San Nicolas, Santa Barbara, Santa Monica and San

- Pedro Basins, California. *Geochimica et Cosmochimica Acta*, **43**, 1791-1801.
- DeLAPPE, B.W., R.W. RISEBROUGH and W. WALKER II (1983) A large-volume sampling assembly for the determination of synthetic organic and petroleum compounds in the dissolved and particulate phases of seawater. *Canadian Journal of Fisheries and Aquatic Science*, **40** (suppl.2), 322-336.
- DRUFFEL, E.R.M. and P.M. WILLIAMS (1990) Identification of a deep marine source of particulate organic carbon using bomb ^{14}C . *Nature, London*, **347**, 172-174.
- DRUFFEL, E.R.M., P.M. WILLIAMS and Y. SUZUKI (1989) Concentrations and radiocarbon signatures of dissolved organic matter in the Pacific Ocean. *Geophysical Research Letters*, **16**, 991-994.
- DRUFFEL, E.R.M., P.M. WILLIAMS, K.J. ROBERTSON, S. GRIFFIN, A.J.T. JULL and D. DONAHUE (1989) Radiocarbon in dissolved organic and inorganic carbon from the central North Pacific. *Radiocarbon*, **31**, 523-532.
- EGANHOUSE, R.P. (1982) *Organic matter in municipal wastes and storm runoff: characterization and budget to the coastal waters off southern California*. PhD dissertation, University of California, Los Angeles, Los Angeles, CA, 230pp.
- EGANHOUSE, R.P. and I.R. KAPLAN (1988) Depositional history of recent sediments from San Pedro shelf, California: reconstruction using elemental abundance, isotopic composition and molecular markers. *Marine Chemistry*, **24**, 163-191.
- EMERY, K.O. (1960) *The Sea off Southern California*. John Wiley, New York, 366pp.
- EMERY, K.O. and E.E. BRAY (1962) Radiocarbon dating of California basin sediments. *Bulletin of the American Association of Petroleum Geologists*, **46**, 1839-1856.
- EPPLEY, R.W., E.H. RENGGER and P.R. BETZER (1983) The residence time of particulate organic carbon in the surface layer of the ocean. *Deep-Sea Research*, **30**, 311-323.
- FINNEY, B. and C-A. HUH (1989) High resolution sedimentary records of heavy metals from the Santa Monica and San Pedro Basins, California. *Marine Pollution Bulletin*, **20**, 181-187.
- FRY, B. (1988) Food web structure on Georges Bank from stable C, N and S isotopic compositions. *Limnology and Oceanography*, **33**, 1182-1190.
- FRY, B. and E.B. SHERR (1984) ^{13}C measurements as indicators of carbon flow in marine and freshwater ecosystems. *Contributions to Marine Science*, **27**, 13-47.
- GNAIGER, E. and G. BITTERLICH (1984) Proximate biochemical composition and caloric content calculated from elemental CHN analysis: A stoichiometric concept. *Oecologia*, **62**, 289-298.
- HUBBS, C.L., G.S. BIEN and H.E. SUESS (1965) La Jolla natural radiocarbon measurements IV. *Radiocarbon*, **7**, 66-117.
- HUH, C-A. and T.M. BEASLEY (1987) Profiles of dissolved and particulate thorium isotopes in the water column of coastal Southern California. *Earth and Planetary Science Letters*, **85**, 1-10.
- HUH, C-A., D.L. ZAHNLE and L.F. SMALL (1986) Budgets and behaviors of uranium and thorium series isotopes in Santa Monica Basin sediments. In: *West Coast Environmental Program California Basin Study (CaBS), Progress Report III*, December 1986.
- HUH, C-A., L.F. SMALL, S. NIEMNIL, B.P. FINNEY, B.M. HICKEY, N.B. KACHEL, D.S. GORSLINE and P.M. WILLIAMS (1990) Sedimentation dynamics in the Santa Monica - San Pedro Basin off Los Angeles: radiochemical, sediment trap and transmissometer studies. *Continental Shelf Research*, **10**, 137-164.
- JACKSON, G.A., F. AZAM, A.F. CARLUCCI, R.W. EPPLEY, B. FINNEY, D.S. GORSLINE, B. HICKEY, C-A. HUH, R.A. JAHNKE, I.R. KAPLAN, M.R. LANDRY, L.F. SMALL, M.I. VENKATESAN, P.M. WILLIAMS and K.M. WONG (1989) Elemental cycling and fluxes off Southern California. *EOS, Transactions American Geophysical Union*, **70**, 146-149; 154-155.
- JAHNKE, R.A. (1990) Early diagenesis and recycling of biogenic debris at the seafloor, Santa Monica Basin, California. *Journal of Marine Research*, **48**, 413-436.
- JEFFREY, A.W.A., R.C. PFLAUM, J.M. BROOKS and W. SACKETT (1983) Vertical trends in particulate organic carbon ^{13}C : ^{12}C ratios in the upper water column. *Deep-Sea Research*, **30**, 971-983.
- JULL, A.J.T., D.J. DONAHUE, A. HATHEWAY, T.W. LINICK and J.J. TOOLIN (1986) Production of graphite targets by deposition from CO/H_2 for precision accelerator ^{14}C measurements. *Radiocarbon*, **28**, 191-197.
- LAIRD, J.C., D.P. JONES and C.S. YENTSCH (1967) A submersible filtering unit. *Deep-Sea Research*, **14**, 251-252.
- LANDRY, M.R., W.K. PETERSON and C.C. ANDREWS (1992) Particulate flux in the water column overlying Santa Monica Basin. *Progress in Oceanography*, **30**, 167-195.
- LEVIN, I., B. KROMER, H. SCHOCH-FISCHER, M. BRUNS, M. MUNNICH, D. BERDAN, J.C. VOGEL and

- K.O. MUNNICH (1985) Twenty-five years of tropospheric ^{14}C observations in Central Europe. *Radiocarbon*, **27**, 1-19.
- LINDROTH, P. and K. MOPPER (1979) High performance liquid chromatographic determination of sub picomole amounts of amino acids by precolumn fluorescence derivatization with o-phthalaldehyde. *Analytical Chemistry*, **51**, 1667-1674.
- LINICK, T.W. (1975) *Uptake of Bomb-produced carbon-14 by the Pacific Ocean*. PhD dissertation, University of California, San Diego, Department of Chemistry.
- LIU, K.-K. and I.R. KAPLAN (1982) Nitrous oxide in the sea off Southern California. In: *The Environment of the Deep-Sea*. W.G. ERNST and J.G. MORIN, editors, Prentice Hall, New York, 73-92.
- LIU, K.-K. and I.R. KAPLAN (1989) The eastern tropical Pacific as a source of ^{15}N -enriched nitrate in seawater off southern California. *Limnology and Oceanography*, **34**, 820-830.
- MALOUTA, D.N., D.S. GORSLINE and S.E. THORNTON (1981) Processes and rates of recent (Holocene) basin filling in an active transform margin: Santa Monica Basin, California continental borderland. *Journal of Sedimentary Petrology*, **51**, 1077-1095.
- MANNING, M.R., D.C. LOWE, W.H. MELHUISE, R.J. SPARKS, G. WALLACE, C.A.M. BREENINKMEIJER and R.C. MCGILL (1990) The use of radiocarbon measurements in atmospheric studies. *Radiocarbon*, **32**, 37-58.
- MCCAVE, I.N. (1984) Size spectra and aggregation of suspended particles in the deep ocean. *Deep-Sea Research*, **31**, 329-352.
- MINAGAWA, M., D.A. WINTER and I.R. KAPLAN (1984) Comparison of Kjeldahl and combustion methods for measurement of nitrogen isotope ratios in organic matter. *Analytical Chemistry*, **56**, 1859-1861.
- MULLIN, M.M., G.H. RAU and R.W. EPPLEY (1984) Stable nitrogen isotopes in zooplankton: some geographical and temporal variation in the North Pacific. *Limnology and Oceanography*, **29**, 1267-1273.
- MYERS, E.P. (1974) *The concentration and isotopic composition of carbon in marine sediments affected by a sewage discharge*. PhD dissertation, California Institute of Technology, Pasadena, CA, 179pp.
- NELSON, J.R., J.R. BEERS, R.W. EPPLEY, G.A. JACKSON, J.J. MCCARTHY and A. SOUTAR (1987) A particle flux study in the Santa Monica - San Pedro Basin off Los Angeles: particle flux, primary production, and transmissometer survey. *Continental Shelf Research*, **7**, 307-328.
- NISSENBAUM, A. and I.R. KAPLAN (1972) Chemical and isotopic evidence for the *in situ* origin of marine humic substances. *Limnology and Oceanography*, **17**, 570-582.
- NYDAL, R., K. LOVESETH and F.H. SKOGSETH (1980) Transfer of bomb ^{14}C to the ocean surface. *Radiocarbon*, **22**, 626-635.
- ÖSTLUND, H.G. and M. STUIVER (1980) GEOSECS Pacific radiocarbon. *Radiocarbon*, **22**, 25-53.
- PETERS, K.E., R.E. SWEENEY and I.R. KAPLAN (1978) Correlation of carbon and nitrogen stable isotope ratios in sedimentary organic matter. *Limnology and Oceanography*, **23**, 598-604.
- RAU, G.H., T. TAKAHASHI and D.J. DES MARAIS (1989) Latitudinal variations in plankton $\delta^{13}\text{C}$: Implications for CO_2 and productivity in past oceans. *Nature, London*, **341**, 516-518.
- RAU, G.H., R.E. SWEENEY, I.R. KAPLAN, A.J. MEARNES and D.R. YOUNG (1981) Differences in animal ^{13}C , ^{15}N and D abundance between a polluted and an unpolluted coastal site: likely indicators of sewage uptake by a marine food web. *Estuarine, Coastal and Shelf Science*, **13**, 701-707.
- RISEBROUGH, R. and A.L. BURLINGAME (1978) Water column hydrocarbon chemistry (dissolved and particulate). In: *Southern California Baseline Study and Analysis, 1975-1976*. Vol. II. Integrated Summary Report. La Jolla, California: Science Applications Inc. 375-389.
- ROBINSON, S.W. and G. THOMPSON (1981) ^{14}C corrections for marine shell dates with application to southern Pacific NW coast prehistory. *Syesis*, **14**, 45-57.
- SAINO, T. and A. HATTORI (1987) Geographic distribution of suspended particulate organic nitrogen and its ^{15}N natural abundance in the Pacific and its marginal seas. *Deep-Sea Research*, **34**, 807-827.
- SHARP, J.H. (1974) Improved analysis for "particulate" organic carbon and nitrogen from seawater. *Limnology and Oceanography*, **19**, 984-989.
- SMALL, L.F., M.R. LANDRY, R.W. EPPLEY, F. AZAM and A.F. CARLUCCI (1989) Role of plankton in the carbon and nitrogen budgets of Santa Monica Basin, California. *Marine Ecology Progress Series*, **56**, 57-74.
- SMITH, P.E. and R.W. EPPLEY (1982) Primary production and the anchovy population in the Southern California Bight: comparison of time series. *Limnology and Oceanography*, **27**, 1-17.
- SOUTAR, A. (1979) Collection of benthic sediment samples. In: *Southern California Outer Continental Shelf*

-
- Environmental Baseline Study, 1976/1977, (Second Year)*, Volume II, Technical Reports, Report 2. BLM contract to SAI AA550-CT6-40, NTIS PB80-166093.
- STUIVER, M. and H.A. POLACH (1977) Discussion: Reporting of ^{14}C data. *Radiocarbon*, **19**, 355-363.
- SUGIMURA, Y. and Y. SUZUKI (1988) A high temperature catalytic oxidation method of non-volatile dissolved organic carbon in seawater by direct injection of liquid sample. *Marine Chemistry*, **24**, 105-131.
- SWEENEY, R.E. and K.R. KAPLAN (1980a) Natural abundances of ^{15}N as a source indicator for near-shore sedimentary and dissolved nitrogen. *Marine Chemistry*, **9**, 81-94.
- SWEENEY, R.E. and I.R. KAPLAN (1980b) Tracing flocculent industrial and domestic sewage transport on San Pedro Shelf, Southern California, by nitrogen and sulfur isotope ratios. *Marine Environmental Research*, **3**, 215-224.
- SWEENEY, R.E., E.K. KALIL and I.R. KAPLAN (1980) Characterization of domestic and industrial sewage in Southern California coastal sediments using nitrogen, carbon, sulfur and uranium tracers. *Marine Environmental Research*, **3**, 225-243.
- THOMPSON, S. and G. EGLINTON (1978) The fractionation of a recent sediment for organic geochemical analysis. *Geochimica et Cosmochimica Acta*, **42**, 199-207.
- VENKATESAN, M.I. and I.R. KAPLAN (1990) Sedimentary coprostanol as an index of sewage addition in Santa Monica Basin, Southern California. *Environmental Science and Technology*, **24**, 208-214.
- VENKATESAN, M.I. and I.R. KAPLAN (1992) Vertical and lateral transport and the carbon budget in Santa Monica Basin, California. *Progress in Oceanography*, **30**, 291-312.
- VENKATESAN, M.I., S. BRENNER, E. RUTH, J. BONILLA and I.R. KAPLAN (1980) Hydrocarbons in age-dated sediment cores from two basins in the Southern California Bight. *Geochimica et Cosmochimica Acta*, **44**, 789-802.
- WELSCHMEYER, N.A. and C.J. LORENZEN (1985) Chlorophyll budgets: zooplankton grazing and phytoplankton growth in a temperate fjord and the Central Pacific Gyres. *Limnology and Oceanography*, **30**, 1-21.
- WILLIAMS, P.M. (1986) Chemistry of the dissolved and particulate phases in the water column. In: *Plankton Dynamics of the Southern California Bight*. R.W. EPPLEY, editor, Springer-Verlag, New York, 53-82.
- WILLIAMS, P.M. and E.R.M. DRUFFEL (1987) Radiocarbon in dissolved organic matter in the central North Pacific Ocean. *Nature, London*, **330**, 246-248.
- WILLIAMS, P.M. and E.R.M. DRUFFEL (1988) Dissolved organic matter in the ocean: Comments on a controversy. *Oceanography Magazine*, **1**, 14-17.
- WILLIAMS, P.M. and L.I. GORDON (1970) Carbon-13:carbon-12 ratios in dissolved and particulate organic matter in the sea. *Deep-Sea Research*, **17**, 19-27.
- WILLIAMS, P.M., A.F. CARLUCCI and R. OLSON (1980) A deep profile of some biologically important properties in the central North Pacific gyre. *Oceanologica Acta*, **3**, 471-476.
- WILLIAMS, P.M., J.A. MCGOWAN and M. STUIVER (1970) Bomb carbon-14 in deep-sea organisms. *Nature, London*, **227**, 375-376.