

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California
Contract No. W-7405-eng-48

THE $l = 2$ S-WAVE π - π SCATTERING LENGTH
Janos Kirz, Joseph Schwartz, and Robert D. Tripp

November 7, 1961

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

THE I=2 S-WAVE π - π SCATTERING LENGTH

Janos Kirz, Joseph Schwartz, and Robert D. Tripp

Lawrence Radiation Laboratory
University of California
Berkeley, California

November 7, 1961

ABSTRACT

Through investigation of the reaction $\pi^+ p \rightarrow \pi^+ \pi^+ n$ at an incident pion kinetic energy of 357 Mev, we have found the I = 2 π - π scattering length $|a_2| \approx 0.15 \lambda_\pi$. This work is compared with other experiments bearing on this scattering length.

THE I=2 S-WAVE π - π SCATTERING LENGTH*

Janos Kirz, Joseph Schwartz, and Robert D. Tripp

Lawrence Radiation Laboratory
University of California
Berkeley, California

November 7, 1961

A number of recent experimental and theoretical papers relate directly or indirectly to a_2 , the isotopic spin=2 S-wave π - π scattering length. These are discussed in the latter part of this article. Because these papers are frequently widely divergent in their conclusions, we have undertaken a more definitive measurement of this scattering length by investigating the production of pions in the reaction $\pi^+ + p \rightarrow \pi^+ + \pi^+ + n$ in the 72-inch Lawrence Radiation Laboratory hydrogen bubble chamber.

We see no clear evidence for a low-energy π - π interaction in the isotopic spin 2 state. An upper limit $|a_2| \leq 0.15 \lambda_\pi$ is indicated by this experiment.

The pion beam kinetic energy was 357 ± 5 Mev at the chamber center with a total energy loss in the chamber fiducial region of 40 Mev. The energy chosen was as high as possible, in order to assure a reasonable production cross section while still permitting visual separation (on the basis of track ionization) of this reaction from much more common elastic scattering. The film was scanned twice, giving a scanning efficiency $> 95\%$. The events were measured on digital measuring projectors, and kinematic fits were obtained by using IBM 709 programs PANG and KICK. 213 events of the type $\pi^+ + \pi^+ + n$ were observed among a total of 67,000 interactions. Normalizing to a total $\pi^+ - p$ cross section of 40.5 mb,¹ we obtain $\sigma(\pi^+ + p \rightarrow \pi^+ + \pi^+ + n) = 0.12 \pm 0.01$ mb. Figure 1 shows this point and that of Willis² at 500 Mev, the only other measure of this cross section. The solid curve shows the cross section as calculated by Schnitzer³

using a 1-pion peripheral-collision model and including corrections due to re-scattering of one of the pions (illustrated in Figs. 2a and 2b). The total cross section in this model is proportional to a_2^2 , where a_2 is the S-wave π - π scattering length in the $I=2$ state.⁴ The value used for a_2 was $0.16 \lambda_\pi$. To fit our measurement this model would require $a_2 = 0.08 \lambda_\pi$. The dotted curve shows the dependence of three-body phase space on energy, normalized to our cross section.

Figure 3 shows the number of dipions vs relative pion momentum in the center of mass of the dipion. The curve, normalized to the total number of events, represents phase space. A strong final-state interaction between the pions might be expected to distort this in the manner suggested by Watson.⁵ We see no statistically significant deviation from phase space. The interaction between the π^+ and π^- is quite small in this experiment, since their relative momentum is always below the $3/2, 3/2$ resonance and furthermore the charge state is predominantly $I=1/2$; averaged over all momenta it is about 20 mb and virtually independent of the relative dipion momentum. A plot similar to Fig. 3 but with the π^+ - π^- relative momentum as abscissa is more sensitive to the π - π interaction. This plot has been found to fit phase space very well.

Figure 4 is a plot of the events as a function of p^2 , the square of non-relativistic momentum transfer to the nucleon in units of m_π^2 . The solid curve represents phase space, whereas the dotted curve is the distribution to be expected from a peripheral collision model.^{6,7} The curves are normalized to give the same maximum as the data. The data clearly disagree with phase space and, for $p^2 > 4$, with the peripheral collision model. To study the latter in more detail we have made a Chew-Low extrapolation,⁷ shown in Fig. 5. The data are divided into two ω^2 intervals, $\omega^2 = 4$ to 6 and 6 to 8.5 (where $\omega =$ total dipion energy in its center of mass in units of m_π). The quantity

$(p^2 + 1)^2 \partial^2 \sigma / \partial p^2 \partial \omega^2 = 2 \{ (f^2 / 2\pi) p^2 \omega [(\omega^2 / 4) - 1]^{1/2} / q^2 \} \sigma_{\pi\pi}$ is constructed for various p^2 intervals. f^2 is the π - n coupling constant and q is the incident pion momentum (laboratory system). A good linear fit is made for $p^2 < 4$, yielding extrapolated cross sections at $p^2 = -1$ of $\sigma_{\pi\pi} = 1 \pm 6$ mb and -2 ± 7.5 mb for the lower and higher ω^2 intervals respectively.

Additional uncertainty arises from the assumption of linearity in the extrapolation. This uncertainty is difficult to evaluate, but from Fig. 5 it seems unlikely that $\sigma_{\pi\pi}$ could exceed 10 mb. With $\sigma_{\pi\pi} = 8\pi a_2^2$ this leads to $|a_2| \lesssim 0.15 \lambda_\pi$.

Although one expects the extrapolation curve to show deviations from linearity for large p^2 , it is rather surprising to see the deviations occur as abruptly as indicated in Fig. 5. This again suggests that the $I=2$ π - π interaction is extremely small, and perhaps the dominant mechanism for this reaction is not via peripheral collisions. It should be remarked that calculations by Rodberg based on a static model (containing no π - π interaction),⁸ although yielding too small a cross section for $\pi^- + p \rightarrow \pi^- + \pi^+ + n$, are in agreement with this experiment so far as total cross section is concerned.⁹

Other evidence bearing on a_2 generally involves the difference $a_2 - a_0$, where a_0 is the $I=0$ scattering length. They are:

1. The analysis by Khuri and Treiman¹⁰ of τ -meson decay, considering all deviations from phase space in τ decay to be due to final-state π - π interactions, gives for the most recent data¹¹ $a_2 - a_0 = +0.9 \pm 0.1 \lambda_\pi$.

2. Experiments by Batusov et al.¹² on the reaction $\pi^- + p \rightarrow \pi^- + \pi^+ + n$ near threshold, and using the theory of Anselm and Gribov,¹³ yield $a_2 - a_0 = -0.35 \pm 0.30 \lambda_\pi$.

3. Either of the above two results combined with the interpretation of the experiment by Abashian, Booth, and Crowe¹⁴ on the anomaly in He^3 production

from $p + d$ collisions as a strong final-state $n-\pi$ interaction in the $I = 0$ state ($a_0 \sim 2.5 \lambda_n$) yields a very large a_2 , inconsistent with our result.

4. As remarked previously, calculations of pion production in pion nucleon collisions indicated a_2 to be small and $a_0 \approx 0.5$ to $0.65 \lambda_n$,³ considerably less than that of Reference 14.

Theoretical calculations by Desai¹⁵ based on crossing symmetry relate the $n-\pi$ scattering lengths in the three isotopic spin states. His calculations lead to our result for a negative $n-\pi$ coupling constant λ whose magnitude is less than 0.15. This value of the coupling constant would yield an $a_0 \leq 1.9 \lambda_n$.

We thank Professor Luis W. Alvarez for his encouragement and support, Professor Frank S. Crawford, Jr., for use of his pion beam, and Professor Geoffrey F. Chew and Professor Charles J. Goebel for theoretical advice.

Footnotes and References

- * This work was done under the auspices of the U. S. Atomic Energy Commission.
1. N. P. Klepikov, V. A. Meshcheryakov, and S. N. Sokolov, Joint Institute for Nuclear Research Report D-584, 1960.
 2. W. J. Willis, Phys. Rev. 116, 753 (1959).
 3. Howard J. Schnitzer, Pion-Pion Scattering and Low-Energy Pion Production, University of Rochester preprint; Charles J. Goebel and Howard J. Schnitzer, Phys. Rev. 123, 1021 (1961).
 4. Figure 2b includes a contribution which is proportional to a_1 . This should, however, be small in our energy region.
 5. K. M. Watson, Phys. Rev. 88, 1163 (1952).
 6. C. Goebel, Phys. Rev. Letters 1, 337 (1958).
 7. G. F. Chew and F. E. Low, Phys. Rev. 113, 1640 (1959).
 8. L. S. Rodberg, Phys. Rev. 106, 1090 (1957).
 9. Other calculations using the static model but with different approximations result in a larger cross section than we observe. For a comparison of the various approximations and other references see E. Kazes, Phys. Rev. 107, 1131 (1957).
 10. N. N. Khuri and S. B. Treiman, Phys. Rev. 119, 1115 (1960).
 11. M. Ferro-Luzzi, D. H. Miller, J. J. Murray, A. H. Rosenfeld, and R. D. Tripp, Nuovo cimento (to be published).
 12. Yu. A. Batusov, S. A. Bunyatov, V. M. Sidorov, and V. A. Yarba, in Proceedings of the 1960 Annual International Conference on High Energy Physics at Rochester (Interscience Publishers, Inc., New York, 1960), p 79.
 13. A. A. Anselm and V. N. Gribov, Soviet Physics--JETP 37, 501 (1959).
 14. N. E. Booth, A. Abashian, and K. M. Crowe, Phys. Rev. Letters 7, 35 (1961).
 15. E. R. Desai, Phys. Rev. Letters 6, 497 (1961).

Figure Legends

Fig. 1. Cross sections for $\pi^+ + p \rightarrow \pi^+ + \pi^+ + n$. The point at 500 Mev is that of Willis (2). Our point is shown at 357 Mev. The solid curve is the expected energy dependence calculated by Schnitzer (Reference 3), the dashed curve is three-body phase space scaled to our point.

Fig. 2. Diagrams for the peripheral collision model: (a) one-pion exchange and (b) rescattering of one of the pions.

Fig. 3. Histogram of the number of dipions vs relative pion momentum in the dipion center-of-mass system. The curve represents phase space expected on the statistical model.


Fig. 4. Histogram of the number of events as a function of the square of the nonrelativistic momentum transfer to the nucleon (units of pion mass squared). The solid curve represents phase space, the dotted curve is the dependence expected from the peripheral collision model. Both curves are normalized to give the same maximum as the data.

Fig. 5. Chew-Low extrapolation for

(a) $4 < \omega^2 < 6$, (b) $6 < \omega^2 < 8.5$.


The linear fits are made for the points with $p^2 < 4$. The extrapolated values at $p^2 = -1$ are

(a) 1 ± 6 mb, (b) -2 ± 7.5 mb.



MU-25071

Fig. 1. Cross sections for $\pi^+ + p \rightarrow \pi^+ + \pi^+ + n$. The point at 500 Mev is that of Willis (2). Our point is shown at 357 Mev. The solid curve is the expected energy dependence calculated by Schnitzer (Reference 3), the dashed curve is three-body phase space scaled to our point.



MU-25072

Fig. 2. Diagrams for the peripheral collision model: (a) one-pion exchange and (b) rescattering of one of the pions.






Fig. 3. Histogram of the number of dipions vs relative pion momentum in the dipion center-of-mass system. The curve represents phase space expected on the statistical model.



MU-25074

Fig. 4. Histogram of the number of events as a function of the square of the nonrelativistic momentum transfer to the nucleon (units of pion mass squared). The solid curve represents phase space, the dotted curve is the dependence expected from the peripheral collision model. Both curves are normalized to give the same maximum as the data.



MU-25075

Fig. 5. Chew-Low extrapolation for
(a) $4 < \omega^2 < 6$, (b) $6 < \omega^2 < 8.5$.
The linear fits are made for the points with $p^2 < 4$. The
extrapolated values at $p^2 = -1$ are
(a) 1 ± 6 mb, (b) -2 ± 7.5 mb.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.