

UCLA

UCLA Previously Published Works

Title

Puberty enables oestradiol-induced progesterone synthesis in female mouse hypothalamic astrocytes

Permalink

<https://escholarship.org/uc/item/23q0h3r2>

Journal

Journal of Neuroendocrinology, 34(6)

ISSN

0953-8194

Authors

Mohr, Margaret A

Keshishian, Tina

Falcy, Brennan A

et al.

Publication Date

2022-06-01

DOI

10.1111/jne.13082

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NoDerivatives License, available at <https://creativecommons.org/licenses/by-nd/4.0/>

Peer reviewed

Progesterone receptors in AVPV kisspeptin neurons are sufficient for positive feedback induction of the LH surge

Margaret A. Mohr¹, Lourdes A. Esparza², Paige Steffen², Paul E Micevych^{1*}, and Alexander S. Kauffman^{2*}

¹Dept. of Neurobiology, David Geffen School of Medicine at UCLA, Los Angeles, Los Angeles, CA

²Dept. of OBGYN and Reproductive Sciences, University of California, San Diego, La Jolla, CA

*co-senior authors

Disclosure Statement: The authors have nothing to disclose.

Correspondence:

Dr. Alexander S. Kauffman

Department of OBGYN and Reproductive Sciences

University of California, San Diego

9500 Gilman Drive, #0674

La Jolla, CA 92093

Email: akauffman@ucsd.edu

and

Dr. Paul E Micevych

Dept. of Neurobiology

University of California, Los Angeles

10833 Le Conte Ave.

Los Angeles, CA 90095-1763

Email: pmicevych@mednet.ucla.edu

Abstract

Kisspeptin, encoded by *Kiss1*, stimulates GnRH neurons to govern reproduction. In female rodents, estrogen-sensitive kisspeptin neurons in the rostral anteroventral periventricular (AVPV) hypothalamus are thought to mediate estradiol (E_2)-induced positive feedback induction of the preovulatory luteinizing hormone (LH) surge. AVPV kisspeptin neurons co-express estrogen and progesterone receptors (PGR) and are activated during the LH surge. While E_2 effects on kisspeptin neurons have been well-studied, progesterone's regulation of kisspeptin neurons is less understood. Using transgenic mice lacking PGR exclusively in kisspeptin cells (termed KissPRKOs), we previously demonstrated that progesterone action specifically in kisspeptin cells is essential for ovulation and normal fertility. Unlike control females, KissPRKO females did not generate proper LH surges, indicating that PGR signaling in kisspeptin cells is required for proper positive feedback. However, since PGR was knocked out from all kisspeptin neurons in the brain, that study was unable to determine the specific kisspeptin population mediating PGR action on the LH surge. Here, we used targeted Cre-mediated AAV technology to re-introduce PGR selectively into AVPV kisspeptin neurons of adult KissPRKO females, and tested whether this rescues occurrence of the LH surge. We found that targeted upregulation of PGR in kisspeptin neurons exclusively in the AVPV is sufficient to restore proper E_2 -induced LH surges in KissPRKO females, suggesting that this specific kisspeptin population is a key target of the necessary progesterone action for the surge. These findings further highlight the critical importance of progesterone signaling, along with E_2 signaling, in the positive feedback induction of LH surges and ovulation.

Introduction

In female mammals, ovulation is triggered by an estradiol (E_2)-induced preovulatory luteinizing hormone (LH) surge, which is driven by a surge in gonadotropin-releasing hormone (GnRH) secretion from the brain (i.e., E_2 positive feedback) (1-3). Kisspeptin, encoded by *Kiss1*, potently stimulates GnRH secretion and is essential for the GnRH/LH surge, as female mice lacking *Kiss1* or its receptor, *Kiss1r*, fail to exhibit LH surges or concurrent GnRH neuron activation (3-5). The hypothalamus contains two populations of kisspeptin-synthesizing neurons, one in the arcuate nucleus (ARC) and another rostrally in the continuum spanning the anteroventral periventricular nucleus and neighboring periventricular nucleus (collectively termed here the AVPV; also called the RP3V) (6-8). E_2 's positive feedback effects on the LH surge are thought to be mediated specifically by AVPV *Kiss1* neurons, because 1) E_2 elevates *Kiss1* levels in the AVPV (8-11), 2) AVPV *Kiss1* neurons express $ER\alpha$ (11), 3) AVPV *Kiss1* mRNA and neuronal activation increase in a circadian manner coincident with the LH surge (10, 12-14), and 4) *Kiss1* and kisspeptin levels in the AVPV are sexually dimorphic (greater in females), correlating with the LH surge occurring only in female rodents (8, 15).

While most positive feedback studies focus on E_2 , P4 and its receptor (PGR) are also critical contributors to the LH surge. Like $ER\alpha$ knockout (KO) mice, global PR KO female mice are infertile and unable to produce LH surges (16, 17). Converging pieces of evidence indicate that E_2 -induced local synthesis of P4 in the hypothalamus is critical for the rodent LH surge (18-20). Specifically, E_2 appears to act in $ER\alpha$ -expressing astrocytes in the hypothalamic AVPV to induce the synthesis and secretion of P4 ("neuroP"), which then acts paracrinely to promote the LH surge (18-23). However, the specific neural population(s) that neuroP4 acts in for this process remains unknown. In immortalized cells representing adult female AVPV kisspeptin neurons (mHypoA51), neuroP augmented E_2 action on the expression and release of kisspeptin (24, 25). Because AVPV kisspeptin neurons *in vivo* express PGR (4, 26-28), we previously tested whether PGR signaling in kisspeptin cells is critical for the LH surge. Using transgenic mice with selective KO of PGR specifically in kisspeptin cells ("KissPRKO"), we demonstrated that P4 signaling directly in kisspeptin cells is necessary for the E_2 -induction of the LH surge and ovulation ((27); also see (29)). Moreover, AVPV *Kiss1* cells of E_2 -treated KissPRKOs had normal *Kiss1* mRNA levels but reduced neuronal activation (27), indicating that P4 signaling in *Kiss1* cells is critical for activation of AVPV kisspeptin neurons during the LH surge, perhaps due to P4-induced calcium release, as observed in immortalized kisspeptin cells (25).

While our previous studies implicated that PGR signaling in kisspeptin cells as essential for the LH surge, a limitation was that kisspeptin cells are present in several brain regions, including the AVPV, ARC, and extra-hypothalamic areas like the medial amygdala (MeA) and bed nucleus of the stria terminalis (BNST) (30-33). KissPRKOs lack PGR in all these kisspeptin populations, precluding a definitive conclusion of the specific population that is the direct target for P4 in the LH surge process. Therefore, we tested here whether selective reintroduction of PGR in just one kisspeptin population, the AVPV, of KissPRKOs could rescue the occurrence of the LH surge even with PGR still absent in all other kisspeptin cells.

Materials and Methods

Animals

The same mouse line was used as in our prior study (27), including both KissCre⁺ PR^{fl/fl} females (termed “KissPRKO”; PGR knocked out in *Kiss1* cells) and KissCre⁻ PR^{fl/fl} controls (termed “WT”; PR still present in all cells, including *Kiss1* cells) (**Figure 1**). Our prior study confirmed proper Cre mediated excision of *Pgr* DNA in the AVPV and ARC of KissPRKOs but not in Cre- controls or in tissues known to lack *Kiss1* expression (27). Genotypes and occasional occurrence of germline recombination were determined via PCR analysis of tail DNA; germline-recombined (global PRKO) mice were excluded. Female mice were weaned at 3 weeks and housed 2-3/cage under a 12-12 light-dark cycle (lights off at 1800h). All experiments were approved by the UCSD IACUC.

AAV treatment to induce PR in AVPV kisspeptin cells and mCherry IHC validation

We previously demonstrated that KissPRKO females have impaired ovulatory function and LH surges. Here, we determined if we could rescue the LH surge in KissPRKOs by selectively re-introducing PGR into just AVPV kisspeptin neurons, while all other kisspeptin cells still lack PGR. AAVs were used to deliver a Cre-dependent *Pgr* construct or a control construct to AVPV kisspeptin neurons (which express Cre in Kiss1Cre⁺ mice) (**Figure 1**). 2-3 month old KissPRKO females (Cre⁺) and WT controls (Cre⁻) were

anesthetized with isoflurane and site-specific AVPV-targeted (from bregma: A/P: +0.5 mm; D/V: -5.5 mm, M/L +/-0.3 mm) bilateral injections were made. 500 nL of either AAV1-CMV-DIO-mPGR-2A-mCherry or AAV2/1-CAG-DIO-mCherry (Vector Biolabs) were infused at a flow rate of 100 nL/min; the former AAV, termed “Pgr”, allows for *Pgr* and mCherry expression in Cre-expressing cells while the latter AAV, termed “mCh”, was a control AAV with the mCherry reporter but no *Pgr* construct. After AAV infusion, the needle was left in place for 10 min to allow diffusion and prevent backflow. To validate the AAV technique, several non-experimental mice were perfused and examined for proper targeting via immunohistochemistry (IHC) analysis of mCherry induction in the AVPV region of Cre+ but not WT (Cre-) mice. Because fixed tissue was needed for IHC, these validation mice were separate from the experimental mice evaluated for LH surges and *Kiss1+Pgr* mRNA co-expression (which required fresh frozen tissue for *in situ* hybridization). IHC was performed on fixed AVPV-containing brain slices using an mCherry primary antibody [1:5000; Abcam ab167453 (34)] and AlexaFluor 594 donkey anti-rabbit secondary [1:2000; Jackson 711-585-152 (35)], following previous methods (36). For assessment of LH surges and *Pgr* mRNA induction after AAV treatment, there were 4 experimental groups: KissPRKO^{Pgr}, KissPRKO^{mCh}, WT^{Pgr}, and WT^{mCh}. Mice were given 4 weeks to recover prior to LH surge testing and brain collection. Double-label ISH evaluation of *Pgr+Kiss1* co-expression was used to confirm successful anatomical targeting of AAV to AVPV *Kiss1* neurons; 1 animal with missed targeting was excluded from the study.

Estradiol-induced LH surge

Four weeks after AAV infusion, mice were anesthetized with isoflurane, ovariectomized (OVX), and given subcutaneous Silastic implants (inner diameter: 0.078 in, outer diameter: 0.125 in) containing 0.75 µg of 17-β E₂ dissolved in sesame oil. This E₂ implant is commonly used to induce LH surges and produces elevated serum E₂ levels ~16-24 pg/ml, resembling mouse proestrus levels (5, 37). This positive feedback paradigm produces an LH surge two days later before lights off (5, 13, 14, 37-39). OVX+E₂ mice from all 4 treatment groups (KissPRKO^{Pgr}, KissPRKO^{mCh}, WT^{Pgr}, and WT^{mCh}) were killed 2 days after E₂ implantation, either in the AM (0900-1000h, control time when the LH surge does not occur; n=4-6/group) or in the PM just before lights off (17:30-18:00h, time of the expected LH surge; n=7-10/group) (**Figure 1**). Blood was collected at sacrifice and the serum isolated and stored at -20°C. Serum LH was measured by the University of Virginia Ligand Assay Core using a sensitive LH sandwich radioimmunoassay (limit of detectability 0.04 ng/ml) (40, 41). LH surges were defined as >0.70 ng/ml.

Brains were collected fresh frozen onto dry ice and stored at -80°C . Brains were sectioned on a cryostat into alternating sets of 20- μm sections onto Superfrost-plus slides and stored at -80°C .

Double-label In Situ Hybridization (ISH) for Pgr expression in Kiss1 cells

Double-label ISH was used to measure *Kiss1* and *Pgr* co-expression levels in the AVPV. One set of AVPV brain sections was assayed using well-established radio-labeled (^{33}P) *Pgr* (0.04 pmol/ml) and digoxigenin-labeled (DIG; 1:500) *Kiss1* antisense riboprobes, following our standard double-label ISH protocol (14, 27, 42-44). ISH slides were analyzed “blindly” with respect to treatment using microscopy linked to an automated imaging processing system (Dr. Don Clifton, Univ. Washington) that counts the number of DIG-containing (*Kiss1* mRNA) cells under fluorescence microscopy and then quantifies the number of silver grains (*Pgr* mRNA) overlying each DIG cell under dark-field microscopy. A minimum of 3 AVPV sections (mean: 5.5 sections/animal) and 145 *Kiss1* cells (mean: 290 cells/animal) was counted for each animal. Signal-to-background ratios of *Pgr* expression levels for individual DIG (*Kiss1*) cells were auto-calculated by the program, and a *Kiss1* cell considered double-labeled with *Pgr* if its ratio was >3 (13, 14, 43-46).

Statistical Analyses

All data are expressed as mean \pm SEM. Differences were analyzed by ANOVA, followed by post-hoc comparisons via Fisher's Protected LSD test. Statistical significance was set at $p < 0.05$.

Results

mCherry localization in the AVPV of Cre+ mice and Pgr upregulation in Kiss PRKO females

Pgr or mCh AAV was infused into the AVPV of adult Cre+ (*KissPRKO*) females and Cre- (WT) controls (**Figure 1**). We first confirmed proper localization and targeting of the AAV injection, using IHC to identify proper mCherry expression in the AVPV of Cre+ mice but not Cre- controls (**Figure 2A, B**). We also confirmed no spread of mCherry expression in the ARC (**Figure 2C**).

Next, to verify that our *Pgr* AAV treatment upregulated *Pgr* levels in *Kiss1* cells of Cre+ mice, *Pgr* + *Kiss1* co-expression levels in the AVPV were determined with double ISH. *Pgr* was co-expressed with *Kiss1* at similar levels between WT^{Pgr} and WT^{mCh} (**Figure 3**), as expected for Cre- controls. As we previously reported, KissPRKOs normally have much less *Pgr* co-expression than WTs, evidenced by significantly lower % *Pgr*+*Kiss1* co-expression in KissPRKO^{mCh} versus either WT^{Pgr} and WT^{mCh} controls ($p < 0.05$ for each; **Figure 3**). Confirming the AAV-mediated upregulation of *Pgr*, KissPRKO^{Pgr} had significantly increased % *Pgr*+*Kiss1* co-expression versus KissPRKO^{mCh} ($p < 0.05$; **Figure 3**), with similar % co-expression as the WT control groups (**Figure 3**). Mean levels of *Pgr* mRNA in *Kiss1* neurons were also similar between KissPRKO^{Pgr} and WT^{Pgr} controls (13.5±1.5 vs 12.0±1.8 grains/cell, $p=0.53$). Thus, the experimental paradigm effectively re-introduced *Pgr* expression in the AVPV of KissPRKOs that normally lack PR in kisspeptin neurons.

Positive feedback induction of the LH surge is rescued in KissPRKO^{Pgr} females

We hypothesized that the previously-observed loss of LH surges in KissPRKOs (27) reflects an impairment in the AVPV kisspeptin system caused by absent PGR signaling in this specific neural population. We tested whether selective re-introduction of PGR into AVPV kisspeptin neurons, while leaving PGR absent in all other kisspeptin cells, could rescue the LH surge in OVX+E₂ females. All 4 groups exhibited low LH in the AM, as expected for this non-surge time, with no group differences (**Figure 4C**). In the PM, the overall occurrence of an LH surge was greatly increased in KissPRKO^{Pgr} compared with KissPRKO^{mCh} females and was not different between KissPRKO^{Pgr} and WT^{Pgr} and WT^{mCh} control groups (**Figure 4A**). Indeed, virtually all PM KissPRKO^{Pgr} females demonstrated a surge, whereas no PM KissPRKO^{mCh} females surged (**Figure 4B**).

In the PM, mean LH was significantly elevated in both WT control groups versus AM values, whereas mean LH of PM KissPRKO^{mCh} females was lower ($p < 0.05$ versus each PM WT group; **Figure 4C**) and did not differ significantly from AM groups, similar to our previous report (27). By contrast, PM KissPRKO^{Pgr} females exhibited elevated mean LH versus all AM groups and versus PM KissPRKO^{mCh} females ($p < 0.05$ for each; **Figure 4C**). The mean LH of PM KissPRKO^{Pgr} females was not significantly different than PM WT^{mCh} controls but was lower than WT^{Pgr} (**Figure 4C**), due primarily to one very high WT^{Pgr} value (**Figure 4B**; not significantly different if that value excluded). Collectively, these results demonstrate that re-introduction of *Pgr* selectively in AVPV kisspeptin cells rescues the circadian-timed LH surge.

Discussion

Ovulation is dependent on sex steroid-induced positive feedback induction of GnRH/LH surges. Ample evidence implicates the AVPV kisspeptin system in mediating the positive feedback effects of E_2 on the LH surge (reviewed in (3, 9). Along with E_2 , P4 is also critical for the LH surge, and we and another group recently reported that P4 action specifically in kisspeptin cells is necessary for females to generate proper LH surges (27, 29). However, neither *in vivo* study determined which exact kisspeptin population mediates this necessary action of PGR. AVPV kisspeptin neurons co-express PGR (27, 28), suggesting they are key targets for P4 action. Supporting this, mHypoA51 cells (representing AVPV kisspeptin cells) respond to P4 by increasing *Kiss1* expression and *in vitro* kisspeptin release (24, 25). To formally test whether AVPV kisspeptin cells are critical for positive feedback induction of the LH surge, we re-introduced *Pgr* expression selectively into AVPV kisspeptin neurons of KissPRKO females using targeted AAV technology. We demonstrate that such upregulation of PGR in kisspeptin neurons specifically within the AVPV is sufficient to rescue the E_2 -induced LH surge in KissPRKO females, suggesting these neurons are targets of the endogenous P4 that promotes the surge.

Female KissPRKO mice were previously shown to display impacted fertility, including fewer litters and a notable reduction in litter size (27). The reduced fecundity was traced to diminished ovulatory events in KissPRKO females, reflected by fewer corpora lutea and diminished or absent LH surges in response to elevated E_2 (positive feedback) (27, 29). These sequelae mirrored rats in which hypothalamic P4 synthesis was blocked (19). Moreover, we previously demonstrated that P4 stimulated the release of intracellular calcium in mHypoA51 cells, mirroring our prior *in vivo* results that KissPRKO AVPV kisspeptin neurons were not properly activated by E_2 , correlating with impaired LH surges. Here, we both verified and extended that finding to show that the deficit is likely in kisspeptin neurons specifically in the AVPV, as selective re-introduction of PGR in just AVPV kisspeptin neurons was sufficient to rescue both the occurrence and magnitude of LH surges. These findings support prior evidence linking the AVPV kisspeptin population to the circadian-timed E_2 -induced LH surge and also supports recent findings that P4 can act *in vitro* in AVPV mHypoA51 kisspeptin cells to activate MAPK and Src kinases and induce kisspeptin release (24). Importantly, our present data also indicate that PGR is not required in other kisspeptin populations, such as the ARC, MeA, or BNST, for proper LH surges, though we cannot rule out if PGR action in those populations is sufficient for surge induction.

Two KissPRKO^{Pgr} females did not show an LH surge, but this was not due to insufficient *Pgr* induction, as *Pgr+Kiss1* co-expression was not different between those 2 non-surgings mice and the other 8 surging KissPRKO^{Pgr} mice or surging WT controls (data not shown). Rather, in typical mouse LH surge studies, a small percentage (10-20%) of normal animals sometimes do not surge (for reasons unknown) and the 2 non-surgings KissPRKO^{Pgr} animals likely reflect that normal outcome, similar to the 2 non-surgings WT^{Pgr} (Figure 4B).

Our findings demonstrate that signaling of endogenous P4 directly in AVPV kisspeptin cells is essential for the positive feedback induction of LH surges. These findings further highlight the critical importance of P4 signaling, along with E₂ signaling, in positive feedback induction of LH surge. We have proposed that the source of P4 in this LH surge process is neural rather than ovarian, derived in astrocytes of the AVPV region, under the influence of elevated E₂ (18-22). Indeed, because all our mice were OVX, any requisite P4 for the surge was non-ovarian in origin, further underlining the role of neuroP in E₂ triggered LH surges and ovulation.

Accepted Manuscript

Acknowledgements: The authors thank Ruby Parra, Angela Wong, Tina Keshishian, Melinda Mittelman-Smith, and Shannon Stephens for technical and experimental support.

Grant Support: This research was supported by NIH grants R01 HD090161, R01 HD100580, and R01 HD042635. MAM was supported by NIH F32 HD097965. Additional support provided by NICHD grants P50 HD012303 (U.C. San Diego) and R24 HD1020614 (University of Virginia Ligand Core).

Data Availability Statement: Some or all data generated during and/or analyzed during the current study are not publicly available but are available from the corresponding author on reasonable request.

Accepted Manuscript

References

1. **Freeman ME** 2006 Neuroendocrine control of the ovarian cycle in the rat. In: Neill JD ed. *Physiology of Reproduction*, 3rd Edition; 2283-2326
2. **Wintermantel TM, Campbell RE, Porteous R, Bock D, Grone HJ, Todman MG, Korach KS, Greiner E, Perez CA, Schutz G, Herbison AE** 2006 Definition of estrogen receptor pathway critical for estrogen positive feedback to gonadotropin-releasing hormone neurons and fertility. *Neuron* 52:271-280
3. **Khan AR, Kauffman AS** 2012 The role of kisspeptin and RFamide-related peptide-3 neurones in the circadian-timed preovulatory luteinising hormone surge. *J Neuroendocrinol* 24:131-143
4. **Clarkson J, d'Anglemont de Tassigny X, Moreno AS, Colledge WH, Herbison AE** 2008 Kisspeptin-GPR54 signaling is essential for preovulatory gonadotropin-releasing hormone neuron activation and the luteinizing hormone surge. *J Neurosci* 28:8691-8697
5. **Dror T, Franks J, Kauffman AS** 2013 Analysis of multiple positive feedback paradigms demonstrates a complete absence of LH surges and GnRH activation in mice lacking kisspeptin signaling. *Biol Reprod* 88:146
6. **Clarkson J, Herbison AE** 2006 Postnatal development of kisspeptin neurons in mouse hypothalamus; sexual dimorphism and projections to gonadotropin-releasing hormone neurons. *Endocrinology* 147:5817-5825
7. **Gottsch ML, Cunningham MJ, Smith JT, Popa SM, Acohido BV, Crowley WF, Seminara S, Clifton DK, Steiner RA** 2004 A role for kisspeptins in the regulation of gonadotropin secretion in the mouse. *Endocrinology* 145:4073-4077
8. **Kauffman AS, Gottsch ML, Roa J, Byquist AC, Crown A, Clifton DK, Hoffman GE, Steiner RA, Tena-Sempere M** 2007 Sexual differentiation of Kiss1 gene expression in the brain of the rat. *Endocrinology* 148:1774-1783
9. **Herbison AE** 2008 Estrogen positive feedback to gonadotropin-releasing hormone (GnRH) neurons in the rodent: the case for the rostral periventricular area of the third ventricle (RP3V). *Brain research reviews* 57:277-287
10. **Adachi S, Yamada S, Takatsu Y, Matsui H, Kinoshita M, Takase K, Sugiura H, Ohtaki T, Matsumoto H, Uenoyama Y, Tsukamura H, Inoue K, Maeda K** 2007 Involvement of anteroventral periventricular metastin/kisspeptin neurons in estrogen positive feedback action on luteinizing hormone release in female rats. *J Reprod Dev* 53:367-378
11. **Smith JT, Cunningham MJ, Rissman EF, Clifton DK, Steiner RA** 2005 Regulation of Kiss1 gene expression in the brain of the female mouse. *Endocrinology* 146:3686-3692

12. **Smith JT, Popa SM, Clifton DK, Hoffman GE, Steiner RA** 2006 Kiss1 neurons in the forebrain as central processors for generating the preovulatory luteinizing hormone surge. *J Neurosci* 26:6687-6694
13. **Robertson JL, Clifton DK, de la Iglesia HO, Steiner RA, Kauffman AS** 2009 Circadian Regulation of Kiss1 Neurons: Implications for Timing the Preovulatory GnRH/LH Surge. *Endocrinology* 150:3664–3671
14. **Poling MC, Luo EY, Kauffman AS** 2017 Sex Differences in Steroid Receptor Coexpression and Circadian-Timed Activation of Kisspeptin and RFRP-3 Neurons May Contribute to the Sexually Dimorphic Basis of the LH Surge. *Endocrinology* 158:3565-3578
15. **Homma T, Sakakibara M, Yamada S, Kinoshita M, Iwata K, Tomikawa J, Kanazawa T, Matsui H, Takatsu Y, Ohtaki T, Matsumoto H, Uenoyama Y, Maeda KI, Tsukamura H** 2009 Significance of Neonatal Testicular Sex Steroids to Defeminize Anteroventral Periventricular Kisspeptin Neurons and the GnRH/LH Surge System in Male Rats. *Biol Reprod* 81:1216-1225
16. **Chappell PE, Lydon JP, Conneely OM, O'Malley BW, Levine JE** 1997 Endocrine defects in mice carrying a null mutation for the progesterone receptor gene. *Endocrinology* 138:4147-4152
17. **Chappell PE, Schneider JS, Kim P, Xu M, Lydon JP, O'Malley BW, Levine JE** 1999 Absence of gonadotropin surges and gonadotropin-releasing hormone self-priming in ovariectomized (OVX), estrogen (E2)-treated, progesterone receptor knockout (PRKO) mice. *Endocrinology* 140:3653-3658
18. **Micevych P, Sinchak K** 2011 The Neurosteroid Progesterone Underlies Estrogen Positive Feedback of the LH Surge. *Front Endocrinol (Lausanne)* 2:90
19. **Micevych P, Sinchak K** 2008 Estradiol regulation of progesterone synthesis in the brain. *Mol Cell Endocrinol* 290:44-50
20. **Micevych P, Sinchak K, Mills RH, Tao L, LaPolt P, Lu JK** 2003 The luteinizing hormone surge is preceded by an estrogen-induced increase of hypothalamic progesterone in ovariectomized and adrenalectomized rats. *Neuroendocrinology* 78:29-35
21. **Sinchak K, Mohr MA, Micevych PE** 2020 Hypothalamic Astrocyte Development and Physiology for Neuroprogesterone Induction of the Luteinizing Hormone Surge. *Front Endocrinol (Lausanne)* 11:420
22. **Mohr MA, Wong AM, Tamm RJ, Soma KK, Micevych PE** 2019 Pubertal development of estradiol-induced hypothalamic progesterone synthesis. *Horm Behav* 111:110-113
23. **Mittelman-Smith MA, Rudolph LM, Mohr MA, Micevych PE** 2017 Rodent Models of Non-classical Progesterone Action Regulating Ovulation. *Front Endocrinol (Lausanne)* 8:165
24. **Mittelman-Smith MA, Wong AM, Micevych PE** 2018 Estrogen and Progesterone Integration in an in vitro Model of RP3V Kisspeptin Neurons. *Neuroendocrinology* 106:101-115

25. **Mittelman-Smith MA, Wong AM, Kathiresan AS, Micevych PE** 2015 Classical and membrane-initiated estrogen signaling in an in vitro model of anterior hypothalamic kisspeptin neurons. *Endocrinology* 156:2162-2173
26. **Zhang J, Yang L, Lin N, Pan X, Zhu Y, Chen X** 2014 Aging-related changes in RP3V kisspeptin neurons predate the reduced activation of GnRH neurons during the early reproductive decline in female mice. *Neurobiology of aging* 35:655-668
27. **Stephens SB, Tolson KP, Rouse ML, Jr., Poling MC, Hashimoto-Partyka MK, Mellon PL, Kauffman AS** 2015 Absent Progesterone Signaling in Kisspeptin Neurons Disrupts the LH Surge and Impairs Fertility in Female Mice. *Endocrinology* 156:3091-3097
28. **Stephens SBZ, Kauffman AS** 2021 Estrogen regulation of the molecular phenotype and active transcriptome of AVPV kisspeptin neurons. *Endocrinology*
29. **Gal A, Lin PC, Cacioppo JA, Hannon PR, Mahoney MM, Wolfe A, Fernandez-Valdivia R, Lydon JP, Elias CF, Ko C** 2016 Loss of Fertility in the Absence of Progesterone Receptor Expression in Kisspeptin Neurons of Female Mice. *PLoS One* 11:e0159534
30. **Kim J, Semaan SJ, Clifton DK, Steiner RA, Dhamija S, Kauffman AS** 2011 Regulation of Kiss1 expression by sex steroids in the amygdala of the rat and mouse. *Endocrinology* 152:2020-2030
31. **Di Giorgio NP, Semaan SJ, Kim J, Lopez PV, Bettler B, Libertun C, Lux-Lantos VA, Kauffman AS** 2014 Impaired GABAB receptor signaling dramatically up-regulates Kiss1 expression selectively in nonhypothalamic brain regions of adult but not prepubertal mice. *Endocrinology* 155:1033-1044
32. **Stephens SB, Chahal N, Munaganuru N, Parra RA, Kauffman AS** 2016 Estrogen Stimulation of Kiss1 Expression in the Medial Amygdala Involves Estrogen Receptor-alpha But Not Estrogen Receptor-beta. *Endocrinology* 157:4021-4031
33. **Stephens SBZ, Di Giorgio NP, Liaw RB, Parra RA, Yang JA, Chahal N, Lux-Lantos VA, Kauffman AS** 2018 Estradiol-Dependent and -Independent Stimulation of Kiss1 Expression in the Amygdala, BNST, and Lateral Septum of Mice. *Endocrinology* 159:3389-3402
34. RRID: AB_2571870. <https://antibodyregistry.org/search?q=ab167453>
35. RRID: AB_2340621. <https://antibodyregistry.org/search?q=711-585-152>
36. **Falcy BA, Mohr MA, Micevych PE** 2020 Immunohistochemical amplification of mCherry fusion protein is necessary for proper visualization. *MethodsX* 7:100946
37. **Christian CA, Mobley JL, Moenter SM** 2005 Diurnal and estradiol-dependent changes in gonadotropin-releasing hormone neuron firing activity. *Proc Natl Acad Sci U S A* 102:15682-15687

38. **Stephens SBZ, Rouse ML, Tolson KP, Liaw RB, Parra RA, Chahal N, Kauffman AS** 2017 Effects of Selective Deletion of Tyrosine Hydroxylase from Kisspeptin Cells on Puberty and Reproduction in Male and Female Mice. *eNeuro* 4
39. **Dror T, Franks J, Kauffman AS** 2013 Analysis of Multiple Positive Feedback Paradigms Demonstrates a Complete Absence of LH Surges and GnRH Activation in Mice Lacking Kisspeptin Signaling. *Biol Reprod* 88:1-8
40. RRID: AB_2665514. https://antibodyregistry.org/search.php?q=AB_2665514
41. RRID: AB_2665513. https://antibodyregistry.org/search.php?q=AB_2665513
42. **Yang JA, Hughes JK, Parra RA, Volk KM, Kauffman AS** 2018 Stress rapidly suppresses in vivo LH pulses and increases activation of RFRP-3 neurons in male mice. *J Endocrinol* 239:339-350
43. **Semaan SJ, Murray EK, Poling MC, Dhamija S, Forger NG, Kauffman AS** 2010 BAX-Dependent and BAX-Independent Regulation of Kiss1 Neuron Development in Mice. *Endocrinology* 151:5807-5817
44. **Semaan SJ, Kauffman AS** 2015 Daily successive changes in reproductive gene expression and neuronal activation in the brains of pubertal female mice. *Mol Cell Endocrinol* 401:84-97
45. **Luo E, Stephens SB, Chaing S, Munaganuru N, Kauffman AS, Breen KM** 2016 Corticosterone Blocks Ovarian Cyclicity and the LH Surge via Decreased Kisspeptin Neuron Activation in Female Mice. *Endocrinology* 157:1187-1199
46. **Esparza LA, Schafer D, Ho BS, Thackray VG, Kauffman AS** 2020 Hyperactive LH Pulses and Elevated Kisspeptin and NKB Gene Expression in the Arcuate Nucleus of a PCOS Mouse Model. *Endocrinology* 161

Figure Legends

Figure 1: Experimental design and timeline for virally re-expressing PGR selectively in AVPV kisspeptin neurons of KissPRKO females and then assessing their ability to generate E₂-induced LH surges.

Figure 2: Double-label IHC verification of proper mCherry expression in AVPV kisspeptin cells of **A)** Cre⁺ mice (KissPRKO) but not **B)** Cre⁻ mice (WT controls) after targeted AAV injection into the AVPV. Kisspeptin staining is yellow; mCherry is pseudo-colored magenta. White arrowheads in **A** denote kisspeptin neurons. Importantly, there was no spread of AVPV-infused AAV to the ARC region, indicated by **C)** complete absence of mCherry in kisspeptin cells in the ARC of Cre⁺ animals. Scale bars are shown in white for each image.

Figure 3: Very low co-expression of *Pgr* in AVPV *Kiss1* neurons of KissPRKO^{mCh} females and up-regulation of *Pgr* co-expression in KissPRKOs after “Pgr AAV” treatment, as assessed with double-label ISH. **A)** Representative microscope images of *Pgr* (white silver grains) and *Kiss1* (red fluorescence) co-expression in AVPV neurons of OVX+E₂ females 4 weeks after treatment with either Pgr AAV or control mCh AAV. Yellow arrowheads denote example *Kiss1* cells with *Pgr* co-expression; blue arrowheads denote example non-double labeled *Kiss1* cells. **B)** Mean % of *Kiss1*+*Pgr* co-expression in the AVPV of all 4 PM groups (n=7-10/group). Different letters above the bars signify significantly different ($p < 0.05$) from each other.

Figure 4: Selective re-introduction of PR into AVPV kisspeptin neurons rescues the LH surge in OVX+E₂ females. **A)** The occurrence of LH surges in each group, represented by the % of females demonstrating a surge after treatment with a validated E₂ positive feedback regimen known to elicit an LH surge in the PM but not AM time-point (n=7-10/PM group and n=4-6/AM group). The LH surge occurrence was very low in KissPRKO^{mCh} females and increased to WT levels in KissPRKO^{Pgr} females. **B)** Individual dot plot of PM LH values showing the high proportion of animals achieving an LH surge in both WTs and KissPRKO^{Pgr} (Cre⁺ Pgr) females but not KissPRKO^{mCh} (Cre⁺ mCh) females. Dashed horizontal line designates LH surge threshold (>0.70 ng/ml). **C)** Mean serum LH levels in OVX+E₂ females at the AM and PM time-periods demonstrating elevated surge-levels of LH in PM KissPRKO^{Pgr} but not PM KissPRKO^{mCh} females (n=7-

10/PM group and n=4-6/AM group). Different letters above the bars signify significantly different ($p < 0.05$) from each other.

Accepted Manuscript

Figure 1

Figure 1




Figure 2

Figure 2




Figure 3

Figure 3





Figure 4

Figure 4

