

SEMIPARAMETRIC MIXTURE OF BINOMIAL REGRESSION WITH A DEGENERATE COMPONENT

J. Cao and W. Yao

Simon Fraser University and Kansas State University

Abstract: Many datasets contain a large number of zeros, and cannot be modeled directly using a single distribution. Motivated by rain data from a global climate model, we study a semiparametric mixture of binomial regressions, in which both the component proportions and the success probabilities depend on the predictors nonparametrically. An EM algorithm is proposed to estimate this semiparametric mixture model by maximizing the local likelihood function. We also consider a special case in which the component proportions are constant while the component success probabilities still depend on the predictors nonparametrically. This model is estimated by a one-step backfitting procedure, and the estimates are shown to achieve the optimal convergence rates. The asymptotic properties of the estimates for both models are established. The proposed procedures are applied to rain data from a global climate model and historical rain data from Edmonton, Canada. Simulation studies show that satisfactory estimates are obtained for the proposed models for finite samples.

Key words and phrases: climate change, EM algorithm, weather data

1 Introduction

It is of great interest to study the evidence and impacts of climate change from weather data over periods of time that range from decades to millions of years (Parmesan and Yohe (2003), IPCC (2007), Tebaldi and Sansó (2009), and Smith et al. (2009)). While historical weather data are often limited, massive amounts of data for future weather can be generated from a global climate model.

Global climate models are mathematical models of the general circulation of a planetary atmosphere or ocean. There are about 25 versions of global climate models developed in different research centers across the world. Global climate models are commonly used for simulating the atmosphere or ocean of the earth

using complex computer programs. They are widely used for weather forecasting, understanding the climate, and projecting climate changes. The Geophysical Fluid Dynamics Laboratory (GFDL) in the U.S.A. developed one global climate model and implemented some computer simulations (Delworth et al. (2006), Gnanadesikan et al. (2006), Wittenberg et al. (2006) and Stouffer et al. (2006)).




Figure 1: The number of rain days per week in the Edmonton area, Canada in three periods 1981-2000, 2046-2065, and 2081-2100, based on the outputs of the GFDL’s global climate model.

Figure 1 displays the number of rain days per week in the Edmonton area, Canada in three periods 1981-2000, 2046-2065, and 2081-2100, based on the outputs of the GFDL’s global climate model. Although such count data might typically be modelled using a binomial distribution, preliminary analysis indicates that this variable does not follow a binomial distribution since too many weeks have no rain days.

Motivated by this example, we propose a semiparametric mixture of binomial regression model

$$f(X(t) | \pi_1(t), p(t)) = \pi_1(t)\text{Bin}(X(t); N, 0) + \pi_2(t)\text{Bin}(X(t); N, p(t)), \quad (1)$$

where $\text{Bin}(X; N, p)$ is the probability mass function of a binomial distribution for the variable X based on N trials with success probability p . The first component is a degenerate distribution with mass 1 on 0. To make the model identifiable, we assume $N \geq 2$; see Teicher (1961) and Lindsay (1995). Two nonparametric functions, $\pi_1(t)$ and $\pi_2(t)$, are the proportions of zero component and binomial component, respectively, under the constraint $\pi_1(t) + \pi_2(t) = 1$. The nonparametric function, $p(t)$, is the success probability in the binomial component. The semiparametric mixture model (1) can be used to model many data with extra numbers of zeros, for example, the number of days per week having a forest fire, since zeros are commonplace in winter.

Mixtures of binomial distributions

$$\pi_1 \text{Bin}(x, p_1) + \dots + \pi_m \text{Bin}(x, p_m) \quad (2)$$

were first used by Pearson (1915) to model yeast cell count data. Since then, mixtures of binomial distributions have been used in such fields as medicine (Farewell and Sprott (1988)), biology (Brooks et al. (1997)) and veterinary science (Böhning et al. (1998)). If the component-specific p_j 's depend on some predictors parametrically, then (2) is called a ‘‘mixture of binomial regression’’. Mixtures of binomial regression models have wide applications in medicine (Zhu and Zhang (2004)), biology (Follmann and Lambert (1989)), marketing research (Wedel and DeSarbo (1933) and De Soete and DeSarbo (1991)), genetics (Zhang and Merikangas (2000)), medical research (Lwin and Martin (1989)), the economics of labor markets (Geweke and Keane (1999)), and agriculture (Wang and Puterman (1998)). However, conventional mixtures of binomial regression models require strong parametric assumptions about the p_j 's and cannot account for the dependence of π_j 's on the predictors.

The semiparametric mixture model (1) removes the parametric assumptions about the component proportions and success probabilities. The functions $\pi_1(t)$ and $p(t)$ in the semiparametric mixture model are estimated using nonparametric smoothing methods such as kernel regression. We propose an EM algorithm (Dempster, Laird, and Rubin (1977)) to maximize the local likelihood function, and prove that the EM algorithm monotonically increases the local likelihood function. The convergence rate of the consistent estimates and their asymptotic normality are also established.

In some cases, one has prior knowledge that the component proportions are constant over t . Therefore, we also consider the model

$$f(X(t) | \pi_1, p(t)) = \pi_1 \text{Bin}(X(t); N, 0) + \pi_2 \text{Bin}(X(t); N, p(t)). \quad (3)$$

This model requires a local estimate of $p(t)$ and a global estimate of π_1 . Then it is difficult to estimate both $p(t)$ and π_1 efficiently. We propose a one-step backfitting algorithm, in which π_1 is first estimated globally given an initial consistent estimate of $p(t)$, and then $p(t)$ is updated given the global estimate of π_1 . We show that the one-step backfitting estimates for both $p(t)$ and π_1 achieve the optimal convergence rates; the computation is much more efficient than performing multiple iterations.

The rest of the paper is organized as follows. In Section 2, we introduce the estimation procedure for both proposed semiparametric mixture of binomial regression models. The asymptotic properties of the proposed procedures are established in Section 3. The bandwidth selection is discussed in Section 4. In Section 5, we demonstrate the proposed procedures by modeling the rain data from GFDL's global climate model and the historical rain data from Edmonton, Canada. In Section 6, we use simulations to compare the finite sample performance of the proposed semiparametric mixture models with a non-mixture model.

2 Methods

In this section, we introduce the estimation procedures and algorithms for the proposed semiparametric mixture models (1) and (3). In (1), both the mixing proportion $\pi_1(t)$ and the success probability $p(t)$ depend on the predictor t nonparametrically. We propose to use a local likelihood criterion to estimate them. An EM algorithm is also proposed to maximize the local likelihood. In (3), the component proportion π_1 is a constant parameter while the success probability $p(t)$ depends on the predictor t nonparametrically. Therefore, we need to estimate π_1 using global data and $p(t)$ using local data. We propose a one-step backfitting procedure to achieve the optimal convergence rates for both the estimates of π_1 and $p(t)$.

2.1 Semiparametric mixture model with time-varying proportions

The semiparametric mixture model (1) has two nonparametric functions to estimate: the proportion of zero component $\pi_1(t)$ and the success probability $p(t)$. Kernel regression is applied to estimate these nonparametric functions, though one could use such other nonparametric smoothing methods as local polynomial, spline smoothing, and wavelets.

The two nonparametric functions, at any point t_0 , are estimated by maximizing the local log-likelihood (Tibshirani and Hastie (1987) and Fan and Gijbels (1996))

$$\begin{aligned} \ell(\boldsymbol{\theta}(t_0)) = & \frac{1}{n} \sum_{i=1}^n K_h(t_i - t_0) \log \left[\pi_1(t_0) I(x_i = 0) \right. \\ & \left. + \{1 - \pi_1(t_0)\} \binom{N}{x_i} p(t_0)^{x_i} \{1 - p(t_0)\}^{N-x_i} \right], \end{aligned} \quad (4)$$

where $\boldsymbol{\theta}(t_0) = \{\pi_1(t_0), p(t_0)\}^T$, x_i is the measurement or observation for $X(t)$ at t_i , $i = 1, \dots, n$, and $K_h(\cdot) = h^{-1}K(\cdot/h)$ is a rescaling of the kernel function $K(\cdot)$ with the bandwidth h . We use the Gaussian kernel for $K(\cdot)$ for the data analysis in Section 5 and our simulation study in Section 6. The choice of bandwidth h is discussed in Section 4.

Note that there is no explicit solution to the maximization of (4), we propose using an EM algorithm. Define a vector of component indicator $\mathbf{z}_i = (z_{i1}, z_{i2})^T$, where

$$z_{ij} = \begin{cases} 1, & \text{if } (x_i, t_i) \text{ is from the } j\text{-th component;} \\ 0, & \text{otherwise.} \end{cases}$$

Then the complete local log-likelihood function for the complete data $\{(x_i, \mathbf{z}_i), i = 1, \dots, n\}$, after omitting some irrelevant constants, is

$$\begin{aligned} l_c(\boldsymbol{\theta}(t_0)) = & \sum_{i=1}^n K_h(t_i - t_0) [I(x_i = 0) z_{i1} \log \pi_1(t_0) \\ & + z_{i2} \{\log(1 - \pi_1(t_0)) + x_i \log p(t_0) + (N - x_i) \log(1 - p(t_0))\}]. \end{aligned}$$

Let $\mathbf{y} = \{(x_1, t_1), \dots, (x_n, t_n)\}$, and $\boldsymbol{\theta}^{(k)}(t_0)$ be the value of $\boldsymbol{\theta}(t_0)$ after the k -th EM iteration. The E-step at the $(k + 1)$ -st iteration requires one to find

$E_{\boldsymbol{\theta}^{(k)}(t_0)}(l_c(\boldsymbol{\theta}(t_0)) \mid \mathbf{y})$. Since $l_c(\boldsymbol{\theta}(t_0))$ is a linear function of z_{ij} , the E-step is equivalent to finding the classification probabilities $p_{ij}^{(k+1)} = E_{\boldsymbol{\theta}^{(k)}(t_0)}(Z_{ij} \mid \mathbf{y})$, where Z_{ij} is the random variable corresponding to z_{ij} . The M step at the $(k+1)$ -st iteration requires one to maximize $E_{\boldsymbol{\theta}^{(k)}(t_0)}(l_c(\boldsymbol{\theta}(t_0)) \mid \mathbf{y})$ with respect to $\boldsymbol{\theta}(t_0)$, for which there are explicit solutions.

Let $\pi_1^{(k)}(t_0)$ and $p^{(k)}(t_0)$ be the value of $\pi_1(t_0)$ and $p(t_0)$ at the k -th iteration.

E step : find the classification probabilities, given the current estimate,

$$\begin{aligned} p_{i1}^{(k+1)} &= \frac{\pi_1^{(k)}(t_0)\text{Bin}(x_i; N, 0)}{\pi_1^{(k)}(t_0)\text{Bin}(x_i; N, 0) + \{1 - \pi_1^{(k)}(t_0)\}\text{Bin}(x_i; N, p^{(k)}(t_0))}, \\ p_{i2}^{(k+1)} &= 1 - p_{i1}^{(k+1)}, i = 1, \dots, n. \end{aligned}$$

M step : update $\{\pi_1(t_0), p(t_0)\}$ by

$$\begin{aligned} \pi_j^{(k+1)}(t_0) &= \frac{\sum_{i=1}^n K_h(t_i - t_0) p_{ij}^{(k+1)}}{\sum_{i=1}^n \sum_{j=1}^2 K_h(t_i - t_0) p_{ij}^{(k+1)}}, j = 1, 2, \\ p^{(k+1)}(t_0) &= \frac{\sum_{i=1}^n K_h(t_i - t_0) p_{i2}^{(k+1)} x_i}{N \sum_{i=1}^n K_h(t_i - t_0) p_{i2}^{(k+1)}}. \end{aligned} \quad (5)$$

The algorithm monotonically increases the local log-likelihood (4) after each iteration, as shown in the following theorem.

Theorem 2.1. *At each iteration of the E and M steps, $\ell(\boldsymbol{\theta}^{(k+1)}(t_0)) \geq \ell(\boldsymbol{\theta}^{(k)}(t_0))$ for all k , where $\boldsymbol{\theta}(t_0) = (\pi_1(t_0), p(t_0))$ and $\ell(\cdot)$ is defined at (4).*

The proof of Theorem 2.1 is given in the supplementary file.

2.2 Semiparametric mixture model with constant proportions

For the semiparametric model (3), the success probability $p(t)$ needs to be estimated locally while the constant proportion π_1 can be estimated globally. It is then not easy to estimate both $p(t)$ and π_1 efficiently. In this section, we propose a one-step backfitting procedure to estimate the model (3): π_1 is estimated globally given an initial consistent estimate of $p(t)$; we update the estimate for $p(t)$ given the root n consistent estimate of π_1 .

We first consider the global estimation of it. Let $\hat{p}(t)$ and $\hat{\pi}_1$ denote consistent estimates of $p(t)$ and π_1 , respectively; these can be obtained by maximizing the local log-likelihood (4). Since $\hat{\pi}_1$ is a local estimator, it is not root n consistent. To improve efficiency, π_1 can be estimated globally by maximizing the log-likelihood (6) using the EM algorithm after replacing $p(t)$ in (3) by $\hat{p}(t)$:

$$\ell_1(\pi_1) = \frac{1}{n} \sum_{i=1}^n \log \left[\pi_1 I(x_i = 0) + \pi_2 \binom{N}{x_i} \hat{p}(t_i)^{x_i} \{1 - \hat{p}(t_i)\}^{N-x_i} \right] \quad (6)$$

Let $\tilde{\pi}_1$ be the maximizer of (6). We prove the root n consistency of $\tilde{\pi}_1$ in Section 3.2.

The EM algorithm to maximize (6) at the $(k+1)$ th step is as follows

E step : find the classification probability given the current estimates

$$p_{i1}^{(k+1)} = \frac{\pi_1^{(k)} \text{Bin}(x_i; N, 0)}{\pi_1^{(k)} \text{Bin}(x_i; N, 0) + \pi_2^{(k)} \text{Bin}(x_i; N, \hat{p}(t_i))},$$

$$p_{i2}^{(k+1)} = 1 - p_{i1}^{(k+1)}, i = 1, \dots, n.$$

M step : update (π_1, π_2) by

$$\pi_j^{(k+1)} = \frac{\sum_{i=1}^n p_{ij}^{(k+1)}}{n}, j = 1, 2.$$

We can next improve the efficiency of the estimator for $p(t)$ given $\tilde{\pi}_1$. Replacing π_1 in (3) by $\tilde{\pi}_1$, we propose to estimate $p(t_0)$, for any given t_0 , by maximizing the local likelihood function

$$\ell_2(p(t_0)) = \frac{1}{n} \sum_{i=1}^n K_h(t_i - t_0) \log \left[\tilde{\pi}_1 I(x_i = 0) + (1 - \tilde{\pi}_1) \binom{N}{x_i} p(t_0)^{x_i} (1 - p(t_0))^{N-x_i} \right]. \quad (7)$$

Let $\tilde{p}(t_0)$ be the resulting estimate of $p(t_0)$. Since $\tilde{\pi}_1$ is a root n consistent estimate of π_1 , the $\tilde{p}(t_0)$ has the same efficiency as if π_1 were known. Therefore, $\tilde{p}(t_0)$ is more efficient than $\hat{p}(t_0)$, which needs to account for the uncertainty of $\hat{\pi}_1$, since $\hat{p}(t_0)$ and $\hat{\pi}_1$ are estimated locally simultaneously in (4). See Theorem 3.3 and 3.4 for more details.

An EM algorithm is proposed to maximize (7). The estimate for $p(t_0)$ is updated at the $(k+1)$ th step as follows

E step : find the classification probability given the current estimate

$$p_{i1}^{(k+1)} = \frac{\tilde{\pi}_1 \text{Bin}(x_i; N, 0)}{\tilde{\pi}_1 \text{Bin}(x_i; N, 0) + (1 - \tilde{\pi}_1) \text{Bin}(x_i; N, p^{(k)}(t_0))}$$

$$p_{i2}^{(k+1)} = 1 - p_{i1}^{(k+1)}, i = 1, \dots, n.$$

M step : update $p(t_0)$ by

$$p^{(k+1)}(t_0) = \frac{\sum_{i=1}^n K_h(t_i - t_0) p_{i2}^{(k+1)} x_i}{N \sum_{i=1}^n K_h(t_i - t_0) p_{i2}^{(k+1)}}.$$

The ascending property of the above EM algorithm can be established along the lines of Theorem 2.1, and we omit this here.

One might further employ the backfitting procedures with a full iteration between estimating π_1 and $p(t)$ (see, for example, Buja, Hastie and Tibshirani (1989), Hastie and Tibshirani (1990), and Opsomer and Ruppert (1999)), or with a profile likelihood approach (Severini and Staniswalis (1994)), to improve efficiency. However, we prove in Section 3.2 that the one-step backfitting procedure achieves the optimal convergence rate, and with less computation.

The idea of one-step estimate has been used by many authors to simplify the computation procedure. See, for example, Carroll et al. (1997) for the generalized partially linear single-index model, and Li and Liang (2008) for the generalized partially linear model.

3 Statistical Theory

In this section, we investigate the asymptotic properties of the proposed estimates for the semiparametric mixture models at (1) and (3).

3.1 Semiparametric mixture model with time-varying proportions

Convergence rates of the estimates $\hat{\pi}_1(t)$ and $\hat{p}(t)$ for (1) are in Theorem 3.1, and asymptotic normality results are in Theorem 3.2.

Theorem 3.1. *Under Conditions A–D in the Appendix, there exists a consistent maximizer $\hat{\theta}(t_0) = (\hat{\pi}_1(t_0), \hat{p}(t_0))^T$ for the local log-likelihood function (4) such*

that

$$\left\| \hat{\boldsymbol{\theta}}(t_0) - \boldsymbol{\theta}_0(t_0) \right\| = O_p \left\{ (nh)^{-1/2} + h^2 \right\},$$

where $\| \cdot \|$ is the Euclidian norm, and $\boldsymbol{\theta}_0(t_0)$ is the true value of $\boldsymbol{\theta}(t_0) = (\pi_1(t_0), p(t_0))^T$.

The proof of Theorem 3.1 is given in the Appendix.

For asymptotic normality of the estimates, we need some notations. Let

$$f(x, \boldsymbol{\theta}) = \pi_1 \text{Bin}(x; N, 0) + \pi_2 \text{Bin}(x; N, p),$$

where $\boldsymbol{\theta} = (\pi_1, p)$. Let $l(x, \boldsymbol{\theta}) = \log f(x, \boldsymbol{\theta})$ and

$$l_1(x, \boldsymbol{\theta}) = \frac{\partial}{\partial \boldsymbol{\theta}} l(x, \boldsymbol{\theta}), \quad l_2(x, \boldsymbol{\theta}) = \frac{\partial^2}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} l(x, \boldsymbol{\theta});$$

$$G(t) = E\{l_1(X, \boldsymbol{\theta}_0(t_0)) \mid T = t\}, \quad \mathcal{I}(t) = -E\{l_2(X, \boldsymbol{\theta}_0(t_0)) \mid T = t\}. \quad (8)$$

The moments of K and K^2 are denoted, respectively, by

$$\mu_j = \int t^j K(t) dt \quad \text{and} \quad \nu_j = \int t^j K^2(t) dt.$$

Theorem 3.2. *Under Conditions A – D in the Appendix, the estimate $\hat{\boldsymbol{\theta}}(t_0) = (\hat{\pi}_1(t_0), \hat{p}(t_0))^T$ satisfies*

$$\sqrt{nh} \left\{ \hat{\boldsymbol{\theta}}(t_0) - \boldsymbol{\theta}_0(t_0) - b(t_0)h^2 + o_p(h^2) \right\} \xrightarrow{D} N \left\{ 0, g^{-1}(t_0) \mathcal{I}^{-1}(t_0) \nu_0 \right\},$$

where $g(t)$ is the marginal density of the time random variable T , and

$$b(t_0) = \mathcal{I}^{-1}(t_0) \left\{ \frac{G'(t_0)g'(t_0)}{g(t_0)} + \frac{1}{2}G''(t_0) \right\} \mu_2, \quad (9)$$

The proof of Theorem 3.2 is given in the Appendix.

3.2 Semiparametric mixture model with constant proportions

Let $\tilde{\pi}_1$ and $\tilde{p}(t_0)$ denote the estimates for the semiparametric mixture of binomial regression model (3) using the one-step backfitting procedure introduced in Section 2.2. We first establish the root n consistency of $\tilde{\pi}_1$ and give its asymptotic distribution in Theorem 3.3. The asymptotic distribution of $\tilde{p}(t_0)$ is given in Theorem 3.4.

Theorem 3.3. *Under Conditions A – D in the Appendix, if $nh^4 \rightarrow 0$ and $nh^2 \log(1/h) \rightarrow \infty$, then*

$$\sqrt{n}(\tilde{\pi}_1 - \pi_1) \rightarrow N(0, \mathcal{I}_{\pi_1}^{-2}\Sigma),$$

where

$$\begin{aligned} \mathcal{I}_{\pi_1} &= -E \left\{ \frac{\partial^2 f(x, \pi_1, p(t))}{\partial \pi_1^2} \right\}, \\ \Sigma &= \text{var} \left\{ \frac{\partial f(x, \pi_1, p(t))}{\partial \pi_1} - \mathcal{I}_{\pi_1 p}(t) \psi(t, x) \right\}, \\ \mathcal{I}_{\pi_1 p}(t) &= -E \left\{ \frac{\partial^2 f(X, \pi_1, p(t))}{\partial \pi_1 \partial p} \middle| T = t \right\}, \end{aligned}$$

and $\psi(t, x)$ is the second entry of $\mathcal{I}(t)^{-1}l_1(x, \boldsymbol{\theta}(t))$.

The proof of Theorem 3.3 is given in the Appendix. That $h = o(n^{-1/4})$ while the optimal bandwidth for kernel smoothing usually satisfies $h = O(n^{-1/5})$ means that undersmoothing is required for $\tilde{p}(t)$ in order to get the root n convergence rate for $\tilde{\pi}_1$. This is consistent with what has been found by Carroll et al. (1997) and Li and Liang (2008).

Theorem 3.4. *Under Conditions A – D in the Appendix,*

$$\sqrt{nh} \left\{ \tilde{p}(t_0) - p(t_0) - \tilde{b}(t_0)h^2 \right\} \xrightarrow{D} N(0, g(t_0)^{-1} \mathcal{I}_p(t_0)^{-1} \nu_0),$$

where

$$\begin{aligned} \tilde{b}(t_0) &= \frac{1}{2g(t_0)\mathcal{I}_p(t_0)} \left\{ \Gamma''(t_0)g(t_0) + 2\Gamma'(t_0)g'(t_0) \right\} \mu_2, \\ \Gamma(t) &= E \left\{ \frac{\partial f(x, \pi_1, p(t_0))}{\partial p} \middle| T = t \right\}, \\ \mathcal{I}_p(t) &= \text{var} \left\{ \frac{\partial f(x, \pi_1, p(t_0))}{\partial p} \middle| T = t \right\}. \end{aligned}$$

The proof of the Theorem 3.4 is given in the Appendix. Note that $\mathcal{I}_p(t)$ is the (2, 2) element of $\mathcal{I}(t)$ and $\Gamma(t)$ is the second entry of $G(t)$, where $\mathcal{I}(t)$ and $G(t)$ are defined in (8). If $\mathcal{I}^{22}(t)$ is the (2, 2) element of $\mathcal{I}^{-1}(t)$, $1/\mathcal{I}_p(t_0) \leq \mathcal{I}^{22}(t_0)$. Comparing the results in Theorem 3.2 and 3.4, we see that the one-step backfitting estimator $\tilde{p}(t_0)$ has smaller asymptotic bias and variance than does $\hat{p}(t_0)$.

4 Bandwidth Selection

The nonparametric functions in (1) and (3) are estimated using kernel regression with some bandwidth h . The optimal bandwidth can be obtained by minimizing the asymptotic weighted mean square error; a practical data-driven bandwidth selector can be based on the Nadaraya-Watson estimator. For simplicity, we focus on (1). The methods are easily adaptive for (3).

Based on Theorem 3.2, one can see that the asymptotic bias of $\hat{\boldsymbol{\theta}}$ is $b(t_0)h^2$ and the asymptotic covariance is $(nh)^{-1}g^{-1}(t_0)\mathcal{I}^{-1}(t_0)\nu_0$. An optimal bandwidth for estimating $\boldsymbol{\theta}(t_0)$ can be obtained by minimizing the asymptotic weighted mean squared error (MSE)

$$\begin{aligned} & \mathbb{E} \left[\left\{ \hat{\boldsymbol{\theta}}(t_0) - \boldsymbol{\theta}(t_0) \right\}^T W \left\{ \hat{\boldsymbol{\theta}}(t_0) - \boldsymbol{\theta}(t_0) \right\} \right] \\ &= b(t_0)^T W b(t_0) h^4 + \frac{\nu_0}{g(t_0)nh} \text{tr} \{ \mathcal{I}^{-1}(t_0) W \} + o_p(a_n), \end{aligned}$$

where $a_n = \{h^4 + (nh)^{-1}\}$, $\text{tr}(A)$ is the trace of A , and W is a weight function. Then the optimal local bandwidth is

$$\hat{h}_{opt}(t_0) = \left[\frac{\text{tr} \{ \mathcal{I}^{-1}(t_0) W \} \nu_0}{4b(t_0)^T W b(t_0) g(t_0)} \right]^{1/5} n^{-1/5}, \quad (10)$$

where $b(t_0)$ is given in (9). If our main interest is in $p(t_0)$, the weight function W can be $\text{diag}\{0, 1\}$; if we are interested in both $\pi_1(t_0)$ and $p(t_0)$, we can take $W = \mathcal{I}(t_0)$, which is proportional to the inverse of the asymptotic variance of $\hat{\boldsymbol{\theta}}(t_0)$.

Based on the asymptotic bias and variance, we can also derive a global bandwidth selector by minimizing the asymptotic integrated weighted mean squared error (MSE)

$$\int \mathbb{E} \left[\left\{ \hat{\boldsymbol{\theta}}(t) - \boldsymbol{\theta}(t) \right\}^T W \left\{ \hat{\boldsymbol{\theta}}(t) - \boldsymbol{\theta}(t) \right\} \right] w(t) dt,$$

where $w(t)$ is any weight function, for example $g(t)$, or 1. Then the optimal global bandwidth is

$$\hat{h}_{opt} = \left[\frac{\nu_0 \int \text{tr} \{ \mathcal{I}^{-1}(t) W \} g^{-1}(t) w(t) dt}{4 \int b(t)^T W b(t) w(t) dt} \right]^{1/5} n^{-1/5}. \quad (11)$$

Note that there are some unknown quantities in (10) and (11). To proceed, one can use the Plug-In idea (see, for example, Ruppert, Sheather, and Wand (1995)) and replace the unknown quantities by estimates of them. In addition, one can also use cross-validation to choose the bandwidth, with a little more computation.

Noting that (5) is a conventional Nadaraya-Watson estimator if p_{i2} is either 0 or 1, we can also employ the existing bandwidth selector for the Nadaraya-Watson estimator; see, for example, Rice (1984) or Hurvich, Simonoff, and Tsai (1998), based on some initial partition of the data into different components. The initial classification probabilities, p_{ij} , can be estimated by assuming $\pi_1(t)$ and $p(t)$ are constant or polynomial functions of t . This simple initial fit cannot guarantee a consistent estimate, but is easy to implement and generally works well. This idea of initial parametric fit has been used by Fan and Gijbels (1996, Sec 4.2). Note that the nonzero observation x_i must be from the second component; the only uncertainty is the zero observations of x . Therefore, the impact of misspecification of $\pi_1(t)$ and $p(t)$ is very small. One can also iterate the above procedure several times to get a refined bandwidth. We use this bandwidth selection method in our simulations and data applications.

5 Applications

5.1 Rain data from GFDL's Global Climate Model

GFDL's computer simulation, based on their global climate model, generated rain data at 128 grid points in longitude and 64 grid points in latitude over the whole earth in three time periods: 1981-2000, 2046-2065, and 2081-2100. We chose a grid point close to Edmonton, Canada, and analyzed the rain data at this grid point. Figure 1 displays the number of rain days per week during the three time periods; a rain day is defined as a day with more than 1 millimeter of rainfall. Model (1) with $N = 7$ was used to analyze the rain data from each of the three time periods. We used the bandwidth introduced at the end of Section 4, and the Gaussian kernel for $K(\cdot)$ for our model (1). Similar choices were used for the other examples.

It is of great interest to estimate the trend of extreme weather when studying




Figure 2: The top and bottom panels show the estimated probabilities of having zero and seven rain days in one week within the three time periods 1981-2000 (solid lines), 2046-2065 (dashed lines) and 2081-2100 (thick dash-dotted lines), respectively.

the evidence of climate change. For the rain data, extreme weather includes having zero rain days in one week (too dry) or having seven rain days in one week (too wet). The probabilities of having zero and seven rain days in one week are calculated as $P(X(t) = 0) = \hat{\pi}_1(t) + \{1 - \hat{\pi}_1(t)\}\{1 - \hat{p}(t)\}^7$ and $P(X(t) = 7) = \{1 - \hat{\pi}_1(t)\}\hat{p}(t)^7$, respectively. Figure 2 displays the estimated $P(X(t) = 0)$ and $P(X(t) = 7)$ in the three time periods. The time period 2081-2100 has a high probability of having zero rain days in one week 22.0% on average. The average probability of having zero days in one week in the time period 2081-2100 increases 27.4% and 6.2% from the time periods 1981-2000 and 2046-2065, respectively.

The time period 2081-2100 also has a high probability of having seven rain days in one week, 0.26% on average. The time period 2081-2100 has the average probability of having seven rain days in one week increasing 9.4% and 84.0% from the time periods 1981-2000 and 2046-2065, respectively. This is suggestive of more extreme weather in the time period 2081-2100.

5.2 Historical Rain Data in Edmonton




Figure 3: The number of rain days per week in Edmonton, Canada during 1961-1993.

Figure 3 displays the number of rain days per week in Edmonton, Canada during 1961-1993. We fit the rain data with Model (1) with $N = 7$.

Figure 4 displays the estimates of the probabilities of having zero and seven rain days in one week. The time period 1986-1987 was extremely dry, having the largest probability of zero rain days in one week (21.1%) and the smallest probability of seven rain days in one week (0.042%). On the other hand, 1973 had extreme rainfall, with the smallest probability of having zero rain days in one week (7.4%) and the largest probability of having seven rain days in one week (0.200%).

We also fit the rain data with the non-mixture model

$$f(X(t) | p(t)) = \text{Bin}(X(t); N, p(t)). \quad (12)$$

This model ignores the degenerate zero component and is equivalent to Model (1) when $\pi_1(t) \equiv 0$. As a result, it seriously underestimates the probability of having zero and seven rain days in one week, which is around 5.9% and 0.046% for the whole time period. In Figure 4, one can also see that the estimates from




Figure 4: The top and bottom panels display the estimated probabilities of having zero and seven rain days in one week in Edmonton, Canada during 1961-1993, respectively. The solid and dashed lines are the estimates from Model (1) and Model (12), respectively. The dotted curves are the corresponding 95% confidence intervals for the estimates from Model (1).

the non-mixture model are almost flat; the non-mixture model does not have enough flexibility to capture the variation in the data.

The parametric bootstrap is applied to obtain 95% confidence intervals for the probabilities of having zero and seven rain days in one week, implemented as follows. Simulated data are generated from Model (1) with $N = 7$, where $\pi_1(t)$ and $p(t)$ are then to be the estimates from the rain data. The semiparametric mixture model is estimated from the simulated data in 1000 simulation replicates. The probabilities of having zero and seven rain days in one week are calculated from the 1000 estimates of the semiparametric mixture model. We then obtain 95% confidence interval by calculating the 2.5% and 97.5% quantiles of the 1000 probabilities of having zero and seven rain days in one week. Figure 4 displays the intervals. Note that the probability of having zero rain days in one week, as estimated from the non-mixture model, is under the lower confidence bound.

6 Simulations

Simulation studies were implemented to evaluate the finite sample performance of our estimators for Models (1) and (3); We compare them with the non-mixture model (12).




Figure 5: The mean of the estimates for $p(t)$ and $\pi_1(t)$ in Model (1), both are plotted in solid lines; the dashed lines are the true functions $p(t)$ and $\pi_1(t)$. The dash-dotted line is the mean of the estimates for $p(t)$ in the non-mixture model (12).

The simulated data were generated based on Model (1) and (3). Both cases took the true $p(t) = 0.3(1.5 + \cos(2\pi t))$. In the first case, the true $\pi_1(t)$ was the time-varying function $\pi_1(t) = 0.2(1.5 + \sin(2\pi t))$; in the second case, the true $\pi_1 = 0.4$. We evaluated the finite sample performance by varying the sample size as small ($n = 50$), medium ($n = 100$), and large ($n = 200$). The times were n equally-spaced grid points in $[0,1]$. Models (1) and (12) were estimated from the simulated data in the first case, and Models (3) and (12) were estimated in the second case. The simulation was done with 100 replicates.

Figure 5 displays the mean of the estimates for $\pi_1(t)$ and $p(t)$ in the semi-parametric mixture model (1) in the first scenario. For comparison, we also add the mean of the estimates for $p(t)$ in the non-mixture model (12). From Figure

5, one can see that the mean estimates of both $\pi_1(t)$ and $p(t)$ are very close to the true functions, while the mean estimate of $p(t)$ in the non-mixture model is smaller than the true $p(t)$, and this becomes more serious where the true $\pi_1(t)$ is large.

When the data are simulated based on the first scenario, the estimates for Model (1) and Model (12) are summarized in Table 1. The average absolute values of biases of $\hat{p}(t)$ using the semiparametric mixture model were 12%, 10%, and 7% of those using the non-mixture model when the sample size was 50, 100, and 200, respectively. The estimates using the semiparametric mixture model had slightly smaller average standard deviations for $p(t)$ than those using the non-mixture model. The semiparametric mixture model also reduced the average RMSE of $\hat{p}(t)$ by 53%, 60%, and 68% over the non-mixture model when the sample size was 50, 100, and 200, respectively.

When the data are simulated based on the second scenario, the mixture model (3) was estimated using the one-step estimator and the traditional full iterative backfitting algorithm. Table 2 gives the summary of the estimates. The two algorithms have almost the same quality of estimates for $p(t)$; the backfitting algorithm had a slightly smaller RMSE for $\hat{\pi}_1$ than the one-step method, but the one-step method took less than half the time. The non-mixture model (12) was also fitted to the same simulated data. The mixture model (3) reduced the average RMSE of $\hat{p}(t)$ by 62%, 70%, and 77% over the non-mixture model when the sample size was 50, 100, and 200, respectively.

Appendix

In this section, we provide a sketch of the proof of Theorems 2.1 and 3.3. The proof of Theorems 3.1, 3.2, and 3.4 are standard and are omitted here. Refer to the supplementary file for more detail.

Let $g(t)$ be the density function for t . The following technical conditions are to be imposed. They are not the weakest possible conditions, but they facilitate the proofs.

Technical Conditions:

Table 1: The summary of the estimates for Model (1) and (12) when the simulated data were generated based on Model (1). The true success probability function $p(t) = 0.3(1.5 + \cos(2\pi t))$, and the true $\pi_1 = 0.2(1.5 + \sin(2\pi t))$. The last three columns are the absolute values of bias, standard deviation (SD), and root mean squared error (RMSE) of the estimates for Models (1) and (12), averaged over n equally spaced points in $[0,1]$. Model (12) was estimated with the Penalized Iteratively Reweighted Least Squares (PIRLS) method (see e.g. Wood (2000)) using the “mgcv” package in R (R Development Core Team (2010)).

n	Model		BIAS	SD	RMSE
50	Mixture	$\pi_1(t)$	0.013	0.164	0.165
		$p(t)$	0.017	0.080	0.082
	Non-Mixture	$p(t)$	0.137	0.100	0.173
100	Mixture	$\pi_1(t)$	0.014	0.133	0.134
		$p(t)$	0.013	0.059	0.061
	Non-Mixture	$p(t)$	0.132	0.069	0.152
200	Mixture	$\pi_1(t)$	0.013	0.099	0.100
		$p(t)$	0.009	0.044	0.045
	Non-Mixture	$p(t)$	0.132	0.047	0.142

Table 2: The summary of the estimates for Model (3) and (12) when the simulated data were generated based on Model (1). The true success probability function was $p(t) = 0.3(1.5 + \cos(2\pi t))$, and the true $\pi_1 = 0.4$. “|BIAS|”, “SD”, “RMSE” are the absolute values of bias, standard deviation (SD) and root mean squared error (RMSE) of the estimates for Models (3) and (12), averaged over n equally spaced points in $[0,1]$. Model (12) was estimated with the Penalized Iteratively Reweighted Least Squares (P-IRLS) method (see e.g. Wood (2000)) using the “mgcv” package in R (R Development Core Team 2010). The last column is the computing time in seconds for 100 Simulations.

n	Model		Method	BIAS	SD	RMSE	Time
50	Mixture	π_1	One-Step	0.032	0.074	0.081	1.656
			Backfitting	0.011	0.075	0.076	3.890
		$p(t)$	One-Step	0.010	0.077	0.078	1.656
			Backfitting	0.009	0.079	0.079	3.890
	Non-Mixture	$p(t)$	P-IRLS	0.178	0.104	0.207	7.407
	100	Mixture	π_1	One-Step	0.032	0.055	0.064
Backfitting				1.9e-5	0.060	0.060	7.412
$p(t)$			One-Step	0.011	0.057	0.059	3.374
			Backfitting	0.009	0.058	0.059	7.412
Non-Mixture		$p(t)$	P-IRLS	0.182	0.073	0.197	8.625
200		Mixture	π_1	One-Step	0.026	0.037	0.045
	Backfitting			0.001	0.038	0.038	15.637
	$p(t)$		One-Step	0.009	0.042	0.044	7.710
			Backfitting	0.008	0.043	0.044	15.637
	Non-Mixture	$p(t)$	P-IRLS	0.181	0.053	0.190	11.625

A $\pi_1(t)$ and $p(t)$ have continuous second derivatives at t_0 , $0 < \pi_1(t_0) < 1$ and $0 < p(t_0) < 1$. (For Model (3), we make the same assumption for $p(t)$ and assume $0 < \pi_1 < 1$.)

B $g(t)$ has a continuous second derivative at the point t_0 and $g(t_0) > 0$.

C $K(\cdot)$ is a symmetric (about 0) kernel density with compact support $[-1, 1]$.

D The bandwidth h tends to zero such that $nh \rightarrow \infty$.

Proof of Theorem 2.1.

Note that

$$\ell(\boldsymbol{\theta}) = \frac{1}{n} \sum_{i=1}^n K_h(t_i - t_0) \log f(x_i, \boldsymbol{\theta}).$$

Hence,

$$\begin{aligned} \ell(\boldsymbol{\theta}^{(k+1)}) - \ell(\boldsymbol{\theta}^{(k)}) &= \sum_{i=1}^n \log \left\{ \frac{\pi_1^{(k)} B(x_i, N, 0) \pi_1^{(k+1)} B(x_i, N, 0)}{f(x_i, \boldsymbol{\theta}^{(k)}) \pi_1^{(k)} B(x_i, N, 0)} \right. \\ &\quad \left. + \frac{\pi_2^{(k)} B(x_i, N, p^{(k)}) \pi_2^{(k+1)} B(x_i, N, p^{(k+1)})}{f(x_i, \boldsymbol{\theta}^{(k)}) \pi_2^{(k)} B(x_i, N, p^{(k)})} \right\} K_h(x_i - x_0) \\ &= \sum_{i=1}^n \log \left\{ r_{i1}^{(k+1)} \frac{\pi_1^{(k+1)} B(x_i, N, 0)}{\pi_1^{(k)} B(x_i, N, 0)} + r_{i2}^{(k+1)} \frac{\pi_2^{(k+1)} B(x_i, N, p^{(k+1)})}{\pi_2^{(k)} B(x_i, N, p^{(k)})} \right\} K_h(x_i - x_0) \end{aligned}$$

By Jensen's inequality,

$$\begin{aligned} \ell(\boldsymbol{\theta}^{(k+1)}) - \ell(\boldsymbol{\theta}^{(k)}) &\geq \sum_{i=1}^n \left[r_{i1}^{(k+1)} \log \left\{ \frac{\pi_1^{(k+1)} B(x_i, N, 0)}{\pi_1^{(k)} B(x_i, N, 0)} \right\} K_h(x_i - x_0) \right. \\ &\quad \left. + r_{i2}^{(k+1)} \log \left\{ \frac{\pi_2^{(k+1)} B(x_i, N, p^{(k+1)})}{\pi_2^{(k)} B(x_i, N, p^{(k)})} \right\} K_h(x_i - x_0) \right] \end{aligned}$$

Based on (5), we have $\ell(\boldsymbol{\theta}^{(k+1)}) - \ell(\boldsymbol{\theta}^{(k)}) \geq 0$.

Proof of Theorem 3.3.

Let

$$f(x_i, \pi_1, \hat{p}(t_i)) = \log \left[\pi_1 I(x_i = 0) + \pi_2 \binom{N}{x_i} \hat{p}(t_i)^{x_i} (1 - \hat{p}(t_i))^{N-x_i} \right].$$

Based on a Taylor expansion of (6), we have that $\sqrt{n}(\tilde{\pi}_1 - \pi_1) = B_n^{-1}A_n + o_p(1)$, where

$$A_n = \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{\partial f(x_i, \pi_1, \hat{p}(t_i))}{\partial \pi_1} \text{ and } B_n = -\frac{1}{n} \sum_{i=1}^n \frac{\partial^2 f(x_i, \pi_1, \hat{p}(t_i))}{\partial \pi_1^2}$$

It can be shown that

$$B_n = -\mathbb{E} \left\{ \frac{\partial^2 f(x_i, \pi_1, p(t_i))}{\partial \pi_1^2} \right\} + o_p(1) = \mathcal{I}_{\pi_1} + o_p(1).$$

And that

$$\begin{aligned} A_n &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{\partial f(x_i, \pi_1, p(t_i))}{\partial \pi_1} + \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{\partial^2 f(x_i, \pi_1, p(t_i))}{\partial \pi_1 \partial p} \{\hat{p}(t_i) - p(t_i)\} + O_p(d_{1n}) \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{\partial f(x_i, \pi_1, p(t_i))}{\partial \pi_1} + S_{n1} + O_p(d_{1n}). \end{aligned}$$

where $d_{1n} = n^{-1/2} \|\tilde{\pi}_1 - \pi_1\|_\infty^2 = o_p(1)$. From the proof of Theorem 3.2 (see the Supplement), we have

$$\hat{\boldsymbol{\theta}}(t_i) - \boldsymbol{\theta}(t_i) = \frac{1}{n} g(t_i)^{-1} \mathcal{I}(t_i)^{-1} \sum_{j=1}^n K_h(t_j - t_i) l_1(x_j, \boldsymbol{\theta}(t_j)) + O_p(d_{n2}).$$

Similar to Li and Liang (2008) and Carroll et al. (1997), one can prove that $n^{1/2}d_{n2} = o_p(1)$ uniformly in t_i if $nh^2/\log(1/h) \rightarrow \infty$. Let $\psi(t_j, x_j)$ be the second entry of $\mathcal{I}(t_j)^{-1} l_1(x_j, \boldsymbol{\theta}(t_j))$. Since $p(t_i) - p(t_j) = O(t_i - t_j)$ and $K(\cdot)$ is symmetric about 0, we have

$$\begin{aligned} S_{n1} &= \frac{1}{n^{-3/2}} \sum_{j=1}^n \sum_{i=1}^n \frac{\partial^2 f(x_i, \pi_1, p(t_i))}{\partial \pi_1 \partial p} g(t_i)^{-1} \psi(t_j, x_j) K_h(t_j - t_i) + O_p(n^{1/2}h^2) \\ &= S_{n2} + O_p(n^{1/2}h^2). \end{aligned}$$

It can be shown, by calculating the second moment, that $S_{n2} - S_{n3} = o_p(1)$, where $S_{n3} = -n^{-1/2} \sum_{j=1}^n \xi(t_j, x_j)$, with

$$\xi(t_j, x_j) = -\mathbb{E} \left\{ \frac{\partial^2 f(x, \pi_1, p(t_j))}{\partial \pi_1 \partial p} \mid t = t_j \right\} \psi(t_j, x_j) = \mathcal{I}_{\pi_1 p}(t_j) \psi(t_j, x_j).$$

Using $nh^4 \rightarrow 0$,

$$A_n = n^{-1/2} \sum_{i=1}^n \left\{ \frac{\partial f(x_i, \pi_1, p(t_i))}{\partial \pi_1} - \xi(t_i, x_i) \right\} + o_p(1).$$

We can show that $E(A_n) = 0$. With

$$\Sigma = \text{var}(A_n) = \text{var} \left\{ \frac{\partial f(x, \pi_1, p(t))}{\partial \pi_1} - \xi(t, x) \right\},$$

the Central Limit Theorem gives us $\sqrt{n}(\tilde{\pi}_1 - \pi_1) \rightarrow N(0, \mathcal{I}_{\pi_1}^{-2}\Sigma)$.

Acknowledgment

The authors would like to thank Dr. Peter Hall, an associate editor, and two referees for their constructive suggestions and comments that greatly improved the paper. They also thank Dr. Claudia Tebaldi for providing the rain data from global climate models. JC's research is partly supported by a discovery grant from the Natural Sciences and Engineering Research Council of Canada (NSERC).

References

- Brooks, S. P., Morgan, B. J. T., Ridout, M. S., and Pack, S. E. (1997). Finite mixture models for proportions. *Biometrics* 53, 1097-1115.
- Buja, A., Hastie, T. J., and Tibshirani, R. J. (1989). Linear smoothers and additive models (with discussion). *Annals of Statistics* 17, 453-555.
- Böhning, D. (1998). Zero-inflated Poisson models and C.A.MAN: A tutorial collection of evidence. *Biometrical Journal* 40, 833-843.
- Carroll, R. J. Fan, J., Gijbels, I., and Wang, M. P. (1997). Generalized partially linear single-index models. *Journal of the American Statistical Association* 92, 477-489.
- De Soete, G. and DeSarbo, W. S. (1991). A latent class probit model for analyzing pick any/N data. *Journal of Classification* 8, 45-63.
- Delworth et al. (2006). GFDL's CM2 global coupled climate models - Part 1: Formulation and simulation characteristics, *Journal of Climate* 19, 643-674.

- Dempster, A. P., Laird, N. M., and Rubin, D. B. (1977). Maximum likelihood estimation from incomplete data via the EM algorithm (with discussion). *Journal of the Royal Statistical Society B* 39, 1-38.
- Fan, J. and Gijbels, I. (1996). *Local Polynomial Modelling and Its Applications*. Chapman and Hall, London.
- Farewell, V. T. and Sprott, D. A. (1988). The use of a mixture model in the analysis of count data. *Biometrics* 44, 1191-1194.
- Follmann, D. A. and Lambert, D. (1989). Generalizing logistic regression by nonparametric mixing. *Journal of the American Statistical Association* 84, 295-300.
- Fong, D. Y. T. and Yip, P. S. F. (1993). An EM algorithm for a mixture model of count data. *Statistics & Probability Letters* 17, 53-60.
- Fong, D. Y. T. and Yip, P. S. F. (1993). A note on informatino loss in analyzing a mixture model for count data. *Communications in Statistics-Theory and Methods* 24, 3197-3209.
- Geweke, J. and Keane, M. (1999). Mixture of normals probit models. In *Analysis of Panels and Limited Dependent Variables* edited by C. Hsiao, M. H. Pearson, K. L. Lahiri, and L. F. Lee, 49-78. Cambridge University Press, Cambridge.
- Gnanadesikan et al. (2006). GFDL's CM2 global coupled climate models - Part 2: The baseline ocean simulation, *Journal of Climate* 19, 675-697.
- Hastie, T. and Tibshirani, R. J. (1990). *Generalised Additve Models*. Chapman and Hall, London.
- Hurvich, C. M., Simonoff, J. S., and Tsai, C. L. (1998). Smoothing parameter selection in nonparametric regression using an improved Akaike information criterion. *Journal of the Royal Statistical Society B* 60, 271-293.
- IPCC (2007). Climate change 2007: the physical science basis (summary for policy makers). IPCC.
- Lambert, D. (1992). Zero-inflated Poisson regression with an application to defects in manufacturing. *Technometrics* 34, 1-14.

- Li, R. and Liang, H. (2008). Variable selection in semiparametric regression modeling. *The Annals of Statistics* 36, 261-286.
- Lindsay, B. G., (1995), *Mixture Models: Theory, Geometry, and Applications*, NSF-CBMS Regional Conference Series in Probability and Statistics v 5. Institute of Mathematical Statistics, Hayward, CA.
- Lwin, T. and Martin, P. J. (1989). Probits of mixtures. *Biometrics* 45, 721-732.
- Meng, X. L. (1997). The EM algorithm and medical studies: a historical link. *Statistical Methods in Medical Research* 6, 3-23.
- Nadaraya, E. A. (1964). On estimating regression. *Theory of Applied Probability* 10, 186-190.
- Opsomer, J. D. and Ruppert, D. (1999). A root-n consistent backfitting estimator for semiparametric additive modeling. *Journal of Computational and Graphical Statistics* 8, 715-732.
- Parmesan, C. and Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature* 421, 37-42.
- Pearson, K. (1915). On certain types of compound frequency distributions in which the components can be individually described by binomial series. *Biometrika* 11, 139-144.
- R Development Core Team (2010). *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Vienna, Austria, ISBN 3-900051-07-0, <http://www.R-project.org>.
- Rice, J. (1984). Bandwidth choice for nonparametric regression. *Annals of Statistics* 12, 1215-1230.
- Smith, R.L., C. Tebaldi, D. Nychka, and L. O. Mearns (2009). Bayesian modeling of uncertainty in ensembles of climate models. *Journal of the American Statistical Association* 104, 97-116.
- Severini, T. A. and Staniswalis, J. G. (1994). Quasi-likelihood estimation in semiparametric models. *Journal of the American Statistical Association* 89, 501-511.

- Stouffer et al. (2006). GFDL's CM2 global coupled climate models - Part 4: Idealized climate response, *Journal of Climate* 19, 723-740.
- Tebaldi, C. and Sansó, B. (2009). Joint projections of temperature and precipitation change from multiple climate models: a hierarchical Bayesian approach. *Journal of the Royal Statistical Society A* 1, 83-106.
- Tibshirani, R. J. and Hastie, T. J. (1987). Local Likelihood Estimation. *Journal of the American Statistical Association* 82, 559-567.
- Tierney, H. (1961). Identifiability of mixtures. *The Annals of Mathematical Statistics* 32, 244-248.
- Wang, P. and Puterman, M. L. (1998). Mixed logistic regression models. *Journal of Agricultural, Biological, and Environmental Statistics* 3, 175-200.
- Watson, G. S. (1964). Smooth regression analysis. *Sankhya A* 26, 359-372.
- Wedel, M. and DeSarbo, W. S. (1933). A latent class binomial logit methodology for the analysis of paired comparison data. *Decision Sciences* 24, 1157-1170.
- Wittenberg et al. (2006). GFDL's CM2 global coupled climate models - Part 3: Tropical Pacific climate and ENSO, *Journal of Climate* 19, 698-722.
- Wood, S. N. (2000). Modelling and smoothing parameter estimation with multiple quadratic penalties. *Journal of the Royal Statistical Society, B* 62, 413-428.
- Yin, P. (1988). Inference about the mean of a Poisson distribution in the presence of a nuisance parameter. *Australian Journal of Statistics* 30, 299-306.
- Yin, P. (1991). Conditional inference for a mixture model for the analysis of count data. *Communications in Statistics-Theory and Methods* 20, 2045-2057.
- Zhu, H.-T. and Zhang, H. (2004). Hypothesis testing in mixture regression models. *Journal of the Royal Statistical Society B* 66, 3-16.
- Zhang, H. P. and Merikangas, K. (1964). A frailty model of segregation analysis: Understanding the familial transmission of alcoholism. *Biometrics* 56, 815-823.

J. Cao

Department of Statistics & Actuarial Science, Simon Fraser University

8888 University Drive, Burnaby, BC, V5A1S6, Canada

E-mail: jca76@sfu.ca

W. Yao

Department of Statistics, Kansas State University

101 Dickens Hall, Manhattan, KS, 66506-0802, USA

E-mail: wxyao@ksu.edu