UC Santa Barbara

UC Santa Barbara Previously Published Works

Title

Understanding the development and evolution of novel floral form in Aquilegia.

Permalink

<https://escholarship.org/uc/item/245173f7>

Journal

Current opinion in plant biology, 17(1)

ISSN

1369-5266

Authors

Sharma, Bharti
Yant, Levi
Hodges, Scott A
et al.

Publication Date

2014-02-01

DOI

10.1016/j.pbi.2013.10.006

Peer reviewed

Understanding the development and evolution of novel floral form in *Aquilegia*

Bharti Sharma¹, Levi Yant¹, Scott A Hodges² and Elena M Kramer¹

Flowers of the lower eudicot *Aquilegia* (columbine) possess morphological innovations, namely elaborate petal spurs and a fifth distinct organ identity, the staminodium, that are well suited to the investigation of key questions in developmental evolution. The recent evolution of these characteristics combined with a growing set of genetic and genomic resources has provided insight into how the traits arose and diversified. The petal spur appears to represent a key innovation that diversified largely via modification of specific aspects of cell expansion. In the case of the staminodium, gene duplication has played a role in allowing a novel organ identity to be carved out of the traditional ABC program.

Addresses

¹ Department of Organismic and Evolutionary Biology, Harvard University, 16 Divinity Avenue, Cambridge, MA 02138, USA

² Department of Ecology, Evolution & Marine Biology, University of California, Santa Barbara, Santa Barbara, CA 93106-9610, USA

Corresponding authors: Kramer, Elena M (ekramer@oeb.harvard.edu)

Current Opinion in Plant Biology 2014, 17:22–27

This review comes from a themed issue on **Growth and development**

Edited by **David R Smyth** and **Jo Ann Banks**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 15th November 2013

1369-5266/\$ – see front matter, © 2013 Elsevier Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.pbi.2013.10.006>

Introduction

The established plant genetic models have provided us with a detailed understanding of the genetic programs controlling the development of structures such as lateral organs, including the floral organs (reviewed [1,2]). However, we must recognize that these models capture only a very limited component of the enormous diversity in floral morphology. Commonly known as columbine, *Aquilegia* is a member of the basal eudicot family Ranunculaceae whose morphology and evolutionary history make it a powerful tool for studying the evolution of novel morphology [3,4]. The genus encompasses 70 recently diversified species, which although morphologically varied (Figure 1), remain largely interfertile [5]. In addition to this interesting recent history, the phylogenetic position of *Aquilegia*, roughly intermediate between the core eudicots and grasses, provides an attractive data point for deeper comparisons across the flowering plants. *Aquilegia* possesses five distinct types of floral organs: petaloid sepals in the first whorl;

nectiferous, spurred petals in the second whorl; four to seven whorls of 10 stamens each; a whorl of 10 sterile organs termed the staminodia; and an innermost whorl of four to seven carpels (Figure 2a,d). Although the floral bauplans of the diverse *Aquilegia* species are very similar, the flowers differ in color, many aspects of spur morphology, and, in one case, the presence of staminodia. Several groups have been using a growing set of genetic and genomic tools to investigate the genetic basis of the novel floral features of *Aquilegia*. These tools include a high-quality Sanger-sequenced genome for *A. × coerulea* ‘Goldsmith’ [6] with detailed gene annotation utilizing multiple deeply sequenced expressed sequence tag (EST) and RNAseq libraries (e.g., <http://compbio.dfci.harvard.edu/tgi/cgi-bin/tgi/gimain.pl?gudb=aquilegia>), resequencing of 13 additional species as well as the sister genus *Semiaquilegia* (Figure 1), two bacterial artificial chromosome (BAC) libraries [7], and a well-established virus induced gene silencing (VIGS) protocol [8].

There are a number of immediate questions suggested by *Aquilegia*’s floral morphology in relation to the otherwise conserved ABC model of floral organ identity. The ABC model suggests that floral organ identity is established by the interaction of three classes of gene activity: A class alone determines sepals; A + B, petals; B + C, stamens; and C alone, carpels [9]. This model fails to explain, however, how a flower could produce two whorls of morphologically distinct petaloid organs, as in the sepals and petals of *Aquilegia*, or even more dramatically, an entirely novel fifth organ such as the staminodium. Expression and functional studies of A, B and C gene homologs in *Aquilegia* have shed light on these questions. In regard to A function, it has previously been demonstrated that this role is not well conserved, even within the core eudicots [10,11]. Consistent with this, the *Aquilegia* homologs of the Arabidopsis A gene *APETALA1* (*AP1*), *AqFL1a* and *AqFL1b*, control leaf morphogenesis and inflorescence structure but do not contribute significantly to flower development [12••]. In contrast, C class function, represented by homologs of *AGAMOUS* (*AG*), is generally well-conserved across the angiosperms in terms of promoting stamen and carpel identity as well as floral meristem determinacy. Ongoing functional studies of the *Aquilegia* *AG* paralogs *AqAG1* and *AqAG2* suggest that this is equally true for the loci, although a role in novel staminodium identity requires further investigation (B Sharma *et al.*, unpublished data).

This leaves us to focus on the role of the B class genes, *APETALA3* (*AP3*) and *PISTILLATA* (*PI*) in Arabidopsis.

Figure 1

Simplified phylogeny of *Aquilegia* species relative to the sister genus *Semiaquilegia* (based on [22,30]). All species represented have been the subject of genome resequencing or, in the case of *A. jonesii*, deep RNA-seq. The evolution of several features is represented by colored bars, specifically the initial gain and subsequent loss of both petal nectar spurs and staminodia. The geographic distribution of the species is also indicated to the right of the phylogeny. EU = European, AS = Asian, and NA = North American. Panels (a)–(l) are photographs of the species, as indicated by the corresponding superscript letters. (a) *A. aurea*. (b) *A. vulgaris*. (c) *A. oxysepala*. (d) *A. ecalcarata*. (e) *A. formosa*. (f) *A. coerulea*. (g) *A. pinetorum*. (h) *A. longissima*. (i) *A. canadensis*. (j) *A. jonesii*. (k) *A. flabellata*. (l) *Semiaquilegia*. All photographs by SA Hodges except (c) and (k) by Hong-Xing Xiao.

They offer an obvious starting point since the novel features of *Aquilegia* involve both different forms of petaloidy and derivatives of the stamens. Early studies found that there are three major paralogous lineages of *AP3* in the Ranunculid order and, in *Aquilegia*, due to a more recent duplication, the genus actually possesses four *AP3*-like genes: *AqAP3-1*, *AqAP3-2*, *AqAP3-3* and *AqAP3-3b* [13,14]. These proteins all appear to function as heterodimers with the protein of the single *PI* homolog, *AqPI*, but each have distinct expression patterns [14]. *AqAP3-1* expression is initially broad but rapidly becomes restricted to the staminodia, *AqAP3-2* is also broad at early stages but is only consistently expressed in stamens, and both *AqAP3-3* and *AqAP3-3b* are entirely restricted to the petals, with *AqAP3-3b* expressed at much lower levels. Consistent with the dependence of each *AP3* protein on *AqPI*, the latter is broadly expressed across the petals, stamens and staminodia. Furthermore, transient RNAi targeting *AqPI* demonstrates that the B functional domain in *Aquilegia* is expanded to include the staminodia as well as the canonical petals and stamens (Figure 2e) [14]. Following from these data, the questions that have received recent attention are: Do the *AP3* paralogs, in fact, show subfunctionalization and neofunctionalization as suggested by their expression patterns? and How has both petal and staminodium morphology diversified down-stream of organ identity?

Petal identity and diversification

Work in *Aquilegia* has also informed questions about the evolution of petal identity itself. The diversity of Ranunculid petal form, combined with a variability even of their presence within the family, prompted botanists to conclude that they evolved from stamens many times independently [15,16]. With the advent of gene expression studies, however, Rasmussen *et al.* [17] suggested that this diversity was instead controlled by a commonly inherited identity program, which was lost in some cases rather than gained multiple times. More recent work in *Aquilegia* has supported this hypothesis: the MADS-domain *AqAP3-3/3b* paralogs have been shown to specifically promote petal identity with no contribution to stamen or staminodium development (Figure 2f) [18]. Furthermore, similar studies in the distantly related Ranunculaceae genus *Nigella* have recovered the same result [19^{••},20[•]]. While this evidence does seem to argue for a conserved petal identity controlled by *AqAP3-3* orthologs, the question remained: What happens to these genes when petals are naturally lost? This point has been addressed by a study that examined multiple pairs of closely related genera in which one taxon had petals while the other lacked them [19^{••}]. In every apetalous genus, expression of the *AqAP3-3* ortholog is dramatically reduced, if not completely eliminated, and the genomic loci show a variety of evidence of pseudogenization. All of these data

Figure 2

Novel floral organs of *Aquilegia* and homeotic RNAi phenotypes that affect their identity. **(a)** Flowers of *A. formosa*. On the left, a pre-anthesis flower with all stamens removed. The petaloid sepals (sep) and spurred petals (pet) are visible along with the inner staminodia (std) surrounding the carpels (car). Note that the staminodia are free from each other. On the right, a post-anthesis flower in which the sepals, petals and stamens have naturally abscised while the persistent staminodia are fused to form a continuous sheath that surrounds the carpels. Photograph by SA Hodges. **(b)** The spurred petal of *A. x coerulea* 'Goldsmith'. **(c)** The staminodial sheath of *A. x coerulea* 'Goldsmith'. Photographs in (b) and (c) by B Sharma. **(d)** Floral diagram for a wildtype (WT) flower showing five distinct organ identities: sepals (sep), petals (pet), stamens (sta), staminodia (std), and carpels (car). **(e)** Phenotype of *AqPI*-silenced flower (*aqpi*) in which petals are replaced by sepals and both stamens and staminodia by carpels. **(f)** In *AqAP3-3*-silenced flowers (*aqap3-3*), the petals are replaced by sepals with no effect on the other floral organs [18]. **(g)** *AqAP3-2*-silenced flowers (*aqap3-2*) show stunted, sterilized stamens, indicated in grey [32]. **(h)** *AqAP3-1*-silenced flowers (*aqap3-1*) display strong transformation of staminodia into carpels and some perturbation of inner stamen development [32]. **(i)** Double silencing of *AqAP3-1* and *AqAP3-2* (*aqap3-1/2*) results in complete transformation of all stamens and staminodia into carpels with no effect on petal identity but some reduction in size [32].

are, therefore, consistent with a model in which petal identity was commonly inherited but turned off in many separate instances. Of course, one may also ask why the loss of petals is so commonly tolerated in the family. One likely explanation is the presence of petaloid sepals in

almost all of the taxa. This transference of primary attractive function from the second to the first whorl could have released constraint on the petals, allowing them to diversify in morphology or, on occasion, be lost [17,19••].

Having established the genetic basis of petal identity, we can now turn our attention to the elaboration of these organs. *Aquilegia* flowers feature a diverse array of petal nectar spurs, the shape of which is tightly associated with specialized pollinators that range from bees to hummingbirds to hawkmoths (Figure 1; [21,22]). Tight correspondence between a pollinator's tongue and the spur length in any particular species is considered a product of selection for pollen removal and receipt, driving spur length over a 16-fold range and reproductive isolation between species [22]. Combined with the explosive worldwide radiation of *Aquilegia* species after the evolution of the nectar spur, this novelty is thought to be the key innovation that enabled *Aquilegia* to become a textbook example of adaptive radiation [23]. Despite the clear ecological and evolutionary significance of spur length, however, almost nothing was known about *Aquilegia* spur morphogenesis until very recently.

Early hypotheses posited that spur development is caused primarily by 'meristematic knobs' thought to flank the spur attachment point, adding one cell at a time until the final spur shape is achieved [24]. Recently, however, spur growth has been shown to be an extreme example of organ curvature, progressing in two dynamic phases. During phase I, diffuse cell divisions occur throughout petal primordium but when the nascent spur cup reaches ~1 mm in length, the division domain contracts in a wave that begins at both the blade and base, progressing eventually toward the future nectary [25**]. The result is that cell divisions persist in the spur region for longer than in the blade and base, which promotes the initial out-pocketing of the spur. This localized domain of cell division continues to contract until the spur reaches ~5 mm, when it enters phase II. This consists of highly anisotropic cell elongation that drives the bulk of spur growth and continues until the spur reaches its final length. The inference that spur shape ontogeny is governed by these two simple phases was confirmed by modeling: spurs were computationally 'grown' using thousands of experimental measurements of cell area and anisotropy to achieve spur profiles that were in agreement with natural profiles and shapes [25**]. Another critical finding of this study is that the considerable variation in spur length across various *Aquilegia* species appears to be entirely controlled by variation in anisotropic cell elongation, rather than cell number. The degree of elongation is, in turn, controlled by the duration of the elongation period, which suggests that a macro-morphological feature — spur length — is controlled by a cell level parameter, which is itself mediated by heterochronic shifts. A clear target of future work is to understand the genetic basis of this morphological novelty, which will take advantage of genome sequence resources that cover a wide range of species with various spur shapes and lengths, as well as the secondarily spurless *A. ecalcarata* (Figure 1d). This natural variation is a powerful tool

because the species are interfertile, which allows genetic dissection through segregation studies [26] and there are even natural hybrid zones where association mapping is promising [27].

***Aquilegia* staminodia, more than sterilized stamens**

Unlike the staminodia of many other angiosperms, which may simply be aborted stamens, the staminodia of *Aquilegia* are a continuous whorl of sterile, laterally expanded organs that surround the carpel whorl (Figure 2a,c). After anthesis, when all of the outer floral organs abscise, the staminodia stay attached to the receptacle and undergo late congenital fusion to form a cylindrical sheath [14]. The current hypothesis is that this reflects an herbivory deterrence mechanism protecting the early developing fruits [28], but this has yet to be tested in the field. In *Semiaquilegia*, the sister genus to *Aquilegia* (Figure 1l), similar organs are present in the same position, but they are more variable in number and morphology [29]. No other genus in the Ranunculaceae has similar staminodia in this position, suggesting that they evolved recently in a stepwise fashion, becoming sterilized in the last common ancestor of *Aquilegia* + *Semiaquilegia* ~8 mya and further elaborated in the lineage leading to *Aquilegia* ~6 mya [30*].

As described above, expression studies implicated the *AP3* paralogs *AqAP3-1* and *AqAP3-2* in the differential control of staminodium and stamen identity, respectively. Using VIGS, a transient RNAi approach, each of these paralogs has been silenced individually and in combination [32**]. In *AqAP3-2* silenced plants, stamens show broad anther necrosis and, in the most severe phenotypes, anthers were highly reduced to yield naked sterile filaments (Figure 2g), reminiscent of the underdeveloped staminodia seen in *Semiaquilegia*. There was no effect on the actual staminodia or other floral organs in *AqAP3-2*-silenced flowers, however. In the *AqAP3-1* knockdowns, the primary phenotype was homeotic conversion of the staminodia toward carpel identity with some weak necrosis or transformation to carpel identity in the inner stamens (Figure 2h). These findings would appear to suggest that *AqAP3-2* is essential to the proper development and fertility of the stamens while the main role of *AqAP3-1* is staminodium identity with some gradient of influence on the innermost stamens. These conclusions are supported by the double silencing phenotype, in which all of the stamens and staminodia were strongly transformed to carpels (Figure 2i). There was also some reduction in petal size but no obvious effect on identity. These findings reveal a complex picture of a modified ABC model following gene duplication. While *AqAP3-3* experienced a relatively ancient subfunctionalization to control only petal identity, more recent evolutionary changes to the *AqAP3-1* and *AqAP3-2* paralogs have allowed the definition of a new organ identity derived from the innermost

whorl of stamens. Both *AqAP3-1* and *AqAP3-2* are broadly expressed throughout the stamen/staminodium domain at very early stages and it appears that this early *AqAP3-1* expression is sufficient to promote some degree of stamen identity. Because of the later establishment of differential expression, we see that proper stamen development requires *AqAP3-2* while staminodium identity depends on *AqAP3-1*. Again, the available genomic resources for *Aquilegia* as well as its sister genus *Semiaquilegia*, will hopefully facilitate a better understanding of how this differential expression pattern evolved, as well as how the entire staminodium identity program was derived from the ancestral stamen program.

Conclusion

Aquilegia's recent evolutionary history makes it a particularly useful model for studying traits related to the evolution of floral novelty. That being said, there are many other developmental features of interest where *Aquilegia*, along with its genomic resources, will represent an important model, including compound leaves, cymose inflorescence structure, flower color, perennial life-form and the requirement for vernalization in controlling flowering time. For instance, the genus has been used to demonstrate that the role of *CUC* homologs in the dissection of leaf margins is deeply conserved across eudicots [31]. As our tools continue to improve, these comparisons will grow even more powerful.

Acknowledgements

Work described here was supported by NSF award IOS 0720240 and 1121005 to EMK, NSF award EF-0412727 to SAH and EMK and a Ruth L Kirschstein National Research Service Award from the National Institutes of Health to LY (1 F32 GM096699).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Moon J, Hake S: **How a leaf gets its shape.** *Curr Opin Plant Biol* 2011, **14**:24-30.
2. Causier B, Schwarz-Sommer Z, Davies B: **Floral organ identity: 20 years of ABCs.** *Sem Cell Dev Biol* 2010, **21**:73-79.
3. Kramer EM: ***Aquilegia*: a new model for plant development, ecology, and evolution.** *Ann Rev Plant Biol* 2009, **60**:261-277.
4. Kramer EM, Hodges SA: ***Aquilegia* as a model system for the evolution and ecology of petals.** *Philos Trans R Soc Lond B Biol Sci* 2010, **365**:477-490.
5. Hodges SA, Derieg NJ: **Adaptive radiations: from field to genomic studies.** *Proc Natl Acad Sci U S A* 2009, **106**:9947-9954.
6. Joint Genome Institute, Phytozome v9.1. <http://www.phytozome.net/>.
7. Fang GC, Blackmon BP, Henry DC, Staton ME, Saski CA, Hodges SA, Tomkins JP, Luo H: **Genomic tools development for *Aquilegia*: construction of a BAC-based physical map.** *BMC Gen* 2010, **11**:621.
8. Gould B, Kramer EM: **Virus-induced gene silencing as a tool for functional analyses in the emerging model plant *Aquilegia* (columbine, Ranunculaceae).** *Plant Methods* 2007, **3**:6.
9. Coen ES, Meyerowitz EM: **The war of the whorls: genetic interactions controlling flower development.** *Nature* 1991, **353**:31-37.
10. Litt A: **An evaluation of A-function: evidence from the *APETALA1* and *APETALA2* gene lineages.** *Int J Plant Sci* 2007, **168**:73-91.
11. Davies B, Cartolano M, Schwarz-Sommer Z: **Flower development: the *Antirrhinum* perspective.** *Adv Bot Res* 2006, **44**:279-321.
12. Pabón-Mora N, Sharma B, Holappa L, Kramer EM, Litt A: **The *Aquilegia FRUITFULL*-like genes play key roles in leaf morphogenesis and inflorescence development.** *Plant J* 2013, **74**:197-212.
- The authors demonstrate that the closest *AP1* homologs of *Aquilegia*, *AqFL1a/b*, do not significantly contribute to floral development but, instead, play novel roles in compound leaf development, particularly marginal lobing.
13. Kramer EM, Di Stilio VS, Schluter P: **Complex patterns of gene duplication in the *APETALA3* and *PISTILLATA* lineages of the Ranunculaceae.** *Int J Plant Sci* 2003, **164**:1-11.
14. Kramer EM, Holappa L, Gould B, Jaramillo MA, Setnikov D, Santiago P: **Elaboration of B gene function to include the identity of novel floral organs in the lower eudicot *Aquilegia* (Ranunculaceae).** *Plant Cell* 2007, **19**:750-766.
15. Prantl K: **Beiträge zur Morphologie und Systematik der Ranunculaceen.** *Bot Jahrb Syst* 1888, **9**:225-273.
16. Tamura M: **Morphology, ecology, and phylogeny of the Ranunculaceae IV.** *Sci Rep, Osaka Univ* 1965, **14**: 53-71.
17. Rasmussen DE, Kramer EM, Zimmer EA: **One size fits all? Molecular evidence for a commonly inherited petal identity program in the Ranunculales.** *Am J Bot* 2009, **96**:1-14.
18. Sharma B, Guo C, Kong H, Kramer EM: **Petal-specific subfunctionalization of an *APETALA3* paralog in the Ranunculales and its implications for petal evolution.** *New Phytol* 2011, **190**:870-883.
19. Zhang R, Guo C, Zhang W, Wang P, Li L, Duan X, Du Q, Zhao L, Shan H, Hodges SA *et al.*: **Disruption of the petal identity gene *APETALA3-3* is highly correlated with loss of petals within the buttercup family (Ranunculaceae).** *Proc Natl Acad Sci U S A* 2013, **110**:5074-5079.
- Detailed studies of multiple pairs of petalous and apetalous genera in the Ranunculaceae find that while *AP3-3* orthologs are specifically expressed in the petals of the former, their expression has been consistently lost in the latter. In addition, the genomic loci of apetalous genera show a range of signs of pseudogenization including complete deletion, major indels and increased rates of sequence evolution.
20. Goncalves B, Nogue O, Jabbour F, Ridet C, Morin H, Laufs P, Manicacci D, Damerval C: **An *APETALA3* homolog controls both petal identity and floral meristem patterning in *Nigella damascena* L. (Ranunculaceae).** *Plant J* 2013, **6**:223-235.
- In the first reverse genetic study of *Nigella*, the authors show that in addition to being required for petal identity, the *AP3-3* ortholog appears to contribute to the maintenance of AG homolog expression in outer stamen whorls.
21. Fulton M, Hodges SA: **Floral isolation between *Aquilegia formosa* and *Aquilegia pubescens*.** *Proc R Soc Lond B* 1999, **266**:2247-2252.
22. Whittall JB, Hodges SA: **Pollinator shifts drive increasingly long nectar spurs in columbine flowers.** *Nature* 2007, **447**:706-710.
23. Hodges SA, Arnold ML: **Spurring plant diversification: are floral nectar spurs a key innovation?** *Proc R Soc Lond B* 1995, **262**:343-348.
24. Tepfer SS: **Floral anatomy and ontogeny in *Aquilegia formosa* var. *truncata* and *Ranunculus repens*.** *Univ Calif Publ Bot* 1953, **25**:513-648.
25. Puzey JR, Gerbode S, Hodges SA, Mahadevan L, Kramer EM: **Evolution of spur-length diversity in *Aquilegia* petals is achieved solely through cell-shape anisotropy.** *Proc R Soc Lond B* 2012, **279**:1640-1645.

Cell level studies of petal spur development find that spur growth is primarily mediated by anisotropic cell elongation and, further, that variation in this one parameter can explain the full range of spur length differences seen within the genus.

26. Hodges SA, Whittall JB, Fulton M, Yang JY: **Genetics of floral traits influencing reproductive isolation between *Aquilegia formosa* and *Aquilegia pubescens***. *Am Nat* 2002, **159**(Suppl.):S51-S60.
 27. Cooper EA, Whittall JB, Hodges SA, Nordborg M: **Genetic variation at nuclear loci fails to distinguish two morphologically distinct species of *Aquilegia***. *PLoS ONE* 2010, **5**:e8655.
 28. Voelckel C, Borevitz J, Kramer EM, Hodges SA: **Within and between whorls: comparative transcriptional profiling of *Aquilegia* and *Arabidopsis***. *PLoS ONE* 2010, **5**:e9735.
 29. Tucker SC, Hodges SA: **Floral ontogeny of *Aquilegia*, *Semiaquilegia*, and *Isopyrum* (Ranunculaceae)**. *Intl J Plant Sci* 2005, **166**:557-574.
 30. Fior S, Li MG, Oxelman B, Viola R, Hodges SA, Ometto L, Varotto C: **Spatiotemporal reconstruction of the *Aquilegia* rapid radiation through next-generation sequencing of rapidly evolving cpDNA regions**. *New Phytol* 2013, **198**: 579-592.
- A well resolved phylogenetic analysis of the *Aquilegia* radiation, examining both biogeographic patterns and the timing of major divergences.
31. Blein T, Pulido A, Vialette-Guiraud A, Nikovics K, Morin H, Hay A, Johansen IE, Tsiantis M, Laufs P: **A conserved molecular framework for compound leaf development**. *Science* 2008, **322**:1835-1839.
 32. Sharma B, Kramer EM: **Sub- and neofunctionalization of *APETALA3* paralogs have contributed to the evolution of novel floral organ identity in *Aquilegia* (columbine, Ranunculaceae)**. *New Phytol* 2013, **197**:949-957.
- Functional demonstration the subfunctionalization and neofunctionalization of the *AqAP3-1* and *AqAP3-2* paralogs. Data also suggest potential auto-regulatory or cross-regulatory interactions among the *AP3* paralogs.