UC Riverside

UC Riverside Previously Published Works

Title

Robust fitting of mixtures of factor analyzers using the trimmed likelihood estimator

Permalink

<https://escholarship.org/uc/item/24j2d57q>

Journal

Communications in Statistics: Simulation and Computation, 46(2)

ISSN

0361-0918

Authors

Yang, L
Xiang, S
Yao, W

Publication Date

2017-02-07

DOI

10.1080/03610918.2014.999088

Peer reviewed

Robust Fitting of Mixtures of Factor Analyzers Using the Trimmed Likelihood Estimator

Li Yang^{*}, Sijia Xiang[†] and Weixin Yao[‡]

Abstract

Mixtures of factor analyzers have been popularly used to cluster the high dimensional data. However, the traditional estimation method is based on the normality assumptions of random terms and thus is sensitive to outliers. In this article, we introduce a robust estimation procedure of mixtures of factor analyzers using the trimmed likelihood estimator (TLE). We use a simulation study and a real data application to demonstrate the robustness of the trimmed estimation procedure and compare it with the traditional normality based maximum likelihood estimate.

Key words: EM algorithm, Factor analysis, Mixture models, Robustness, Trimmed likelihood estimator.

1 Introduction

Factor analysis is a statistical dimension reduction technique for modeling the covariance structure of high dimensional data using a small number of latent variables (Ghahramani

^{*}Department of Statistics, Kansas State University. E-mail: liy@k-state.edu.

[†]Corresponding Author, School of Mathematics and Statistics Zhejiang University of Finance and Economics. E-mail: fxbxsj@gmail.com.

[‡]Department of Statistics, University of California, Riverside. E-mail: weixin.yao@ucr.edu.

and Hinton, 1997). It can be extended by allowing different local factor models in different regions of the input space. This results in a model which performs clustering and dimension reduction at the same time, and can be thought of as a reduced dimension mixture of Gaussians. Ghahramani and Hinton (1997) and Hinton et al. (1997) originally proposed mixtures of factor analyzers (MFA) model. They used this model to visualize high dimensional data in a lower dimensional space to explore the grouping structure. Tipping and Bishop (1997, 1999) and Bishop (1998) considered the related model of mixtures of principal component analysers for the same purpose. MFA model is in fact a nonlinear model which can be considered as a combination of traditional [factor analysis](#) (FA) model and the finite mixture models. Therefore, MFA model offers a way to overcome the linear limitation of the traditional FA model. In recent years, MFA model has received considerable interest. See, for example, Fokoué and Titterington (2003), Yung (1997), Dolan and VanderMaas (1998), and Arminger et al. (1999). McLachlan et al. (2003) discussed the application of mixtures of factor analyzers to density estimation and the clustering of high-dimensional data.

MFA has been traditionally fitted using the maximum likelihood estimator (MLE) based on the normality assumptions of the random terms. Ghahramani and Hinton (1997) introduced an exact Expectation-Maximization (EM) algorithm to compute the MLE of MFA. However, it is well known that the normal based MLE can be very sensitive to outliers. In fact, even a single outlier can make an enormous impact on the MLE, which in mixture models means that at least one of the component parameter estimates might be arbitrarily large.

In this article, a robust fitting of mixtures of factor analyzers is introduced based on the idea of trimmed likelihood estimator (TLE) (Neykov et al., 2007). The TLE is designed to fit the majority of the data, whereas the remaining data will be considered as outliers and thus will not be used for parameter estimation. We use a simulation

study and a real data application to demonstrate the robustness of the new estimation procedure and compare it with the traditional normality based maximum likelihood estimate.

The rest of the paper is organized as follows. In Section 2, we briefly introduce the EM algorithm for a **factor analysis** (FA) and the mixture of factor analyzers (MFA). Section 3 presents the robust fitting of the mixture of factor analyzers using the trimmed likelihood estimator (TLE). Simulation results and a real data application are presented in Section 4. A discussion section ends the article.

2 Mixtures of Factor Analyzers

2.1 Factor analysis

Let $\mathbf{y}_1, \dots, \mathbf{y}_n$ be a random sample of size n on a p -dimensional random vector. A typical **factor analysis** model is given by:

$$\mathbf{y}_i = \boldsymbol{\mu} + \Lambda \mathbf{z}_i + \mathbf{e}_i, i = 1, \dots, n, \quad (2.1)$$

where $\boldsymbol{\mu}$ is the mean of $\mathbf{y}_i$, $\mathbf{z}_i$ is a q -dimensional ($q < p$) vector of latent or unobservable variables called factors, and Λ ($p \times q$) is a factor loading matrix. The factors $\mathbf{z}_i$ are assumed to be i.i.d. $\mathcal{N}_q(\mathbf{0}, \mathbf{I}_q)$, independent of the errors $\mathbf{e}_i$, which are assumed to be i.i.d. $\mathcal{N}_p(\mathbf{0}, \Psi)$ with Ψ a diagonal matrix $\Psi = \text{diag}(\sigma_1^2, \dots, \sigma_p^2)$. The marginal density of $\mathbf{y}_i$ is then $\mathcal{N}_p(\boldsymbol{\mu}, \Lambda \Lambda^T + \Psi)$. For the purpose of classifying and reducing data, the traditional **factor analysis** is a useful tool for reducing a mass of information to an efficient description and grouping interdependent variables into descriptive categories. In statistics, it is a method used for explaining data, in particular, correlations between variables in multivariate observations.

The [factor analysis](#) model (2.1) can be fitted by maximizing the log-likelihood:

$$\ell(\boldsymbol{\theta}) = \sum_{i=1}^n \log\{(2\pi)^{p/2} |\Lambda\Lambda^T + \Psi|^{-1/2} \exp[-\frac{1}{2}(\mathbf{y}_i - \boldsymbol{\mu})^T(\Lambda\Lambda^T + \Psi)^{-1}(\mathbf{y}_i - \boldsymbol{\mu})]\},$$

with $\boldsymbol{\theta} = (\boldsymbol{\mu}^T, \Lambda^T, \Psi^T)^T$, which can be computed iteratively via the EM algorithm if $\mathbf{z}_i$ is considered the missing data.

E-step: Given the current estimator $\boldsymbol{\theta}^{(k)}$, calculate the following conditional expectation given the observed data $\mathbf{y}$:

$$\begin{aligned} \mathbf{a}_i^{(k)} &= E(\mathbf{z}_i | \mathbf{y}_i, \boldsymbol{\theta}^{(k)}) = \Lambda^{(k)T} (\Psi^{(k)} + \Lambda^{(k)} \Lambda^{(k)T})^{-1} \mathbf{y}_i, \\ \mathbf{b}_i^{(k)} &= E(\mathbf{z}_i \mathbf{z}_i^T | \mathbf{y}_i, \boldsymbol{\theta}^{(k)}) = \mathbf{I} - \Lambda^{(k)T} (\Psi^{(k)} + \Lambda^{(k)} \Lambda^{(k)T})^{-1} \Lambda^{(k)} \\ &\quad + \{\Lambda^{(k)T} (\Psi^{(k)} + \Lambda^{(k)} \Lambda^{(k)T})^{-1} \mathbf{y}_i\} \{\Lambda^{(k)T} (\Psi^{(k)} + \Lambda^{(k)} \Lambda^{(k)T})^{-1} \mathbf{y}_i\}^T. \end{aligned}$$

M-step: Calculate

$$\begin{aligned} \boldsymbol{\mu}^{(k+1)} &= \sum_{i=1}^n (\mathbf{y}_i - \Lambda^{(k)} \mathbf{a}_i^{(k)}), \\ \Lambda^{(k+1)} &= \left\{ \sum_{i=1}^n \mathbf{y}_i \mathbf{a}_i^{(k)T} \right\} \left\{ \sum_{i=1}^n \mathbf{b}_i^{(k)} \right\}^{-1}, \\ \Psi^{(k+1)} &= \frac{1}{n} \text{diag} \left\{ \sum_{i=1}^n (\mathbf{y}_i \mathbf{y}_i^T - \Lambda^{(k+1)} \mathbf{a}_i^{(k)} \mathbf{a}_i^{(k)T}) \right\}. \end{aligned}$$

2.2 Mixtures of factor analyzers

Although the [factor analysis](#) model (2.1) provides a global linear model for the presentation of the data in a lower-dimensional subspace, its application is limited when the data is not homogenous. The mixture of factor analyzers model (MFA), which allows different local factor models in different regions of the input space, is a natural extension of the [factor analysis](#). Assume we have a mixture of m factor analyzers with mixing

71 proportion π_j , $j = 1, \dots, m$. The marginal density of $\mathbf{y}$ is given by:

$$f(\mathbf{y}; \boldsymbol{\theta}) = \sum_{j=1}^m \pi_j \mathcal{N}_p(\mathbf{y}; \boldsymbol{\mu}_j, \Lambda_j \Lambda_j^T + \Psi), \quad (2.2)$$

where $\boldsymbol{\theta} = (\boldsymbol{\pi}^T, \boldsymbol{\mu}^T, \Lambda^T, \Psi^T)^T$, $\boldsymbol{\pi} = (\pi_1, \dots, \pi_{m-1})^T$, $\boldsymbol{\mu} = (\boldsymbol{\mu}_1^T, \dots, \boldsymbol{\mu}_m^T)^T$, $\Lambda = (\Lambda_1^T, \dots, \Lambda_m^T)^T$. Here, $\boldsymbol{\mu}_j$ is the mean of the j^{th} component, Λ_j is the factor loading matrix of the j^{th} component, and Ψ is the diagonal matrix of the error terms. It will be useful in the estimation equations to have a definition of the mixture factor analyzers in terms of conditional densities. For the j^{th} component, the conditional density function is:

$$f_j(\mathbf{y}|\mathbf{z}) = \mathcal{N}_p(\mathbf{y}; \boldsymbol{\mu}_j + \Lambda_j \mathbf{z}, \Psi).$$

Within each component of the mixture, we have the following joint density of $\mathbf{y}$ and $\mathbf{z}$:

$$\begin{bmatrix} \mathbf{y} \\ \mathbf{z} \end{bmatrix} \sim \mathcal{N}_{p+q} \left(\begin{bmatrix} \boldsymbol{\mu}_j \\ 0 \end{bmatrix}, \begin{bmatrix} \Lambda_j \Lambda_j^T + \Psi & \Lambda_j \\ \Lambda_j^T & \mathbf{I}_q \end{bmatrix} \right).$$

Similar to the factor analysis, the mixture of factor analyzers can be estimated by maximizing the following likelihood:

$$\ell(\boldsymbol{\theta}) = \sum_{i=1}^n \log \sum_{j=1}^m \pi_j \left[(2\pi)^{p/2} |\Lambda_j \Lambda_j^T + \Psi|^{-1/2} \exp \left\{ -\frac{1}{2} (\mathbf{y}_i - \boldsymbol{\mu}_j)^T (\Lambda_j \Lambda_j^T + \Psi)^{-1} (\mathbf{y}_i - \boldsymbol{\mu}_j) \right\} \right]. \quad (2.3)$$

72 However, there is no explicit solution for the above maximizer. Ghahramani and Hinton
73 (1997) introduced an EM algorithm to maximize (2.3). More specifically, let ω_{ij} be an

74 indicator variable indicating which component $\mathbf{y}_i$ comes from. That is,

$$\omega_{ij} = \begin{cases} 1, & \text{if } \mathbf{y}_i \text{ is from } j^{\text{th}} \text{ component,} \\ 0, & \text{otherwise.} \end{cases} \quad (2.4)$$

Then the complete log-likelihood for $\{(\mathbf{y}_i, \mathbf{z}_i, \omega_{ij}), i = 1, \dots, n, j = 1, \dots, m\}$ is

$$\ell_c(\boldsymbol{\theta}) = \sum_{i=1}^n \log \prod_{j=1}^m \pi_j^{\omega_{ij}} \left[(2\pi)^{p/2} |\Psi|^{-1/2} \exp \left\{ -\frac{1}{2} (\mathbf{y}_i - \boldsymbol{\mu}_j - \Lambda_j \mathbf{z}_i)^T \Psi^{-1} (\mathbf{y}_i - \boldsymbol{\mu}_j - \Lambda_j \mathbf{z}_i) \right\} \right]^{\omega_{ij}}.$$

The EM algorithm iterates between E-step, which computes the expected complete log-likelihood given current parameter estimates, and M-step, which maximizes the expected completed log-likelihood calculated in the E-step. We summarize the EM algorithm to maximize (2.3) as follows:

E-step: Given the current estimator $\boldsymbol{\theta}^{(k)}$, calculate the following conditional expectation given the observed data $\mathbf{y}$:

$$E(\omega_{ij} | \mathbf{y}_i, \boldsymbol{\theta}^{(k)}) = \frac{\pi_j^{(k)} \mathcal{N}_p(\mathbf{y}_i; \boldsymbol{\mu}_j^{(k)}, \Lambda_j^{(k)} \Lambda_j^{(k)T} + \Psi^{(k)})}{\sum_{j=1}^m \pi_j^{(k)} \mathcal{N}_p(\mathbf{y}_i; \boldsymbol{\mu}_j^{(k)}, \Lambda_j^{(k)} \Lambda_j^{(k)T} + \Psi^{(k)})} = p_{ij}^{(k)},$$

$$\mathbf{a}_{ij}^{(k)} = E(\mathbf{z}_i | \mathbf{y}_i, \omega_{ij} = 1, \boldsymbol{\theta}^{(k)}) = \Gamma_j^{(k)} (\mathbf{y}_i - \boldsymbol{\mu}_j^{(k)}),$$

$$\mathbf{b}_{ij}^{(k)} = E(\mathbf{z}_i \mathbf{z}_i^T | \mathbf{y}_i, \omega_{ij} = 1, \boldsymbol{\theta}^{(k)}) = I - \Gamma_j^{(k)} \Lambda_j^{(k)} + \Gamma_j^{(k)} (\mathbf{y}_i - \boldsymbol{\mu}_j^{(k)}) \{ \Gamma_j^{(k)} (\mathbf{y}_i - \boldsymbol{\mu}_j^{(k)}) \}^T,$$

75 where $\Gamma_j = \Lambda_j^T (\Psi + \Lambda_j \Lambda_j^T)^{-1}$.

76

M-step: Calculate

$$\pi_j^{(k+1)} = \frac{1}{n} \sum_{i=1}^n p_{ij}^{(k)},$$

$$\boldsymbol{\mu}_j^{(k+1)} = \left\{ \sum_{i=1}^n p_{ij}^{(k)} (\mathbf{y}_i - \Lambda_j^{(k)} \mathbf{a}_{ij}^{(k)}) \right\} \left\{ \sum_{i=1}^n p_{ij}^{(k)} \right\}^{-1},$$

$$\Lambda_j^{(k+1)} = \left\{ \sum_{i=1}^n p_{ij}^{(k)} (\mathbf{y}_i - \boldsymbol{\mu}_j^{(k+1)}) \mathbf{a}_{ij}^{(k)T} \right\} \left\{ \sum_{i=1}^n p_{ij}^{(k)} \mathbf{b}_{ij}^{(k)} \right\}^{-1},$$

$$\Psi^{(k+1)} = \frac{1}{n} \text{diag} \left\{ \sum_{i=1}^n \sum_{j=1}^m p_{ij}^{(k)} (\mathbf{y}_i - \boldsymbol{\mu}_j^{(k+1)} - \Lambda_j^{(k+1)} \mathbf{a}_{ij}^{(k)}) (\mathbf{y}_i - \boldsymbol{\mu}_j^{(k+1)})^T \right\}.$$

3 Robust Fitting of Mixtures of Factor Analyzers Using the Trimmed Likelihood Estimator

The maximum likelihood estimator introduced in Section 2 is easy to implement, but very sensitive to outliers. Even a single outlier can make an enormous impact on the MLE, and make at least one of the component parameters to be arbitrarily large. To overcome this, McLachlan et al. (2007), Andrews et al. (2011), and Baek and McLachlan (2011) proposed mixtures of t -factor analyzers by assuming multivariate t -distributions for component errors and factor distributions. In this section, we apply the idea of trimmed likelihood estimator (TLE), proposed by Neykov et al. (2007), to fit the the mixtures of factor analyzers in a robust way. Compared to the proposed method based on TLE, the mixture of t -distributions has a very small breakdown point and is not robust when the outliers are extreme (Hennig, 2004; Yao et al., 2014).

Suppose a number k ($k \leq n$) of n observations are regular observations in the data, and the remaining $n - k$ observations may be gross or outliers. The basic idea of TLE is removing the $n - k$ observations which do not follow the model, and using only the k observations to fit the model. The combinatorial nature of the TLE can be expressed as:

$$\max_{I \in I_k} \max_{\boldsymbol{\theta}} \sum_{i \in I} \log f(\mathbf{y}_i; \boldsymbol{\theta}),$$

where I_k is the set of all k -subsets of $(1, \dots, n)$ and $f(\mathbf{y}; \boldsymbol{\theta})$ is defined in (2.2). The fact that all possible $\binom{n}{k}$ partitions of the data have to be fitted by the MLE makes

the estimation procedure very computational expensive. To find an approximate TLE solution for large data sets, an algorithm called FAST-TLE was developed by Neykov and Müller (2003). The basic idea behind FAST-TLE algorithm contains two steps: a trial step followed by a refinement step.

(i) Trial step: Randomly select a subsample of size k^* from the data sample and then fit the model to that subsample to get a trial maximum likelihood estimate (MLE).

(ii) Refinement step: This step is based on the so-called concentration procedure.

(a) Starting with the trial MLE, find a combination with the k smallest negative log-likelihoods based on the current estimate.

(b) Obtain an improved estimator by fitting the model to these k cases.

(c) Repeat (a) and (b) until convergence.

At the end of this step, the solution with the largest trimmed likelihood is stored.

This value may not be guaranteed to be the global optimal but would be a close approximation to it.

The choice of trial size k^* and refinement subsample size k are related to the breakdown point (BP). The breakdown point (i.e., the smallest fraction of contamination that can cause the estimator to take arbitrary large values) of TLE was studied by using d -fullness technique. Vandev and Neykov (1993) determined the value of d for the mixtures of normals to be $m(p + 1)$. It was proved that if $\log f(y)$ is d -full, then the BP of TLE is not less than $\frac{1}{n} \min\{n - m + 1, m - d + 1\}$ (Neykov and Müller, 2003). The trial subsample size k^* should be greater than or equal to d for the existence of MLE. The choice of k can be any number within $[d, n]$. When $k = \lfloor (n + d + 1)/2 \rfloor$, the BP of the TLE is maximized (Neykov and Müller, 2003). If the expected percentage of outliers α in the data is a known priori, a recommended choice of k is $\lfloor n(1 - \alpha) \rfloor$ which can increase the efficiency of the TLE.

116 The process of TLE applied particularly to the mixtures of factor analyzers can be
 117 performed as follows:

118 **Input:** A trial subset with sample size equals to k^* and initial parameters $\boldsymbol{\theta}^{(0)} =$
 119 $(\boldsymbol{\pi}^{(0)T}, \boldsymbol{\mu}^{(0)T}, \Lambda^{(0)T}, \Psi^{(0)T})^T$.

120 **Output:** A subset of size k which has the k smallest negative log-likelihoods.

121 At the $(l + 1)^{th}$ iteration:

122 **E-step:** Compute the expectation of component indicators ω_{ij} , latent variable $\mathbf{z}$, and
 123 $\mathbf{z}\mathbf{z}^T$ based on the current subsample of size k .

124 **M-step:** Maximize the complete log-likelihood of subsample of size k with respect to
 125 each unknown parameter and thus get a new parameter

$$\boldsymbol{\theta}^{(l+1)} = (\boldsymbol{\pi}^{(l+1)T}, \boldsymbol{\mu}^{(l+1)T}, \Lambda^{(l+1)T}, \Psi^{(l+1)T})^T.$$

126 **T-step:** Define a new subsample of size k which has the k smallest negative log-
 127 likelihoods with the new parameter $\boldsymbol{\theta}^{(l+1)}$.

128 Repeat **EMT** steps until convergence.

129 4 Simulation Study and Real Data Application

130 4.1 Simulation study

131 In this section, we use a simulation study to assess the performance of the MLE and the
 132 TLE to the mixtures of factor analyzers. For TLE, 20 randomly generated initial values
 133 are used and TLE reports the estimate whose log-likelihood is the biggest. True value
 134 (T) is also used as initial value for MLE and TLE. For the 20 initial values, we first
 135 use the R code “hc” from the R package “mclust” to cluster the randomly generated
 136 subsets of the data and then use the R code “factanal” from the R package “stats” to

do factor analysis for each cluster. The trimming proportion α is set to be 5% and thus $k = \lfloor n(1 - \alpha) \rfloor$ is used for TLE in all examples. We will discuss how to choose α data adaptively in Section 5.

A two-component mixture of factor analyzers are considered in the simulation:

$$f(\mathbf{y}) = \sum_{j=1}^2 \pi_j \mathcal{N}_p(\mathbf{y}; \boldsymbol{\mu}_j, \Lambda_j \Lambda_j^T + \Psi),$$

where the mixing proportions are $\pi_1 = 0.4$ and $\pi_2 = 0.6$. The means $\boldsymbol{\mu}_1$ and $\boldsymbol{\mu}_2$ are $p \times 1$ vectors with all the elements equal to 0 and 5, respectively, and the factor loading matrices Λ_1 and Λ_2 are $p \times 2$ matrices with all the elements equal to 0.5 and 1, respectively. That is,

$$\boldsymbol{\mu}_1 = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}_{p \times 1}, \boldsymbol{\mu}_2 = \begin{pmatrix} 5 \\ \vdots \\ 5 \end{pmatrix}_{p \times 1},$$

$$\Lambda_1 = \begin{pmatrix} 0.5 & 0.5 \\ \vdots & \vdots \\ 0.5 & 0.5 \end{pmatrix}_{p \times 2}, \Lambda_2 = \begin{pmatrix} 1 & 1 \\ \vdots & \vdots \\ 1 & 1 \end{pmatrix}_{p \times 2}.$$

We consider $p = 10, 20$, and 30 . Sample sizes of $n = 200$ and $n = 400$ are conducted over 200 repetitions. To assess the robustness of the estimators, only $(1 - \alpha_0) \times 100\%$ of the observations are generated from the above model with $\alpha_0 = 0, 0.01, 0.03$, and 0.05 , and the remaining $\alpha_0 \times 100\%$ of the data is generated randomly from $U(20, 30)$. The simulation was done through R on a personal laptop with an i7-3610QM CPU and 8GB of RAM. The computation time of the new algorithm (with 20 random initial values) is 45 seconds for $n = 200$ and 61 seconds for $n = 400$.

The performance of the estimates is measured by the miss-classification probability

(MCP), which is defined to be the proportion of observations that are misclassified:

$$\text{MCP} = 1 - \left\{ \sum_{i=1}^n \sum_{j=1}^2 \omega_{ij} I_{p_{ij} > 0.5} \right\} / n,$$

where ω_{ij} , defined in (2.4), indicates which component $\mathbf{y}_i$ comes from, and p_{ij} is the classification probability calculated by

$$p_{ij} = \frac{\hat{\pi}_j \mathcal{N}_p(\mathbf{y}_i; \hat{\boldsymbol{\mu}}_j, \hat{\Lambda}_j \hat{\Lambda}_j^T + \hat{\Psi})}{\sum_{j=1}^2 \hat{\pi}_j \mathcal{N}_p(\mathbf{y}_i; \hat{\boldsymbol{\mu}}_j, \hat{\Lambda}_j \hat{\Lambda}_j^T + \hat{\Psi})}, i = 1, \dots, n, j = 1, 2.$$

Note that for mixture models there are well known label switching issues (Celeux, et al., 2000; Stephens, 2000; Jasra et al., 2005; Yao and Lindsay, 2009; Grün and Leisch, 2009; Yao, 2012a, 2012b). In our simulations, the labels are found by minimizing the MCP.

Tables 1 and 2 report the means and standard deviations of MCP for $n = 200$ and 400 , respectively. Based on the above tables, both TLE(T) and TLE(I) have smaller MCP than MLE for all three p values and both $n = 200$ and $n = 400$. In Tables 3 and 4, we also report the means and standard deviations of the Euclidean distance between the estimates $\hat{\pi}_1$, $\hat{\boldsymbol{\mu}}_1$, and $\hat{\boldsymbol{\mu}}_2$ and their corresponding true values based on 200 repetitions. From the tables, we can see that the TLEs with both true initial values and random initial values have better performance than the MLE when there are outliers, especially for $\boldsymbol{\mu}_2$ and π_1 . The TLEs with randomly generated initial values work almost the same as those with true initial values. In addition, the TLE still works well when the trimming proportion is larger than the proportion of outliers. Furthermore, when there are no outliers ($\alpha = 0$), TLE has comparable performance to the traditional MLE.

Table 1: Average (Std) of MCP, with $n = 200$.

Dimension	Method	$\alpha = 0$	$\alpha = 0.01$	$\alpha = 0.03$	$\alpha = 0.05$
$p = 10$	MLE	0.984(0.012)	0.117(0.032)	0.103(0.031)	0.089(0.029)
	TLE(T)	0.984(0.011)	0.017(0.010)	0.018(0.012)	0.017(0.012)
	TLE(I)	0.982(0.012)	0.019(0.011)	0.020(0.013)	0.020(0.014)
$p = 20$	MLE	0.982(0.012)	0.089(0.030)	0.097(0.029)	0.140(0.029)
	TLE(T)	0.982(0.012)	0.019(0.013)	0.020(0.013)	0.067(0.010)
	TLE(I)	0.980(0.014)	0.022(0.015)	0.022(0.014)	0.070(0.013)
$p = 30$	MLE	0.151(0.354)	0.076(0.025)	0.105(0.031)	0.100(0.032)
	TLE(T)	0.151(0.353)	0.026(0.014)	0.033(0.018)	0.021(0.012)
	TLE(I)	0.145(0.347)	0.029(0.021)	0.040(0.036)	0.026(0.029)

4.2 Real data application

In this example, we consider applying both MLE and TLE of the mixture of factor analyzers to the wine data, which is available at the Machine Learning Repository of the University of California. The data set contains the results of chemical analysis of wines grown in the same region in Italy, but derived from three different cultivars. Therefore, a three component mixture model is suitable to fit the data if we do not use the cultivars of the wines. The analysis determined the quantities of $p = 13$ constituents found in each of $n = 178$ wines. Both MLE and TLE of the mixture of factor analyzers were fitted to this data set. Similar to the simulation study, the trimming proportion is set to be 0.05 for TLE.

Based on McLachlan and Peel (2000), the miss-classification rate is smallest for $q = 2$ and 3. In our analysis, $q = 2$ is used as our reduced dimension. Figure 1 shows the estimated posterior means of the $q = 2$ factors following a three-component mixture of factor analyzers of the wine data, which is actually the $\mathbf{a}_{ij}$ calculated from E-step. These posterior means have been plotted with their true group labels corresponding to

Table 2: Average (Std) of MCP, with $n = 400$.

Dimension	Method	$\alpha = 0$	$\alpha = 0.01$	$\alpha = 0.03$	$\alpha = 0.05$
$p = 10$	MLE	0.986(0.006)	0.125(0.024)	0.123(0.020)	0.130(0.019)
	TLE(T)	0.986(0.006)	0.025(0.007)	0.044(0.006)	0.064(0.006)
	TLE(I)	0.986(0.006)	0.026(0.008)	0.044(0.006)	0.064(0.006)
$p = 20$	MLE	0.986(0.006)	0.110(0.021)	0.123(0.022)	0.131(0.019)
	TLE(T)	0.986(0.006)	0.025(0.007)	0.044(0.006)	0.065(0.007)
	TLE(I)	0.986(0.006)	0.025(0.007)	0.045(0.006)	0.065(0.007)
$p = 30$	MLE	0.984(0.008)	0.096(0.021)	0.124(0.020)	0.091(0.022)
	TLE(T)	0.984(0.009)	0.025(0.006)	0.047(0.008)	0.016(0.007)
	TLE(I)	0.984(0.009)	0.025(0.007)	0.047(0.008)	0.017(0.008)

the three different cultivars displayed. From Figure 1 we can see that mixtures of factor analyzers have been useful here in exploring the grouping structure of the data in a much reduced dimension.

To assess the robustness of the two estimation methods, we also consider the contaminated data by adding 1% and 3% outliers from $U(9, 11)$. Table 5 displays the estimated means $\boldsymbol{\mu}_1$, $\boldsymbol{\mu}_2$, and $\boldsymbol{\mu}_3$ via MLE and TLE when the proportion of outliers are $\alpha_0 = 0$, 0.01, and 0.03, and Table 6 displays the estimated component proportions π_1 and π_2 . The true parameter values are calculated by using true classification labels based on the cultivars of the wines. From both tables, we see that when there are no outliers ($\alpha_0 = 0$), both MLE and TLE can provide comparatively good estimators. When the data is contaminated, however, TLE performs much better than MLE. As the proportion of outliers gets higher, MLE departs further away from the original MLE, while TLE does not change much when the outliers are added to the data.

Figure 1: Wine data: Plot of the estimated posterior means of the $q = 2$ factors (Δ , $\circ$, and $*$ denote true component membership).

5 Discussion

Mixtures of factor analyzers have been popularly used to do dimension reduction and model based clustering for high dimensional data. In this article, we investigate a robust estimation procedure of the mixtures of factor analyzers based on the TLE proposed by Neykov et al. (2007). The simulation study and real data analysis demonstrated the effectiveness of the TLE based robust estimation procedure.

It is well known that the scale estimate by TLE is biased for univariate data. A scale factor is usually needed to make the scale estimate an unbiased consistent estimator. Based on our limited empirical experience, the TLE based covariance estimate for mixtures of factor analyzers are also biased. However, it requires more theoretical studies whether a scale or vector factor could make the TLE based covariance estimator unbiased and consistent.

In our examples, we have fixed the trimming proportion to be 0.05 for TLE. It works well whenever the true proportions of outliers are no more than 5%. However, it requires

203 more research to find a data adaptive optimal or conservative trimming proportion for
 204 TLE in practice. Neykov et al. (2007) recommended a graphical tool to choose the
 205 trimming proportion in their examples. However, based on our limited empirical expe-
 206 rience, such graphical tool was not very successful in choosing the trimming proportion
 207 for mixtures of factor analyzers. There have been many methods proposed for choosing
 208 the trimming proportion for TLE in the non-mixture context. For example, Jurečková
 209 et al. (1994) studied the problem of choosing the trimming proportion for a trimmed L -
 210 estimator of location, and recommended the L -estimators with smooth weight functions.
 211 For the trimmed mean in the location modeling and for the trimmed least-squares esti-
 212 mator in the linear regression model, Dodge and Jurečková (1997) proposed a partially
 213 adaptive estimator of the trimming proportion based on a rank-based decision proce-
 214 dure. Clark and Schubert (2010) studied an adaptive trimmed likelihood estimator of
 215 regression, whose algorithm tends to expose the outliers automatically and provide the
 216 estimators with the outliers removed. It will be interesting to know whether we can
 217 extend the foregoing methods to adaptively choose the trimming proportion for TLE in
 218 the mixture context.

219 References

- 220 Andrews, J. L., McNicholas, P. D. and Subedi, S. (2011). Model-based classification
 221 via mixtures of multivariate t -distribution. *Computational Statistics & Data Analysis*,
 222 55(1), 520-529.
- 223 Arminger, G., Stein, P. and Wittenberg, J. (1999). Mixtures of conditional mean and
 224 covariance structure models. *Psychometrika*, 65, 475-494.
- 225 Baek, J. and McLachlan, G. J. (2011). Mixtures of common t -factor analyzers for clus-
 226 tering high-dimensional microarray data. *Oringnal Paper*, 27(9), 1269-1276.

- 227 Bishop, C. M. (1998). Latent variable models. *Jordan, M.I. (Ed.), Learning in Graphical*
228 *Models*, pp, 371-403.
- 229 Clark, B. R. and Schubert, D. (2010). Adaptive trimmed likelihood estimation in regres-
230 sion. *Probability and Statistics*, 30, 203-219.
- 231 Celeux, G., Hurn, M. and Robert, C. P. (2000). Computational and inferential difficulties
232 with mixture posterior distributions. *Journal of the American Statistical Association*,
233 95, 957-970.
- 234 Dodge, Y. and Jurečková J. (1997). Adaptive choice of proportion in trimmed least-
235 squares estimation. *Statistics & Probability Letters*, 33(2), 167-176.
- 236 Dolan, C. and Van der Mass, H. (1998), Fitting multivariate normal finite mixtures
237 subject to structural equation modelling. *Psychometrika*, 63, 227-253.
- 238 Fokoué, E. and Titterington, D. M. (2003). Mixtures of factor analysers: Bayesian esti-
239 mation and inference by stochastic simulation. *Machine Learning*, 50, 73-94.
- 240 Ghahramani, Z. and Hinton, G. E. (1997). The EM algorithm for mixture of factor
241 analyzers. *University of Toronto Technical Report*, CRG-TR-96-1.
- 242 Grün, B. and Leisch, F. (2009). Dealing with label switching in mixture models under
243 genuine multimodality. *Journal of Multivariate Analysis*, 100, 851-861.
- 244 Hennig, C. (2004). Breakdown points for maximum likelihood-estimators of location-
245 scale mixtures. *Annals of Statistics*, 32, 1313-1340.
- 246 Hinton, G. E., Dayan, P. and Revow, M. (1997). Modeling the manifolds of images of
247 handwritten digits. *IEEE Transaction, Neural Networks*, 8, 65-73.

248 Jasra, A., Holmes, C. C., and Stephens, D. A. (2005). Markov chain Monte Carlo meth-
249 ods and the label switching problem in Bayesian mixture modeling. *Statistical Science*,
250 20, 50-67.

251 Jurečková, J., Koenker, R. and Welsh, A. H. (1994). Adaptive choice of trimming pro-
252 portions. *Annals of the Institute of Statistical Mathematics*, 46(4), 737-755.

253 McLachlan, G. J., Bean, R. W. and Ben-Tovim Jones, L. (2007). Extension of the mix-
254 ture of factor analyzers model to incorporate the multivariate t -distribution. *Com-*
255 *putational Statistics & Data Analysis*, 51, 5327-5338.

256 McLachlan, G. J. and Peel, D. (2000). Finite mixture models. John Wiley & Sons, Inc,
257 New York.

258 McLachlan, G. J., Peel, D. and Bean, R. W. (2003). Modelling high-dimensional data by
259 mixtures of factor analyzers. *Computational Statistics & Data Analysis*, 41, 379-388.

260 Neykov, N., Filzmoser, P., Dimova, R. and Neytchev, P. (2007). Robust fitting of mix-
261 tures using the trimmed likelihood estimator. *Computational Statistics & Data Anal-*
262 *ysis*, 52, 299-308.

263 Neykov, N. M. and Müller, C. H. (2003). Breakdown points of the trimmed likelihood
264 and related estimators in generalized linear models. *Developments in Robust Statistics*,
265 pp, 277-286.

266 Neal, R. M. and Hinton, G. E. (1998). Learning in graphical models. MIT Press.

267 Stephens, M. (2000). Dealing with label switching in mixture models. *Journal of Royal*
268 *Statistical Society, Ser. B*, 62, 795-809.

269 Tipping, M. E. and Bishop, C. M. (1997). Mixtures of probabilistic principal component

- 270 analysers. *Technical Report*, No. NCRG/97/003, Neural Computing Research Group,
271 Aston University, Birmingham
- 272 Tipping, M. E. and Bishop, C. M. (1999). Mixtures of probabilistic principal component
273 analysers. *Neural Computation*, 11, 443-482.
- 274 Vandev, D. L. and Neykov, N. M. (1993). Robust maximum likelihood in the Gaussian
275 case. In Morgenthaler, S., Ronchetti, E. and Stahel, W. A. (eds), *New Directions in*
276 *Data Analysis and Robustness*, 257-264. Basel, Birkhauser Verlag.
- 277 Yao, W. (2012a). Model based labeling for mixture models. *Statistics and Computing*,
278 22, 337-347.
- 279 Yao, W. (2012b). Bayesian mixture labeling and clustering. *Communications in Statistics*
280 - *Theory and Methods*, 41, 403-421.
- 281 Yao, W., Wei, Y., and Yu, C. (2014). Robust Mixture Regression Using T-Distribution.
282 *Computational Statistics and Data Analysis*, 71, 116-127.
- 283 Yao, W. and Lindsay, B. G. (2009). Bayesian mixture labeling by highest posterior
284 density. *Journal of American Statistical Association*, 104, 758-767.
- 285 Yung, Y. F. (1997). Finite mixtures in confirmatory factor analysis models. *Psychome-*
286 *trika*, 62, 297-330.

Table 3: Average (Std) of Euclidean distance, with $n = 200$.

Dimension	Method	$\alpha = 0$	$\alpha = 0.01$	$\alpha = 0.03$	$\alpha = 0.05$
$p = 10$	MLE	μ_1 : 0.023(0.026)	0.051(0.032)	0.042(0.038)	0.044(0.033)
		μ_2 : 0.025(0.034)	1.359(0.469)	2.979(1.505)	6.368(0.825)
		π_1 : 0.001(0.002)	0.021(0.012)	0.021(0.016)	0.030(0.016)
	TLE(T)	μ_1 : 0.024(0.020)	0.023(0.020)	0.021(0.021)	0.025(0.014)
		μ_2 : 0.028(0.030)	0.030(0.035)	0.024(0.028)	0.032(0.035)
		π_1 : 0.001(0.002)	0.001(0.002)	0.002(0.003)	0.003(0.004)
	TLE(I)	μ_1 : 0.026(0.022)	0.025(0.021)	0.021(0.022)	0.030(0.030)
		μ_2 : 0.030(0.034)	0.033(0.038)	0.031(0.066)	0.036(0.038)
		π_1 : 0.001(0.002)	0.001(0.002)	0.002(0.003)	0.003(0.004)
$p = 20$	MLE	μ_1 : 0.022(0.015)	0.046(0.091)	0.042(0.024)	0.042(0.027)
		μ_2 : 0.027(0.029)	0.849(0.298)	2.792(0.479)	5.449(0.690)
		π_1 : 0.001(0.001)	0.013(0.009)	0.020(0.012)	0.028(0.014)
	TLE(T)	μ_1 : 0.023(0.015)	0.026(0.024)	0.023(0.018)	0.025(0.016)
		μ_2 : 0.029(0.030)	0.036(0.046)	0.030(0.030)	0.031(0.036)
		π_1 : 0.001(0.002)	0.001(0.002)	0.002(0.003)	0.003(0.003)
	TLE(I)	μ_1 : 0.024(0.016)	0.029(0.025)	0.027(0.025)	0.029(0.023)
		μ_2 : 0.039(0.057)	0.047(0.068)	0.037(0.037)	0.038(0.040)
		π_1 : 0.001(0.002)	0.002(0.003)	0.002(0.003)	0.003(0.003)
$p = 30$	MLE	μ_1 : 0.004(0.010)	0.034(0.022)	0.040(0.024)	0.018(0.032)
		μ_2 : 0.005(0.021)	0.528(0.213)	2.248(0.392)	1.551(2.216)
		π_1 : 0.001(0.001)	0.008(0.008)	0.019(0.012)	0.010(0.016)
	TLE(T)	μ_1 : 0.004(0.009)	0.024(0.015)	0.024(0.014)	0.010(0.018)
		μ_2 : 0.009(0.043)	0.027(0.033)	0.028(0.031)	0.008(0.020)
		π_1 : 0.001(0.001)	0.002(0.002)	0.002(0.003)	0.001(0.003)
	TLE(I)	μ_1 : 0.004(0.010)	0.047(0.201)	0.079(0.465)	0.044(0.401)
		μ_2 : 0.012(0.063)	0.037(0.048)	0.039(0.049)	0.013(0.036)
		π_1 : 0.001(0.001)	0.002(0.002)	0.003(0.007)	0.001(0.005)

Table 4: Average (Std) of Euclidean distance, with $n = 400$.

Dimension	Method	$\alpha = 0$	$\alpha = 0.01$	$\alpha = 0.03$	$\alpha = 0.05$
$p = 10$	MLE	μ_1 : 0.010(0.007)	0.031(0.020)	0.023(0.015)	0.020(0.013)
		μ_2 : 0.013(0.018)	1.566(0.289)	3.757(0.364)	6.630(0.595)
		π_1 : 0.001(0.001)	0.025(0.011)	0.026(0.010)	0.030(0.011)
	TLE(T)	μ_1 : 0.011(0.008)	0.012(0.009)	0.012(0.009)	0.012(0.008)
		μ_2 : 0.015(0.021)	0.016(0.017)	0.013(0.014)	0.012(0.012)
		π_1 : 0.001(0.001)	0.001(0.001)	0.001(0.001)	0.002(0.002)
	TLE(I)	μ_1 : 0.011(0.009)	0.012(0.009)	0.013(0.009)	0.012(0.009)
		μ_2 : 0.016(0.022)	0.017(0.019)	0.015(0.016)	0.014(0.014)
		π_1 : 0.001(0.001)	0.001(0.001)	0.001(0.001)	0.002(0.002)
$p = 20$	MLE	μ_1 : 0.011(0.006)	0.025(0.013)	0.021(0.012)	0.020(0.014)
		μ_2 : 0.013(0.013)	1.056(0.235)	2.963(0.324)	5.713(0.511)
		π_1 : 0.001(0.001)	0.018(0.008)	0.024(0.010)	0.028(0.010)
	TLE(T)	μ_1 : 0.011(0.007)	0.011(0.006)	0.012(0.008)	0.012(0.008)
		μ_2 : 0.014(0.016)	0.016(0.016)	0.013(0.013)	0.013(0.015)
		π_1 : 0.001(0.001)	0.001(0.001)	0.001(0.001)	0.002(0.002)
	TLE(I)	μ_1 : 0.012(0.008)	0.011(0.006)	0.012(0.008)	0.013(0.014)
		μ_2 : 0.016(0.020)	0.018(0.017)	0.014(0.014)	0.015(0.016)
		π_1 : 0.001(0.001)	0.001(0.001)	0.001(0.001)	0.002(0.002)
$p = 30$	MLE	μ_1 : 0.011(0.008)	0.021(0.013)	0.022(0.014)	0.016(0.014)
		μ_2 : 0.014(0.015)	0.715(0.171)	2.503(0.316)	3.616(2.238)
		π_1 : 0.001(0.001)	0.013(0.008)	0.022(0.010)	0.021(0.016)
	TLE(T)	μ_1 : 0.012(0.009)	0.011(0.007)	0.012(0.007)	0.009(0.008)
		μ_2 : 0.018(0.024)	0.014(0.013)	0.017(0.019)	0.009(0.011)
		π_1 : 0.001(0.001)	0.001(0.001)	0.001(0.002)	0.001(0.002)
	TLE(I)	μ_1 : 0.013(0.009)	0.012(0.008)	0.012(0.007)	0.009(0.008)
		μ_2 : 0.019(0.023)	0.016(0.015)	0.019(0.019)	0.010(0.013)
		π_1 : 0.001(0.001)	0.001(0.001)	0.001(0.002)	0.001(0.002)

Table 5: Wine data: Estimated Means with $\alpha_0 = 0, 0.01$, and 0.03 .

	True	$\alpha_0 = 0$		$\alpha_0 = 0.01$		$\alpha_0 = 0.03$	
		MLE	TLE	MLE	TLE	MLE	TLE
μ_1	13.74	13.66	13.74	13.44	13.74	12.34	13.73
	2.01	1.99	2.01	1.61	2.02	0.21	1.99
	2.46	2.47	2.46	2.09	2.46	0.79	2.43
	17.04	17.49	17.05	16.42	17.18	15.77	17.01
	106.34	107.87	106.30	105.67	106.04	105.95	105.34
	2.84	2.85	2.84	2.50	2.84	1.29	2.84
	2.98	3.00	2.98	2.69	2.98	2.11	2.96
	0.29	0.29	0.29	-0.03	0.29	-1.25	0.28
	1.90	1.92	1.90	1.53	1.90	0.66	1.87
	5.53	5.44	5.52	5.29	5.53	7.09	5.50
	1.06	1.07	1.06	0.71	1.06	-0.40	1.06
	3.16	3.16	3.16	2.78	3.14	1.53	3.14
	1115.71	1097.23	1114.12	1144.08	1115.45	1284.31	1115.80
μ_2	12.28	12.28	12.30	12.34	12.32	12.92	12.30
	1.93	1.95	1.96	1.98	1.95	1.97	1.97
	2.24	2.22	2.25	2.26	2.24	2.33	2.24
	20.24	19.96	20.26	20.21	20.09	18.88	20.08
	94.55	91.86	90.09	94.98	90.07	99.06	91.30
	2.26	2.23	2.23	2.30	2.24	2.51	2.24
	2.08	2.04	2.06	2.14	2.05	2.48	2.07
	0.36	0.37	0.38	0.37	0.37	0.33	0.38
	1.63	1.60	1.55	1.64	1.53	1.75	1.59
	3.09	3.05	3.07	3.17	3.07	4.11	3.06
	1.06	1.05	1.06	1.05	1.05	1.06	1.05
	2.79	2.77	2.79	2.82	2.78	2.95	2.78
	519.51	502.67	496.14	534.54	496.23	777.10	498.36
μ_3	13.15	13.12	13.13	13.12	13.12	13.11	13.12
	3.33	3.31	3.37	3.30	3.30	3.27	3.29
	2.44	2.44	2.43	2.44	2.44	2.43	2.44
	21.42	21.42	21.34	21.42	21.41	21.33	21.41
	99.31	100.03	99.35	100.03	100.04	100.02	100.05
	1.68	1.68	1.65	1.68	1.67	1.68	1.67
	0.78	0.79	0.77	0.79	0.79	0.80	0.79
	0.45	0.44	0.45	0.44	0.44	0.44	0.44
	1.15	1.16	1.12	1.16	1.16	1.15	1.16
	7.40	7.29	7.27	7.28	7.27	7.25	7.25
	0.68	0.69	0.69	0.69	0.69	0.69	0.69
	1.68	1.70	1.68	1.70	1.70	1.69	1.70
	629.90	630.27	629.56	630.53	631.24	627.43	632.32

Table 6: Wine data: Estimated component proportions with $\alpha_0 = 0, 0.01$, and 0.03 .

	True	$\alpha_0 = 0$		$\alpha_0 = 0.01$		$\alpha_0 = 0.03$	
		MLE	TLE	MLE	TLE	MLE	TLE
π_1	0.3315	0.3516	0.3516	0.3049	0.3386	0.0331	0.3201
π_2	0.3989	0.3726	0.3726	0.4190	0.3697	0.6853	0.3869