

UCSF

UC San Francisco Previously Published Works

Title

Cardiovascular health metrics among South Asian adults in the United States: Prevalence and associations with subclinical atherosclerosis

Permalink

<https://escholarship.org/uc/item/25s3k6ns>

Journal

Preventive Medicine, 96

ISSN

0091-7435

Authors

Talegawkar, Sameera A
Jin, Yichen
Kandula, Namratha R
et al.

Publication Date

2017-03-01

DOI

10.1016/j.ypmed.2016.12.017

Peer reviewed



Published in final edited form as:

Prev Med. 2017 March ; 96: 79–84. doi:10.1016/j.ypmed.2016.12.017.

Cardiovascular health metrics among South Asian adults in the United States: Prevalence and associations with subclinical atherosclerosis

Sameera A. Talegawkar, PhD^a, Yichen Jin, MSPH^b, Namratha R. Kandula, MD, MPH^c, and Alka M. Kanaya, MD^d

^aDepartments of Exercise and Nutrition Sciences and Epidemiology and Biostatistics, and the Sumner M. Redstone Global Center for Prevention and Wellness and, Milken Institute School of Public Health, The George Washington University, 950 New Hampshire Ave, NW, 2nd Floor, Washington, DC 20052, USA

^bDepartment of Exercise and Nutrition Sciences, Milken Institute School of Public Health, The George Washington University, 950 New Hampshire Ave, NW, 2nd Floor, Washington, DC 20052, USA

^cDivision of General Internal Medicine and Geriatrics, Department of Medicine, Northwestern University Feinberg School of Medicine, Rubloff Building 10th Floor, 750 N Lake Shore, Chicago IL 6061, USA

^dDivision of General Internal Medicine, Department of Medicine, University of California, San Francisco, 1545 Divisadero, San Francisco CA 94115 USA

Abstract

South Asians, a fast growing ethnic group in the US, have an increased risk for cardiovascular disease compared to the general population. We examined the prevalence and distribution of the American Heart Association's Life's Simple 7 cardiovascular health (CVH) metrics using data from Mediators of Atherosclerosis in South Asians Living in America (MASALA) study, and cross-sectional associations between number of CVH metrics in the ideal range with subclinical atherosclerosis assessed using coronary artery calcium (CAC) measured using cardiac computed tomography and carotid intima media thickness (CIMT) measured using high-resolution B-mode ultrasonography. CAC was modeled as Agatston scores=0, 1–400 and >400; CIMT was examined continuously and as internal CIMT >1.5 mm. In the MASALA cohort (N=875; mean age: 55 y; 53% men; living in greater San Francisco and Chicago areas; October 2010-March 2013) without prevalent coronary heart disease, no participant had all 7 metrics in ideal range; approximately

Address for Correspondence: Sameera A. Talegawkar, PhD, Department of Exercise and Nutrition Sciences, Milken Institute School of Public Health, The George Washington University, 950 New Hampshire Ave, NW, 2nd Floor Washington, DC 20052, Phone: 202-994-8964, Fax: 202-994-3601, sameera.talegawkar@alumni.tufts.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Financial disclosure: No financial disclosures were reported by the authors of this paper.

Conflict of interest: The authors' declare that there are no conflicts of interest.

20% of the participants had at least 5 metrics in ideal range. Higher number of CVH metrics in the ideal range was inversely associated with subclinical atherosclerosis. A 1 unit increase in the number of CVH metrics in the ideal range was associated with 32% lower odds of CAC=1–400 (vs. CAC=0; OR=0.68, 95% CI: 0.60, 0.78) and 28% lower odds of internal CIMT > 1.5mm (OR=0.72, 95% CI: 0.61, 0.85). These data show the prevalence of CVH metrics among South Asians in the US, and provide empirical evidence on inverse associations of meeting ideal levels for higher number of metrics and subclinical atherosclerosis.

Introduction

South Asians, individuals who originate from countries in the Indian sub-continent including India, Pakistan, Bangladesh, Sri Lanka and Nepal, are the sixth largest and fastest growing ethnic group in the United States (US) (Hoeffel et al., 2012). Several studies have underscored the higher risk for cardiovascular disease (CVD) morbidity and mortality among South Asians compared to other racial and ethnic groups indicating the presence of health disparities (Anand et al., 2000; Harding et al., 2008; Jose et al., 2014; Palaniappan et al., 2004). However, there have been few investigations that have focused on elucidating the reasons for the higher risk in this segment of the US population.

In 2010, the American Heart Association (AHA) Strategic Planning Task Force & Statistics Committee set a 10 year goal of improving the cardiovascular health (CVH) of *all* Americans by 20% by the year 2020; to measure progress toward the goal, the AHA defined metrics for ideal, intermediate and poor cardiovascular health for 7 modifiable health behaviors and factors, including smoking, diet, physical activity, body mass index, blood pressure, total cholesterol, and fasting glucose (Lloyd-Jones et al., 2010). As part of their annual surveillance and monitoring, the AHA jointly with the Centers for Disease Control and Prevention and the National Institutes of Health reports the newest nationally representative data on CVD and stroke statistics as well as the current status of CVH for the nation. In 2016 the AHA reported that only 18 % of American adults had 5 or more of these metrics with ideal levels (Mozaffarian et al., 2015). Additionally, studies have also reported the presence of racial and ethnic disparities in the prevalence of ideal CVH metrics. For example, using data from 44, 959 adults from the National Health and Nutrition Examination Survey (NHANES, 1988–1994, 1999–2004, and 2005–2010), Yang et al. (Yang et al., 2012) reported that non-Hispanic whites tended to meet a greater number of CVH metrics as compared to other racial groups. There are, however, a dearth of available data examining the prevalence and distribution of each metric and the overall CVH score for different Asian American subgroups.

Our goal was to determine the prevalence and distribution of the AHA CVH metrics in a cohort of community dwelling, middle-aged and older South Asians in the US. In addition, we also examined the associations between the number of ideal metrics with measures of subclinical atherosclerosis. We postulated that a higher number of CVH metrics in the ideal range would be inversely associated with subclinical atherosclerosis.

Methods

Study population

We used data collected at baseline from the Mediators of Atherosclerosis in South Asians Living in America (MASALA) Study, a community-based sample of South Asians who were free from CVD at baseline. Details regarding recruitment have been published elsewhere (Kanaya et al., 2013). Briefly, using surname-based recruitment methods, a community-based sample of 906 South Asians (age range: 40–84 y, 46% women) were recruited between October 2010 and March 2013 from the two clinical sites including the nine counties of the San Francisco Bay Area (representing the University of California at San Francisco (UCSF) field site) and seven census tracts closest to the Northwestern University (NWU) medical center. For these analyses we excluded 31 participants with missing data for one or more components of the CVH score resulting in a sample of 875 participants. The study protocol and procedures were approved by the institutional review boards of UCSF and NWU. All study participants signed informed consent.

Measures of subclinical atherosclerosis

Coronary Artery Calcium—Cardiac CT scans were performed using a gated-cardiac CT scanner. At UCSF, the system was either the 16D scanner (Philips Medical Systems, Andover, MA) or the MSD Aquilion 64 model (Toshiba Medical Systems, Tustin, CA). At NWU, the technicians used the Sensation Cardiac 64 Scanner (Siemens Medical Solutions, Malvern, PA). All CT scans were sent in batches to the centralized reading center at Harbor-UC Los Angeles Medical Center, where they were interpreted with Rephot Imaging software. Coronary Artery Calcium (CAC) Agatston scores were reported for each of the four major coronary arteries and the summed score was used (A. Kanaya et al., 2014). For all further analyses, the CAC score was modeled categorically as Agatston scores=0, 1–400 and >400.

Carotid Artery Intima Thickness—Right and left internal and common carotid artery intima media thickness (CIMT, in mm) was assessed using high-resolution B-mode ultrasonography. These data were read at the Wake Forest University Medical Center. Wall thickness was defined using the far wall measures. For quality control, a single reader completed 25 repeat CIMT measures. The intra-class correlation coefficient for the internal carotid IMT was 0.96 and 0.78 for the common carotid IMT (A. Kanaya et al., 2014). For all further analyses, CIMT was modeled on: (i) a continuous scale; and (ii) categorically for internal CIMT > 1.5 mm.

Cardiovascular health metrics—Cardiovascular health was operationalized using AHA Life's Simple 7 criteria (LS7) and comprised of seven health behaviors and factors including healthy diet score, physical activity, body mass index (BMI), smoking status, blood pressure, plasma total cholesterol and fasting glucose concentrations. Some of the criteria were adapted based on data available for the MASALA cohort. Specifically, 1) Diet Quality: The Study of Health Assessment and Risk in Ethnic food frequency questionnaire (FFQ) developed and validated for South Asians in Canada was used to assess dietary intake (Kelemen et al., 2003). Participants were given 1 point for meeting each of the dietary goals.

These included: i) Fruit and vegetables: ≥ 4.5 cups/day (potato and potato preparation and fruit and vegetable juices were excluded); ii) Fish: $\geq$ two 3.5 oz servings/week (excluding fried fish); iii) a fiber-to-carbohydrate ratio: ≥ 1.0 g fiber per 10 g carbohydrate/day.; iv) no more than 36 fluid oz/week of sugar-sweetened beverages (including non-diet soda and fruit drinks); v) less than 1500 mg of sodium/day. 2) Physical Activity: The Typical Week's Activity Survey was used to assess levels of various types of physical activity and exercise and the Metabolic Equivalents (METs) of each activity was specified by Ainsworth et al. (Ainsworth et al., 1999). Specifically, we used intentional exercise, including walking for exercise, dance, conditional activities and sports to calculate time spent in moderate and vigorous activities. Time spent in activities identified as either vigorous (>6 METs) or moderate (3–6 METs) were used in derivation. The average time per week spent engaged in all activities at either a vigorous or moderate level was computed for each participant and participants were then categorized based on the AHA criteria (Lloyd-Jones et al., 2010). 3) Smoking status: Smoking status was self-reported and assessed using a questionnaire (Kanaya et al., 2013). 4) BMI: Participant weight was measured on a standard balance-beam scale or a digital weighing scale, and height was measured using a stadiometer. Weight (in kgs) divided by height (in meters) squared was used to calculate BMI. 5) Blood Pressure: Blood pressure was measured thrice seated using an automated blood pressure monitor (V100 Vital Signs Monitor; GE Healthcare, Fairfield, CT) with the average of the last two readings used for analysis (Kanaya et al., 2013). Blood samples in the cohort were obtained after a 12-hour fast, and 6) Total Cholesterol was measured using enzymatic methods, while 7) Blood Sugar was measured with the use of a hexokinase method. Use of glucose, blood pressure and lipid lowering medications were all assessed using a questionnaire (Kanaya et al., 2013).

Participants were assigned a score of 1 for each metric if they met the ideal level and 0 for intermediate or poor levels. The individual metric scores were summed to derive an ideal CVH score which could range from 0–7 with higher scores representing better adherence to the AHA LS7 criteria. For further analyses, in addition to being modeled on a continuous scale, based on the distribution of the overall score in the cohort, we also categorized the score as having 4 or more metrics in the ideal range, followed by having 3 metrics in the ideal range and lastly having 2 or less metrics in the ideal range.

Covariates

Information on age, sex, length of residence in the US, acculturation status, education, family income, insurance type and alcohol intake was obtained using a structured interview and standard questionnaires (Kanaya et al., 2013). Participants reported highest level of education completed which was categorized as Less than high school/High school or some college/Bachelor's degree/Higher than bachelor's degree. Insurance type was classified as Government/HMO/Other/None. Length of residence in the US was self-reported and modeled on a continuous scale as years. A traditional cultural beliefs scale was used as a marker of acculturation (Mukherjee et al., 2013). Scores on this scale were categorized as strong (scores <12), moderate (scores 12–17) or weak (scores >17) traditional beliefs. Alcohol intake (g/d) was derived from self-report.

Statistical analysis

The prevalence of each CVH metric was examined. Baseline characteristics were reported as mean (standard deviation [SD]) for continuous variables or number (percentage) for categorical variables. Differences across the 3 categories of number of ideal metrics were examined. Unadjusted associations between socio-demographic and health characteristics were examined using t-tests, Kruskal Wallis tests, or chi-squared tests as appropriate. The number of ideal metrics met was modeled on both a continuous and categorical scale. Linear and logistic regression models were used to examine the associations between number of ideal metrics met and internal and common CIMT modeled on a continuous scale or internal CIMT >1.5 mm. For CAC, multinomial logistic regression models were used to assess the associations between number of ideal metrics and Agatston score=0, 1–400 and >400. Sequential models were examined and covariates were determined based on preliminary univariate analyses. Final models included: age, sex, length of residence in the US, traditional cultural beliefs, insurance type, study site and alcohol intake. We also examined whether age, sex, length of residence in the US, traditional cultural beliefs, insurance type and study site modified the associations between ideal CVH and subclinical atherosclerosis. In addition we performed sensitivity analyses, wherein we assigned each CVH metric component a score of 2, 1, 0 for meeting the ideal, intermediate, poor criterion, respectively, and used the overall CVH score (which could range from 0 to 14) by summing up scores across the 7 metrics to assess the risk of subclinical atherosclerosis in the adjusted linear and logistic regression models. Analyses were performed using Stata version 13.1. (StataCorp, 2013)

Results

The distributions of socio-demographic and health characteristics by the number of ideal CVH metrics met are shown in Table 1. The average age (Mean $\pm$ SD) of the 875 MASALA participants included in these analyses was 55 $\pm$ 9 y, included 53% men, and participants had a mean residence in the US of 27 $\pm$ 11 y. Almost 42% percent of the cohort had at least 4 of the AHA metrics in the ideal range, followed by 27.9% with 3 ideal metrics and lastly, 30% had 2 or fewer ideal metrics. Individuals with 4 or more ideal CVH metrics were more likely to be women, younger, have weaker traditional cultural beliefs (were more acculturated), more likely to have private health insurance, hold a bachelor's degree or higher, and to be seen at the UCSF study site compared to those with no more than 2 ideal metrics.

None of the MASALA study participants had all 7 CVH metrics in the ideal range. Approximately 20.5% of the participants had 5 or 6 ideal health metrics with some differences by age and sex: 27.4% for those aged < 55 y vs. 13.2 % for those $\geq$ 55 y; 14.9 % for men vs. 26.6% for women (Table 2). Among the health behaviors, more than 80% of the MASALA study participants did not currently smoke, 65% met the ideal levels for physical activity, 44% had ideal BMI and less than 3% met the ideal criteria for diet quality (Figure 1). Among the other health factors, only 56% of the participants had the ideal levels for fasting glucose, 40% for total cholesterol and 33% for blood pressure.

Using multinomial logistic regression, we examined the associations between the number of CVH metrics in the ideal range and CAC scores. For the multivariate model, a 1 unit

increase in number of metrics in the ideal range was associated with a 32% lower odds of CAC =1–400 (OR=0.68, 95% CI: 0.60, 0.78). When modeled categorically, participants with 4 or more CVH metrics in the ideal range had significantly lower odds of CAC =1–400 (OR=0.30, 95% CI: 0.20, 0.46) than participants meeting the ideal levels for 2 or less of the metrics. Similar, graded and inverse associations were seen for the odds of CAC> 400 (Table 3).

For CIMT (Table 4), for the multivariate model, every additional increase in the number of CVH metrics in the ideal range was associated with a 0.02 mm and a 0.05 mm lower common and internal CIMT respectively. When modeled categorically, participants with 4 or more CVH metrics in the ideal range had 0.07 mm and 0.16 mm lower common and internal CIMT respectively, than participants meeting the ideal levels for 2 or less of the metrics. Similar protective associations for number of metrics in the ideal range were observed for internal CIMT modeled on categorical scale.

Using the recommended BMI cut-points for overweight/obesity in Asians (Misra et al., 2009) 33%, 43% and 24% of the MASALA participants were classified as poor, intermediate and ideal respectively for the BMI metric. This also resulted in changes in the proportion of ideal metrics fulfilled by the participants, but the proportion meeting the ideal levels for 5 or 6 of the metrics remained at 20% and the associations with subclinical atherosclerosis persisted (data not shown). In sensitivity analyses (Supplementary Table 1) overall CVH remained inversely associated with CAC and CIMT.

Discussion

This study yielded several important insights about ideal CVH and subclinical atherosclerosis among middle-aged US South Asian men and women without clinical CVD. Using the AHA metrics and definitions of ideal CVH, none of the MASALA study participants achieved ideal levels for all 7 components. Importantly, there was a graded association with the number of ideal CVH metrics with CAC and both common and internal CIMT, with a higher CVH score associated with lower levels of subclinical atherosclerosis. To our knowledge, this is the first study to examine the prevalence of CVH metrics in a large cohort of US South Asians, and to demonstrate relationships between CVH metrics and measures of subclinical atherosclerosis in this ethnic group.

Since the release of the AHA LS7 metrics in 2010, there have been several reports describing the prevalence of these health metrics in the general US population. Using data from the 2011–2012 National Health and Nutrition Examination Survey, Mozaffarian and colleagues (Mozaffarian et al., 2015) reported that 0.1 % of adults aged 20 years and above had ideal levels for all 7 of the health factors and behaviors. The prevalence estimates for those aged 40–59 y for meeting ideal levels for at least 5 criteria were 11% with those meeting ideal criteria for smoking status (74%), followed by fasting blood glucose (52%), physical activity (41%), blood pressure (34%), total cholesterol (34%), BMI (25%), and lastly diet quality (0.5%). This is similar to what we report in our study, wherein smoking status had the highest and diet quality had the lowest proportion of participants with the ideal levels. A lower proportion of MASALA participants also had lower ideal fasting

glucose and blood pressure metrics than the US population. Interestingly, almost two-thirds of the MASALA study participants met the ideal levels for physical activity, whereas the calculated exercise levels for the cohort are quite low (median MET mins/week: 1027 for men and 840 for women) (A. M. Kanaya et al., 2014). A third of the MASALA study participants reported walking for exercise which combined with other intended exercise activities may explain the relatively high proportion of participants meeting the ideal physical activity levels.

In supplementary analyses we examined the distribution of the CVH metrics by a traditional cultural beliefs scale, a measure of acculturation in this cohort (Supplementary Table 2). Participants with weaker traditional beliefs were more physically active and had better diet quality. Associations of better diet quality with greater acculturation may be due to the components of diet score, which includes foods such as fish not consumed by South Asians following a vegetarian diet for religious reasons. We have previously also demonstrated that dietary intakes in the cohort differed by length of residence in the US, with largely mixed benefits (Talegawkar et al., 2016). These findings suggest that immigration and acculturation are associated with CVH behaviors in South Asians, and should be considered when planning interventions for immigrant populations.

In addition to reporting the prevalence of the CVH metrics, this study also demonstrated protective associations between the number of ideal CVH metrics and measures of subclinical atherosclerosis, including CIMT and CAC. Saleem et.al (Saleem et al., 2015) examined the cross-sectional associations between the number of ideal CVH metrics and CAC in middle-aged, predominantly non-Hispanic White patients from a clinic sample. Patients with favorable scores, defined as meeting ideal levels for 4 or more CVH metrics, had over 50% lower odds of any coronary artery calcium than patients with unfavorable (ideal levels for only 0–2 metrics) profiles and 32% lower odds with intermediate scores. More recently, Robbins et.al. demonstrated similar protective associations between CVH metrics and prevalent calcified atherosclerotic plaque (defined as Agatston scores>100) in the participants of the National Heart, Lung, and Blood Institute Family Heart Study, which was also predominantly non-Hispanic whites (Robbins et al., 2015). Among children and young adults aged 12 to 24 y from the Cardiovascular Risk in Young Finns Study cohort, an average lifetime ideal CVH index based on the arithmetic mean of the number of ideal factors and behaviors at baseline and follow-up visit was associated with a reduced risk of CAC and CIMT in middle age. CIMT and CAC are both significant and well-established predictors of future coronary heart disease events (Detrano et al., 2008; Pletcher et al., 2004; Polak et al., 2011), and are useful as surrogate markers for preventing progression and subsequent incidence of disease.

While the predictive associations between cardiovascular health operationalized using the AHA LS7 and CVD morbidity and mortality are not unexpected, there is also evidence from epidemiological investigations that in addition to its associations with heart disease, CVH is also predictive of functional status (Dhamoon et al., 2015), cognitive decline (Crichton et al., 2014; Reis et al., 2013; Thacker et al., 2014), kidney disease (Muntner et al., 2013) and cancer (Foraker et al., 2015; Rasmussen-Torvik et al., 2013). There has been a worldwide increase in the prevalence of chronic diseases (United Nations, 2011), leading to a

concurrent rise in multimorbidity conditions (Garin et al., 2016; Goodman et al., 2016). Therefore, the associations between CVH and other chronic diseases and health conditions underline the public health importance of achieving the ideal levels of the LS7 metric proposed by the AHA to achieve its 2020 impact goals not just for preventing deaths from CVD and stroke but for overall health, wellbeing and longevity.

Strengths of our study include its population- South Asians in the US are a fast growing and understudied minority growing with high risk for chronic disease including heart disease. Key variables in the MASALA study including diet have been collected using culturally appropriate and/or validated instruments. The study also had several other important sociodemographic covariates to facilitate controlling for confounding. Study limitations include that the MASALA study is observational in study design with the potential for residual confounding by unmeasured factors. In addition, the participants of the MASALA study were middle-aged and older and relatively well educated which limits the generalizability of the study findings to younger and less-educated South Asians in the US or other South Asians in the diaspora.

This study provides important new information on the association between the AHA CVH metrics and subclinical atherosclerosis in US South Asians. The low prevalence of ideal CVH metrics in US South Asians is similar to what has been found in the general US population. Longitudinal follow-up of the MASALA cohort will allow further study of the relationship between the AHA CVH metrics and progression of subclinical atherosclerosis, incident CVD events, and mortality. Effective primordial and primary prevention interventions are needed to preserve and improve CVH in US South Asians, as well as continued research on novel and emerging risk factors (Bharmal et al., 2016; Flowers et al., 2015; Nadimpalli et al., 2016; A. D. Shah et al., 2016; B. M. Shah et al., 2016) that may be contributing to the high burden of CVD in South Asians.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors would like to acknowledge Dr. Darwin R. Labarthe for providing critical comments on an earlier version of this manuscript. The project described was supported by Grant Number R01HL093009 from the National Heart, Lung, And Blood Institute and the National Center for Research Resources and the National Center for Advancing Translational Sciences, National Institutes of Health, through UCSF-CTSI Grant Number UL1 RR0241. The findings and conclusions in this article are those of the authors and not necessarily those of the National Institutes of Health.

Abbreviations used

AHA	American Heart Association
CVD	Cardiovascular Disease
CVH	Cardiovascular Health
CAC	Coronary Artery Calcium

CIMT	Carotid Intima Median Thickness
BMI	Body Mass Index
BP	Blood Pressure
FFQ	Food Frequency Questionnaire
LS7	Life's Simple 7
MASALA	Mediators of Atherosclerosis in South Asians Living in America
MET	Metabolic Equivalents
NHANES	National Health and Nutrition Examination Survey
NWU	Northwestern University
US	Study of Health Assessment and Risk in Ethnic, United States
UCSF	University of California San Francisco

References

- Ainsworth BE, Irwin ML, Addy CL, Whitt MC, Stolarczyk LM. Moderate physical activity patterns of minority women: the Cross-Cultural Activity Participation Study. *Journal of women's health & gender-based medicine*. 1999; 8:805–813.
- Anand SS, Yusuf S, Vuksan V, Devanese S, Teo KK, Montague PA, Kelemen L, Yi C, Lonn E, et al. Differences in risk factors, atherosclerosis, and cardiovascular disease between ethnic groups in Canada: the Study of Health Assessment and Risk in Ethnic groups (SHARE). *Lancet*. 2000; 356:279–284. [PubMed: 11071182]
- Bharmal NH, McCarthy WJ, Gadgil MD, Kandula NR, Kanaya AM. The Association of Religious Affiliation with Overweight/Obesity Among South Asians: The Mediators of Atherosclerosis in South Asians Living in America (MASALA) Study. *J Relig Health*. 2016
- Crichton GE, Elias MF, Davey A, Alkerwi A. Cardiovascular health and cognitive function: the Maine-Syracuse Longitudinal Study. *PloS one*. 2014; 9:e89317. [PubMed: 24595096]
- Detrano R, Guerci AD, Carr JJ, Bild DE, Burke G, Folsom AR, Liu K, Shea S, Szklo M, et al. Coronary calcium as a predictor of coronary events in four racial or ethnic groups. *The New England journal of medicine*. 2008; 358:1336–1345. [PubMed: 18367736]
- Dhamoon MS, Dong C, Elkind MS, Sacco RL. Ideal cardiovascular health predicts functional status independently of vascular events: the Northern Manhattan Study. *Journal of the American Heart Association*. 2015; 4
- Flowers E, Gadgil M, Aouizerat BE, Kanaya AM. Circulating micrornas associated with glycemic impairment and progression in Asian Indians. *Biomark Res*. 2015; 3:22. [PubMed: 26966540]
- Foraker RE, Abdel-Rasoul M, Kuller LH, Jackson RD, Van Horn L, Seguin RA, Safford MM, Wallace RB, Kucharska-Newton AM, et al. Cardiovascular Health and Incident Cardiovascular Disease and Cancer: The Women's Health Initiative. *American journal of preventive medicine*. 2015
- Garin N, Koyanagi A, Chatterji S, Tyrovolas S, Olaya B, Leonardi M, Lara E, Koskinen S, Tobiasz-Adamczyk B, et al. Global Multimorbidity Patterns: A Cross-Sectional, Population-Based, Multi-Country Study. *The journals of gerontology. Series A, Biological sciences and medical sciences*. 2016; 71:205–214.
- Goodman RA, Ling SM, Briss PA, Parrish RG, Salive ME, Finke BS. Multimorbidity Patterns in the United States: Implications for Research and Clinical Practice. *The journals of gerontology. Series A, Biological sciences and medical sciences*. 2016; 71:215–220.

- Harding S, Rosato M, Teyhan A. Trends for coronary heart disease and stroke mortality among migrants in England and Wales, 1979–2003: slow declines notable for some groups. *Heart*. 2008; 94:463–470. [PubMed: 17690159]
- Hoeffel, EM., Rastogi, S., Kim, MO., Hasan, S. Census, U.S.B.o.t. The Asian population: 2010. US Department of Commerce, Economics and Statistics Administration, US Census Bureau; 2012.
- Jose PO, Frank AT, Kapphahn KI, Goldstein BA, Eggleston K, Hastings KG, Cullen MR, Palaniappan LP. Cardiovascular disease mortality in Asian Americans. *Journal of the American College of Cardiology*. 2014; 64:2486–2494. [PubMed: 25500233]
- Kanaya A, Ewing S, Vittinghoff E, Herrington D, Tegeler C, Mills C, Kandula N. Acculturation and Subclinical Atherosclerosis among US South Asians: Findings from the MASALA study. *Journal of clinical and experimental research in cardiology*. 2014; 1
- Kanaya AM, Kandula N, Herrington D, Budoff MJ, Hulley S, Vittinghoff E, Liu K. Mediators of Atherosclerosis in South Asians Living in America (MASALA) Study: Objectives, Methods, and Cohort Description. *Clin Cardiol*. 2013; 36:713–720. [PubMed: 24194499]
- Kanaya AM, Kandula NR, Ewing SK, Herrington D, Liu K, Blaha MJ, Srivastava S, Dave SS, Budoff MJ. Comparing coronary artery calcium among US South Asians with four racial/ethnic groups: The MASALA and MESA studies. *Atherosclerosis*. 2014; 234:102–107. [PubMed: 24632509]
- Kelemen LE, Anand SS, Vuksan V, Yi Q, Teo KK, Devanesen S, Yusuf S. Investigators, S. Development and evaluation of cultural food frequency questionnaires for South Asians, Chinese, and Europeans in North America. *Journal of the American Dietetic Association*. 2003; 103:1178–1184. [PubMed: 12963948]
- Lloyd-Jones DM, Hong Y, Labarthe D, Mozaffarian D, Appel LJ, Van Horn L, Greenlund K, Daniels S, Nichol G, et al. Defining and setting national goals for cardiovascular health promotion and disease reduction: the American Heart Association's strategic Impact Goal through 2020 and beyond. *Circulation*. 2010; 121:586–613. [PubMed: 20089546]
- Misra A, Chowbey P, Makkar BM, Vikram NK, Wasir JS, Chadha D, Joshi SR, Sadikot S, Gupta R, et al. Consensus statement for diagnosis of obesity, abdominal obesity and the metabolic syndrome for Asian Indians and recommendations for physical activity, medical and surgical management. *The Journal of the Association of Physicians of India*. 2009; 57:163–170. [PubMed: 19582986]
- Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, de Ferranti S, Despres JP, Fullerton HJ, et al. Heart disease and stroke statistics--2015 update: a report from the American Heart Association. *Circulation*. 2015; 131:e29–e322. [PubMed: 25520374]
- Mukherjee A, Underwood KC, Stewart AL, Ivey SL, Kanaya AM. Asian Indian views on diet and health in the United States: importance of understanding cultural and social factors to address disparities. *Family & community health*. 2013; 36:311–323. [PubMed: 23986072]
- Muntner P, Judd SE, Gao L, Gutierrez OM, Rizk DV, McClellan W, Cushman M, Warnock DG. Cardiovascular risk factors in CKD associate with both ESRD and mortality. *Journal of the American Society of Nephrology : JASN*. 2013; 24:1159–1165. [PubMed: 23704285]
- Nadimpalli SB, Dulin-Keita A, Salas C, Kanaya AM, Kandula NR. Associations Between Discrimination and Cardiovascular Health Among Asian Indians in the United States. *Journal of immigrant and minority health / Center for Minority Public Health*. 2016
- Palaniappan L, Wang Y, Fortmann SP. Coronary heart disease mortality for six ethnic groups in California, 1990–2000. *Annals of epidemiology*. 2004; 14:499–506. [PubMed: 15310526]
- Pletcher MJ, Tice JA, Pignone M, Browner WS. Using the coronary artery calcium score to predict coronary heart disease events: a systematic review and meta-analysis. *Archives of internal medicine*. 2004; 164:1285–1292. [PubMed: 15226161]
- Polak JF, Pencina MJ, Pencina KM, O'Donnell CJ, Wolf PA, D'Agostino RB Sr. Carotid-wall intima-media thickness and cardiovascular events. *The New England journal of medicine*. 2011; 365:213–221. [PubMed: 21774709]
- Rasmussen-Torvik LJ, Shay CM, Abramson JG, Friedrich CA, Nettleton JA, Prizment AE, Folsom AR. Ideal cardiovascular health is inversely associated with incident cancer: the Atherosclerosis Risk In Communities study. *Circulation*. 2013; 127:1270–1275. [PubMed: 23509058]

- Reis JP, Loria CM, Launer LJ, Sidney S, Liu K, Jacobs DR Jr, Zhu N, Lloyd-Jones DM, He K, et al. Cardiovascular health through young adulthood and cognitive functioning in midlife. *Annals of neurology*. 2013; 73:170–179. [PubMed: 23443990]
- Robbins JM, Petrone AB, Carr JJ, Pankow JS, Hunt SC, Heiss G, Arnett DK, Ellison RC, Gaziano JM, et al. Association of ideal cardiovascular health and calcified atherosclerotic plaque in the coronary arteries: the National Heart, Lung, and Blood Institute Family Heart Study. *American heart journal*. 2015; 169:371–378. e1. [PubMed: 25728727]
- Saleem Y, DeFina LF, Radford NB, Willis BL, Barlow CE, Gibbons LW, Khera A. Association of a favorable cardiovascular health profile with the presence of coronary artery calcification. *Circulation. Cardiovascular imaging*. 2015; 8
- Shah AD, Kandula NR, Lin F, Allison MA, Carr J, Herrington D, Liu K, Kanaya AM. Less favorable body composition and adipokines in South Asians compared with other US ethnic groups: results from the MASALA and MESA studies. *Int J Obes (Lond)*. 2016; 40:639–645. [PubMed: 26499444]
- Shah BM, Shah S, Kandula NR, Gadgil MD, Kanaya AM. Psychosocial Factors Associated with Subclinical Atherosclerosis in South Asians: The MASALA Study. *Journal of immigrant and minority health / Center for Minority Public Health*. 2016
- StataCorp. *Stata Statistical Software: Release. Vol. 13*. College Station, TX: StataCorp LP; 2013.
- Talegawkar SA, Kandula NR, Gadgil MD, Desai D, Kanaya AM. Dietary intakes among South Asian adults differ by length of residence in the USA. *Public health nutrition*. 2016; 19:348–355. [PubMed: 25990446]
- Thacker EL, Gillett SR, Wadley VG, Unverzagt FW, Judd SE, McClure LA, Howard VJ, Cushman M. The American Heart Association Life's Simple 7 and incident cognitive impairment: The REasons for Geographic And Racial Differences in Stroke (REGARDS) study. *Journal of the American Heart Association*. 2014; 3:e000635. [PubMed: 24919926]
- United Nations, GA. *Prevention and Control of Non-communicable Diseases*. United Nations, New York, NY: 2011.
- Yang Q, Cogswell ME, Flanders WD, Hong Y, Zhang Z, Loustalot F, Gillespie C, Merritt R, Hu FB. Trends in cardiovascular health metrics and associations with all-cause and CVD mortality among US adults. *Jama*. 2012; 307:1273–1283. [PubMed: 22427615]

Highlights

- About 20% of South Asians from the MASALA study met ≥ 5 ideal CVH metrics.
- Prevalence of ideal diet was only 1.1 %
- A higher number of ideal CVH metrics was inversely associated with subclinical atherosclerosis




Figure 1. Distribution of Life's Simple 7 metric in the MASALA study participants living in the greater San Francisco and Chicago areas from the US, 2010–2013

Socio-demographic and health characteristics by the number of AHA ideal CVH metrics among MASALA study participants living in the greater San Francisco and Chicago areas from the US, 2010–2013

Table 1

	Overall	Number of ideal metrics fulfilled ^a			P-value
		0–2	3	4–6	
N	875	263	244	368	
Female, %	47.2	36.9	45.5	55.7	<0.001
Age (SD), y	55.3 (9.4)	57.1 (9.0)	56.1 (9.3)	53.5 (9.3)	<0.001
NWU site, %	45.6	51.7	45.1	41.6	0.04
Traditional cultural beliefs, % (n=873)					0.036
- Strong	33.4	38.3	34.0	29.6	
- Intermediate	29.9	23.0	31.6	33.7	
- Weak	36.7	38.7	34.4	36.7	
Length of residence in the US (SD), y	27.4 (10.8)	28.1 (11.3)	27.9 (10.9)	26.6 (10.9)	0.182
Insurance type, %					<0.001
- Government	10.3	13.3	10.7	7.9	
- HMO	76.9	68.4	78.3	82.0	
- Other	4.7	4.6	3.3	5.7	
- None	8.1	13.7	7.8	4.4	
Education, %					<0.001
- Lower than high school	6.9	12.2	7.0	3.0	
- Higher school or some college	5.5	7.6	5.3	4.1	
- Bachelor's degree	29.1	30.0	27.9	29.3	
- Higher than Bachelor's degree	58.5	50.2	59.8	63.6	
Family income ≥ \$75,000, % (n=851)	73.4	64.0	72.6	80.9	<0.001
Current smoker, %	3.2	6.8	3.3	0.5	<0.001
Physical activity, %					<0.001
- Poor	14.7	25.5	16.0	6.3	
- Intermediate	19.8	34.6	19.7	9.2	

	Number of ideal metrics fulfilled ^a				p-value
	Overall	0–2	3	4–6	
-Ideal	65.5	39.9	64.3	84.5	
Diet quality, %					0.269
-Poor	39.1	40.7	43.0	35.3	
-Intermediate	58.5	57.8	54.5	61.7	
-Ideal	2.4	1.5	2.5	3.0	
Lipid lowering medication, %	28.3	46.8	33.6	11.7	<0.001
Blood pressure lowering medication, %	31.1	50.2	35.7	14.2	<0.001
Diabetes medication, %	16.0	33.5	16.8	3.0	<0.001
Alcohol consumption (SD), g/day	2.8 (5.9)	3.5 (6.3)	3.0 (7.1)	2.0 (4.6)	0.006
BMI (SD), kg/m ²	26.0 (4.3)	28.1 (3.7)	26.6 (5.0)	24.2 (3.4)	<0.001
Total cholesterol (SD), mg/dL	187 (36.9)	189 (41.0)	187 (39.3)	187 (31.8)	0.712
Systolic blood pressure (SD), mm Hg	124 (15.7)	130 (13.8)	127 (16.3)	119 (14.6)	<0.001
Diastolic blood pressure (SD), mm Hg	73.4 (9.7)	75.8 (9.3)	74.6 (10.2)	70.9 (9.1)	<0.001
Fasting plasma glucose (SD), mg/dL ^b	103(24.7)	117(31.0)	104(24.3)	92.4 (10.5)	<0.001
CAC score, % (n=869)					<0.001
-0	58.0	40.9	53.3	73.2	
-1–400	34.8	47.5	39.3	22.7	
->400	7.2	11.6	7.4	4.1	
Common CIMT (SD, n=874), mm	0.88 (0.23)	0.93 (0.28)	0.89 (0.21)	0.83 (0.19)	<0.001
Internal CIMT (SD, n=873), mm	1.21 (0.44)	1.33 (0.51)	1.20 (0.43)	1.12 (0.37)	<0.001

^aNo participant had all 7 Life's Simple 7 metrics in the ideal range

^bKruskal-Wallis test

AHA, American Heart Association; MASALA, Mediators of Atherosclerosis in South Asians Living in America; BMI, Body Mass Index; CVH, Cardiovascular Health

Distribution of ideal CVH metrics by age and sex, among MASALA study participants living in the greater San Francisco and Chicago areas from the US, 2010–2013

Table 2

No. of Ideal Health Metrics Present	Total Sample [number(%)] (n=875)	Men [number (%)]			Women [number (%)]		
		Age < 55 (n=219)	Age ≥ 55 (n=243)	p-value	Age < 55 (n=225)	Age ≥ 55 (n=188)	p-value
7	0 (0)	0 (0)	0 (0)		0 (0)	0 (0)	
6	41 (4.7)	11 (5.0)	4 (1.7)	0.05	21 (9.3)	5 (2.7)	0.01
5	138 (15.8)	35 (16.0)	19 (7.8)	0.01	55 (24.4)	29 (15.4)	0.02
4	189 (21.6)	45 (20.6)	49 (20.2)	0.92	55 (24.4)	40 (21.3)	0.46
3	244 (27.9)	56 (25.6)	77 (31.7)	0.15	56 (24.9)	55 (29.3)	0.32
2	176 (20.1)	41 (18.7)	60 (24.7)	0.12	30 (13.3)	45 (23.9)	0.01
1	73 (8.3)	26 (11.9)	26 (10.7)	0.68	8 (3.6)	13 (6.9)	0.13
0	14 (1.6)	5 (2.3)	8 (3.3)	0.52	0 (0)	1 (0.5)	^a

^aNo female under 55y when ideal health metrics=0, therefore age-groups cannot be compared.

CVH, Cardiovascular Health; MASALA, Mediators of Atherosclerosis in South Asians Living in America

Table 3

Odds Ratios (95% CI) for associations between number of CVH metrics in the ideal range and CAC among MASALA study participants living in the greater San Francisco and Chicago areas from the US, 2010–2013

Number of metrics in ideal range	CAC category			
	CAC 1–400 vs. CAC=0		CAC >400 vs. CAC=0	
	OR	95 % CI	OR	95% CI
Crude model				
Continuous	0.64	0.52, 0.72	0.58	0.47, 0.70
Categorical				
0–2	Ref		Ref	
3	0.64	0.44, 0.92	0.49	0.26, 0.93
4–6	0.27	0.19, 0.38	0.20	0.10, 0.38
Multivariate adjusted ^a				
Continuous	0.68	0.60, 0.78	0.57	0.44, 0.74
Categorical				
0–2	Ref		Ref	
3	0.65	0.42, 0.99	0.46	0.21, 1.00
4–6	0.30	0.20, 0.46	0.20	0.09, 0.44

CAC, Coronary Artery Calcium; CVH, Cardiovascular Health; MASALA, Mediators of Atherosclerosis in South Asians Living in America;

^a Multivariate model has been adjusted for sex, age, length of residence in the U.S., traditional cultural beliefs, insurance type, alcohol intake and study site

Table 4

Regression coefficients and Odds Ratios (95% CI) for associations between the number of CVH metrics in the ideal range and CIMT among MASALA study participants living in the greater San Francisco and Chicago areas from the US, 2010–2013

Number of metrics in ideal range	Common CIMT		Internal CIMT		Internal CIMT > 1.5 mm	
	Beta	95% CI	Beta	95% CI	OR	95% CI
Crude model						
Continuous	-0.03	-0.05, 0.02	-0.07	-0.09, -0.05	0.70	0.61, 0.81
Categorical						
0–2	Ref		Ref		Ref	
3	-0.05	-0.09, -0.004	-0.14	-0.22, -0.05	0.55	0.35, 0.85
4–6	-0.11	-0.15, -0.07	-0.21	-0.28, -0.14	0.32	0.21, 0.50
Multivariate model ^a						
Continuous	-0.02	-0.03, -0.01	-0.05	-0.07, -0.03	0.72	0.61, 0.85
Categorical						
0–2	Ref		Ref		Ref	
3	-0.04	-0.08, 0	-0.13	-0.20, -0.06	0.50	0.31, 0.80
4–6	-0.07	-0.11, -0.04	-0.16	-0.23, -0.09	0.35	0.21, 0.57

CIMT, Carotid Intima Media Thickness; MASALA, Mediators of Atherosclerosis in South Asians Living in America

^a Multivariate model has been adjusted for sex, age, length of residence in the US, traditional cultural beliefs score, insurance type, alcohol intake, and study site