Lawrence Berkeley National Laboratory

Recent Work

Title

Stabilization and Structure of the cis Tautomer of a Free-Base Porphyrin.

Permalink

<https://escholarship.org/uc/item/25v7422s>

Journal

Angewandte Chemie (International ed. in English), 56(34)

ISSN

1433-7851

Authors

Thomas, Kolle E
McCormick, Laura J
Vazquez-Lima, Hugo
et al.

Publication Date

2017-08-01

DOI

10.1002/anie.201701965

Peer reviewed

Stabilization and Structure of the *Cis* Tautomer of a Free-Base Porphyrin

Kolle E. Thomas,^[a] Laura J. McCormick,^[b] Hugo Vazquez-Lima^[a] and Abhik Ghosh^{*[a]}

Abstract. Single-crystal X-ray analysis of the $\square$ -heptakis(trifluoromethyl)-meso-tetrakis(*p*-fluorophenyl)porphyrin, $H_2[(CF_3)_7TpFPP]$, has revealed the first example of a stable *cis* tautomer of a free-base porphyrin, the long-postulated intermediate of porphyrin tautomerism. The stability of the unique molecule appears to reflect a dual origin: a strongly saddled porphyrin skeleton, which alleviates electrostatic repulsion between the two NH protons, and two polarization-enhanced, transannular N-H...O-H...N hydrogen bond chains, each involving a molecule of water. DFT calculations suggest that the observed tautomer has a lower energy than the alternative, doubly hydrated *trans* tautomer by some 6.6 kcal/mol. A fascinating prospect thus exists that $H_2[(CF_3)_7TpFPP] \cdot 2H_2O$ and cognate structures may act as supramolecular synthons, which, given their chirality, may even be amenable to resolution into optically pure enantiomers.

The fundamental geometrical disposition and dynamics of the central NH protons of free-base porphyrins were elucidated in a series of seminal studies during the latter part of the last century.^[1] Early X-ray^[2,3,4,5] crystal structures and *ab initio* calculations^[6,7,8] confirmed a linear N-H...H-N arrangement with each NH proton engaged in symmetric, bifurcated hydrogen bonding with two neighboring nitrogen atoms. Detailed kinetic studies further established that the degenerate tautomerism of free-base porphyrins occurs asynchronously, most likely via the *cis* free-base tautomer.^[9,10,11,12] *Ab initio* and density functional theory (DFT) calculations established the *cis* tautomer as a true minimum, with exact C_{2v} symmetry for unsubstituted porphyrin.^[13,14] No direct, experimental characterization of the *cis* tautomer, however, has been reported until now. Here we describe the serendipitous isolation and single-crystal X-ray structure determination of the *cis* tautomer of a free-base porphyrin. The structure reveals an unusual combination of factors that result in the existence of the *cis* tautomer as the energetically preferred form for the free-base porphyrin in question.

In the course of our continuing studies of electron-deficient, highly perfluoroalkylated porphyrinoid systems,^[15,16,17] acid-induced demetalation of an inseparable mixture of $Cu[(CF_3)_7TpFPP]$ and $Cu[(CF_3)_8TpFPP]$ led to the correspond-

Table 1. Crystal and structure refinement data for $H_2[(CF_3)_7TpFPP] \cdot 2H_2O$.

Chemical Formula	$C_{51}H_{23}F_{25}N_4O_2$
[a] Dr. Kolle E. Thomas, Dr. Hugo Vazquez-Lima, Professor Dr. Abhik Ghosh*	
Department of Chemistry	
UiT – The Arctic University of Norway	
9037 Tromsø, Norway	
E-mail: abhik.ghosh@uit.no (AG)	
[b] Dr. Laura J. McCormick	
Advanced Light Source	
Lawrence Berkeley National Laboratory	
Berkeley, CA 94720-8229, USA	

<i>b</i> (Å)	27.9206(18)
<i>c</i> (Å)	11.9969(11)
α (°)	90
β (°)	104.213(6)
γ (°)	90
<i>Z</i>	4
<i>V</i> (Å ³)	4671.0(6)
Temperature (K)	100(2)
ρ (g/cm ³)	1.705
Measured reflections	36249
Unique reflections	7439
Parameters	840
Restraints	56
<i>R</i> _{int}	0.0398
θ range (°)	2.068 - 26.566
<i>R</i> 1, <i>wR</i> 2 all data	0.1103, 0.2762
<i>S</i> (GooF) all data	1.042
Max/min res. Dens. (e/Å ³)	1.452/-1.123

ing free-base porphyrins $H_2[(CF_3)_7TpFPP]$ and $H_2[(CF_3)_8TpFPP]$, which could be readily separated by means of column chromatography.^[18] Of the two products, $H_2[(CF_3)_7TpFPP]$ proved amenable to single-crystal X-ray structure analysis, revealing the first example of a stable, *cis* tautomer of a free-base porphyrin (Figure 1 and Tables 1 and 2). As expected on the basis of its sterically hindered character,^[19,20,21] the porphyrin core is strongly saddled, with the two central NH groups pointing above and below the mean N_4 plane. Each NH group acts as a hydrogen bond donor to a water molecule, which in turn acts as a hydrogen bond donor to an unprotonated nitrogen across the porphyrin. Both the N-H...O and O-H...N hydrogen bonds are approximately linear and ~2.0 Å in length.^[22,23,24,25,26,27] Figure 2 shows the location of the peaks of electron density corresponding to the N- and O-bound hydrogen atoms. The assignment and modeling of hydrogen atom positions is described in detail in the Supporting Information.

The unique stability of the *cis* tautomer appears to result from a dual origin, comprising both strong saddling and the hydrogen bond network. By increasing their spatial separation, strong saddling alleviates the electrostatic repulsion between the central NH protons of the *cis* tautomer. Additional stability accrues via two so-called homodromic,^[24,25,28,29] polarization-enhanced N-H...O-H...N hydrogen bond chains. DFT calculations (PBE0^[30,31]-D3/STO-TZ2P; ADF^[32] 2016) calculations on the *cis* and *trans* tautomers of a series free-base porphyrins.³³

Figure 1. Thermal ellipsoid plots (33%) of $\text{H}_2[(\text{CF}_3)_7\text{TpFPP}]\cdot 2\text{H}_2\text{O}$: (a) top view and (b) close-up of the porphyrin core with hydrogen bond distances (Å).

Table 2. Hydrogen bond metrical parameters (Å, °) for $\text{H}_2[(\text{CF}_3)_7\text{TpFPP}]\cdot 2\text{H}_2\text{O}$.

D-H...A	d(D-H)	d(H...A)	d(D...A)	<DHA
N(2)-H(2)...O(2W)	0.73(6)	2.12(7)	2.847(6)	177(7)
O(2W)-H(3W)...N(4)	0.82(5)	2.22(6)	2.983(6)	156(8)
N(1)-H(1)...O(1W)	0.70(7)	2.19(8)	2.873(7)	166(8)
O(1W)-H(2W)...N(3)	0.84(5)	2.14(6)	2.909(6)	153(9)

are consistent with the dual stabilization of the *cis* tautomer. (The term homodromic refers to a hydrogen-bonded chain or ring where the constituent D-H...A units all have the same directionality.) Thus, the energy of the *cis* tautomer relative to the *trans* tautomer (in kcal/mol) decreases along the series $\text{H}_2[\text{TPP}]$ (8.74), $\text{H}_2[\text{Br}_8\text{TPP}]$ (4.52), $\text{H}_2[\text{I}_8\text{TPP}]$ (2.93),^[34] and $\text{H}_2[(\text{CF}_3)_8\text{TPP}]$ (1.45), paralleling the steep increase in saddling of the optimized structures. Hydrogen bonding with two water molecules then tips the energy balance, favoring the *cis* tautomer for $\text{H}_2[(\text{CF}_3)_8\text{TPP}]\cdot 2\text{H}_2\text{O}$ by 6.62 kcal/mol relative to the *trans* tautomer. As elsewhere,^[24,25] the homodromic hydrogen bond network found in the *cis* tautomer appears to confer much greater stability than the antidromic network assumed for the *trans* tautomer.

The calculated stability of the *cis* tautomer of $\text{H}_2[(\text{CF}_3)_8\text{TPP}]\cdot 2\text{H}_2\text{O}$ by a clear margin of energy potentially presages the emergence of *cis* porphyrin dihydrates as a new supramolecular synthon.^[35,36] Furthermore, the chiral, approximately C_2 -symmetric nature of the compounds may allow successful resolution of the enantiomers, ushering in a new, inherently chiral chromophore.^[37,38] Another interesting aspect of $\text{H}_2[(\text{CF}_3)_7\text{TpFPP}]$ and $\text{H}_2[(\text{CF}_3)_8\text{TpFPP}]$ is that both molecules exhibit strong Q bands extending well into the NIR region (see Figure S3 under Supporting Information), potentially suggesting applications in photodynamic therapy and as NIR dyes.^[39]

In conclusion, a detailed X-ray analysis of $\square$ -heptakis(trifluoromethyl)-*meso*-tetrakis(*p*-fluorophenyl)porphyrin, $\text{H}_2[(\text{CF}_3)_7\text{TpFPP}]$, has yielded unambiguous proof of the existence of a long-postulated

reactive intermediate – the *cis* tautomer of a free-base porphyrin. The stability of the unique molecule has a dual origin, viz. a strongly saddled geometry and a hydrogen bond network involving two water molecules of crystallization per porphyrin unit.

Experimental Section

Synthetic protocols and spectroscopic data have been presented in the Supporting Information, with the present section focusing on the X-ray structure analysis. X-ray data for $\text{H}_2[(\text{CF}_3)_7\text{TpFPP}]\cdot 2\text{H}_2\text{O}$ were collected on beamline 11.3.1 at the Advanced Light Source, using a Bruker D8 diffractometer equipped with a PHOTON100 CMOS detector operating in shutterless mode. The crystal was coated in protective oil prior to being mounted on a MiTeGen[®] kapton micromount and placed under a nitrogen stream at 100(2) K provided by an Oxford Cryostream 800 Plus low temperature apparatus. Diffraction data were collected with synchrotron radiation monochromated using silicon(111) to a wavelength of 0.7749(1) Å. An approximate full-sphere of data was collected using a combination of $\square$ and $\square$ scans with scan speeds of 4° per second for the $\square$ scans, and 1 second per degree for the $\square$ scans at $2\theta = 0$ and -45° , respectively. The structures were solved by intrinsic phasing (SHELXT)^[40] and refined by full-matrix least squares on F^2 (SHELXL-2014).^[41] Absorption corrections were applied using SADABS.^[42] Additional crystallographic information has been summarized in Tables 1 and 2, and full details can be found in the crystallographic information files provided in the Supplementary Information.

Table 3. Selected PBE0 and experimental geometry parameters (Å, deg); relative energies (kcal/mol) of the *cis* tautomers.*

Molecule	<i>trans</i>					<i>cis</i>							E_{cis} (rel)
	$\square$	$\square$	$\square$	<i>a</i>	<i>j</i>	$\square$	$\square$	$\square$	<i>a</i>	<i>b</i>	<i>c</i>	<i>j</i>	
H ₂ [TPP]	13.9	110.7	105.6	2.915	2.303	3.3	110.3	105.9	3.170	2.697	3.083	1.914	8.2
H ₂ [Br ₈ TPP]	63.6	112.8	107.6	2.913	2.351	74.8	112.5	107.7	3.042	2.864	3.009	2.378	5.1
H ₂ [I ₈ TPP]	71.9	112.7	107.5	2.906	2.377	84.8	112.3	107.5	3.027	2.876	2.994	2.431	4.9
H ₂ [(CF ₃) ₈ TPP]	129.6	113.2	107.6	2.979	2.735	134.1	112.9	107.3	3.049	2.956	2.981	2.726	1.1
H ₂ [(CF ₃) ₈ TPP]·2H ₂ O	133.8	112.4	107.6	2.988	2.786	132.6	111.9	107.3	3.070	3.033	3.010	2.778	-8.3
H ₂ [(CF ₃) ₇ TPP]·2H ₂ O (expt)						117.6(11)	110.7(5), 111.3(5)	108.7(4), 107.8(4)	3.047(7)	3.011(6), 3.053(6)	2.992(5)		

* Additional PBE0 hydrogen bond metrical parameters (Å) for the two tautomers of H₂[(CF₃)₈TPP]·2H₂O.

<i>trans</i>					<i>cis</i>				
<i>h</i>	<i>k</i>	<i>m</i>	<i>n</i>	<i>p</i>	<i>h</i>	<i>k</i>	<i>m</i>	<i>n</i>	<i>p</i>
1.019	2.022	2.179	0.971	0.967	1.032	1.841	1.995	0.984	0.965

Figure 2. Electron density map surrounding the central porphyrin unit of $H_2[(CF_3)_7TpFPP] \cdot 2H_2O$. Peripheral CF_3 and C_6H_4F groups have been omitted for clarity. Threshold for the electron density map has been set at $0.2 \text{ e}/\text{\AA}^3$. The yellow and green solid polyhedra represent peaks of electron density. The $F_{\text{observed}} - F_{\text{calculated}}$ electron density map is shown as either green or red polyhedral surfaces for positive and negative $F_{\text{obs}} - F_{\text{calc}}$ values, respectively.

Figure 3. Unit cell of $H_2[(CF_3)_7TpFPP] \cdot 2H_2O$.

All full-occupancy non-hydrogen atoms were refined anisotropically. A detailed discussion of the disorder of the CF_3 groups and the modeling of the N- and O-bound hydrogen atoms is included in the Supporting Information. Hydrogen atoms attached to the carbon atoms belonging to the porphyrin unit were included at their geometrically estimated positions and refined as riding atoms. Once the final model of the porphyrin unit was obtained, peaks of electron density corresponding to the location of the hydrogen atoms belonging to the porphyrin pyrrole groups and water molecules of crystallization were observed. Electron density maps generated using shelXle^[43] (Figure 2) show that the electron density is localized around two cis porphyrin nitrogen atoms (namely N1 and N2). Finally, note that each $H_2[(CF_3)_7TpFPP] \cdot 2H_2O$ unit is chiral and that

there are four such, symmetry-related units, two of each handedness, within each unit cell (Figure 3). The porphyrin units are organized into layers such that all the units within a given layer have the same handedness.

Acknowledgement

This work was supported by projects 231086 and 262229 and of the Research Council of Norway (AG) and by the Advanced Light Source, Berkeley, California. The Advanced Light Source is supported by the Director, Office of Science, Office of Basic Energy Sciences, of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231.

Supporting Information

Details of spectroscopic, electrochemical, and crystallographic data. The crystallographic information files have been deposited at the Cambridge Crystallographic Data Centre and assigned the deposition numbers CCDC 1002320-1002322.

Keywords: porphyrin • tautomer • homodromic • hydrogen bonding • supramolecular

1. H. H. Limbach, J. M. Lopez, M. Kohen, *Phil. Trans. R. Soc. B* 2006, 361, 1399-1415.
2. S. J. Silvers, A. Tulinsky A J. *Am. Chem. Soc.* **1967**, 89, 3331-3337.
3. P. W. Coddling, A. Tulinsky, A J. *Am. Chem. Soc.* **1972**, 94, 4151-4157.
4. B. M. L. Chen, A. Tulinsky, *J. Am. Chem. Soc.* **1972**, 94, 4144-4151.
5. A. Tulinsky, *Ann. N.Y. Acad. Sci.* **1973**, 206, 47-69.
6. J. Almlöf, *Int. J. Quantum Chem.* **1974**, 8, 915-924.
7. A. Ghosh, J. Almlöf, P. G. Gassman, *Chem. Phys. Lett* **1991**, 186, 113-118.
8. J. Almlöf, T. H. Fischer, P. G. Gassman, A. Ghosh, M. Häser, *J. Phys. Chem.* **1993**, 97, 10964-10970.
9. T. J. Butenhoff, C. B. Moore, *J. Am. Chem. Soc.* **1988**, 110, 8336-8341.
10. T. Butenhoff, R. Chuck, H.-H. Limbach, C. B. Moore, *J. Phys. Chem.* **1990**, 94, 7847-7851.
11. H.-H. Limbach, J. Hennig, D. Gerritzen, H. Rumpel, *Faraday Discuss. Chem. Soc.* **1982**, 74, 229-243.
12. J. Braun, M. Schlabach, B. Wehrle, M. Köcher, E. Vogel, H.-H. Limbach, *J. Am. Chem. Soc.* **1994**, 116, 6593-6604.
13. A. Ghosh, J. Almlöf, *J. Phys. Chem.* **1995**, 99, 1073-1075.
14. D. K. Maity, R. L. Bell, T. N. Truong, *J. Am. Chem. Soc.* **2000**, 122, 897-906.
15. K. E. Thomas, I. H. Wasbotten, A. Ghosh *Inorg. Chem.* **2008**, 47, 10469-10478.
16. K. E. Thomas, J. Conradie, L. K. Hansen, A. Ghosh, *Eur. J. Inorg. Chem.* **2011**, 12, 1865-1870.
17. K. E. Thomas, C. M. Beavers, A. Ghosh, *Mol. Phys.* 2012, 110(19-20):2439-2444.
18. Abbreviations: TpFPP = meso-tetrakis(p-fluorophenyl)porphyrin; (CF₃)_{7/8} denotes b-heptakis/octakis(trifluoromethyl) substitution.
19. J. A. Shelnutt, X.-Z. Song, J.-G. Ma, S.-L. Jia, W. Jentzen, C. J. Medforth, *Chem. Soc. Rev.* **1998**, 27, 31-42.
20. C. J. Medforth, M. O. Senge, K. M. Smith, L. D. Sparks, J. A. Shelnutt, *J. Am. Chem. Soc.* **1992**, 114, 9859-9869.
21. I. V. Sazanovich, V. A. Galievsky, A. van Hoek, T. J. Schaafsma, V. L. Malinovskii, D. Holten, V. S. Chirvony, *J. Phys. Chem. B* **2001**, 105, 7818-7829.
22. C. L. Perrin, J. B. Nielson, *Annu. Rev. Phys. Chem.* **1997**, 48, 511-544.
23. G. A. Jeffrey, *An Introduction to Hydrogen Bonding*. Oxford, 1997.
24. S. Scheiner, *Hydrogen Bonding: A Theoretical Perspective*. Oxford, 1997.
25. T. Steiner, *Angew. Chem. Int. Ed* **2002**, 41, 48-76.
26. Y. Marechal, *The Hydrogen Bond and the Water Molecule*. Elsevier, 2006.
27. An explanatory note may be in order in the short N-H distances, 0.73(6) and 0.70(7) Å, observed here. Although it is customary to restrain N-H bond distances to 0.88 Å or 0.91 Å at 100 K, we have not done so here, in order to underscore that the NH hydrogens were assigned and refined in their current positions and that no attempts have been made to influence their relative positions.
28. W. Saenger, *Nature* **1979**, 279, 343-344.
29. W. Saenger, K. Lindner, *Angew. Chem. Int. Ed. Engl.* 1980, 19, 398-399.
30. (a) J. P. Perdew, K. Burke, M. Ernzerhof, *Phys. Rev. Lett.* **1996**, 77, 3865-3868. (b) J. P. Perdew, K. Burke, M. Ernzerhof, *Phys. Rev. Lett.* **1997**, 78, 1396-1396. (c) J. P. Perdew, M. Ernzerhof, K. Burke, *J. Chem. Phys.* **1996**, 105, 9982-9985.
31. F. Weigenda, R. Ahlrichs, *Phys. Chem. Chem. Phys.* **2005**, 7, 3297-3305.
32. (a) G. te Velde, F. M. Bickelhaupt, S. J. A. van Gisbergen, C. Fonseca Guerra, E. J. Baerends, J. G. Snijders, T. Ziegler, *J. Comput. Chem.*, **2001**, 22, 931-967; (b) C. Fonseca Guerra, J. G. Snijders, G. te Velde, E. J. Baerends, *Theor. Chem. Acc.* **1998**, 99, 391-403.
33. The PBE0 hybrid functional has often been used to study hydrogen-bonded systems. The results obtained, particularly the energetics and key geometry parameters, here were also checked against those obtained with other functionals such as OLYP and B3LYP; no significant differences between the different functionals could be discerned.
34. I. K. Thomassen, H. Vazquez-Lima, K. J. Gagnon, A. Ghosh, *Inorg. Chem.* **2015**, 54, 11493-11497.
35. G. R. Desiraju, *Angew. Chem. Int. Ed. Engl.* **1995**, 34, 2311-2327.
36. G. R. Desiraju, J. J. Vittal, A. Ramanan, *Crystal Engineering: A Textbook*. World Scientific, 2011.
37. A. Moscowitz, *Adv. Chem. Phys.* **1962**, 4, 67-112.
38. For a recent review of inherently chiral chromophores, see: M. Kwit, P. Skowronek, J. Gawronski, J. Frelek, M. Woznica, A. Butkiewicz, in *Comprehensive Chiroptical Spectroscopy, Vol. 2: Applications in Stereochemical Analysis of Synthetic Compounds, Natural Products, and Biomolecules* (Eds.: Beroova N, Polavarapu PL, Nakanishi K, Woody RW, Wiley, 2012), Ch 2, pp. 39-72.
39. M. Ethirajan, Y. Chen, P. Joshi, R. K. Pandey, *Chem. Soc. Rev.* **2011**, 40, 340-362.
40. G. M. Sheldrick, *Acta Cryst.* **2015**, A71, 3-8.
41. G. M. Sheldrick, *Acta Cryst.* **2015**, C71, 3-8.
42. SADABS: Area-Detector Absorption Correction; Siemens Industrial Automation, Inc.: Madison, WI, 1996.
43. C. B. Hübschle, G. M. Sheldrick, B. Dittrich, *J. Appl. Cryst.* **2011**, 44, 1281-1284.

Table of Contents Entry

Long the white whale of porphyrin researchers, the elusive *cis* tautomer of a free-base porphyrin has been isolated and structurally characterized for the first time.