

UC Irvine

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health

Title

A Pain in the Wrist: Stingray Envenomation

Permalink

<https://escholarship.org/uc/item/2623r2rt>

Journal

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 13(1)

ISSN

1936-900X

Authors

Tse, Adele E
Evans, David P
Counselman, Frank

Publication Date

2012

DOI

10.5811/westjem.2011.8.6878

License

<https://creativecommons.org/licenses/by-nc/4.0/> 4.0

Peer reviewed

A Pain in the Wrist: Stingray Envenomation

Adele E. Tse, MD
David P. Evans, MD
Francis L. Counselman, MD, CPE

Eastern Virginia Medical School, Department of Emergency Medicine, Norfolk, Virginia

Supervising Section Editor: Sean Henderson, MD

Submission history: Submitted August 10, 2011; Accepted August 22, 2011

Reprints available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.2011.8.6878

[West J Emerg Med. 2012;13(1):80–81.]

A 43-year-old man presented to the emergency department after he was swimming in the ocean and felt a sudden sting followed by a burning pain and muscle spasms in his right

hand. Physical exam was remarkable for tachycardia and the foreign body (Figure 1). Radiograph of his right wrist is also shown (Figure 2).



Figure 1. Patient in emergency department.



Figure 2. Radiograph of hand.

DIAGNOSIS

Stingray envenomations: Stingrays are a cartilaginous bottom-dwelling fish with a hard tail and 2 or more hard barbs, each containing a venom sac. Commonly, envenomation occurs when the victim unintentionally steps on a buried stingray and reflexes cause the ray to lash out with its tail. The venom has vasoconstrictive properties, causing possible necrosis and poor wound healing. Common presentations include local pain, muscle cramps, vomiting, and diarrhea, and rare complications include artery laceration and compartment syndrome. Since the venom is heat sensitive, immediate management includes irrigation and soaking the wound in hot water (110 to 115°F) for 30 to 90 minutes to inactivate the protein.¹ A radiograph should be obtained of the area to evaluate for embedded spines. One must be careful when removing the serrated spine. The best strategy is to open the wound with a scalpel along the nonjagged edges of the spine without disturbing the barbs. These wounds are to be closed by delayed primary closure. According to 1 study, due to the rate of infection, these patients are to be started on a quinolone and updated on tetanus.²

Address for Correspondence: David P. Evans, MD, Eastern Virginia Medical School, Department of Emergency Medicine, Rm 304, Raleigh Building, 600 Gresham Dr, Norfolk, VA 23507. E-mail: evansdp@me.com.

Conflicts of Interest: By the *WestJEM* article submission agreement, all authors are required to disclose all affiliations, funding, sources, and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

REFERENCES

1. Guenin DG, Auerbach PS. Trauma and envenomations from marine fauna. In: Tintinalli JE, Stapczynski S, Cline DM, et al, eds. *Emergency Medicine: A Comprehensive Study Guide*. 7th ed. New York, NY: McGraw-Hill; 1996:868–873.
2. Clark RF, Girard RH, Rao D, et al. Stingray envenomation: a retrospective review of clinical presentation and treatment in 119 cases. *J Emerg Med*. 2007;33:33–37.