

# UCSF

## UC San Francisco Previously Published Works

### Title

Genetic variants of Adam17 differentially regulate TGF $\beta$  signaling to modify vascular pathology in mice and humans

### Permalink

<https://escholarship.org/uc/item/26j5506g>

### Journal

Proceedings of the National Academy of Sciences of the United States of America, 111(21)

### ISSN

0027-8424

### Authors

Kawasaki, Kyoko  
Freimuth, Julia  
Meyer, Dominique S  
et al.

### Publication Date

2014-05-27

### DOI

10.1073/pnas.1318761111

Peer reviewed

# Genetic variants of *Adam17* differentially regulate TGF $\beta$ signaling to modify vascular pathology in mice and humans

Kyoko Kawasaki<sup>a</sup>, Julia Freimuth<sup>a</sup>, Dominique S. Meyer<sup>a</sup>, Marie M. Lee<sup>a</sup>, Akiko Tochimoto-Okamoto<sup>a</sup>, Michael Benzinou<sup>a</sup>, Frederic F. Clermont<sup>a</sup>, Gloria Wu<sup>a</sup>, Ritu Roy<sup>b</sup>, Tom G. W. Letteboer<sup>a,c</sup>, Johannes Kristian Ploos van Amstel<sup>c</sup>, Sophie Giraud<sup>d</sup>, Sophie Dupuis-Girod<sup>d</sup>, Gaeten Lesca<sup>d</sup>, Cornelius J. J. Westermann<sup>e</sup>, Robert J. Coffey, Jr.<sup>f</sup>, and Rosemary J. Akhurst<sup>a,g,h,1</sup>

<sup>a</sup>Helen Diller Family Comprehensive Cancer Center (HDFCCC) and Department of Anatomy, University of California, San Francisco, CA 94158-9001; <sup>b</sup>HDFCCC Biostatistical Core Facility, University of California, San Francisco, CA 94143; <sup>c</sup>Department of Medical Genetics, University Medical Centre, KC04.084.2, Utrecht, The Netherlands; <sup>d</sup>French Hereditary Hemorrhagic Telangiectasia Reference Center, Department of Medical Genetics, Lyon University Hospital, 69622 Villeurbanne, France; <sup>e</sup>St. Antonius Hospital, 3430 EM, Nieuwegein, The Netherlands; <sup>f</sup>Vanderbilt Ingram Cancer Center, Vanderbilt University, Nashville, TN 37232; and <sup>g</sup>Department of Anatomy, and <sup>h</sup>Institute of Human Genetics, University of California, San Francisco, CA 94143

Edited by Mina J. Bissell, E. O. Lawrence Berkeley National Laboratory, Berkeley, CA, and approved April 8, 2014 (received for review November 2, 2013)

**Outcome of TGF $\beta$ 1 signaling is context dependent and differs between individuals due to germ-line genetic variation. To explore innate genetic variants that determine differential outcome of reduced TGF $\beta$ 1 signaling, we dissected the modifier locus *Tgfbm3*, on mouse chromosome 12. On a NIH/OlaHsd genetic background, the *Tgfbm3*<sup>C57</sup> haplotype suppresses prenatal lethality of *Tgfb1*<sup>-/-</sup> embryos and enhances nuclear accumulation of mothers against decapentaplegic homolog 2 (*Smad2*) in embryonic cells. Amino acid polymorphisms within a disintegrin and metalloprotease 17 (*Adam17*) can account, at least in part, for this *Tgfbm3b* effect. ADAM17 is known to down-regulate *Smad2* signaling by shedding the extracellular domain of TGF $\beta$ RI, and we show that the C57 variant is hypomorphic for down-regulation of *Smad2/3*-driven transcription. Genetic variation at *Tgfbm3* or pharmacological inhibition of ADAM17, modulates postnatal circulating endothelial progenitor cell (CEPC) numbers via effects on TGF $\beta$ RI activity. Because CEPC numbers correlate with angiogenic potential, this suggests that variant *Adam17* is an innate modifier of adult angiogenesis, acting through TGF $\beta$ RI. To determine whether human ADAM17 is also polymorphic and interacts with TGF $\beta$  signaling in human vascular disease, we investigated hereditary hemorrhagic telangiectasia (HHT), which is caused by mutations in TGF $\beta$ /bone morphogenetic protein receptor genes, *ENG*, encoding endoglin (HHT1), or *ACVRL1* encoding ALK1 (HHT2), and considered a disease of excessive abnormal angiogenesis. HHT manifests highly variable incidence and severity of clinical features, ranging from small mucocutaneous telangiectases to life-threatening visceral and cerebral arteriovenous malformations (AVMs). We show that ADAM17 SNPs associate with the presence of pulmonary AVM in HHT1 but not HHT2, indicating genetic variation in ADAM17 can potentiate a TGF $\beta$ -regulated vascular disease.**

The TGF $\beta$  signaling pathway is a therapeutic target for numerous disease applications because elevated levels of ligands and signaling components have been shown to drive several pathological states, especially cancer and fibrosis (1). However, this signaling pathway is essential for formation and maintenance of the vascular system, as manifested by the vascular phenotypes of *Tgfb1*, *Tgfb1*, *Tgfb2*, *Eng*, and *Acvrl1* gene knockout mice (2), and by congenital vasculopathies caused by mutations in orthologous human genes (3–5). Therefore, therapeutic downmodulation of TGF $\beta$  signaling has the potential to cause undesirable vascular outcomes (6), making investigation of the molecular mechanisms underlying such phenotypes clinically important.

Studies in mice have shown that, when the TGF $\beta$ 1 pathway is disrupted, innate genetic differences between strains result in variable penetrance and severity of vascular phenotypes (7). Human Mendelian disorders of TGF $\beta$  and bone morphogenetic

protein (BMP) signaling also show variable penetrance and severity that may be attributed, at least in part, to genetic variation within modifier loci (8, 9). One example is hereditary hemorrhagic telangiectasia (HHT), which is caused by loss-of-function mutations in *ENG*, *ALK1*, or *SMAD4* that encode components of the TGF $\beta$ /BMP signaling pathways (4, 5, 10). HHT patients can suffer vascular malformations in multiple organ systems. They initially present with recurrent epistaxis but subsequently develop multiple cutaneous and mucosal telangiectases. Bleeding from gastrointestinal telangiectases can lead to chronic, debilitating, and treatment-refractory anemia. Some patients develop more severe arteriovenous malformations (AVMs) in the lung, brain, and/or liver, which can lead to severe complications. Lung AVMs occur in ~30–50% of patients with HHT, causing right-to-left shunting due to a compromised pulmonary capillary filter, and potentially leading to life-threatening stroke (11).

Here, we characterized a TGF $\beta$  modifier locus on proximal mouse chromosome 12 (12) and found that, like mouse modifiers of common disease phenotypes (13, 14), *Tgfbm3* is genetically complex with positive and negative elements that potentiate or suppress lethal prenatal *Tgfb1*<sup>-/-</sup> vascular dysgenesis. A C57

## Significance

Down-regulation of TGF $\beta$  signaling is a therapeutic strategy for several disease applications, particularly cancer and fibrosis. However, TGF $\beta$  blockade may have adverse effects because TGF $\beta$  signaling is necessary for formation and maintenance of normal blood vessels. Because side effects may vary between individuals due to innate differences in genetic makeup, it is important to find which interindividual genetic variants regulate responses to reduced TGF $\beta$  signaling and how these operate at the molecular level. Here we identified polymorphic variants of mouse a disintegrin and metalloprotease 17 that differentially regulate TGF $\beta$  signaling output and influence the severity of *Tgfb1*-dependent vascular pathology. This has relevance to risk assessment for clinical manifestations in TGF $\beta$ -driven diseases, as well as for prediction of desirable and undesirable responses to anti-TGF $\beta$  therapy.

Author contributions: K.K., F.F.C., T.G.W.L., C.J.J.W., and R.J.A. designed research; K.K., J.F., D.S.M., M.M.L., A.T.-O., F.F.C., G.W., and T.G.W.L. performed research; J.K.P.V.A., S.G., S.D.-G., G.L., C.J.J.W., and R.J.C. contributed new reagents/analytic tools; K.K., J.F., D.S.M., M.M.L., A.T.-O., M.B., F.F.C., R.R., T.G.W.L., and R.J.A. analyzed data; R.J.A. directed collaborations; and K.K., J.F., G.L., C.J.J.W., R.J.C., and R.J.A. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

<sup>1</sup>To whom correspondence should be addressed. E-mail: rakhurst@cc.ucsf.edu.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1318761111/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1318761111/-DCSupplemental).

versus NIH haplotype over a subinterval of *Tgfbm3*, termed *Tgfbm3b*, functions to support *Tgfb1*-independent developmental angiogenesis and, in adults, potentiates circulating endothelial progenitor cell (CEPC) numbers, a surrogate marker for angiogenic capacity (15). We show that *Tgfbm3b* encodes a hypomorphic variant of a disintegrin and metalloprotease 17 (ADAM17) that, due to ineffectual down-regulation of TGF $\beta$ RI signaling, potentiates TGF $\beta$ -mothers against decapentaplegic homologs 2 and 3 (Smad2/3) signaling downstream of the ligand. We also provide evidence that genetic variation within human *ADAM17* associates with the presence of potentially life-threatening pulmonary AVMs in patients with HHT1 carrying *ENG* mutations, giving credence to the notion that genetic variation at *ADAM17* may determine severity of TGF $\beta$  pathway-regulated disease in humans.

## Results


***Tgfbm3b*<sup>C57</sup> Is a Suppressor of *Tgfb1*<sup>-/-</sup> Prenatal Lethality.** *Tgfb1*<sup>-/-</sup> mice are not viable. They either die prenatally from vascular dysgenesis 9.5–10.5 days postcoitum (16), or postnatally from lethal multifocal inflammation (17). Previously we mapped variant genomic loci that determine variable penetrance of *Tgfb1*<sup>-/-</sup> prenatal lethality (7, 9, 12, 18). We showed that a C57 *Tgfbm3*<sup>C57</sup> haplotype on proximal chromosome 12 accentuates prenatal lethality of NIH.*Tgfb1*<sup>-/-</sup> embryos, albeit that the *Tgfbm3*<sup>NIH</sup> allele, acting alone, is insufficient to suppress C57.*Tgfb1*<sup>-/-</sup> embryo lethality (12). To further dissect *Tgfbm3*, we used marker-assisted backcross to generate *Tgfb1*<sup>+/-</sup> congenic sublines carrying different extents of C57 over the *Tgfbm3* locus, which were defined by mapping using SSR and SNP markers (Fig. 1 A and B and Table S1). As seen within other genetic modifier loci (13, 14, 19), we found that *Tgfbm3*<sup>C57</sup> is complex and possesses two modifying elements, a proximal enhancer (*Tgfbm3a*<sup>C57</sup>; centromere to *Lpin1* gene) and a distal suppressor (*Tgfbm3b*<sup>C57</sup>; *Ntsr2* to *Grhl1*) of *Tgfb1*<sup>-/-</sup> prenatal lethality (Fig. 1C).

**Genetic Variation at *Tgfbm3b* Modulates Nuclear Localization of SMAD2.** We focused on analysis of *Tgfbm3b*, because the ability to suppress *Tgfb1*<sup>-/-</sup> embryo-lethality suggests that this locus acts


downstream of TGF $\beta$ 1 to compensate for reduced ligand levels, whereas potentiation of a lethal phenotype might be achieved through a variety of mechanisms, not necessarily related to TGF $\beta$  signaling. Indeed, classical forward genetic screens for suppressors of gene knockout phenotypes have been used extensively in lower organisms to identify novel components of specific signaling pathways (20). To investigate the effect of variant *Tgfbm3b* on downstream canonical TGF $\beta$  signaling, we examined the nucleocytoplasmic distribution of endogenous Smad2 in wild-type primary mouse embryo fibroblasts (MEFs) derived from NIH.*Tgfbm3*<sup>C57</sup> congenic lines 4 and 6, which differ over an ~1.6-Mb interval of unique gene-rich DNA within *Tgfbm3b* (Fig. 1 A and B). Wild-type NIH and line 4 MEFs, homozygous for the *Tgfbm3b*<sup>NIH</sup> allele, showed predominantly cytoplasmic localization of Smad2, which relocated to the nucleus upon TGF $\beta$  treatment (Fig. 2 A–D, and G). In contrast, wild-type line 6 MEFs, which bear the C57 *Tgfbm3b*<sup>C57</sup> allele, exhibited nuclear Smad2 staining before addition of TGF $\beta$ . The differential in nuclear Smad2 staining between line 4 and line 6 mice was further accentuated after TGF $\beta$  treatment (Fig. 2 E–G). The C57 allele of *Tgfbm3b* appeared to act in a recessive manner, because wild-type MEFs generated by intercrossing NIH.C57-*Tgfbm3* line 4 with line 6 mice (NIH.*Tgfbm3b*<sup>C57/NIH</sup>), showed the same Smad2 distribution as those from line 4 MEFs (NIH.*Tgfbm3b*<sup>NIH/NIH</sup>) that bear only the NIH allele. *Tgfbm3b*<sup>C57</sup> therefore appears to be a recessive *Tgfb1* modifier that acts downstream of TGF $\beta$ 1 to modulate Smad2 levels.

***Tgfbm3b* Is a Modifier of Circulating Endothelial Progenitor Cell Numbers.** Because *Tgfb1* and *Tgfbm3b* interact to regulate vascular development in utero (Fig. 1C), we postulated that these loci may interact to regulate postnatal angiogenesis. In the adult, angiogenesis is uncommon except during wound healing and in disease states, such as cancer and retinopathy (21). Postnatal angiogenesis occurs predominantly by sprouting from existing blood vessels, but can be supported by endothelial progenitor cells (EPCs) that home to the angiogenic site after release from the bone marrow (22) or that appear by vasculogenesis local to the site of injury (23). Several studies have shown a strong association between CEPCs, active angiogenesis, and vascular outcomes in mice and humans (24–26). Importantly, angiogenic capacity and CEPC numbers correlate, and both are determined by innate genetic variation (15). However, as yet no variant genetic loci that modulate CEPC levels have been characterized.

Because *Tgfb1* has been shown to be necessary for a robust CEPC response to injury and associated reparative angiogenesis (26), we postulated that variants of *Tgfbm3b* may interact with *Tgfb1* to regulate CEPC levels during postnatal angiogenesis. FACS analysis was used to monitor adult CEPC numbers in NIH and congenic NIH.C57-*Tgfbm3* lines 4 and 6 following s.c. implantation of a highly angiogenic syngeneic carcinoma cell line. All three mouse lines showed elevation in CEPC (CD45<sup>-</sup>, CD117<sup>+</sup>, CD13<sup>+</sup>) and circulating endothelial cell (CEC) (CD45<sup>-</sup>, CD117<sup>-</sup>, CD13<sup>+</sup>) numbers, 1–2 d following administration of this angiogenic stimulus (Fig. 3 A and B). NIH and line 4 mice, both homozygous NIH for the major *Tgfbm3b*<sup>NIH</sup> allele, showed similar profiles of CEPC and CEC induction, with only a weak and transient increase in CEPC numbers. In contrast, line 6 mice, homozygous for the *Tgfbm3b*<sup>C57</sup> allele, which suppresses *Tgfb1*<sup>-/-</sup> embryo lethality, elicited a much greater CEPC response than NIH or line 4 mice (Fig. 3 A and B). Significantly, differences in CEPC numbers between mice that harbor the two alternative *Tgfbm3b* genetic variants were not paralleled by differentials in total circulating CD45<sup>+</sup> cells, or specifically in CD45<sup>+</sup> CD11b<sup>+</sup> F4.80<sup>+</sup> Gr1<sup>-</sup> monocytes or CD45<sup>+</sup> CD11b<sup>+</sup> Gr1<sup>+</sup> myeloid-derived suppressor cells (Fig. S1), implying that the effects observed were specific to EPCs. Importantly, loss of a single *Tgfb1* allele severely reduced the accentuated CEPC and CEC responses to the angiogenic stimulus seen in line 6 mice (NIH.*Tgfbm3b*<sup>C57/C57</sup>) (Fig. 3 C and D), demonstrating that TGF $\beta$ 1 is required for the activity of *Tgfbm3b*<sup>C57</sup> in this assay and



**Fig. 1.** *Tgfbm3* is a complex locus harboring an enhancer (*Tgfbm3a*) and suppressor (*Tgfbm3b*) of *Tgfb1*<sup>-/-</sup> prenatal lethality. (A) Cartoon of *Tgfbm3* on proximal mouse chromosome 12, indicating positions of genes within “islands” of gene-rich unique DNA (black blocks) interspersed by extensive regions (megabases) of repetitive DNA (white blocks). The human *TGFBM3* syntenic region (2p23–25) lacks repetitive DNA and is thus far shorter (Fig. S3). Asterisks mark genes with amino acid polymorphisms between NIH and C57. (B) C57 regions within NIH.C57-*Tgfbm3* congenic lines 4 and 6, aligned to A. Black blocks indicate C57; white blocks indicate NIH genomic DNA. The C57→NIH transition in line 4 occurs between rs49671069 and rs3724468, located just distal to *Lpin1*, and that of line 6 occurs between rs29180890 and rs31626421, located just distal to *Taf1b*. (C) Congenic test cross data generated by intraline NIH.*Tgfb1*<sup>+/-</sup> intercrosses of NIH, NIH.C57-*Tgfbm3a* (line 4), and NIH.C57-*Tgfbm3a-Tgfbm3b* (line 6). Survival to birth (STB) rates were estimated from the ratio of *Tgfb1*<sup>-/-</sup> neonates to *Tgfb1*<sup>+/+</sup> neonates on the day of birth. \**P* = 0.07; \*\**P* < 0.01.



**Fig. 2.** Early passage NIH MEFs homozygous for *Tgfbm3b*<sup>C57</sup> show higher basal and inducible nuclear Smad2 levels than those with *Tgfbm3b*<sup>NIH</sup>. (A–F) Smad2 staining of MEFs derived from wild-type mice of a parental NIH background (A and B), or on the line 4 (C and D) or line 6 (E and F) congenic backgrounds, before (A, C and E) and 1 h after (B, D, and F) stimulation with TGFβ 0.5 ng/mL. (G) Quantification of nuclear Smad2 intensity. \**P* ≤ 0.05. Each experiment was independently reproduced three times with three technical replicates within each experiment.

giving support to a model of genetic interaction between *Tgfb1* and *Tgfbm3b* in regulating CEPC levels. We surmise that genetic variation within *Tgfbm3b* regulates CEPC numbers in response to an angiogenic stimulus. We believe that this is the first report of an endogenous polymorphic genetic modifier of CEPC number in vivo.


**Innate Genetic Variation of *Adam17* Within *Tgfbm3b* Modifies TGFβ Signaling.** Whole genome SNP analysis showed that lines 4 and 6 were both ≥95% homozygous NIH, including the *Tgfbm1* and *Tgfbm2* loci (Table S1). It is therefore unlikely but not impossible that differences in TGFβ biology observed between congenic lines 4 and 6 were influenced by genetic variation at unlinked loci. Definitive proof that *Tgfbm3b* directly influences TGFβ signaling therefore required analysis of the variant genes within that locus. We sequenced the coding regions of candidates located within *Tgfbm3b* (Fig. 1A). Three genes, *Greb1*, *E2f6*, and *Adam17*, each showed amino acid polymorphisms between C57 and NIH mice (Table S2). *E2f6* possesses a single *p.Leu10Gln* polymorphism, but comparative functional analysis of the two protein variants showed no significant influence on TGFβ signaling in vitro.

*Adam17* possesses two nonsynonymous coding SNPs. This gene encodes a disintegrin and metalloprotease 17 (ADAM17), also termed tumor necrosis factor α converting enzyme (TACE), a transmembrane metalloprotease involved in cell surface shedding and activation of a number of growth factors, including TNFα, EGF-R ligands, Notch1, and Notch4 (27). A *p.Asp113Asn* variant in the C57 allele introduces a charge change at an evolutionarily conserved residue within the ADAM17 Pro domain. This domain is involved in ADAM17 polypeptide processing, subcellular trafficking, and activation (Fig. 4A and B). The ADAM17 *p.Asp113Asn* variant has only been observed in one other *Mus musculus domesticus* strain, NZW/LacJ, and two wild-derived *Mus musculus* subspecies, *Mus musculus molossinus* and *Mus musculus musculus* (Fig. 4B). The C57 ADAM17 *p.Ileu613Val* variant (Fig. 4A) is located in the cysteine-rich domain. This is a relatively common variant in laboratory mouse strains and is not evolutionarily conserved (Fig. 4B).

ADAM17 has been shown to cleave the TGFβ type I receptor, effectively reducing Smad2 signaling (28). We therefore tested the comparative activity of ectopically expressed ADAM17 variants to modulate TGFβ responsiveness in NIH 3T3 fibroblasts using a CAGA<sub>12</sub>-luciferase assay that reports on Smad2/3 transcriptional activity. As predicted, ectopic expression of the major


(NIH) ADAM17 isoform reduced TGFβ-dependent Smad2/3 transcriptional output (Fig. 4C), presumably by shedding TβRI (28). Ectopic C57 ADAM17 protein levels were similar to those of the NIH variant (Fig. S2), but showed a 50% attenuation in the ability to reduce TGFβ-induced CAGA<sub>12</sub>-luciferase activity (Fig. 4C), suggesting that this isoform is hypomorphic with respect to TβRI sheddase activity. Interestingly, ADAM17 polypeptides harboring single C57 amino acid variants showed intermediate levels of CAGA<sub>12</sub>-luciferase activity between that of the C57 and NIH alleles (Fig. 4C), suggesting that both amino acid variants contribute to the overall decrease in TβRI sheddase activity. We conclude that the C57 genome encodes a recessive *Adam17* variant that is hypomorphic for down-regulation of TβRI, resulting in a hyperactivated state of TβRI-Smad2/3 signaling that, in the presence of TGFβ2, TGFβ3, or maternal TGFβ1, may contribute to the rescue of line 6 *Tgfb1*<sup>-/-</sup> embryos from lethal vascular dysgenesis (Fig. 5).

**ADAM17 Inhibition Potentiates CEPC Levels by Enhancing TGFβ Signaling.** We postulated that genetic variation of *Adam17* is also responsible for the modulatory effects of *Tgfbm3b* on CEPC levels. Based on the observation that *Tgfbm3b*<sup>C57</sup>, harboring the hypomorphic *Adam17* allele, potentiated CEPC levels in response to the angiogenic stimulus, we hypothesized that pharmacological ADAM17 inhibition would accentuate release of CEPCs from the bone marrow. Indeed, systemic treatment of mice with an ADAM17-specific small molecule inhibitor, TMI-005 (29), enhanced CEPC and CEC numbers 16- to 30-fold (Fig. 6A–C). This TMI-005-induced increase in CEPC response was observed even in mice carrying the hypomorphic *Adam17*<sup>C57</sup> allele (line 6, Fig. 6B and C), consistent with the fact that the hypomorphic C57 variant retains significant residual ADAM17 activity. Based on our genetic interaction analysis, we hypothesized that the effect of hypomorphic ADAM17 on CEPC and CEC responses in vivo was mediated by reduced shedding and consequent increased activation of the TGFβ type I receptor. In concordance with this hypothesis, the large induction in CEPC numbers



**Fig. 3.** *Tgfbm3b* is a TGFβ1-dependent modifier of adult CEPC response to angiogenesis. Three mouse strains: (i) NIH, (ii) congenic NIH.*Tgfbm3a* line 4, and (iii) NIH.*Tgfbm3ab* line 6, were implanted with syngeneic CarB cells ( $2 \times 10^5$ ) expressing high VEGF and SDF1 levels, to provide an angiogenic stimulus. Blood was harvested at the indicated time points post-CarB implantation, and assayed by FACS analysis for (A) CD45<sup>+</sup>, CD13<sup>+</sup>, CD117<sup>-</sup> CECs, and (B) (CD45<sup>+</sup>, CD13<sup>+</sup>, and CD117<sup>+</sup>) CEPCs. Similar experiments were performed using NIH.*Tgfbm3ab* line 6 congenic mice that were either wild type or haploinsufficient for *Tgfb1* (C and D). Each time point represents data from four mice per genotype, and each experiment was independently reproduced three times.





**Fig. 4.** *Adam17* polymorphic variants differentially regulate canonical TGFβ-SMAD signaling output. (A) Position of C57 amino acid substitutions within the ADAM17 protein. (B) Evolutionary conservation of ADAM17 amino acid residues 113 and 594/613 from chick to human. (C) TGFβ1/Smad2/3-mediated transcriptional responses of NIH 3T3 fibroblasts transfected with expression vectors encoding NIH-*Adam17*, *Adam17*<sup>Asp113Asn</sup>, *Adam17*<sup>Ileu613Val</sup>, or C57. *Adam17*<sup>Asp113Asn,Ileu613Val</sup>. Cells were transfected with the indicated *Adam17* expression constructs together with a Smad2/3-responsive pCAGA-luciferase construct. Note that exogenous ADAM17 expression was in vast excess to that of endogenous NIH 3T3 *Adam17* (Fig. S2). Activation of luciferase was assayed 24 h after addition, or not, of 1 ng/mL TGFβ. \**P* ≤ 0.05, \*\**P* ≤ 0.01. Each experiment was independently reproduced three times with three technical replicates within each experiment.

observed in the presence of TMI-005 was completely ablated by Ly2109761 (Fig. 6C), a small molecule inhibitor of TβRI kinase (30). We therefore conclude that the stimulatory effect of ADAM17 inhibition on CEPC levels is mediated by enhanced TβRI signaling.

**Genetic Variation Within Human ADAM17 Associates with Pulmonary Vascular Lesions in HHT.** Because *Adam17* interacts with *Tgfb1* to modulate prenatal vascular development and adult CEPC levels in mice, we postulated that genetic variation in *ADAM17* may modify TGFβ-regulated vascular disorders in humans. HHT is such a disorder, caused by haploinsufficiency of *ENG*, *ACVRL1*, or *SMAD4*, and considered to be a disease of excess angiogenesis and/or altered endothelial cell fate, resulting from perturbations in the TGFβ/BMP signaling pathways (31–33). Although HHT is rare, affecting 1 in 5–10,000 people worldwide, studies of this disease are valuable to our overall understanding of human vascular biology. We hypothesized that genetic variation within human *ADAM17* may influence clinical severity of HHT, one feature of which is development of pulmonary AVMs. We therefore investigated genetic association between SNPs within *ADAM17* and the presence of pulmonary AVM in HHT mutation carriers.


We genotyped six common independent *ADAM17* tagSNPs (*r*<sup>2</sup> < 0.7) in a cohort of 401 Dutch and Dutch Antillean HHT mutation carriers, together with 235 unaffected first degree relatives. Gamete competition analysis (34), a modified version of the transmission distortion test, revealed that three of the six *ADAM17* SNPs showed significant association between the minor allele and presence of pulmonary AVM in HHT mutation carriers (*P* < 0.05, Table S3: *rs10495565*, *rs12474540*, and *rs17524594*). When Dutch patients were stratified by HHT subtype, significant association with pulmonary AVM was observed for HHT1 (mutant *ENG*), but not for HHT2 (mutant *ACVRL1*). Importantly, of 175 additional tagSNPs that spanned the human 2p25 genomic interval syntenic to *Tgfbm3b* (750 kb proximal and 1 Mb distal from *ADAM17*; Fig. S3) only one other SNP (*rs2304401*) showed significant genetic association to lung AVM in the Dutch cohort. In humans, a positive genetic association

requires confirmation in an independent population. We therefore genotyped *rs10495565* and *rs12474540* (*ADAM17*), as well as *rs2304401* in an additional French cohort of patients with HHT (*n* = 222). Both *ADAM17* SNPs showed a strong trend toward genetic association between minor alleles with pulmonary AVM in HHT1 (*n* = 75, Table S4; *P* = 0.067) but not HHT2 mutation carriers (*n* = 147), with the statistical power being limited by the small size of this French HHT cohort. *rs2304401* showed no genetic association. In conclusion, despite the statistical power constraints of working with small populations with a rare genetic disorder, the data strongly suggest that differential interactions between *ADAM17* variants and the TGFβ signaling pathway can influence severity of vascular pathology in humans.


**Discussion**

In the current study, we report the presence of a polymorphic genetic locus in mice that regulates both *Tgfb1*-dependent prenatal and postnatal vascular biology. We show that the C57 allele of *Tgfbm3b*, a component of the larger *Tgfbm3* locus on mouse chromosome 12, is a suppressor of *Tgfb1*<sup>-/-</sup> prenatal lethality and modulates CEPC levels in adults, a surrogate for angiogenic capacity. Although these genetic data might be confounded by the presence of a low level of genetic heterogeneity elsewhere in the genome (9), we provide molecular and pharmacological evidence that genetic variation of the *Adam17* gene within *Tgfbm3b* is at least in part responsible for this modifier effect.

It is notable that the C57 mouse strain has a low angiogenic potential and low CEPC levels (15), as well as manifesting complete penetrance of prenatal lethality of *Tgfb1*<sup>-/-</sup> embryos from vascular dysgenesis (7, 9). It is therefore counterintuitive that the C57 allele of *Tgfbm3b* would be proangiogenic, as manifested by enhanced adult CEPC numbers and suppression of *Tgfb1*<sup>-/-</sup> prenatal lethality in line 6 compared with line 4 mice. This may be explained if, in pure C57 mice, the proangiogenic activity of *Tgfbm3b*<sup>C57</sup> is masked by stronger angiogenesis suppressor genes located elsewhere in the C57 genome, such as within *Tgfbm3a*. This paradox highlights the complexity of genetic interactions between endogenous variants observed in polygenic traits, such as angiogenesis, as has previously been documented in cancer susceptibility loci (35–37). In the current study, the confounding effects of unlinked modifier genes are largely, although not entirely, eliminated by utilization of the NIH congenic lines that differ only across *Tgfbm3*.



**Fig. 5.** Model depicting effect of ADAM17 variant on the canonical TGFβ signaling pathway. TGFβ ligand induces heterologomerization of TβRII and TβRI, activation of TβRI by TβRII, and phosphorylation of Smad2 by TβRI kinase. P-Smad2 oligomerizes with Smad4 to shuttle to the nucleus and instigate a Smad-mediated transcriptional response. The model proposes differential shedding and consequent inactivation of TβRI by the NIH versus C57 variants of ADAM17. The C57 ADAM17 isoform has relatively weak inhibitory activity against TβRI, consequently elevating nuclear P-Smad2/3 and Smad2/3-mediated transcription in cells harboring this variant.



**Fig. 6.** T $\beta$ RI kinase activity mediates the effect of hypomorphic ADAM17 on CEPC numbers in vivo. Mice were treated, or not, with 50 mg/kg of a pharmacological ADAM17 inhibitor, TMI-005, and/or 100mg/kg of an orally available T $\beta$ RI kinase inhibitor, Ly2109761, before and during induction of angiogenesis, as described in the legend to Fig. 3. Mice were dosed twice a day with drug or vehicle, from one day before CarB implantation. Blood samples were collected before tumor implantation (0 h) and 24 h post implantation. CEPC and CEC numbers were assayed as described in Fig. 3. (A) CEC and (B) CEPC numbers, in response to inhibition of ADAM17 by TMI-005. Elevated induction of (C) CEPCs in congenic mice was neutralized by treatment with the T $\beta$ RI kinase inhibitor, Ly2109761. Each time point reports the mean of four mice per experiment, and each experiment was replicated three times. \* $P \leq 0.05$ , \*\* $P \leq 0.01$ , \*\*\* $P \leq 0.001$ .

It seems counterintuitive that MEFs from pure NIH mice, which have a relatively high *Tgfb1*<sup>-/-</sup> embryo survival rate, have lower nuclear Smad2 levels compared with line 6 MEFs (NIH. *Tgfbm3b*<sup>C57</sup>). Moreover, adult NIH mice did not demonstrate an accentuated CEPC response to the angiogenic stimulus, despite a high *Tgfb1*<sup>-/-</sup> embryo survival rate. These facts might be reconciled if NIH alleles within the proximal *Tgfbm3a* haplotype support prenatal *Tgfb1*<sup>-/-</sup> development in a Smad2-independent fashion that does not influence CEPC levels in the adult. In the current study we restricted our analysis to *Tgfbm3b* because this locus not only modulated adult CEPC levels, but also suppressed *Tgfb1*<sup>-/-</sup> embryo lethality, suggesting a direct influence of *Tgfbm3b* on the TGF $\beta$  signaling pathway.

We present evidence that functional polymorphism within ADAM17 is a major contributor to the modifier effects of *Tgfbm3b*. ADAM17 is a sheddase that cleaves many substrates and, by so doing, is involved in processing and activating TNF $\alpha$ , Notch, EGFR, and amphiregulin, as well as suppressing T $\beta$ RI signaling (27, 28). We found that the C57 *Adam17* allele harbors two amino acid polymorphisms, *p.Asp113*→*Asn* and *p.Ileu613*→*Val*, which together result in a protein that is hypomorphic in down-modulating TGF $\beta$  signaling, culminating in higher levels of nuclear Smad2 signaling downstream of T $\beta$ RI and enhanced Smad2/3-dependent transcriptional reporter activity. This effect may compensate somewhat for lack of TGF $\beta$ 1 ligand in *Tgfb1*<sup>-/-</sup> embryos, allowing more of them to undergo normal embryonic development, possibly supported by TGF $\beta$ 2, TGF $\beta$ 3, or maternal TGF $\beta$ 1.

Intriguingly, we found that inheritance of the hypomorphic C57 *Adam17* variant or pharmacological inhibition of ADAM17 results in higher CEPC numbers in mice. Many studies have found that CEPC numbers correlate with productive angiogenesis and vascular stabilization. CEPC numbers have been

positively associated with a reduced risk of vascular disease, as well as a better outcome after a cardiovascular incident (38). In stroke victims, CEPC numbers are lower compared with control subjects, whereas higher CEPC numbers correlate with a more favorable outcome (24, 25). An inverse correlation between CEPC number and aneurysm size has been reported in patients with idiopathic thoracic ascending aortic aneurysm (TAAA) (39), an intriguing observation because TAAD (TAAA with dissections) is a major health concern in patients who have mutations in TGF $\beta$  signaling pathway genes (40–42). The fact that genetic reduction of TGF $\beta$ 1 or pharmacological inhibition of T $\beta$ RI signaling neutralized the effects of ADAM17 hypoactivity on CEPC numbers, not only further illustrates the importance of interaction between ADAM17 activity and TGF $\beta$  signaling in vivo, but suggests that many of the effects of ADAM17 on angiogenesis are mediated via TGF $\beta$ RI.

It is likely that these amino acid substitutions in C57 ADAM17 alter microdomains that differentially affect substrate specificities. In fact, the *p.Ileu613Val* variant is located within the cysteine-rich substrate recognition domain (43); thus, variation in *Adam17 p.Asn113-Val613* may preferentially affect T $\beta$ RI signaling over that of other substrates, such as TNF $\alpha$ . If the C57 ADAM17 variant reduced activation of EGFR, TGF $\alpha$  or TNF $\alpha$  processing, wild-type C57 mice might be expected to present with dramatic phenotypes, such as perinatal lethality (44) or inflammatory skin and bowel disease (45), observed in mice or humans lacking functional ADAM17, or with wavy fur and open eyelids, as observed in a more severe murine ADAM17 hypomorph (46), which is clearly not the case.

Notch is another ADAM17 substrate, and is a vascular quiescence factor with a known involvement in TGF $\beta$  and ALK1 signaling (32, 33). It is thus possible that reduction in Notch activation, due to reduced ADAM17 activity, may also contribute to hyperelevation of CEPC numbers and increased angiogenesis observed in the NIH.C57-*Adam17* and TMI-005-treated mice (32). Nevertheless, the fact that specific pharmacological inhibition of T $\beta$ RI completely suppressed the TMI-005-induced CEPC response, argues in favor of a major role of TGF $\beta$  signaling in mediating the effects of hypomorphic ADAM17 on CEPC number. We conclude that TGF $\beta$ 1-T $\beta$ RI-Smad2/3 signaling plays an important proangiogenic role not only during embryogenesis, but in release of adult bone-marrow-derived CEPCs to the blood, and that these processes are modulated by interaction between T $\beta$ RI and genetic variants of *Adam17*.

Finally, in the current study, we provide evidence that genetic variation in human *ADAM17* may influence clinical outcome of a TGF $\beta$ -regulated disease, HHT. We show genetic association between *ADAM17* SNPs and the presence of pulmonary AVMs in patients with HHT, suggesting a contribution to the risk of lung AVM. The human study is limited by the small cohort size necessitated by working with a rare disease, and the SNPs that drive association between *ADAM17* and PAVM in HHT remain to be identified. However, the fact that three of six independent SNPs ( $r^2 < 0.7$ ) showed significant genetic association in the Dutch population as well as marginal association in French HHT is compelling. Human variation in *ADAM17* at the specific amino acid residues orthologous to those of the C57 variant (113 and 613) has not been reported to date. However, clusters of “potentially detrimental” amino acid variants lie close to the orthologous polypeptide positions of the C57 mouse variants, between residues 97–120 (Pro domain) and 505–610 (cysteine-rich domain) (Polyphen and SIFT analysis in dbSNP). These protein variants may contribute to differences in TGF $\beta$  signaling within the human population. Intriguingly, in both the Dutch and French populations, genetic association was only observed within HHT1 but not in HHT2. This may be of biological significance, because *ACVRL1*, mutated in HHT2, encodes a serine threonine kinase signaling receptor that is primarily activated by BMP9 rather than TGF $\beta$ , whereas *ENG*/endoglin, mutated in HHT1, is involved in presentation of various ligands to their respective type II receptors, including TGF $\beta$ 1 and -3, activins, and BMPs.

In conclusion, here we show that genetic variants of mouse *Adam17* can profoundly influence the outcome of vascular phenotypes in vivo, both pre- and postnatally. This explains one component of the genetic complexity that determines individualized responses to reduced TGF $\beta$ 1 signaling. It is likely that human genetic variation in *ADAM17*, by differential effects on T $\beta$ RI, will also determine risks and outcomes of more common human diseases, as well as clinical responses to TGF $\beta$  pathway-targeted therapies (1).

## Materials and Methods

The parental mouse strains under study were NIH/OlaHsd and C57BL/6NTac, each carrying a *Tgfb1*<sup>Tm1n</sup> allele (17). NIH.C57-*Tgfbm3* congenic mouse lines were generated by repeated backcross (more than seven generations) of NIH.C57.F1 mice to NIH/OlaHsd, with selection for the *Tgfb1*<sup>Tm1n</sup> allele on chromosome (Chr) 7 and for C57 SSR markers at *Tgfbm3*. Congenic lines

were genotyped at *Tgfbm1* (Chr 5) and *Tgfbm2* (Chr 1) to confirm homozygosity for NIH and exclude the possibility of confounding effects due to genetic variation at these *Tgfbm* loci. Further methods are available in *SI Materials and Methods*.

**ACKNOWLEDGMENTS.** We thank Mamie Higgins and Minh Thu Luu for outstanding technical assistance and Jonathan Yingling for supplying Ly2109761. Human genotyping was performed by the Helen Diller Family Comprehensive Cancer Center Genome Core Facility. This work was funded by National Institutes of Health Grants HL078564, GM60514, CA116019, and an award from the HHT International Foundation (to R.J.A.). K.K. received a fellowship from the Uehara Memorial Foundation, Japan. F.F.C. received a fellowship from the Belgian American Educational Foundation; F.F.C. and M.B. were both recipients of American Heart Association fellowships; D.S.M. was the recipient of a Swiss National Foundation fellowship, and T.G.W.L. was the recipient of a fellowship from the Ter Meulen Fund, The Netherlands.

- Akhurst RJ, Hata A (2012) Targeting the TGF $\beta$  signalling pathway in disease. *Nat Rev Drug Discov* 11(10):790–811.
- Pardali E, Goumans MJ, ten Dijke P (2010) Signaling by members of the TGF- $\beta$  family in vascular morphogenesis and disease. *Trends Cell Biol* 20(9):556–567.
- Loeys BL, et al. (2006) Aneurysm syndromes caused by mutations in the TGF- $\beta$  receptor. *N Engl J Med* 355(8):788–798.
- McAllister KA, et al. (1995) Six novel mutations in the endoglin gene in hereditary hemorrhagic telangiectasia type 1 suggest a dominant-negative effect of receptor function. *Hum Mol Genet* 4(10):1983–1985.
- Johnson DW, et al. (1996) Mutations in the activin receptor-like kinase 1 gene in hereditary haemorrhagic telangiectasia type 2. *Nat Genet* 13(2):189–195.
- Anderton MJ, et al. (2011) Induction of heart valve lesions by small-molecule ALK5 inhibitors. *Toxicol Pathol* 39(6):916–924.
- Bonyadi M, et al. (1997) Mapping of a major genetic modifier of embryonic lethality in TGF  $\beta$  1 knockout mice. *Nat Genet* 15(2):207–211.
- Kang HC, et al. (2013) Multiple self-healing squamous epithelioma (MSSE): Rare variants in an adjacent region of chromosome 9q22.3 to known *TGFBF1* mutations suggest a digenic or multilocus etiology. *J Invest Dermatol* 133(7):1907–1910.
- Benzinou M, et al. (2012) Mouse and human strategies identify *PTPN14* as a modifier of angiogenesis and hereditary haemorrhagic telangiectasia. *Nat Commun* 3:616.
- Gallione CJ, et al. (2006) SMAD4 mutations found in unselected HHT patients. *J Med Genet* 43(10):793–797.
- Showlin CL (2010) Hereditary haemorrhagic telangiectasia: Pathophysiology, diagnosis and treatment. *Blood Rev* 24(6):203–219.
- Tang Y, et al. (2005) Epistatic interactions between modifier genes confer strain-specific redundancy for *Tgfb1* in developmental angiogenesis. *Genomics* 85(1):60–70.
- Morel L, Blenman KR, Croker BP, Wakeland EK (2001) The major murine systemic lupus erythematosus susceptibility locus, *Sle1*, is a cluster of functionally related genes. *Proc Natl Acad Sci USA* 98(4):1787–1792.
- Samuelson DJ, et al. (2007) Rat *Mcs5a* is a compound quantitative trait locus with orthologous human loci that associate with breast cancer risk. *Proc Natl Acad Sci USA* 104(15):6299–6304.
- Shaked Y, et al. (2005) Genetic heterogeneity of the vasculogenic phenotype parallels angiogenesis: Implications for cellular surrogate marker analysis of antiangiogenesis. *Cancer Cell* 7(1):101–111.
- Dickson MC, et al. (1995) Defective haematopoiesis and vasculogenesis in transforming growth factor- $\beta$  1 knock out mice. *Development* 121(6):1845–1854.
- Kulkarni AB, et al. (1993) Transforming growth factor  $\beta$  1 null mutation in mice causes excessive inflammatory response and early death. *Proc Natl Acad Sci USA* 90(2):770–774.
- Tang Y, et al. (2003) Genetic modifiers interact with maternal determinants in vascular development of *Tgfb1*(*-/-*) mice. *Hum Mol Genet* 12(13):1579–1589.
- Subramanian S, et al. (2005) Epistatic suppression of systemic lupus erythematosus: Fine mapping of *Sle1* to less than 1 mb. *J Immunol* 175(2):1062–1072.
- St Johnston D (2002) The art and design of genetic screens: Drosophila melanogaster. *Nat Rev Genet* 3(3):176–188.
- Potente M, Gerhardt H, Carmeliet P (2011) Basic and therapeutic aspects of angiogenesis. *Cell* 146(6):873–887.
- Gao D, et al. (2009) Bone marrow-derived endothelial progenitor cells contribute to the angiogenic switch in tumor growth and metastatic progression. *Biochim Biophys Acta* 1796(1):33–40.
- Tongers J, Roncalli JG, Losordo DW (2010) Role of endothelial progenitor cells during ischemia-induced vasculogenesis and collateral formation. *Microvasc Res* 79(3):200–206.
- Sobrinho T, et al. (2007) The increase of circulating endothelial progenitor cells after acute ischemic stroke is associated with good outcome. *Stroke* 38(10):2759–2764.
- Yip HK, et al. (2008) Level and value of circulating endothelial progenitor cells in patients after acute ischemic stroke. *Stroke* 39(1):69–74.
- Wara AK, et al. (2011) TGF- $\beta$ 1 signaling and Krüppel-like factor 10 regulate bone marrow-derived proangiogenic cell differentiation, function, and neovascularization. *Blood* 118(24):6450–6460.
- Scheller J, Chalaris A, Garbers C, Rose-John S (2011) ADAM17: A molecular switch to control inflammation and tissue regeneration. *Trends Immunol* 32(8):380–387.
- Liu C, Xu P, Lamouille S, Xu J, Derynck R (2009) TACE-mediated ectodomain shedding of the type I TGF- $\beta$  receptor downregulates TGF- $\beta$  signaling. *Mol Cell* 35(1):26–36.
- Thabet MM, Huizinga TW (2006) Drug evaluation: Apratastat, a novel TACE/MMP inhibitor for rheumatoid arthritis. *Curr Opin Investig Drugs* 7(11):1014–1019.
- Li HY, et al. (2008) Optimization of a dihydropyridopyrazole series of transforming growth factor- $\beta$  type I receptor kinase domain inhibitors: Discovery of an orally bioavailable transforming growth factor- $\beta$  receptor type I inhibitor as antitumor agent. *J Med Chem* 51(7):2302–2306.
- Dupuis-Girod S, et al. (2012) Bevacizumab in patients with hereditary hemorrhagic telangiectasia and severe hepatic vascular malformations and high cardiac output. *JAMA* 307(9):948–955.
- Larrivée B, et al. (2012) ALK1 signaling inhibits angiogenesis by cooperating with the Notch pathway. *Dev Cell* 22(3):489–500.
- Kim JH, Peacock MR, George SC, Hughes CC (2012) BMP9 induces EphrinB2 expression in endothelial cells through an Alk1-BMPRII/ActRII-ID1/ID3-dependent pathway: Implications for hereditary hemorrhagic telangiectasia type II. *Angiogenesis* 15(3):497–509.
- Lange K, Sinsheimer JS, Sobel E (2005) Association testing with Mendel. *Genet Epidemiol* 29(1):36–50.
- Moore JH, Williams SM (2009) Epistasis and its implications for personal genetics. *Am J Hum Genet* 85(3):309–320.
- Samuelson DJ, Aperavich BA, Haag JD, Gould MN (2005) Fine mapping reveals multiple loci and a possible epistatic interaction within the mammary carcinoma susceptibility quantitative trait locus, *Mcs5*. *Cancer Res* 65(21):9637–9642.
- Nagase H, Mao JH, de Koning JP, Minami T, Balmain A (2001) Epistatic interactions between skin tumor modifier loci in interspecific (spretus/musculus) backcross mice. *Cancer Res* 61(4):1305–1308.
- Werner N, et al. (2007) Endothelial progenitor cells correlate with endothelial function in patients with coronary artery disease. *Basic Res Cardiol* 102(6):565–571.
- Parietti E, et al. (2011) Presence of circulating endothelial progenitor cells and levels of stromal-derived factor-1 $\alpha$  are associated with ascending aorta aneurysm size. *Eur J Cardiothorac Surg* 40(1):e6–e12.
- Lindsay ME, et al. (2012) Loss-of-function mutations in *TGFB2* cause a syndromic presentation of thoracic aortic aneurysm. *Nat Genet* 44(8):922–927.
- Doyle AJ, et al. (2012) Mutations in the TGF- $\beta$  repressor *SKI* cause Shprintzen-Goldberg syndrome with aortic aneurysm. *Nat Genet* 44(11):1249–1254.
- Loeys BL, et al. (2005) A syndrome of altered cardiovascular, craniofacial, neurocognitive and skeletal development caused by mutations in *TGFBF1* or *TGFBF2*. *Nat Genet* 37(3):275–281.
- Takeda S, Igarashi T, Mori H, Araki S (2006) Crystal structures of VAP1 reveal ADAMs' MDC domain architecture and its unique C-shaped scaffold. *EMBO J* 25(11):2388–2396.
- Peschon JJ, et al. (1998) An essential role for ectodomain shedding in mammalian development. *Science* 282(5392):1281–1284.
- Blaydon DC, et al. (2011) Inflammatory skin and bowel disease linked to *ADAM17* deletion. *N Engl J Med* 365(16):1502–1508.
- Hassemer EL, et al. (2010) The waved with open eyelids (*woe*) locus is a hypomorphic mouse mutation in *Adam17*. *Genetics* 185(1):245–255.