

UCSF

UC San Francisco Previously Published Works

Title

Pilot study of extended-release lorcaserin for cocaine use disorder among men who have sex with men: A double-blind, placebo-controlled randomized trial

Permalink

<https://escholarship.org/uc/item/26m1r9ff>

Journal

PLOS ONE, 16(7)

ISSN

1932-6203

Authors

Santos, Glenn-Milo

Ikeda, Janet

Coffin, Phillip

et al.

Publication Date

2021

DOI

10.1371/journal.pone.0254724

Peer reviewed


RESEARCH ARTICLE

Pilot study of extended-release lorcaserin for cocaine use disorder among men who have sex with men: A double-blind, placebo-controlled randomized trial

Glenn-Milo Santos^{1,2*}, Janet Ikeda², Phillip Coffin^{2,3}, John E. Walker², Tim Matheson², Matthew McLaughlin², Jennifer Jain⁴, Eric Vittinghoff⁵, Steven L. Batki^{4,6}

1 Department of Community Health Systems, University of California, San Francisco, CA, United States of America, **2** Center on Substance Use and Health, San Francisco Department of Public Health, San Francisco, CA, United States of America, **3** Division of HIV, Infectious Disease & Global Medicine, University of California, San Francisco, CA, United States of America, **4** Department of Psychiatry and Behavioral Sciences, University of California, San Francisco, CA, United States of America, **5** Department of Epidemiology and Biostatistics, University of California, San Francisco, CA, United States of America, **6** San Francisco VA Health Care System (SFVAHCS), San Francisco, CA, United States of America

* Glenn-Milo.Santos@ucsf.edu



OPEN ACCESS

Citation: Santos G-M, Ikeda J, Coffin P, Walker JE, Matheson T, McLaughlin M, et al. (2021) Pilot study of extended-release lorcaserin for cocaine use disorder among men who have sex with men: A double-blind, placebo-controlled randomized trial. *PLoS ONE* 16(7): e0254724. <https://doi.org/10.1371/journal.pone.0254724>

Editor: Bernard Le Foll, Centre for Addiction and Mental Health, CANADA

Received: March 11, 2021

Accepted: June 28, 2021

Published: July 15, 2021

Copyright: © 2021 Santos et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: This study involves sensitive data related to illicit substance use among a small sample size. To protect the privacy of study participants, interested parties will be granted access the data after receiving IRB approval for use from their research institutions, the University of California San Francisco IRB (IRB@ucsf.edu) and contacting PI (GMS at GlennMilo.Santos@gmail.com).

Abstract

Objective

To determine if men who have sex with men (MSM) with cocaine use disorder (CUD) and actively-using cocaine could be enrolled and retained in a pharmacologic intervention trial of lorcaserin—a novel 5-HT_{2c}R agonist—and determine the degree to which participants would adhere to study procedures.

Methods

This was a phase II randomized, double-blind, placebo-controlled pilot study with 2:1 random parallel group assignment to daily extended-release oral lorcaserin 20 mg versus placebo ([clinicaltrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT03192995) identifier-NCT03192995). Twenty-two of a planned 45 cisgender MSM with CUD were enrolled and had weekly follow-up visits during a 12-week treatment period, with substance use counseling, urine specimen collection, and completion of audio-computer assisted self-interview (ACASI) behavioral risk assessments. Adherence was measured by medication event monitoring systems (MEMS) caps and self-report. This study was terminated early because of an FDA safety alert for lorcaserin's long-term use.

Results

Eighty-six percent completed the trial, with 82% of weekly study follow-up visits completed. Adherence was 55.3% (lorcaserin 51.6% vs. placebo 66.2%) by MEMS cap and 56.9% (56.5% vs. placebo 57.9%) by self-report and did not differ significantly by treatment assignment. Intention-to-treat analyses (ITT) did not show differences in cocaine positivity by urine screen between the lorcaserin and placebo groups by 12 week follow-up (incidence risk

Funding: This study was funded by the National Institutes of Drug Abuse (NIDA), grant number R34DA042161-01A1 (PI: G.-M. Santos).

Competing interests: Authors declare no competing interest. Dr Coffin has led National Institutes of Health-funded studies receiving donated ledipasvir-sofosbuvir from Gilead Sciences (2016-2017) outside the submitted work. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

ratio [IRR]: 0.96; 95%CI = 0.24–3.82, $P = 0.95$). However, self-reported cocaine use in time-line follow-back declined more significantly in the lorcaserin group compared to placebo (IRR: 0.66; 95%CI = 0.49–0.88; $P = 0.004$).

Conclusion

We found that it is feasible, acceptable, and tolerable to conduct a placebo-controlled pharmacologic trial for MSM with CUD who are actively using cocaine. Lorcaserin was not associated with significant reductions in cocaine use by urine testing, but was associated with significant reductions in self-reported cocaine use. Future research may be needed to continue to explore the potential utility of 5-HT_{2C}R agonists.

Introduction

Cocaine use in the United States (US) is an unrelenting public health problem with serious negative medical, societal, and economic impacts [1–3]. Its use in the US has also increased during the COVID-19 pandemic [4]. In both powder (cocaine hydrochloride) and crack (free base) forms, cocaine is an addictive stimulant; it is estimated that up to one in six cocaine users may develop a moderate to severe use disorder in their lifetime [2, 5, 6]. Cocaine is associated with a range of negative health consequences, including direct cardiac toxicity, myocardial infarctions, and sudden cardiac death [7–11].

Additionally, cocaine use is a major public health issue among men who have sex with men (MSM). National HIV Behavioral Surveillance (NHBS) data indicate that 37% of MSM in the US used cocaine in the past 12 months, which is 15 times more prevalent than the general population [12, 13]. In San Francisco, cocaine use (past 12 months) climbed from 19% in 2003–2005 to 35% in 2014 [14, 15]. This increase corresponded with a significant decrease (test for trend $p < 0.001$) in methamphetamine use among San Francisco MSM (from 22% to 13%) [15, 16].

In addition, cocaine use plays an important role in the HIV epidemic, particularly for MSM, whom disproportionately account for new HIV infections. It is associated with HIV-related risks (e.g., sexual risk behaviors [17–23] and needle-sharing during injection drug use [24, 25]) as well as new HIV infections in multiple longitudinal studies [26–28]. Moreover, among individuals living with HIV, cocaine use has been associated with poor adherence to antiretroviral treatment (ART), HIV disease progression, lower CD4 count, and higher mortality [29–32].

Given the health hazards associated with cocaine use, there is an urgent need to develop effective treatments, particularly for those with cocaine use disorder (CUD) and for populations with higher prevalence of use and associated harm, including MSM. Unlike alcohol and opioid use disorders, there are currently no FDA-approved medications for CUD, which limits treatment options for this pressing public health issue to behavioral interventions alone [33]. Although some positive findings have been associated with certain pharmacologic agents to treat cocaine use, an effective medication for treatment compared to placebo remains elusive in clinical trials [33]. Lorcaserin, a novel selective serotonin (5-HT) receptor agonist that was FDA-approved for weight management, was, at the time of study design, considered to be a promising agent to treat CUD [34]. Cocaine use increases levels of dopamine (DA), the neurochemical responsible for the reinforcing effects of the drug, and data suggest that DA activity is modulated by the central 5-HT systems [35–37], particular via the 5-HT_{2C} receptor (5-HT_{2C}R), which has been identified as the primary 5-HT receptor subtype that inhibits tonic

and cocaine-evoked DA activity [37]. 5-HT_{2C}R agonists can reduce dopamine release in the nucleus accumbens (NAcc) and frontal neocortex [35, 37]. Preclinical data also support the potential of lorcaserin in reducing cocaine use [36, 38–48]. 5-HT_{2C}R selective agonists, including lorcaserin, have been associated with reductions in cocaine self-administration in mice [49, 50] and non-human primates [47, 51]. The only human cocaine trial with lorcaserin was a laboratory study that found that lorcaserin compared to placebo significantly reduced craving for cocaine and delayed cocaine administration [52]. Further support for studying lorcaserin was based on the possibility that lorcaserin had the potential to reduce cocaine use by reducing impulsivity. Excess impulsivity has long been associated with drug use disorders and cocaine use induces greater impulsivity [53]. 5-HT_{2C}R agonists have reduced impulsive action [53, 54] and lorcaserin has specifically decreased nicotine-induced impulsive action in animal studies [55]. We hypothesize that lorcaserin's *dual effects of inhibiting dopamine levels and decreasing impulsivity* may effectively help MSM reduce or stop their cocaine use.

Given the mechanism of action of lorcaserin and the preclinical data supporting its potential for cocaine treatment, we conducted a pilot clinical trial to examine the feasibility, acceptability, and tolerability of lorcaserin among MSM with CUD in an outpatient study. In exploratory analyses, we also examined lorcaserin's effects on cocaine use and behavioral outcomes including sexual behaviors. This study focused on MSM because the prevalence of CUD is high in this population and effective treatments for CUD can also lead to reductions in HIV transmission among MSM.

Methods

Study design and recruitment

This was a randomized, double-blind, placebo-controlled, 12-week parallel group pilot study with 2:1 random assignment to 20 mg of extended-release oral lorcaserin versus placebo. Assignment of two-thirds of participants to the lorcaserin arm was specifically chosen to improve our ability to detect adverse events in the active treatment arm, while preserving our ability to assess recruitment in a study in which randomization to placebo. Participants were recruited via street outreach, recruitment flyers, STD and HIV clinics, needle exchanges, community organizations, MSM bars, online Web sites, and social media. Potential participants completed a brief telephone screen to assess initial eligibility and, if eligible, were scheduled for an in-person screening visit. All participants gave informed consent using UCSF IRB-approved consent forms. A 10-item true/false questionnaire was used to verify participants' understanding of the trial. The target sample size for the study was 45 participants. With this sample size, we estimated that proportions for our feasibility and acceptability outcomes would be estimated within margins of sampling error (MSEs; i.e., half widths of 95% confidence intervals) of ≤ 14.4 percentage points, and means with MSEs of 0.30 standard deviations, both typical for a small pilot study. We also estimated that for the sample size of 45, the minimum detectable effects of the intervention on the cocaine use outcomes would need to be substantial, as is typical of a pilot study. In our prior trial for MSM actively using stimulants [56], the within-subject correlation of the repeated measures for urine positivity was 0.45, the urine positivity in the control group was 55%, and the retention was 93% at the end of the study. On this basis, we estimated that the sample would provide 80% power to detect reductions of 29 percentage points in the rate of cocaine urine positivity.

Early study termination

On February 13, 2020, the Food and Drug Administration (FDA) issued a safety alert after concluding the assessment of the long-term safety of lorcaserin in another clinical trial among

6,000 participants who took lorcaserin for over 4.3 years of treatment and 6,000 who took placebo; all participants were overweight or obese, with cardiovascular disease or multiple cardiovascular disease risk factors (Cardiovascular and Metabolic Effects of Lorcaserin in Overweight and Obese Patients—Thrombolysis in Myocardial Infarction 61 [CAMELLIA--TIMI 61] clinical trial) [57]. The FDA's analysis of this clinical trial showed an increased risk of cancer, with 7.1% in placebo group being diagnosed with cancer, compared to 7.7% in lorcaserin group. There was no difference in cancer risk noted in the first months of lorcaserin use, but the difference emerged with years of use. In response to the FDA safety alert, active participants in our study at that time ($n = 2$) were instructed to stop study medication use immediately and all participants (2 active and 20 former) were unblinded of their treatment assignment, informed of the FDA safety alert, and provided a copy of the FDA communication. The FDA's statement did not recommend special screening for patients who have taken lorcaserin; it recommended that standard screening recommendations for cancer should be implemented for individual patients, regardless of prior lorcaserin treatment. In consultation with the FDA and the study sponsor, our study was paused (i.e., screening and enrollment was stopped), while the FDA determined whether it was feasible or worthwhile to develop revised safety guidelines for investigational studies with lorcaserin for short-term treatment. On May 20, 2020, the FDA determined that it would not be feasible for researchers to acquire the medication lorcaserin, even for short-term investigational use, and no formal guidelines would be developed for studies of lorcaserin with short-term use. At this point, the FDA recommended study termination. The Data and Safety Monitoring Board (DSMB) was advised of the FDA recommendation and the DSMB approved the initiation of study closeout and recommended the study proceed to data analysis based on data collected from 22 participants enrolled at the time of the FDA safety alert. Since termination of the study, a meta-analysis of 4 clinical trials comprising a cumulative sample of 19,771 participants published in December 2020 did not identify a significant increase in risk for cancer with lorcaserin treatment compared to placebo [58].

Study participants

Participants were eligible if they reported at least one day of active cocaine use during the previous 30 days confirmed by urinalysis or sweat patch at screening (at least one sample positive for cocaine metabolites within two screening and two run-in visits); were seeking treatment for cocaine use and met Diagnostic and Statistical Manual of Mental Disorders (DSM-V) criteria for mild to severe CUD; were HIV negative by rapid antibody test and HIV-pooled RNA test or HIV-positive with medical record of HIV infection and CD4 count; were 18–65 years of age; identified as a cisgender or transgender male who has sex with men; did not have any acute medical or psychiatric illnesses; and had baseline safety labs without clinically significant abnormalities.

We excluded individuals for any psychiatric (e.g., depression with suicidal ideation) or medical condition that would preclude safe study participation or compliance to procedures; any severe current substance use disorder (other than severe cocaine, nicotine, alcohol, or cannabis use disorder) according to DSM-V; testing positive for HIV at screening if previously unaware of HIV infection; a known allergy or previous adverse reaction to lorcaserin; having a current CD4 count < 100 cells/mm³ or HIV viral load greater than 50 copies/mL; moderate to severe liver disease (AST, ALT > 3 times upper limit of normal); severely impaired renal function (creatinine clearance $< \text{or} = 30$ mL/min); use of exclusionary medications that affect the serotonergic neurotransmitter system (e.g., selective serotonin reuptake inhibitors (SSRIs), selective serotonin-norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants, or

monoamine oxidase inhibitors (MAOIs)); self-reporting a predisposition to priapism; currently participating in other intervention studies or recently participating in the last 30 days; anticipated use of agents that are associated with valvulopathy and/or pulmonary hypertension; currently receiving maintenance treatment with an opioid medication (buprenorphine or methadone); currently being in court-mandated drug treatment; any physical condition affecting drug absorption (e.g., gastrectomy); a 12-lead electrocardiogram (ECG) demonstrating QTc > 450 or a QRS interval > 120 msec at screening; current alcohol use disorder that is judged to require supervised medical detoxification (Clinical Institute Withdrawal Assessment for Alcohol, Revised (CIWA-Ar) score > or = 10); being pregnant; or having a body mass index (BMI) < 15, a BMI \geq 30 with desire to use weight management medication, or a BMI > 35. The exclusion criteria for BMI \geq 30 was included because lorcaserin was an FDA-approved treatment for weight loss indicated for obese patients (BMI \geq 30) and it would have been unethical to enroll individuals who are interested in a weight loss medication because this study was placebo-controlled (i.e., they could have been randomized to placebo). Participants screened with BMI \geq 30 with desire to use weight management medication were encouraged to speak with their primary health care provider about treatment options for weight loss by study clinicians.

The exclusion criteria for current CD4 count was revised from <200 cells/mm³ to the criteria above of <100 cells/mm³ and approved by the UCSF-IRB on January 29, 2020. The exclusion criteria for CIWA-Ar was added to the protocol after enrollment started, and approved by the UCSF-IRB on February 11, 2019.

Study procedures

All study procedures were conducted at the San Francisco Department of Public Health (25 Van Ness Avenue, San Francisco, CA). At screening visits, after completing informed consent, participants completed a medical history, vitals, physical, complete blood count, a comprehensive metabolic panel, and a 12-lead ECG. To confirm active cocaine use during screening, the rapid qualitative urine test Medtox Verdict II (Medtox Diagnostics, Burlington, NC) and tamper-evident sweat patches (PharmChek®, PharmChem, Inc., Fort Worth, Tx) were used. Participants who reported being HIV negative or did not know their HIV status received HIV rapid testing (OraQuick HIV 1/2, OraSure Technologies, Inc., Bethlehem, PA) and HIV pooled RNA testing; HIV-positive participants received CD4 count testing. Participants received HIV risk-reduction or "Prevention with Positives" counseling, as appropriate, based on Centers for Disease Control and Prevention (CDC) guidelines [59]. Participants were evaluated for any psychiatric (e.g., depression with suicidal ideation) or medical condition that would preclude safe study participation, as well as any severe current substance use disorder (other than cocaine, nicotine, alcohol, or cannabis use disorder) according to DSM-V. Staff collected extensive participant contact information and two back-up contacts. Eligible participants were scheduled for an enrollment visit.

At enrollment, treatment was assigned using double-blinded block randomization generated using a computer program. The study used randomly selected block sizes of 3 and 6. The study statistician provided the randomization codes to the Safeway compounding pharmacy, which prepared kits corresponding with the treatment assignment (20 mg extended-release lorcaserin or matching placebo) from the randomization code. The study clinicians, staff and participants were all unaware of the treatment assignments for the study drug kits, hence the study was double-blinded. Study clinicians provided training and instructions on dosing of medication during enrollment. Participants were instructed to take one pill each day of extended-release lorcaserin 20mg or placebo. Medications were dispensed in bottles with

MEMS caps (AARDEX Group, Liège, Belgium), which are wireless medication monitoring devices that record each opening as a real-time medication event [60]. All participants were asked about potential adverse events at each follow-up visit; symptom-driven physical exams and safety laboratory monitoring were done at weeks 4, 8, and 12. Adverse events were classified using the Division of AIDS (DAIDS) Table for Grading Severity of Adult Adverse Experiences for HIV Prevention Trials Network [61].

Participants were seen every week for substance use counseling and urine tests for cocaine metabolites. Trained staff, supervised by a clinical psychologist, administered brief (20–30 minutes) substance use counseling at follow-up visits, which was modified from a standardized, manual-driven psychosocial substance use counseling program using cognitive behavioral therapy [62] and motivational interviewing techniques [63, 64] incorporating the Stages of Change Model [65]. This platform has been used in brief substance use behavioral interventions and has had high acceptability among MSM who use substances in prior trials [66–69]. Audio-computer assisted self-interviews (ACASI) were used to standardize data collection and minimize reporting bias [70, 71].

HIV risk-reduction counseling and testing were repeated for HIV-negative participants at the final visit. Participants were paid \$20 for each screening visit, \$10 for run-in visits, \$40 for enrollment, \$15 for weekly visits, \$35 for month 1 and 2 visits, and \$50 for the final visit. The study was conducted under the FDA Investigational New Drug Application (IND #134951). Procedures were approved by University of California San Francisco's Human Research Protection Program (IRB Number 17–21502), and the trial was registered at clinicaltrials.gov (Identifier-NCT03192995).

Outcome measures

For feasibility outcome measures, we computed the proportions of participants eligible and enrolled among those recruited and screened, the proportion of scheduled visits completed, and the proportion of participants retained to the end of the study to assess feasibility of enrolling and retaining MSM with CUD in a 12-week, randomized, double-blind study of lorcaserin versus placebo. We also tabulated the proportion of participants in each group who correctly guessed their treatment assignment to determine if there was significant evidence of unblinding at the end of the trial.

Acceptability measures were computed from questions on attitudes about trial participation and level of satisfaction with trial procedures. We calculated the proportion of participants who were satisfied/highly satisfied with the study, who would participate in a similar study in the future, and the proportion who would recommend the study to a friend. We evaluated lorcaserin use acceptability by examining the frequency of taking the study drug as measured by the number of MEMS cap openings. Cumulative percent adherence was calculated by dividing the frequency of openings at a given time point divided by the number of days since enrollment. For tolerability measures, we computed the proportion of those experiencing adverse events, both overall and by type, to explore the safety of lorcaserin versus placebo among individuals with CUD who are active users.

To measure cocaine use outcomes at each weekly visit, we used qualitative urine screens and self-reported use in the past week, recognizing the differences in detection windows between urine screens and self-reported use. Prior cocaine treatment studies have also documented greater sensitivity in self-report of cocaine compared to qualitative cocaine urine screens [72, 73]. At each weekly visit, we tested urine samples using rapid qualitative urine test Medtox Verdict II, which are estimated to detect cocaine use from the past 1 to 3 days [74]. Self-reported cocaine use from each day within the prior week was measured using a modified

timeline followback [75]. We also calculated the proportion of participants who had any cocaine use in either urine screens or timeline followback to create a composite measure accounting for cocaine use from both urine and self-reported measures.

Standardized measures were used to assess other secondary outcomes, including cocaine craving and severity of cocaine dependence, and sexual risk behavior [18, 75–77]. For monthly visits, the participants completed the measures related to impulsivity, including the Balloon Analog Risk Task (BART) [78] and the Barratt Impulsiveness Scale (BIS) [79].

Statistical analyses

We used Wilcoxon rank sum and Fisher's exact tests, as appropriate, to assess the comparability of participants by treatment assignment at baseline, as well as differences in feasibility, acceptability, and tolerability measures. We compared cumulative adherence between treatment groups using the Wilcoxon test.

Using intention to treat (ITT) analyses, we conducted *a priori* exploratory assessments using GEE Poisson models with robust standard errors to account for within-subject correlation, to assess treatment effects on cocaine urine positivity, and self-reported cocaine use using timeline follow-back. The primary ITT analyses excluded cocaine urine test results from week 1, whereas in a sensitivity analysis, we include those results. In additional sensitivity analyses, we included results from missing urine tests as positive for cocaine use, and analyzed treatment effects while adjusting for baseline cocaine use. In all models, the effect of the intervention was estimated by the interaction between treatment assignment and a linear term in time. For GEE diagnostic procedures, departures from linear trends as well as degree to which the within-subject correlation of the outcomes declines over time were evaluated. Although the robust standard errors in our models accounted for declines in within-subject correlation, models with unstructured correlations were also fitted as a sensitivity analysis. Additionally, we examined treatment effects on any cocaine use based on urine positivity or timeline follow-back. In addition, we conducted as-treated analyses accounting for a time-dependent variable for cumulative medication adherence, calculated from MEMS data, in the analyses examining treatment effects on cocaine urine positivity and self-report; in this analysis, the as-treated effect is estimated by the group-by-adherence interaction. Finally, to evaluate for consecutive weeks of continued abstinence, we compared the “number of beyond-threshold weeks of success” (NOBWOS), defined as the number of consecutive weeks of abstinence lasting through the end of the study [80], using the Wilcoxon test.

We also evaluated treatment effects on cocaine-associated sexual risk behaviors using GEE Poisson, binomial, and negative binomial models using ITT analyses. We used GEE models to explore whether lorcaserin reduced impulsivity as measured by BART and BIS. We also assessed whether lorcaserin reduced cocaine craving and severity of cocaine dependence, using similar methods. Analyses were conducted with STATA 15.0 (StataCorp, College Station, TX).

Results

Screening and enrollment

Fig 1 shows results for screening, recruitment, assignment, and retention for the study period from February 2018 to February 2020. One hundred and six people were assessed for eligibility and consented to participate, of whom 66 were deemed ineligible. The most common reasons for ineligibility were other severe substance use disorders ($n = 36$); high BMI ($n = 8$); inability to provide cocaine-positive urine/sweat samples ($n = 4$); and having ECG values outside the normal range ($n = 4$). In addition, 15 people were lost to follow-up during screening, 1




Fig 1. Study participant flow in screening, randomization, follow-up and analysis.

<https://doi.org/10.1371/journal.pone.0254724.g001>

declined participation, and 2 were in the screening phase when the study was terminated. At the time of study termination, 22 participants (21% of those screened) were randomized (16 to lorcaserin, 6 to placebo; see Fig 1).

Participant characteristics

We recruited a diverse sample of MSM (41% White, 14% Hispanic/Latino, 23% Black, and 18% Mixed or Other race), of whom 18% were HIV positive (Table 1). All participants were cisgender men. No transgender men enrolled in our trial. The median age of participants was 39 (IQR: 32–55). Baseline demographic characteristics were similarly distributed in both groups.

Retention

Overall, the mean percentage of weekly study follow-up visits completed was 82% (SD: 23%). Mean percentage of weekly follow-up visits were similar ($P = 0.91$) between the lorcaserin (83% [SD: 22]) and the control group (81% [SD: 26]). Overall, 19 participants (86%) were retained at the end of the study, and this study completion rate was similar between groups (lorcaserin 13/16 [81%], placebo 6/6 [100%] $P = 0.53$).

Table 1. Baseline demographic and behavioral characteristics of trial participants.

	Control	Treatment	Total
	N = 6	N = 16	N = 22
Age, median (IQR)	40 (39–62)	36 (29–51)	39 (32–55)
Race/Ethnicity			
White	2 (33%)	7 (44%)	9 (41%)
Asian	0 (0%)	1 (6%)	1 (5%)
Black	0 (0%)	5 (31%)	5 (23%)
Latino	2 (33%)	1 (6%)	3 (14%)
Other	2 (33%)	2 (12%)	4 (18%)
Education			
high school or less	1 (17%)	5 (31%)	6 (27%)
some college	3 (50%)	4 (25%)	7 (32%)
college or above	2 (33%)	7 (44%)	9 (41%)
Income			
under \$20,000	1 (17%)	7 (44%)	8 (36%)
\$20–39,999	3 (50%)	2 (12%)	5 (23%)
\$40,000 or more	2 (33%)	7 (44%)	9 (41%)
Employment status			
not employed	1 (17%)	5 (31%)	6 (27%)
full-time	2 (33%)	9 (56%)	11 (50%)
part-time	2 (33%)	2 (12%)	4 (18%)
employed student	1 (17%)	0 (0%)	1 (5%)
Powder cocaine, past 4 weeks			
no use	1 (17%)	3 (19%)	4 (18%)
any use	5 (83%)	13 (81%)	18 (82%)
Crack/Rock Cocaine, past 4 weeks			
no use	4 (67%)	10 (62%)	14 (64%)
any use	2 (33%)	6 (38%)	8 (36%)
Ever hospitalized for cocaine use			
No	5 (83%)	14 (88%)	19 (86%)
Yes	1 (17%)	2 (12%)	3 (14%)
Ever received cocaine treatment			
No	4 (67%)	10 (62%)	14 (64%)
Yes	2 (33%)	6 (38%)	8 (36%)
Craving for Cocaine—Visual Analog Scale	30 (14–45)	40 (2–72)	35 (5–63)
HIV Status			
HIV-positive	0 (0%)	4 (25%)	4 (18%)
HIV-negative	6 (100%)	12 (75%)	18 (82%)
Has regular health care provider			
No	0 (0%)	2 (12%)	2 (9%)
Yes	6 (100%)	14 (88%)	20 (91%)
Depressive Symptoms—Center for Epidemiological Studies Depression (CESD), median (IQR)	10 (7–14)	10 (5–15)	10 (6–14)

<https://doi.org/10.1371/journal.pone.0254724.t001>

Acceptability of procedures

Overall, 74% of participants reported being satisfied/highly satisfied with their study participation. In addition, 89% reported that they would be interested in participating in a similar study in the future, and 89% reported that they would likely recommend participation in the study to a friend. Acceptability of study procedures was similar between the treatment groups.

Adherence

Overall, the mean percentage of study medication taken by participants as measured by MEMS cap data was 55.3% (SD: 26.6) and similar between groups (lorcaserin 51.6% [SD: 27.7]; placebo 66.2% [SD: 21.7]; $P = 0.32$). The mean percentage adherence by self-report was 56.9% (SD: 19.3) overall and similar between treatment groups (lorcaserin 56.5% [21.8]; placebo 57.9% [12.2]; $P = 0.83$).

Assessment of unblinding

Treatment guessing accuracy between participants in the two groups did not differ significantly ($P = 0.63$). In the lorcaserin group, 23% guessed they were on lorcaserin. In the placebo group, 50% guessed they were on placebo.

Tolerability

The overall frequency of adverse events (AEs) was similar between the two groups. There were two serious adverse events (SAEs) in the study that were both deemed unrelated to lorcaserin. One SAE occurred with a participant from the placebo group and involved increased AST (Grade 3 AE) and increased ALT (Grade 2 AE) at month 1 of treatment (resolving to Grade 2 and Grade 1 AEs at the end of study participation). The other SAE was observed in a participant in the lorcaserin group who developed an abscess on his left hip that resolved after a course of antibiotics and was determined to not be related to lorcaserin. The most frequently observed AEs in both groups were hyperglycemia ($n = 4$; lorcaserin = 3; placebo = 1; $P = 0.99$), coughing (lorcaserin = 2, placebo = 1; $P = 0.8$), and abdominal pain (lorcaserin = 2, placebo = 1; $P = 0.8$). The most frequently observed AEs only present in the lorcaserin group were: rash ($n = 3$) and diarrhea ($n = 3$), however, there were no statistically significant differences in the occurrence of these AEs compared to the placebo group (both $P = 0.53$).

Exploratory efficacy analyses of cocaine use and secondary outcomes

The proportion of urine-positive samples for the placebo group was 0% (0/6) at baseline, and 20% (1/5) at week 12 visit. In the lorcaserin group, 50% (8/16) of samples were positive at baseline, and 58% (7/12) at week 12 visit. The proportion of urine-positive samples at baseline were not statistically significantly different between groups ($P = 0.051$). In intention-to-treat analyses (ITT) summarized in [Table 2](#), changes in cocaine positivity by urinalyses were similar between lorcaserin and placebo by 12 week follow-up (incidence risk ratio [IRR]: 0.96; 95% CI = 0.24–3.82, $P = 0.95$), with no evidence of departures from linearity ($P = 0.37$). The within-subject correlation for urine positivity declined over time. Results were similar for sensitivity analyses including week 1 results, imputing positive results from missing urines, using GEE models with an unstructured correlation, and adjusting for baseline urine-positivity.

The proportion of past-week cocaine use self-reported from timeline followback for the placebo group was 100% (6/6) at baseline, and 100% (6/6) at week 12 visit. In the lorcaserin group, 75% (12/16) self-reported past-week cocaine use at baseline, and 54% (7/13) at week 12 visit. The proportion of self-reported past-week cocaine use data at baseline were not statistically significantly different between groups ($P = 0.541$). In ITT analyses, self-reported cocaine use in timeline follow-back declined significantly in the lorcaserin group compared to placebo (IRR: 0.66; 95%CI = 0.49–0.88; $P = 0.004$), with no evidence of departure from linearity ($P = 0.56$). The within-subject correlation for urine positivity declined over time. Results were similar and remain statistically significant in favor of lorcaserin in sensitivity analyses

Table 2. Exploratory efficacy analyses of primary and secondary outcome with sensitivity analyses.

Primary Outcome: Cocaine Urine Positivity			Incidence	(95% CI)	P-value	
			Risk Ratio (IRR)			
Intention-to-treat analyses	Treatment effect		0.96	(0.24–3.82)	0.954	
	Sensitivity Analyses	Inclusion of week 1 results	1.14	(0.16–8.10)	0.899	
		Imputing of positive results from missing urines	0.70	(0.24–2.03)	0.507	
		Treatment effect using unstructured correlation	2.87	(0.46–17.87)	0.258	
		Adjusting for baseline urine positivity	3.11	(0.74–13.11)	0.122	
As-treated analyses	Cumulative adherence and treatment effects for urine positivity		1.01	(0.99–1.03)	0.61	
Primary Outcome: Cocaine Use by self-report in timeline followback			IRR	(95% CI)	P-value	
Intention-to-treat analyses	Treatment effect		0.66	(0.49–0.88)	0.004	
	Sensitivity analysis	Inclusion of week 1 results	0.66	(0.500.89)	0.005	
		Treatment effect using unstructured correlation	0.60	(0.45–0.81)	0.001	
		Adjusting for cocaine use week before enrollment	0.68	(0.51–0.91)	0.009	
		As-treated analyses	Cumulative adherence and treatment effects for self-reported cocaine use		1.01	(0.99–1.02)
Primary Outcome: Any cocaine use based on urine positivity or timeline followback			IRR	(95% CI)	P-value	
Intention-to-treat analysis	Treatment effect		0.76	(0.55–1.04)	0.085	
Secondary Outcomes			Coefficient	(95% CI)	P-value	
Cocaine craving—Visual Analog Scale			10.39	(-1.76–22.53)	0.09	
Severity of Dependence Scale for Cocaine			0.28	(-3.43–3.99)	0.88	
Balloon Analog Risk Task (BART)			-6.03	(-17.1–5.04)	0.29	
Barratt Impulsiveness Scale Score			3.40	(-4.50–11.29)	0.40	
Depression Score (CES-D)			2.86	(-4.95–10.67)	0.47	
Sexual Behavior Outcomes			Risk Ratio (RR)	(95% CI)	P-value	
Number of male partners			0.03	(0.00–1.24)	0.065	
Number of male partners who use cocaine with			0.01	(0.00–6.58)	0.152	
Anal intercourse with HIV serodiscordant partners			0.54	(0.17–1.67)	0.283	
Condomless anal intercourse with HIV serodiscordant partners			0.05	(0.00–1.93)	0.107	
Other Secondary Outcomes			Lorcaserin	Placebo	Overall	P-value
MEMS Caps Percent Adherence, mean (SD)			51.64 (27.73)	66.17 (21.70)	55.3 (26.6)	0.77
Self-reported Percent Adherence, mean (SD)			56.51 (21.76)	57.99 (12.18)	56.9 (19.3)	0.83
Number of beyond-threshold weeks (NOBWOS)*, median (IQR)			0 (0–0.5)	1.5 (0–5)	0 (0–1)	0.27

<https://doi.org/10.1371/journal.pone.0254724.t002>

including week 1 results, using GEE models with an unstructured correlation, and adjusting for baseline urine-positivity.

In ITT analyses on the composite outcome for any cocaine use based on urine positivity or reported use in timeline followback, results were similar between the lorcaserin and placebo groups (IRR: 0.76; 95% CI = 0.55–1.04; P = 0.085). We did not observe significant treatment effects on the secondary outcomes of impulsivity, depression, craving, sexual risk behaviors, and NOBWOS (see Table 2; all P>0.05). In as-treated analyses accounting for cumulative adherence, cocaine urine positivity (IRR: 1.01; 95% CI = 0.99–1.03; P = 0.61) and self-report (IRR: 1.01; 95% CI = 0.99–1.02; P = 0.34) were also similar between treatment groups.

Discussion

This study demonstrated that it was feasible to enroll and retain cisgender MSM with CUD who are actively using cocaine in a pharmacologic intervention trial with excellent rates of participation in study visits, procedures, and follow-up evaluations. In addition, our participants reported high acceptability of study procedures and moderate levels of study medication adherence.

Due to its early termination, this study's final sample is much too small to estimate the treatment effect reliably enough for sample size determination for a larger efficacy trial. Nevertheless, this pilot may provide important information on appropriate modeling techniques given the distribution of the data (e.g., based on the pattern of treatment effect, the intraclass correlation of the repeated measures, and whether and how steeply the within-subject correlation declines over time, and the loss to follow-up rate). In our data, we observed an increasing treatment effect with no evidence of departure from a linearly increasing effect, suggesting that this model was appropriate. Additionally, we observe some decline in the within-subject correlation over time. However, as noted, fitting alternate GEE models using unstructured correlations did not change overall findings for our cocaine use outcomes. Finally, the lost to follow-up rate of 14% may also be encouraging for feasibility of future trials. These findings may suggest that it is viable to test interventions to address CUD among people who are actively using cocaine and have an interest in reducing their cocaine use. Given the morbidity, mortality and HIV risk associated with CUD, especially among MSM, as well as the lack of pharmacological interventions for CUD, these findings provide helpful information in addressing this public health issue.

Our pilot study was not powered to detect treatment effects between lorcaserin and placebo on cocaine use and sexual risk outcomes. In addition, because of the early termination of our study, we did not reach our target sample size, further reducing study power. While we did not observe a statistically significant treatment effect in urine positivity for cocaine, we observed a significant protective effect on cocaine use by self-report in timeline follow-back assessments in favor of lorcaserin. Although the protective effect from lorcaserin on self-reported cocaine use observed may be subject to bias, the double-blind study design may have helped mitigate this concern [81]. Furthermore, as noted, we did not observe evidence of unblinding between the two groups (i.e., because participants didn't know which group they were assigned to, we have no reason to believe that self-report due to social desirability or recall bias would be systematically different between groups). As previously noted, another possible explanation for the different results observed between the two cocaine measures is that the window for cocaine use in the weekly urine tests—estimated to be between 1 to 3 days—was too narrow to detect cocaine use from the prior week [74]. Hence, we may have had non-differential misclassification of cocaine use during weeks when participants used cocaine outside the urine detection window, which would bias the results toward to null [82].

The treatment effect of lorcaserin on the self-reported use of cocaine in our study can be considered in parallel with a human laboratory study finding that lorcaserin compared to placebo significantly reduced craving for cocaine and delayed cocaine administration [52]. However, the negative findings from the recently completed NIDA Clinical Trials Network (CTN) study of lorcaserin for CUD (ClinicalTrials.gov NCT03007394) do not lend support to its utility to achieve abstinence (data not yet peer-reviewed, but overall findings described in [83]). That said, the primary outcome of this CTN lorcaserin trial was “Actual Number of Subjects That Successfully Achieve Abstinence From Cocaine During the Last Three Weeks of Treatment”; it did not report whether lorcaserin was associated with reductions in cocaine use [83]. Nevertheless, because lorcaserin is no longer commercially available (the manufacturer has removed it from the market voluntarily at the request of the FDA), our findings may be viewed more generally as potential evidence in support of future medications with similar mechanisms of action as lorcaserin (i.e., other selective serotonin agonists or other 5-HT_{2C}R agents [84]). As noted, there are currently no FDA-approved pharmacotherapies for CUD. Hence, as novel pharmacologic agents with similar mechanisms as lorcaserin are identified and deemed safe for use, it may be important for future research to examine their therapeutic potential for CUD. Furthermore, given the linkages between cocaine use and HIV-related sexual risk

behaviors and HIV care outcomes, conducting future trials of pharmacotherapies could aid in the development of efficacious strategies for CUD that can also jointly address HIV transmission and acquisition.

This pilot study has several limitations. As mentioned, this study was not originally powered to assess the efficacy of lorcaserin versus placebo, and because the study was terminated early, the study was likely further under-powered in these efficacy analyses. These should be kept in mind while interpreting the null findings in exploratory analyses between treatment groups for cocaine urine-positivity and sexual risk behaviors. Additionally, as noted above the measures on self-reported cocaine use is subject to non-differential misclassification, which is more likely to lead to bias toward the null [82]—it is possible that the effect of lorcaserin may be larger with more precise objective measures. Moreover, quantitative urine analysis could enhance the ability to detect changes in cocaine use that are otherwise missed in qualitative urine screens [85]. Furthermore, given the exploratory nature of the between-group analyses, we did not formally control for multiple hypotheses testing, therefore findings should also be interpreted with caution, particularly those with wide confidence intervals. Additionally, the early termination of our study, in combination with our 2:1 treatment assignment resulted in imbalanced demographic characteristics in the trial, which may hinder external validity. Finally, our small sample size and use of non-probability sampling, and study population of cisgender MSM limits the generalizability of our findings.

Despite these limitations, we found that it is feasible, acceptable, and tolerable to conduct a placebo-controlled pharmacologic trial for MSM with CUD who are actively using, with strong attendance, compliance with study activities, and retention. In this pilot study, lorcaserin, a 5-HT_{2c}R agonist, was not associated with significant reductions in cocaine urine positivity, but it was associated with significant reductions in self-reported cocaine use. Future research is needed to continue to explore the potential utility of 5-HT_{2c}R agonists. Given the public health importance of CUD, the lack of FDA-approved pharmacologic interventions, and cocaine's linkages to HIV, efforts to evaluate other cocaine interventions are urgently needed.

Supporting information

S1 File. Consort checklist. This is the CONSORT checklist for this clinical trial. (DOC)

S2 File. Study protocol. This is the protocol for this clinical trial. (PDF)

Acknowledgments

All authors participated in the revision of the manuscript for important intellectual content and provided approval of the final version of the manuscript.

Author Contributions

Conceptualization: Glenn-Milo Santos, Eric Vittinghoff.

Data curation: John E. Walker, Matthew McLaughlin, Jennifer Jain.

Formal analysis: Glenn-Milo Santos, Jennifer Jain.

Funding acquisition: Glenn-Milo Santos.

Investigation: Glenn-Milo Santos, John E. Walker.

Methodology: Janet Ikeda, Phillip Coffin, Tim Matheson, Eric Vittinghoff, Steven L. Batki.

Project administration: Janet Ikeda, Phillip Coffin, John E. Walker, Tim Matheson.

Supervision: Glenn-Milo Santos, Janet Ikeda.

Validation: Matthew McLaughlin.

Writing – original draft: Glenn-Milo Santos.

Writing – review & editing: Janet Ikeda, Phillip Coffin, John E. Walker, Tim Matheson, Matthew McLaughlin, Jennifer Jain, Eric Vittinghoff, Steven L. Batki.

References

1. Riezzo I, Fiore C, De Carlo D, Pascale N, Neri M, Turillazzi E, et al. Side effects of cocaine abuse: multi-organ toxicity and pathological consequences. *Curr Med Chem*. 2012; 19(33):5624–46. Epub 2012/09/01. doi: CMC-EPUB-20120817-7 [pii]. <https://doi.org/10.2174/092986712803988893> PMID: 22934772.
2. Karila L, Petit A, Lowenstein W, Reynaud M. Diagnosis and consequences of cocaine addiction. *Curr Med Chem*. 2012; 19(33):5612–8. Epub 2012/08/04. doi: CMC-EPUB-20120801-8 [pii]. <https://doi.org/10.2174/092986712803988839> PMID: 22856661.
3. Substance Abuse and Mental Health Services Administration. Drug Abuse Warning Network, 2011: National Estimates of Drug-Related Emergency Department Visits Rockville, MD: HHS, 2013 04/14/2014. Report No.
4. Millennium Health. COVID-19 Special Edition Reveals Significant Changes in Drug Use During the Pandemic [Report]. *Millennium Health*; 2020 [updated July 8, 2020; cited 2021 March 2. 2021]. Available from: <https://www.millenniumhealth.com/news/signalsreportcovid/>.
5. Anthony J, Warner L, Kessler R. Comparative epidemiology of dependence on tobacco, alcohol, controlled substances, and inhalants: Basic findings from the National Comorbidity Survey. *Experimental and Clinical Psychopharmacology*. 1994; 23:244.
6. Degenhardt L, Baxter AJ, Lee YY, Hall W, Sara GE, Johns N, et al. The global epidemiology and burden of psychostimulant dependence: Findings from the Global Burden of Disease Study 2010. *Drug Alcohol Depend*. 2014. Epub 2014/02/25. doi: S0376-8716(14)00023-4 [pii] <https://doi.org/10.1016/j.drugalcdep.2013.12.025> PMID: 24559607.
7. Vandhuick O, Pistorius MA, Jousse S, Ferreira-Maldent N, Guilmot JL, Guias B, et al. [Drug addiction and cardiovascular pathologies]. *Journal des maladies vasculaires*. 2004; 29(5):243–8. Epub 2005/03/02. [https://doi.org/10.1016/s0398-0499\(04\)96768-6](https://doi.org/10.1016/s0398-0499(04)96768-6) PMID: 15738835.
8. Kloner RA, Hale S, Alker K, Rezkalla S. The effects of acute and chronic cocaine use on the heart. *Circulation*. 1992; 85(2):407–19. Epub 1992/02/01. <https://doi.org/10.1161/01.cir.85.2.407> PMID: 1346509.
9. McCord J, Jneid H, Hollander JE, de Lemos JA, Cercek B, Hsue P, et al. Management of cocaine-associated chest pain and myocardial infarction: a scientific statement from the American Heart Association Acute Cardiac Care Committee of the Council on Clinical Cardiology. *Circulation*. 2008; 117(14):1897–907. Epub 2008/03/19. <https://doi.org/10.1161/CIRCULATIONAHA.107.188950> PMID: 18347214.
10. Phillips K, Luk A, Soor GS, Abraham JR, Leong S, Butany J. Cocaine cardiotoxicity: a review of the pathophysiology, pathology, and treatment options. *American journal of cardiovascular drugs: drugs, devices, and other interventions*. 2009; 9(3):177–96. Epub 2009/05/26. <https://doi.org/10.2165/00129784-200909030-00005> PMID: 19463023.
11. Steinhauer JR, Caulfield JB. Spontaneous coronary artery dissection associated with cocaine use: a case report and brief review. *Cardiovascular pathology: the official journal of the Society for Cardiovascular Pathology*. 2001; 10(3):141–5. Epub 2001/08/04. [https://doi.org/10.1016/s1054-8807\(01\)00074-6](https://doi.org/10.1016/s1054-8807(01)00074-6) PMID: 11485859.
12. Sanchez T, Finlayson T, Drake A, Behel S, Cribbin M, Dinenzo E, et al. Human immunodeficiency virus (HIV) risk, prevention, and testing behaviors—United States, National HIV Behavioral Surveillance System: men who have sex with men, November 2003–April 2005. *MMWR Surveill Summ*. 2006; 55(6):1–16. Epub 2006/07/11. doi: ss5506a1 [pii]. PMID: 16826162.
13. SAMHSA. National Survey on Drug Use and Health. Office of Applied Studies. 2004.
14. Raymond HF. Estimated number of stimulant using men-who-have-sex-with-men in San Francisco. In: Santos GM, editor. San Francisco, CA2012.
15. Raymond HF. National HIV Behavioral Surveillance MSM cycle 4 stimulant use data In: Santos GM, editor. San Francisco, CA 2014.

16. Raymond HF, Chen YH, Ick T, Scheer S, Bernstein K, Liska S, et al. A new trend in the HIV epidemic among men who have sex with men, San Francisco, 2004–2011. *J Acquir Immune Defic Syndr*. 2013; 62(5):584–9. Epub 2013/01/22. <https://doi.org/10.1097/QAI.0b013e318285febf> PMID: 23334505.
17. Mimiaga MJ, Reisner SL, Fontaine YM, Bland SE, Driscoll MA, Isenberg D, et al. Walking the line: stimulant use during sex and HIV risk behavior among Black urban MSM. *Drug Alcohol Depend*. 2010; 110(1–2):30–7. Epub 2010/03/26. doi: S0376-8716(10)00059-1 [pii] <https://doi.org/10.1016/j.drugalcdep.2010.01.017> PMID: 20334986; PubMed Central PMCID: PMC3947405.
18. Colfax G, Vittinghoff E, Husnik MJ, McKirnan D, Buchbinder S, Koblin B, et al. Substance use and sexual risk: a participant- and episode-level analysis among a cohort of men who have sex with men. *Am J Epidemiol*. 2004; 159(10):1002–12. Epub 2004/05/07. <https://doi.org/10.1093/aje/kwh135> PMID: 15128613.
19. Ober A, Shoptaw S, Wang PC, Gorbach P, Weiss RE. Factors associated with event-level stimulant use during sex in a sample of older, low-income men who have sex with men in Los Angeles. *Drug Alcohol Depend*. 2009; 102(1–3):123–9. Epub 2009/03/31. doi: S0376-8716(09)00049-0 [pii] <https://doi.org/10.1016/j.drugalcdep.2009.02.002> PMID: 19327917; PubMed Central PMCID: PMC2751657.
20. Mackesy-Amity ME, Fendrich M, Johnson TP. Symptoms of substance dependence and risky sexual behavior in a probability sample of HIV-negative men who have sex with men in Chicago. *Drug Alcohol Depend*. 2010; 110(1–2):38–43. Epub 2010/03/12. doi: S0376-8716(10)00058-X [pii] <https://doi.org/10.1016/j.drugalcdep.2010.01.016> PMID: 20219291.
21. Lambert G, Cox J, Hottes TS, Tremblay C, Frigault LR, Alary M, et al. Correlates of unprotected anal sex at last sexual episode: analysis from a surveillance study of men who have sex with men in Montreal. *AIDS Behav*. 2011; 15(3):584–95. Epub 2009/12/25. <https://doi.org/10.1007/s10461-009-9605-3> PMID: 20033763.
22. Bowers JR, Branson CM, Fletcher JB, Reback CJ. Predictors of HIV Sexual Risk Behavior among Men Who Have Sex with Men, Men Who Have Sex with Men and Women, and Transgender Women. *Int J Sex Health*. 2012; 24(4):290–302. Epub 2012/01/01. <https://doi.org/10.1080/19317611.2012.715120> PMID: 24660042; PubMed Central PMCID: PMC3960284.
23. Santos GM, Coffin PO, Das M, Matheson T, DeMicco E, Raiford JL, et al. Dose-response associations between number and frequency of substance use and high-risk sexual behaviors among HIV-negative substance-using men who have sex with men (SUMSM) in San Francisco. *J Acquir Immune Defic Syndr*. 2013; 63(4):540–4. Epub 2013/04/11. <https://doi.org/10.1097/QAI.0b013e318293f10b> PMID: 23572012.
24. Tyndall MW, Currie S, Spittal P, Li K, Wood E, O'Shaughnessy MV, et al. Intensive injection cocaine use as the primary risk factor in the Vancouver HIV-1 epidemic. *AIDS*. 2003; 17(6):887–93. Epub 2003/03/28. <https://doi.org/10.1097/00002030-200304110-00014> PMID: 12660536.
25. Cepeda JA, Vickerman P, Bruneau J, Zang G, Borquez A, Farrell M, et al. Estimating the contribution of stimulant injection to HIV and HCV epidemics among people who inject drugs and implications for harm reduction: A modeling analysis. *Drug Alcohol Depend*. 2020; 213:108135. Epub 2020/07/01. <https://doi.org/10.1016/j.drugalcdep.2020.108135> PMID: 32603976; PubMed Central PMCID: PMC7829087.
26. Ostrow DG, Plankey MW, Cox C, Li X, Shoptaw S, Jacobson LP, et al. Specific sex drug combinations contribute to the majority of recent HIV seroconversions among MSM in the MACS. *J Acquir Immune Defic Syndr*. 2009; 51(3):349–55. Epub 2009/04/24. <https://doi.org/10.1097/QAI.0b013e3181a24b20> PMID: 19387357; PubMed Central PMCID: PMC3074969.
27. Koblin BA, Husnik MJ, Colfax G, Huang Y, Madison M, Mayer K, et al. Risk factors for HIV infection among men who have sex with men. *AIDS*. 2006; 20(5):731–9. Epub 2006/03/04. <https://doi.org/10.1097/01.aids.0000216374.61442.55> [pii]. PMID: 16514304.
28. Ostrow DG, DiFranceisco WJ, Chmiel JS, Wagstaff DA, Wesch J. A case-control study of human immunodeficiency virus type 1 seroconversion and risk-related behaviors in the Chicago MACS/CCS Cohort, 1984–1992. Multicenter AIDS Cohort Study. Coping and Change Study. *Am J Epidemiol*. 1995; 142(8):875–83. Epub 1995/10/15. <https://doi.org/10.1093/oxfordjournals.aje.a117727> PMID: 7572964.
29. Hinkin CH, Barclay TR, Castellon SA, Levine AJ, Durvasula RS, Marion SD, et al. Drug use and medication adherence among HIV-1 infected individuals. *AIDS Behav*. 2007; 11(2):185–94. Epub 2006/08/10. <https://doi.org/10.1007/s10461-006-9152-0> PMID: 16897351; PubMed Central PMCID: PMC2867605.
30. Carrico AW. Substance use and HIV disease progression in the HAART era: implications for the primary prevention of HIV. *Life Sci*. 2011; 88(21–22):940–7. Epub 2010/10/12. doi: S0024-3205(10)00437-6 [pii] <https://doi.org/10.1016/j.lfs.2010.10.002> PMID: 20934437.
31. Shoptaw S, Stall R, Bordon J, Kao U, Cox C, Li X, et al. Cumulative exposure to stimulants and immune function outcomes among HIV-positive and HIV-negative men in the Multicenter AIDS Cohort Study. *Int J STD AIDS*. 2012; 23(8):576–80. Epub 2012/08/30. doi: 23/8/576 [pii] <https://doi.org/10.1258/ijsa.2012.011322> PMID: 22930295; PubMed Central PMCID: PMC3576843.

32. Cook JA, Burke-Miller JK, Cohen MH, Cook RL, Vlahov D, Wilson TE, et al. Crack cocaine, disease progression, and mortality in a multicenter cohort of HIV-1 positive women. *AIDS*. 2008; 22(11):1355–63. Epub 2008/06/27. <https://doi.org/10.1097/QAD.0b013e32830507f2> [pii]. PMID: 18580615; PubMed Central PMCID: PMC2645902.
33. Chan B, Kondo K, Freeman M, Ayers C, Montgomery J, Kansagara D. Pharmacotherapy for Cocaine Use Disorder—a Systematic Review and Meta-analysis. *J Gen Intern Med*. 2019; 34(12):2858–73. Epub 2019/06/12. <https://doi.org/10.1007/s11606-019-05074-8> PMID: 31183685; PubMed Central PMCID: PMC6854210.
34. Higgins GA, Fletcher PJ, Shanahan WR. Lorcaserin: A review of its preclinical and clinical pharmacology and therapeutic potential. *Pharmacol Ther*. 2020; 205:107417. Epub 2019/10/20. <https://doi.org/10.1016/j.pharmthera.2019.107417> PMID: 31629010.
35. Rothman RB, Blough BE, Baumann MH. Dopamine/serotonin releasers as medications for stimulant addictions. *Prog Brain Res*. 2008; 172:385–406. Epub 2008/09/06. doi: S0079-6123(08)00919-9 [pii] [https://doi.org/10.1016/S0079-6123\(08\)00919-9](https://doi.org/10.1016/S0079-6123(08)00919-9) PMID: 18772043.
36. Higgins GA, Sellers EM, Fletcher PJ. From obesity to substance abuse: therapeutic opportunities for 5-HT_{2C} receptor agonists. *Trends Pharmacol Sci*. 2013; 34(10):560–70. Epub 2013/09/18. doi: S0165-6147(13)00150-8 [pii] <https://doi.org/10.1016/j.tips.2013.08.001> PMID: 24041919.
37. Alex KD, Pehek EA. Pharmacologic mechanisms of serotonergic regulation of dopamine neurotransmission. *Pharmacol Ther*. 2007; 113(2):296–320. Epub 2006/10/20. doi: S0163-7258(06)00152-5 [pii] <https://doi.org/10.1016/j.pharmthera.2006.08.004> PMID: 17049611; PubMed Central PMCID: PMC2562467.
38. Bubar MJ, Cunningham KA. Serotonin 5-HT_{2A} and 5-HT_{2C} receptors as potential targets for modulation of psychostimulant use and dependence. *Curr Top Med Chem*. 2006; 6(18):1971–85. Epub 2006/10/05. <https://doi.org/10.2174/156802606778522131> PMID: 17017968.
39. Bubar MJ, Cunningham KA. Prospects for serotonin 5-HT_{2R} pharmacotherapy in psychostimulant abuse. *Prog Brain Res*. 2008; 172:319–46. Epub 2008/09/06. doi: S0079-6123(08)00916-3 [pii] [https://doi.org/10.1016/S0079-6123\(08\)00916-3](https://doi.org/10.1016/S0079-6123(08)00916-3) PMID: 18772040.
40. Burton CL, Rizos Z, Diwan M, Nobrega JN, Fletcher PJ. Antagonizing 5-HT_{2A} receptors with M100907 and stimulating 5-HT_{2C} receptors with Ro60-0175 blocks cocaine-induced locomotion and zif268 mRNA expression in Sprague-Dawley rats. *Behav Brain Res*. 2013; 240:171–81. Epub 2012/12/04. doi: S0166-4328(12)00759-0 [pii] <https://doi.org/10.1016/j.bbr.2012.11.030> PMID: 23201361.
41. Craige CP, Unterwald EM. Serotonin (2C) receptor regulation of cocaine-induced conditioned place preference and locomotor sensitization. *Behav Brain Res*. 2013; 238:206–10. Epub 2012/10/30. doi: S0166-4328(12)00684-5 [pii] <https://doi.org/10.1016/j.bbr.2012.10.034> PMID: 23103406; PubMed Central PMCID: PMC3513564.
42. Cunningham KA, Fox RG, Anastasio NC, Bubar MJ, Stutz SJ, Moeller FG, et al. Selective serotonin 5-HT_{2C} receptor activation suppresses the reinforcing efficacy of cocaine and sucrose but differentially affects the incentive-salience value of cocaine- vs. sucrose-associated cues. *Neuropharmacology*. 2011; 61(3):513–23. Epub 2011/05/18. doi: S0028-3908(11)00179-1 [pii] <https://doi.org/10.1016/j.neuropharm.2011.04.034> PMID: 21575646; PubMed Central PMCID: PMC3115535.
43. Filip M, Alenina N, Bader M, Przegalinski E. Behavioral evidence for the significance of serotonergic (5-HT) receptors in cocaine addiction. *Addict Biol*. 2010; 15(3):227–49. Epub 2010/05/12. doi: ADB214 [pii] <https://doi.org/10.1111/j.1369-1600.2010.00214.x> PMID: 20456287.
44. Filip M, Spampinato U, McCreary AC, Przegalinski E. Pharmacological and genetic interventions in serotonin (5-HT)_{2C} receptors to alter drug abuse and dependence processes. *Brain Res*. 2012; 1476:132–53. Epub 2012/04/13. doi: S0006-8993(12)00517-3 [pii] <https://doi.org/10.1016/j.brainres.2012.03.035> PMID: 22494568.
45. Katsidoni V, Apazoglou K, Panagis G. Role of serotonin 5-HT_{2A} and 5-HT_{2C} receptors on brain stimulation reward and the reward-facilitating effect of cocaine. *Psychopharmacology (Berl)*. 2011; 213(2–3):337–54. Epub 2010/06/26. <https://doi.org/10.1007/s00213-010-1887-7> PMID: 20577718.
46. Leggio GM, Cathala A, Moison D, Cunningham KA, Piazza PV, Spampinato U. Serotonin_{2C} receptors in the medial prefrontal cortex facilitate cocaine-induced dopamine release in the rat nucleus accumbens. *Neuropharmacology*. 2009; 56(2):507–13. Epub 2008/11/04. doi: S0028-3908(08)00481-4 [pii] <https://doi.org/10.1016/j.neuropharm.2008.10.005> PMID: 18977370; PubMed Central PMCID: PMC3130963.
47. Manvich DF, Kimmel HL, Howell LL. Effects of serotonin 2C receptor agonists on the behavioral and neurochemical effects of cocaine in squirrel monkeys. *J Pharmacol Exp Ther*. 2012; 341(2):424–34. Epub 2012/02/14. doi: jpet.111.186981 [pii] <https://doi.org/10.1124/jpet.111.186981> PMID: 22328576; PubMed Central PMCID: PMC3336818.
48. Pockros LA, Pentkowski NS, Conway SM, Ullman TE, Zwick KR, Neisewander JL. 5-HT_{2A} receptor blockade and 5-HT_{2C} receptor activation interact to reduce cocaine hyperlocomotion and Fos protein

- expression in the caudate-putamen. *Synapse*. 2012; 66(12):989–1001. Epub 2012/08/14. <https://doi.org/10.1002/syn.21592> PMID: 22886755; PubMed Central PMCID: PMC3476845.
49. Grottick AJ, Fletcher PJ, Higgins GA. Studies to investigate the role of 5-HT(2C) receptors on cocaine- and food-maintained behavior. *J Pharmacol Exp Ther*. 2000; 295(3):1183–91. Epub 2000/11/18. PMID: 11082456.
 50. Pelloux Y, Dilleen R, Economidou D, Theobald D, Everitt BJ. Reduced forebrain serotonin transmission is causally involved in the development of compulsive cocaine seeking in rats. *Neuropsychopharmacology*. 2012; 37(11):2505–14. Epub 2012/07/06. doi: npp2012111 [pii] <https://doi.org/10.1038/npp.2012.111> PMID: 22763621; PubMed Central PMCID: PMC3442345.
 51. Kohut S, Bergman J, Mello N. Pretreatment with either a 5-HT2C agonist or 5-HT2A antagonist decreases cocaine self-administration. *The FASEB Journal*. 2014; 28(1, Supplement 848.9).
 52. Pirtle JL, Hickman MD, Boinepally VC, Surineni K, Thakur HK, Grasing KW. The serotonin-2C agonist Lorcaserin delays intravenous choice and modifies the subjective and cardiovascular effects of cocaine: A randomized, controlled human laboratory study. *Pharmacol Biochem Behav*. 2019; 180:52–9. Epub 2019/02/28. <https://doi.org/10.1016/j.pbb.2019.02.010> PMID: 30811963; PubMed Central PMCID: PMC6529237.
 53. Fletcher PJ, Rizos Z, Noble K, Higgins GA. Impulsive action induced by amphetamine, cocaine and MK801 is reduced by 5-HT(2C) receptor stimulation and 5-HT(2A) receptor blockade. *Neuropharmacology*. 2011; 61(3):468–77. Epub 2011/03/16. doi: S0028-3908(11)00105-5 [pii] <https://doi.org/10.1016/j.neuropharm.2011.02.025> PMID: 21402085.
 54. Navarra R, Comery TA, Graf R, Rosenzweig-Lipson S, Day M. The 5-HT(2C) receptor agonist WAY-163909 decreases impulsivity in the 5-choice serial reaction time test. *Behav Brain Res*. 2008; 188(2):412–5. Epub 2008/01/15. doi: S0166-4328(07)00626-2 [pii] <https://doi.org/10.1016/j.bbr.2007.11.016> PMID: 18191235.
 55. Higgins GA, Silenieks LB, Rossmann A, Rizos Z, Noble K, Soko AD, et al. The 5-HT2C receptor agonist lorcaserin reduces nicotine self-administration, discrimination, and reinstatement: relationship to feeding behavior and impulse control. *Neuropsychopharmacology*. 2012; 37(5):1177–91. Epub 2011/12/23. doi: npp2011303 [pii] <https://doi.org/10.1038/npp.2011.303> PMID: 22189292; PubMed Central PMCID: PMC3306879.
 56. Colfax GN, Santos GM, Das M, Santos DM, Matheson T, Gasper J, et al. Mirtazapine to reduce methamphetamine use: a randomized controlled trial. *Arch Gen Psychiatry*. 2011; 68(11):1168–75. Epub 2011/11/09. doi: 68/11/1168 [pii] <https://doi.org/10.1001/archgenpsychiatry.2011.124> PMID: 22065532; PubMed Central PMCID: PMC3437988.
 57. Bohula EA, Wiviott SD, McGuire DK, Inzucchi SE, Kuder J, Im K, et al. Cardiovascular Safety of Lorcaserin in Overweight or Obese Patients. *N Engl J Med*. 2018; 379(12):1107–17. Epub 2018/08/28. <https://doi.org/10.1056/NEJMoa1808721> PMID: 30145941.
 58. de Andrade Mesquita L, Fagundes Piccoli G, Richter da Natividade G, Frison Spiazzi B, Colpani V, Gerchman F. Is lorcaserin really associated with increased risk of cancer? A systematic review and meta-analysis. *Obes Rev*. 2021; 22(3):e13170. Epub 2020/12/02. <https://doi.org/10.1111/obr.13170> PMID: 33258543.
 59. Revised guidelines for HIV counseling, testing, and referral. *MMWR Recomm Rep*. 2001; 50(Rr-19):1–57; quiz CE1-19a1-CE6-a1. Epub 2001/11/23. PMID: 11718472.
 60. Haberer JE, Kahane J, Kigozi I, Emenyonu N, Hunt P, Martin J, et al. Real-time adherence monitoring for HIV antiretroviral therapy. *AIDS Behav*. 2010; 14(6):1340–6. Epub 2010/09/03. <https://doi.org/10.1007/s10461-010-9799-4> PMID: 20809380; PubMed Central PMCID: PMC2974938.
 61. U.S. Department of Health and Human Services NIOH, National Institute of Allergy and Infectious Diseases, Division of AIDS. Table for Grading the Severity of Adult and Pediatric Adverse Events, Version 2.0. 2014.
 62. Brief D, Bollinger A., Horton G. & LoCastro J. S. Relapse Prevention Treatment for Cocaine Addiction: the RPT-C Manual. Bethesda, MD: Medication Development Division, National Institute on Drug Abuse; 1998.
 63. Miller W, Rollnick S. *Motivational Interviewing*. 2nd ed. New York: Guilford Press; 2002.
 64. Miller WR. Motivational interviewing with problem drinkers. *Behavioural Psychotherapy*. 1983; 11:147–72.
 65. DiClemente CC, Prochaska JO, Gilbertini M. Self-efficacy and the stages of self change in smoking. *Cognitive Therapy and Research*. 1985; 9:181–200.
 66. Das M, Santos D, Matheson T, Santos GM, Chu P, Vittinghoff E, et al. Feasibility and acceptability of a phase II randomized pharmacologic intervention for methamphetamine dependence in high-risk men who have sex with men. *AIDS*. 2010; 24(7):991–1000. Epub 2010/04/17. <https://doi.org/10.1097/qad.0b013e328336e98b> PMID: 20397286.

67. Dunn C, Deroo L, Rivara FP. The use of brief interventions adapted from motivational interviewing across behavioral domains: a systematic review. *Addiction*. 2001; 96(12):1725–42. <https://doi.org/10.1046/j.1360-0443.2001.961217253.x> PMID: 11784466.
68. Rollnick S, Miller W. What is motivational interviewing? *Cognit Behav Psychother*. 1995; 25:325–34.
69. Samet JH, Rollnick S, Barnes H. Beyond CAGE. A brief clinical approach after detection of substance abuse. *Arch Intern Med*. 1996; 156(20):2287–93. <https://doi.org/10.1001/archinte.156.20.2287> PMID: 8911235.
70. Koblin B, Chesney M, Coates T, Mayer K, Agredano F, Aguilu E, et al. Effects of a behavioural intervention to reduce acquisition of HIV infection among men who have sex with men: the EXPLORE randomised controlled study. *Lancet*. 2004; 364(9428):41–50. PubMed PMID: ISI:000222392900030. [https://doi.org/10.1016/S0140-6736\(04\)16588-4](https://doi.org/10.1016/S0140-6736(04)16588-4) PMID: 15234855
71. Macalino GE, Celentano DD, Latkin C, Strathdee SA, Vlahov D. Risk behaviors by audio computer-assisted self-interviews among HIV-seropositive and HIV-seronegative injection drug users. *AIDS Educ Prev*. 2002; 14(5):367–78. <https://doi.org/10.1521/aeap.14.6.367.24075> PMID: 12413183.
72. Delucchi KL, Jones RT, Batki SL. Measurement properties of quantitative urine benzoylecgonine in clinical trials research. *Addiction*. 1997; 92(3):297–302. Epub 1997/03/01. PMID: 9219391.
73. Batki SL, Washburn AM, Delucchi K, Jones RT. A controlled trial of fluoxetine in crack cocaine dependence. *Drug Alcohol Depend*. 1996; 41(2):137–42. Epub 1996/06/01. [https://doi.org/10.1016/0376-8716\(96\)01233-1](https://doi.org/10.1016/0376-8716(96)01233-1) PMID: 8809502.
74. Preston KL, Epstein DH, Cone EJ, Wtsadik AT, Huestis MA, Moolchan ET. Urinary elimination of cocaine metabolites in chronic cocaine users during cessation. *J Anal Toxicol*. 2002; 26(7):393–400. Epub 2002/11/09. <https://doi.org/10.1093/jat/26.7.393> PMID: 12422991.
75. Sobell LC, Brown J, Leo GI, Sobell MB. The reliability of the Alcohol Timeline Followback when administered by telephone and by computer. *Drug Alcohol Depend*. 1996; 42(1):49–54. Epub 1996/09/01. [https://doi.org/10.1016/0376-8716\(96\)01263-x](https://doi.org/10.1016/0376-8716(96)01263-x) PMID: 8889403.
76. Tziortzis D, Mahoney JJ 3rd, Kalechstein AD, TF Newton, De La Garza R 2nd. The relationship between impulsivity and craving in cocaine- and methamphetamine-dependent volunteers. *Pharmacol Biochem Behav*. 2011; 98(2):196–202. Epub 2011/01/11. <https://doi.org/10.1016/j.pbb.2010.12.022> PMID: 21215769.
77. Gonzalez-Saiz F, Domingo-Salvany A, Barrio G, Sanchez-Niubo A, Brugal MT, de la Fuente L, et al. Severity of dependence scale as a diagnostic tool for heroin and cocaine dependence. *Eur Addict Res*. 2009; 15(2):87–93. Epub 2009/01/15. <https://doi.org/10.1159/000189787> PMID: 19142008.
78. Hunt MK, Hopko DR, Bare R, Lejuez CW, Robinson EV. Construct validity of the Balloon Analog Risk Task (BART): associations with psychopathy and impulsivity. *Assessment*. 2005; 12(4):416–28. <https://doi.org/10.1177/1073191105278740> PMID: 16244122.
79. Reise SP, Moore TM, Sabb FW, Brown AK, London ED. The Barratt Impulsiveness Scale-11: reassessment of its structure in a community sample. *Psychol Assess*. 2013; 25(2):631–42. Epub 2013/04/03. <https://doi.org/10.1037/a0032161> PMID: 23544402; PubMed Central PMCID: PMC3805371.
80. McCann DJ, Li SH. A novel, nonbinary evaluation of success and failure reveals bupropion efficacy versus methamphetamine dependence: reanalysis of a multisite trial. *CNS Neurosci Ther*. 2012; 18(5):414–8. Epub 2011/11/11. <https://doi.org/10.1111/j.1755-5949.2011.00263.x> PMID: 22070720; PubMed Central PMCID: PMC6493362.
81. Day SJ, Altman DG. Statistics notes: blinding in clinical trials and other studies. *BMJ*. 2000; 321(7259):504. Epub 2000/08/19. <https://doi.org/10.1136/bmj.321.7259.504> PMID: 10948038; PubMed Central PMCID: PMC1118396.
82. Lash TL, Ahern TP. Bias analysis to guide new data collection. *Int J Biostat*. 2012; 8(2). Epub 2012/04/14. <https://doi.org/10.2202/1557-4679.1345> PMID: 22499734.
83. Negus SS, Banks ML. Learning from lorcaserin: lessons from the negative clinical trial of lorcaserin to treat cocaine use disorder. *Neuropsychopharmacology*. 2020; 45(12):1967–73. Epub 2020/08/26. <https://doi.org/10.1038/s41386-020-00815-4> PMID: 32839526; PubMed Central PMCID: PMC7547654.
84. Palacios JM, Pazos A, Hoyer D. A short history of the 5-HT_{2C} receptor: from the choroid plexus to depression, obesity and addiction treatment. *Psychopharmacology (Berl)*. 2017; 234(9–10):1395–418. Epub 2017/03/08. <https://doi.org/10.1007/s00213-017-4545-5> PMID: 28265714.
85. Li SH, Chiang CN, Tai BC, Marschke CK, Hawks RL. Quantitative versus qualitative urinalysis for benzoylecgonine in clinical trials for the assessment of cocaine use. *Psychopharmacol Bull*. 1995; 31(4):671–9. Epub 1995/01/01. PMID: 8851639.