Safety of Vaccines Used for Routine Immunization in the United States: An Updated Systematic Review and Meta-analysis

Courtney Gidengil, M.D., M.P.H.,[a,b] Matthew Bidwell Goetz, M.D.,[c] Sydne Newberry, Ph.D.,[d] Margaret Maglione, M.P.P.,[d] Owen Hall, M.S.,[d] Jody Larkin, M.S.,[d] Aneesa Motala, B.A.,[d, e] Susanne Hempel, Ph.D.[d, e]

[a]RAND Corporation, 20 Park Plaza, Suite 920, Boston, MA, 02116

[b]Boston Children's Hospital, 300 Longwood Avenue, Boston, MA, 02115

[c]VA Greater Los Angeles Healthcare System and David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA, 90073

[d]RAND Corporation, 1776 Main Street, Santa Monica, CA, 90401

[e]Southern California Evidence Review Center, University of Southern California, Keck School of Medicine, 2001 N Soto Street, Los Angeles, CA, 90033

Corresponding author: Courtney Gidengil, MD MPH

RAND Corporation

20 Park Plaza, Suite 920

Boston, MA 02116

Tel: 617-338-2059

Email: gidengil@rand.org

Safety of Vaccines Used for Routine Immunization in the United States: An Updated Systematic Review and Meta-analysis

Courtney Gidengil, M.D., M.P.H.,[a,b] Matthew Bidwell Goetz, M.D.,[c] Sydne Newberry, Ph.D.,[d] Margaret Maglione, M.P.P.,[d] Owen Hall, M.S.,[d] Jody Larkin, M.S.,[d] Aneesa Motala, B.A.,[d,e] Susanne Hempel, Ph.D.[d,e]

[a]RAND Corporation, 20 Park Plaza, Suite 920, Boston, MA, 02116

[b]Boston Children's Hospital, 300 Longwood Avenue, Boston, MA, 02115

[c]VA Greater Los Angeles Healthcare System and David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA, 90073

[d]RAND Corporation, 1776 Main Street, Santa Monica, CA, 90401

[e]Southern California Evidence Review Center, University of Southern California, Keck School of Medicine, 2001 N Soto Street, Los Angeles, CA, 90033

Corresponding author: Courtney Gidengil, MD MPH

RAND Corporation

20 Park Plaza, Suite 920

Boston, MA 02116

Tel: 617-338-2059

Email: gidengil@rand.org

Abstract

Background: Understanding the safety of vaccines is critical to inform decisions about vaccination. Our objective was to conduct a systematic review of the safety of vaccines recommended for children, adults, and pregnant women in the United States.

Methods: We searched the literature in November 2020 to update a 2014 Agency for Healthcare Research and Quality review by integrating newly available data. Comparative studies that reported the presence or absence of key adverse events were eligible. Adhering to Evidence-based Practice Center methodology, we assessed the strength of evidence (SoE) for all evidence statements. The systematic review is registered in PROSPERO (CRD42020180089).

Results: Of 56,603 reviewed citations, 338 studies reported in 518 publications met inclusion criteria. For children, SoE was high for no increased risk of autism following measles, mumps, and rubella (MMR) vaccine. SoE was high for increased risk of febrile seizures with MMR. Rotavirus vaccine was not associated with intussusception at latest time of follow-up (moderate SoE), nor with increased risk of diabetes (high SoE). There was no evidence of increased risk or insufficient evidence for key adverse events for newer vaccines such as 9-valent human papillomavirus and meningococcal B vaccines. For adults, there was no evidence of increased risk (varied SoE) for key adverse events for the new recombinant adjuvanted zoster vaccine and hepatitis B vaccine with novel immunostimulatory adjuvant. We found no evidence of increased risk (varied SoE) for key adverse events among pregnant women following tetanus, diphtheria, and acellular pertussis vaccine, including for preterm labor and stillbirth (moderate SoE).

Conclusions: Across a large body of research we found few associations of vaccines and serious key adverse events; however, rare events are challenging to study. Any adverse events should be weighed against the protective benefits that vaccines provide.

Keywords: vaccine safety; meta-analysis; systematic review; childhood vaccines; pregnancy

Introduction

Vaccines are considered one of the greatest public health achievements and the effectiveness of vaccines in controlling the spread of and even eradicating many infectious diseases is widely acknowledged [1]. Although vaccination rates for children remain high, parents and caregivers still express worries about the safety of childhood vaccines [2-4]. Vaccination rates for adults lag well behind those for children [5]. Only about a third of pregnant women receive both tetanus, diphtheria, and acellular pertussis (Tdap) and influenza vaccines as indicated during their pregnancies [6], due in part to safety concerns [7].

Safety concerns about vaccines have persisted in spite of the rigorous, transparent processes that vaccines must undergo, overseen in the United States by the U.S. Food and Drug Administration (FDA) [8]. Once a vaccine is licensed and recommended for use following clinical trials, multiple systems are in place to ensure ongoing assessments of safety through Phase IV studies [9], including post-licensure safety surveillance [10] and the FDA's Post-Licensure Rapid Immunization Monitoring (PRISM) system [11-13]. Multiple databases contribute to surveillance, such as the Vaccine Adverse Event Reporting System (VAERS) [14], Vaccine Safety Datalink [15, 16], and Clinical Immunization Safety Assessment project [17, 18].

Reassurance of vaccine safety remains critical for population health in the context of an evolving vaccine landscape and notably the emergence of vaccines against the novel severe acute respiratory syndrome coronavirus (SARS-CoV-2). The purpose of this systematic review was to assess the evidence regarding the safety of vaccines routinely recommended for adults, children, and pregnant women in the United States.

Methods

The evidence review assessed and examined adverse events potentially associated with vaccines to determine the safety of vaccines in adults, children, and pregnant women, following the Agency for Healthcare Research and Quality's Methods Guide for Effectiveness and Comparative Effectiveness Reviews [19] (full details can be found in the online Appendix). The list of included vaccines comprises those licensed by the FDA [20] and included in the CDC's immunization schedules as of November 2020 (Table 1) [21, 22].

This update builds on a 2014 report on the safety of vaccines requested by AHRQ [23], supporting the Office of the Assistant Secretary of Health's Office of Infectious Disease and HIV/AIDS Policy (OASH/OIDP). The 2014 report built upon a 2011 Institute of Medicine consensus report [24]; the prior 2014 report did not search for or include studies on vaccines that were covered in the IOM report and published prior to 2011. Similarly, in this update only for vaccines for which there were new indications (or for new vaccines) did we perform targeted searches for research published prior to 2014. The review is registered in PROSPERO (CRD42020180089) [25], and the review protocol is posted on the EHC website [26].

We searched MEDLINE (including TOXLINE), Embase, CINAHL, Cochrane CENTRAL, Web of Science, and Scopus, through November 2020 (see Appendix for full search strategy). We searched broadly and did not rely on filters for adverse events. Instead, all evaluations of vaccines were obtained and the full text screened for information on adverse events. We reference-mined existing systematic reviews and Advisory Committee on Immunization Practices statements; screened Clinicaltrials.gov; reviewed supplemental material from authors and industry submitted to AHRQ; and consulted with content experts. Experimental and observational studies with a concurrent or historic comparator that reported the presence or absence of adverse events (e.g., self-controlled studies such as those conducted by the Vaccine

Safety Datalink [15]) met inclusion criteria. The update allowed for control groups receiving either no vaccine or standard of care (i.e., the previously available vaccine) as comparators. With the assistance of a technical expert panel–comprised of vaccine experts with particular clinical expertise in key populations (children, adults, older adults, and pregnant women), vaccine safety methodologists, and consumers–we determined a set of key adverse events *a priori* to allow an unbiased synthesis across studies.

Two trained reviewers (with Master's degrees and experience in systematic reviews) independently screened the citations and full text publications. Data were abstracted by an experienced subject matter expert with clinical and research expertise in vaccines (C.G.). For each key adverse event, we computed the relative risk (RR) and 95% confidence intervals (CI) of the adverse event among those who received the vaccine of interest compared to controls across all studies. We combined estimates across studies in random effects meta-analyses using Hartung-Knapp correction of standard errors where appropriate. For cases with zero events across studies, we added a constant to the empty cell to enable computation. We determined the most appropriate meta-analysis model (see Appendix), given that for many adverse events only a small number of studies were available, studies reported on rare events, and several studies reported zero events [27-29]. Where studies did not report sufficient detail and could not be combined into the meta-analysis, we reported the risk estimates provided by the authors.

All studies that reported rates of adverse events that could be computed were combined in meta-analyses. When studies could not be combined statistically, we narratively synthesized the findings to inform the strength of evidence (SoE) assessment and ensure that all available evidence was considered. For the synthesis we determined whether there was evidence of an increased risk of adverse events relative to a control group. In addition to the relative effect, we

also documented the actual incidences, sample sizes, and resulting rates of adverse events in the vaccinated and control groups for each individual study where available. We reviewed all instances where the vaccinated group had reported more instances of adverse events in detail. In the absence of evidence of an increased risk across studies, we also reviewed the risk reported in individual studies and documented the observed rates. For estimates that were imprecise–given the small number of reported events and the small number of samples from which conclusions for the true risk could be estimated–the narrative synthesis also reports observed rates to transparently document the available evidence.

We used McHarm [30] for critical appraisal of individual studies, rating studies that reported timing and severity and used standard, precise definitions of adverse events higher than studies that did not. The body of evidence was assessed based on AHRQ Evidence-based Practice Center grading [31]. We used four criteria to grade the SoE: (1) study limitations; (2) consistency; (3) precision; and (4) reporting bias. We differentiated *high*, *moderate*, *low*, and *insufficient* evidence to communicate the confidence for the findings across studies. *High* confidence indicates that the evidence reflects the true effect; further research is very unlikely to change our confidence in the estimate of effect. *Moderate* confidence indicates that the evidence reflects the true effect; further research may change our confidence in the estimate of effect and may change the estimate. *Low* confidence indicates that the evidence reflects the true effect; further research is likely to change our confidence in the estimate of effect and is likely to change the estimate. *Insufficient* indicates that evidence either is unavailable or does not permit a conclusion.

Findings are reported below for the selected key adverse events (adverse events identified in the prior report that were not selected as key adverse events for this update are included in the

Appendix). We report effect estimates (RR and 95% CI) that could be computed for findings of moderate or high SoE across studies; we also report findings that were of low SoE, but not the effect estimates.

Results

Of 56,603 reviewed citations, 189 new studies met inclusion criteria in this update for a total of 338 studies reported in 518 publications across the prior report and update (Figure 1) [32-549]. Study designs included RCTs, cohort studies, pre-post designs, case-control designs, non-randomized controlled clinical trials, and self-controlled studies (either self-controlled risk interval or self-controlled case series analyses). Many studies followed patients for six months or longer, and some for up to 15 years to record emerging adverse events.

The methodological rigor and reporting of the adverse events over 15 assessed domains varied widely across studies (Appendix Table 1; Appendix Figure 1). Most studies reported the timing and frequency of the adverse events assessment, but few reported the qualifications of the outcome assessors.

Full study characteristics can be found in Appendix Tables 2, 3, and 4.

Safety of vaccines included in the routine immunization schedule in children

A summary of the strength of evidence for the findings can be found in Table 2 (all effect estimates and assessments of the quality of the evidence are in Appendix Table 5, followed by synthesis of the SoE across the prior report and update in Appendix Table 5a).

9-valent human papillomavirus vaccine. All but one study compared 9-valent human papillomavirus vaccine to 2- or 4-valent vaccines. We also reviewed studies that combined

children and adults. There was no evidence of increased risk of autoimmune disease, birth defects, death, reproductive system events, seizures, or spontaneous abortion (all low SoE).

13-valent pneumococcal vaccine. Risk estimates were based on comparisons of 13-valent pneumococcal vaccine to 7-valent pneumococcal vaccine, except for death. There was no evidence of increased risk of death (RR 2.02; CI 0.07, 59.88; risk estimate based on 1 study; moderate SoE assessed across all 5 available studies). The risk estimate was imprecise as the sample size was small with only one event (1/193 vs 0/195). There was also no evidence of increased risk of asthma, cardiovascular events, intussusception, meningitis, reproductive system events, or seizures (all low SoE). There was an increased risk of febrile seizures (low SoE). There was insufficient evidence for 23-valent pneumococcal polysaccharide vaccine for the outcomes of interest.

Diphtheria, tetanus, and pertussis (DTaP) vaccine. There was no evidence of increased risk of type 1 diabetes mellitus (moderate SoE, effect estimate N/A). There was no evidence of increased risk of asthma or death (low SoE).

Tetanus, diphtheria & acellular pertussis vaccine. There was no evidence of increased risk of cardiovascular events or death (low SoE).

***Haemophilus influenzae* type b vaccine**. There was no evidence of increased risk of serious adverse events in the short term (moderate SoE; effect estimates N/A).

Hepatitis A vaccine. There was an increased risk of idiopathic thrombocytopenic purpura (moderate SoE, effect estimate N/A).

Hepatitis B vaccine. There was no evidence of increased risk of multiple sclerosis (moderate SoE, effect estimate N/A).

185 **Inactivated poliovirus vaccine.** There was insufficient evidence for conclusions about
186 increased risk of key adverse events.
187 **Quadrivalent influenza vaccines (IIV)**. Quadrivalent IIV was compared to trivalent IIV in
188 all but one of the studies that contributed to risk estimates (this study only contributed to the risk
189 estimate for death). There was no evidence of increased risk of death (RR 1.08; CI 0.02, 53.95;
190 estimate based on 1 study; moderate SoE across 6 studies). The risk estimate was imprecise
191 because it was based on one small study with no deaths (0/99 vs 0/107). There was no evidence
192 of increased risk of anaphylaxis or systemic allergic reaction, asthma, autoimmune disease,
193 cardiovascular events, febrile seizures, or seizures (low SoE).
194 Quadrivalent live attenuated influenza vaccine (LAIV) was compared to placebo or no
195 vaccine in some studies, or another influenza vaccine (trivalent LAIV or IIV) in other studies.
196 There was no evidence of increased risk of death (when compared to trivalent LAIV) or seizures
197 (when compared to placebo or no vaccine) (low SoE).
198 **Measles, mumps, and rubella vaccine**. There was no evidence of an association with autism
199 (RR 0.60; CI 0.09, 4.12; 2 studies; high SoE across studies from 2014 report and update). There
200 was an increased risk of anaphylaxis (in children with allergies; high SoE), febrile seizures (high
201 SoE), and idiopathic thrombocytopenic purpura (moderate SoE) (effect estimates N/A).
202 **Serogroup A, C, W, and Y meningococcal vaccines.** Some studies of serogroup A, C, W,
203 and Y meningococcal vaccines used another meningococcal vaccine as an active comparator,
204 while others used a non-active comparator (placebo or a base treatment received by both
205 intervention and control groups). All estimates below are based on studies of children with a
206 non-active comparator, but studies where an active comparator was used as well as studies of
207 both children and adults also contribute to the SoE. There was no evidence of increased risk of

cardiovascular events (RR 0.34; CI 0.02, 5.46; estimate based on 1 study; moderate SoE across 3 studies), febrile seizures (RR 0.51; CI 0.18, 1.44; based on 1 study; moderate SoE across 4 studies), intussusception (RR 0.46; CI 0.10, 2.03; 1 study; moderate SoE), idiopathic thrombocytopenic purpura (RR 0.17; CI 0.01, 5.09; based on 1 study; moderate SoE across 3 studies), Kawasaki disease (RR 1.37; CI 0.15, 12.22; based on 1 study; moderate SoE across 2 studies), or seizures (RR 1.51; CI 0.05, 44.86; based on 1 study; moderate SoE across 7 studies).

There was no evidence of increased risk of diabetes (RR 1.32; CI 0.00, 21861366; based on 2 studies; moderate SoE across 6 studies) but the risk estimate was imprecise due to small samples and few or no cases in the vaccinated and unvaccinated groups (1/396 vs 0/397; 0/392 vs 0/296).

There was also no evidence of increased risk of acute disseminated encephalomyelitis, asthma, autoimmune disease, death, encephalitis/encephalopathy, meningitis, multiple sclerosis, reproductive system events, or transverse myelitis (low SoE). There was increased risk of anaphylaxis in children with allergies (moderate SoE; effect estimate N/A), but there was no evidence of increased risk among all children (low SoE).

Serogroup B meningococcal vaccine. There was no evidence of increased risk of anaphylaxis or systemic allergic reaction (RR 0.56; CI 0.00, 34735108; 2 studies; moderate SoE), but the risk estimate was imprecise due to no cases in the vaccinated and unvaccinated groups (0/198 vs 0/121; 0/992 vs 0/501). There was no evidence of increased risk of reproductive system events (RR 0.89; CI 0.01, 65.20; 3 studies; moderate SoE); again, the risk estimate was imprecise, in this case due to small samples and few or no cases in the vaccinated and unvaccinated groups (1/198 vs 0/121; 1/174 vs 0/99; 0/374 vs 1/378). There was no evidence of increased risk of asthma, death, or seizures (low SoE).

Rotavirus vaccine. We found no evidence of increased risk of intussusception across all studies that could be combined for an estimate (RR 0.65; CI 0.41, 1.05; based on 19 studies; moderate SoE across 38 studies), though some observational studies indicated increased risk, particularly around the first dose. There was no evidence of increased risk of asthma (RR 1.33; CI 0.65, 2.72; 5 studies; moderate SoE), autoimmune disease (RR 0.65; CI 0.16, 2.67; 2 studies; moderate SoE), death (RR 1.05; CI 0.82, 1.35; based on 14 studies; moderate SoE across 15 studies), diabetes (RR 0.74; CI 0.45, 1.22; based on 3 studies; high SoE across 4 studies), febrile seizures (RR 0.82; CI 0.33, 2.05; based on 7 studies; moderate SoE across 9 studies), or seizures (RR 1.02; CI 0.25, 4.16; based on 5 studies; moderate SoE across 8 studies).

There was no evidence of increased risk of encephalitis/encephalopathy (RR 0.67; CI 0.00, 85995; based on 2 studies; moderate SoE across 4 studies), but the risk estimate was imprecise due to few or no cases in the vaccinated and unvaccinated groups (1/1647 vs 2/1641;1/34904 vs 1/34862). There was no evidence of increased risk of idiopathic thrombocytopenic purpura (RR 0.64; CI 0.00, 1778394; 2 studies; moderate SoE); again, the risk estimate was imprecise due there being few or no cases in the vaccinated and unvaccinated groups (1/34904 vs 0/34862; 0/4359 vs 2/4328). There was no evidence of increased risk of stroke (RR 1.32; CI 0.00, 1459247; 2 studies; moderate SoE). The risk estimate is imprecise given the small number of studies and studies reporting few or no cases in the vaccinated and unvaccinated groups (1/34904 vs 0/34862; 1/1666 vs 1/1667).

There was no evidence of increased risk of anaphylaxis or systemic allergic reaction, autoimmune thyroiditis (Hashimoto's disease), Kawasaki disease, meningitis, or reproductive system events (all low SoE).

Varicella vaccine. There was evidence of increased risk of anaphylaxis (high SoE) and idiopathic thrombocytopenic purpura (among children aged 11-17 years; moderate SoE) (effect estimates N/A).

Safety of vaccines included in the routine immunization schedule in adults

A summary of the strength of evidence for the findings can be found in Table 3 (all effect estimates and assessments of the quality of the evidence are in Appendix Table 6, followed by synthesis of the SoE in Appendix Table 6a).

13-valent pneumococcal conjugate vaccine. Some studies of 13-valent pneumococcal conjugate vaccine used another pneumococcal vaccine as an active comparator, while others used a non-active comparator (placebo or a base treatment received by both intervention and control groups). All risk estimates below are based on studies with a non-active comparator, but studies where an active comparator was used also contribute to the SoE. There was no evidence of increased risk of cardiovascular events (RR 0.97; CI 0.58, 1.64; based on 4 studies; moderate SoE across 6 studies), myocardial infarction (RR 1.76; CI 0.42, 7.39; based on 4 studies; moderate SoE across 6 studies), or reproductive system events (RR 0.59; CI 0.01, 42.46; based on 3 studies; moderate SoE across 5 studies).

There was no evidence of increased risk of herpes zoster (RR 1.49; CI 0.00, 24855526; 2 studies; moderate SoE). The risk estimate was imprecise as only two studies reported on the outcome with few or no cases occurring in the vaccinated and unvaccinated groups (0/576 vs 0/575; 1/42237 vs 0/42255). There was also no evidence of increased risk of or stroke (RR 1.12; CI 0.00, 451; 2 studies; moderate SoE); the risk estimate was imprecise due to no events in one study (0/551 vs 0/560), and a large sample size with a small number of events in the other (9/42237 vs 8/42255).

We found no evidence of increased risk of acute disseminated encephalomyelitis, anaphylaxis or systemic allergic reactions, asthma, autoimmune disease, death, encephalitis/encephalopathy, idiopathic thrombocytopenic purpura, meningitis, or seizures (low SoE).

23-valent pneumococcal polysaccharide vaccine. We found no evidence of increased risk of death (RR 1.45; CI 0.00, 3455; based on 2 studies; moderate SoE across 4 studies). The risk estimate was imprecise due to a very small study with no events (0/19 vs 0/21) and a larger study with few events (6/725 vs 4/724). We also found high SoE for no evidence of increased risk of cardiovascular events (RR 0.46; CI 0.27, 0.76; based on 4 studies; high SoE across 8 studies) or cerebrovascular events (effect estimate N/A) in people aged 65 years and older.

Hepatitis B vaccine. For HEPLISAV-B (which was compared to previously available hepatitis B vaccines), there was no evidence of increased risk for asthma, autoimmune disease, cardiovascular events, death, herpes zoster, reproductive system events, and stroke (low SoE). For all hepatitis B vaccine, there was no evidence of increased risk of diabetes (RR 0.61; CI 0.55, 0.67; based on 1 study comparing hepatitis B vaccines to no vaccine; moderate SoE across 2 studies). For hepatitis B vaccines (not including HEPLISAV-B) there was no increased risk of multiple sclerosis, but there was increased risk of anaphylaxis in patients allergic to yeast (both moderate SoE; effect estimates N/A).

Influenza vaccines (IIV). Influenza vaccines were compared to an active comparator (either trivalent influenza vaccine or another influenza vaccine). For quadrivalent IIV, we identified no evidence of increased risk of asthma, cardiovascular events, death, myocardial infarction, reproductive system events, seizures, or stroke (low SoE). For adjuvanted IIV (either trivalent or quadrivalent), there was no evidence of stroke (RR 1.18; CI 0.00, 33607; based on 2 studies;

moderate SoE across 3 studies). The risk estimate was imprecise as only two studies reported on the outcome with few or no cases occurring in the vaccinated and comparator groups (0/888 vs 1/888, 3/3545 vs 2/3537). There was no evidence of increased risk of asthma, autoimmune disease, cardiovascular events, death, encephalitis/encephalopathy, Guillain-Barré syndrome, idiopathic thrombocytopenic purpura, myocardial infarction, or seizures (low SoE). For quadrivalent recombinant influenza vaccine, there was no evidence of increased risk of cardiovascular events, death, encephalitis/encephalopathy, myocardial infarction, reproductive system events, or stroke (low SoE). There was insufficient evidence for conclusions about increased risk of key adverse events for quadrivalent live attenuated influenza vaccine.

Measles, mumps, rubella vaccine. There was no evidence of increased risk of type 1 diabetes mellitus (moderate SoE, effect estimate N/A).

Serogroups A, C, W and Y meningococcal vaccines. Some studies of serogroup A, C, W, and Y meningococcal vaccines used another meningococcal vaccine as an active comparator, while others used a non-active comparator (placebo or a base treatment received by both intervention and control groups); all risk estimates below are based on studies with a non-active comparator, but studies where an active comparator was used also contribute to the SoE. There was no evidence of increased risk of death (RR 0.99; CI 0.00, 60563320; based on 2 studies; moderate SoE across 4 studies). The risk estimate was imprecise due to two small studies with no events (00/99 vs 0/100; 0/85 vs 0/84). There was no evidence of increased risk of cardiovascular events, myocardial infarction, or stroke (all low SoE).

Tetanus, diphtheria, and acellular pertussis and tetanus and diphtheria vaccines. There was evidence of increased risk of anaphylaxis (high SoE, effect estimate N/A).

Recombinant zoster vaccine. We found moderate SoE of no evidence of increased risk of cardiovascular events (RR 0.89; CI 0.66, 1.21; 3 studies), death (RR 0.93; CI 0.78, 1.11; 4 studies), myocardial infarction (RR 0.89; CI 0.38, 2.05; 3 studies), or reproductive system events (RR 1.04; CI 0.03, 37.17; 2 studies).

For all other adverse events for which there was moderate SoE, the confidence intervals were wide because the risk estimate was based on two studies with few or no events occurring in the vaccinated and non-vaccinated groups: amyotrophic lateral sclerosis (RR 2.60; CI 0.00, 571537; 2/6950 vs 0/6950, 2/7695 vs 1/7710), anaphylaxis or systemic allergic reaction (RR 1.32; CI 0.00, 1463200; 1/6950 vs 1/6950, 1/7695 vs 0/7710), asthma (RR 0.90; CI 0.00, 493; 2/6950 vs 4/6950, 6/7695 vs 5/7710), diabetes (RR 1.00; CI 0.00, 606; 5/6950 vs 6/6950, 3/7695 vs 2/7710), encephalitis/encephalopathy (RR 0.50; CI 0.00, 2867570; 0/6950 vs 1/6950, 0/7695 vs 1/7710), Guillain-Barré syndrome (RR 0.67; CI 0.00, 86459; 1/6950 vs 2/6950, 1/7695 vs 1/7710), idiopathic thrombocytopenic purpura (RR 2.65; CI 0.00, 530690; 1/6950 vs 0/6950, 3/7695 vs 1/7710), meningitis (RR 0.50; CI 0.00, 2867570; 0/6950 vs 1/6950, 0/7695 vs 1/7710), seizures (RR 1.34; CI 0.00, 13492; 2/6950 vs 0/6950, 3/7695 vs 3/7710), or stroke (RR 1.44; CI 0.03, 71.52; 7/6950 vs 6/6950, 19/7695 vs 12/7710).

We found no evidence of increased risk of herpes zoster (RR 0.09; CI 0.02, 0.30; 5 studies; high SoE). There was no evidence of increased risk of acute disseminated encephalomyelitis, angioedema, ataxia, autoimmune disease, or autoimmune thyroiditis (low SoE).

9-valent human papillomavirus vaccine, hepatitis A vaccine, combination hepatitis A and hepatitis B vaccine, serogroup B meningococcal vaccine, and varicella vaccine. Evidence was insufficient to draw conclusions about increased risk of key adverse events based on studies of adults only.

Safety of vaccines included in the routine immunization schedule in pregnant women (both for the woman and her fetus)

A summary of the strength of evidence for the findings is in Table 4 (all effect estimates and assessments of the quality of the evidence are in Appendix Table 7, followed by synthesis of the SoE in Appendix Table 7a).

We found insufficient evidence to draw conclusions about increased risk of key adverse events for hepatitis B vaccine, quadrivalent inactivated influenza vaccines, or quadrivalent recombinant influenza vaccine in pregnant women.

All studies of Tdap compared to either placebo or base treatment also received by the control groups, except for one study that compared Tdap and Td. There was no evidence of increased risk for maternal cardiovascular events (RR 0.86; CI 0.41, 1.84; 6 studies), maternal death (RR 1.52; CI 0.07, 32.25; 4 studies), maternal diabetes (RR 0.98; CI 0.88, 1.10; 4 studies), eclampsia/pre-eclampsia (RR 0.96; CI 0.92, 1.01; 6 studies), preterm labor (RR 0.62; CI 0.46, 0.82; 10 studies), maternal reproductive system events (RR 0.52; CI 0.05, 5.91; 3 studies), stillbirth (RR 0.44; CI 0.11, 1.80; 6 studies), cardiovascular events in infants (RR 0.77; CI 0.50, 1.20; 4 studies), death in infants (RR 0.15; CI 0.00, 8.88; 3 studies), encephalitis/encephalopathy in infants (RR 1.23; CI 0.60, 2.54; 4 studies), or seizures in infants (RR 1.02; CI 0.76, 1.35; 3 studies) (all moderate SoE). There was also no evidence of increased risk of maternal encephalitis/encephalopathy, autism in infants, birth defects in infants, or febrile seizures in infants (low SoE).

Discussion

We assessed the evidence for the safety of vaccines currently used for routine immunization in the United States among children, adults, and pregnant women. We conducted extensive literature searches, screened 56,603 citations, and abstracted 338 studies reported in 518 publications.

Overall, our evidence review found vaccines to be safe across populations with serious adverse events being rare, consistent with other recent systematic reviews of vaccine safety [550]. For adults, there was no evidence of increased risk (varied SoE) or insufficient evidence for the new recombinant adjuvanted zoster vaccine and hepatitis B vaccine with novel immunostimulatory adjuvant. We found either no evidence of increased risk or insufficient evidence among pregnant women following Tdap, including for preterm labor and stillbirth (moderate SoE).

For children, across all studies SoE was high for no increased risk of autism following measles, mumps, and rubella (MMR) vaccine. SoE was high for increased risk of febrile seizures with MMR. There was no evidence of increased risk (varied SoE) or insufficient evidence for key adverse events for the newer vaccines such as 9-valent human papillomavirus and meningococcal B vaccines. We found high SoE for no increased risk of diabetes following rotavirus vaccine, and moderate SoE for no increased risk of other adverse events, such as autoimmune disease and idiopathic thrombocytopenic purpura. We also found no evidence of increased risk of intussusception following rotavirus vaccine at the latest time of follow-up across studies that could be pooled, consistent with a recent meta-analysis [551]. However, there were mixed findings across other studies, which included pre-post studies, cohort studies, and self-controlled case series, particularly related to the risk following the first dose. While intussusception is a known possible side effect of rotavirus vaccination (listed in the package

inserts for both vaccines and also in the Vaccine Injury Table as a condition covered under the National Vaccine Injury Compensation Program) [552] the finding that there is no increased risk with the longest-term follow-up from clinical trials is noteworthy.

Our study had some limitations. While our literature search procedures were extensive, some unpublished data may not have been identified, although we mitigated this by searching trial registries. The importance of trial registries has increased dramatically since reporting of results has become mandatory. Clinicaltrials.gov is set up to capture results that can be used in systematic reviews and meta-analyses, including data on severe adverse events, serious adverse events, and mortality. In general, the harms data in Clinicaltrials.gov have been found to be more complete than in the corresponding publications,[553, 554] although we note that the database tends to better capture the presence of reported adverse events than the absence of such events.

However, trials often have insufficient sample sizes to identify rare adverse events and may not follow participants long enough to identify long-term sequelae; even in studies with generous follow-up times, timing of events is not always optimally reported. Indeed, many of the harms we assessed as key adverse events (e.g., acute disseminated encephalomyelitis, Guillain-Barré syndrome, transverse myelitis, anaphylaxis) are quite rare and the number of studies that reported on the events for a vaccine was often small. As a result, despite our extensive searches for data that could be combined across studies, our confidence intervals are often wide and the SoE often low or insufficient. Given the limitations of controlled trials, we included post-marketing surveillance and self-controlled analyses (if they met inclusion criteria) when grading the SoE. For example, in the United States the CDC's Vaccine Safety Datalink uses data obtained from eight large health care organizations, enabling studies that may be particularly useful for identifying safety signals and/or investigating concerns for rare serious adverse events. Such

innovative methodologic approaches have improved the analysis of rare adverse events, particularly in the post-marketing phase.

We also may have missed studies due to the challenging nature of assessing harms (as contrasted with assessing effectiveness); however, we screened the full text of all identified vaccine intervention studies, and our search terms did not include safety terms in order not to miss relevant data. Wherever possible, we used data that could be combined in meta-analyses to estimate the relative risk based on all available research studies. When we could not combine data in pooled estimates, we integrated findings (including from the prior 2014 report) in a narrative synthesis to inform the SoE.

This review excluded studies of vaccines not currently in use in the United States and cannot make evidence statements for other vaccine schedules. We also excluded non-English language studies. Although we considered only vaccines approved for use in the United States, it is possible relevant epidemiological studies have been published in non-English journals.

Careful consideration should be given to research gaps, including where the evidence was insufficient to assess the potential associations between some vaccines and particular adverse events and/or where confidence intervals around risk estimates were extremely wide. However, when deciding whether studies are warranted, important factors to consider include the severity and frequency of the adverse event being studied and the challenges of conducting sufficiently powered studies when investigating rare events. Given the rare nature of some of the serious adverse events of interest (e.g., anaphylaxis, immune thrombocytopenia, Guillain-Barré syndrome), ongoing studies of large populations and post-marketing surveillance of vaccines after FDA licensure as noted earlier are needed to identify uncommon adverse events. Future

vaccine research will also need to take into account the expanding landscape of new vaccines and vaccine technologies, in particular the new COVID-19 vaccines [555].

Conclusion

Across a large body of research, we found few instances in which vaccines are rarely associated with serious adverse events; however, potential risks for rare adverse events should be weighed carefully against the protective benefits that those vaccines provide.

Acknowledgments

The authors gratefully acknowledge the following individuals for their contributions to this project: Kim Wittenberg, Stephanie Chang, Christine Chang, Thomas Acciani, David Kim, Kara Elam, Tammy Beckham, Mark Helfand, and Ethan Balk for helpful comments; Jeremy Miles for statistical consultation; Peggy Chen, Alicia Ruelaz Maher, Kelsey O'Hollaren, Nabeel Qureshi, Keller Scholl, Olamigoke Akinniranye, Tim M. Kim, Oluwafemi Jimoh, Lea Xenakis, Weilong (David) Kong, Zichun Xu, and Sangita Baxi for research assistance; and Judy Bearer for administrative assistance.

In addition, the authors would like to thank the Technical Expert Panel while acknowledging that the study questions, design, methodologic approaches, and/or conclusions do not necessarily represent the views of individual technical and content experts. The list of technical experts who provided input for the AHRQ report are as follows: Meghan Baker, Sarah Coles, Frank Destefano, Janet McElhaney, Rebecca Reindel, Thomas Yoshikawa, and Ousseny Zerbo. The authors also sought additional peer review for the AHRQ report and would like to thank the following individuals: Haitao Chu, Tamera Coyne-Beasley, Janet Cragan, Matthew Daley, Cody Meissner, and Barbara Mulach.

***Conflict of interest*:** The authors have no potential conflicts of interest to report, including financial interests, activities, relationships, and affiliations.

Funding source: This study was supported by a contract from the Agency for Healthcare Research and Quality (AHRQ), Rockville, MD (Contract No. 290-2015-00010I). The funder had no role in the design, interpretation of results, nor the decision to submit this manuscript. The views expressed in this article are those of the authors and do not necessarily represent the views of the US government; AHRQ; or the US Department of Health and Human Services.

***Author Contributions*:** All authors attest that they meet the ICMJE criteria for authorship.

References

[1] Centers for Disease Control Prevention. Ten great public health achievements--United States, 1900-1999. MMWR Morb Mortal Wkly Rep. 1999;48:241-3.

[2] Gidengil C, Chen C, Parker AM, Nowak S, Matthews L. Beliefs around childhood vaccines in the United States: A systematic review. Vaccine. 2019;37:6793-802.

[3] Kempe A, Saville AW, Albertin C, Zimet G, Breck A, Helmkamp L, et al. Parental Hesitancy About Routine Childhood and Influenza Vaccinations: A National Survey. Pediatrics. 2020;146.

[4] Szilagyi PG, Albertin CS, Gurfinkel D, Saville AW, Vangala S, Rice JD, et al. Prevalence and characteristics of HPV vaccine hesitancy among parents of adolescents across the US. Vaccine. 2020;38:6027-37.

[5] Centers for Disease Control and Prevention. Vaccination Coverage Among Adults in the United States, National Health Interview Survey, 2017. 2018.

[6] Kahn KE, Black CL, Ding H, Williams WW, Lu PJ, Fiebelkorn AP, et al. Influenza and Tdap Vaccination Coverage Among Pregnant Women - United States, April 2018. MMWR Morb Mortal Wkly Rep. 2018;67:1055-9.

[7] Lindley MC, Kahn KE, Bardenheier BH, D'Angelo DV, Dawood FS, Fink RV, et al. Vital Signs: Burden and Prevention of Influenza and Pertussis Among Pregnant Women and Infants - United States. MMWR Morb Mortal Wkly Rep. 2019;68:885-92.

[8] Marshall V, Baylor NW. Food and Drug Administration regulation and evaluation of vaccines. Pediatrics. 2011;127 Suppl 1:S23-30.

[9] Salmon DA, Pavia A, Gellin B. Editors' introduction: Vaccine safety throughout the product life cycle. Pediatrics. 2011;127 Suppl 1:S1-4.

[10] U.S Food and Drug Administration. Postmarketing Requirements and Commitments: Introduction. 2016.

[11] Nguyen M, Ball R, Midthun K, Lieu TA. The Food and Drug Administration's Post-Licensure Rapid Immunization Safety Monitoring program: strengthening the federal vaccine safety enterprise. Pharmacoepidemiol Drug Saf. 2012;21 Suppl 1:291-7.

[12] Mini-Sentinel. 2011.

[13] U.S Food and Drug Administration. Vaccines. 2020.

[14] Sentinel. Vaccines, Blood, & Biologics Assessments.

[15] Centers for Disease Control and Prevention. Vaccine Safety Datalink (VSD). 2019.

[16] DeStefano F, Vaccine Safety Datalink Research G. The Vaccine Safety Datalink project. Pharmacoepidemiol Drug Saf. 2001;10:403-6.

[17] Centers for Disease Control and Prevention. Clinical Immunization Safety Assessment (CISA) Project. 2020.

[18] Williams SE, Klein NP, Halsey N, Dekker CL, Baxter RP, Marchant CD, et al. Overview of the Clinical Consult Case Review of adverse events following immunization: Clinical Immunization Safety Assessment (CISA) network 2004-2009. Vaccine. 2011;29:6920-7.

[19] Agency for Healthcare Research and Quality. Methods Guide for Effectiveness and Comparative Effectiveness Reviews. Rockville (MD)2008.

[20] U.S Food and Drug Administration. Vaccines Licensed for Use in the United States.

[21] Centers for Disease Control and Prevention. Immunization Schedules: Table 1. Recommended Adult Immunization Schedule for ages 19 years or older, United States, 2021.

[22] Centers for Disease Control and Prevention. Immunization Schedules: Table 1. Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2021.
[23] Maglione MA, Gidengil C, Das L, Raaen L, Smith A, Chari R, et al. Safety of Vaccines Used for Routine Immunization in the United States. Evid Rep Technol Assess (Full Rep). 2014:1-740.
[24] Institute of Medicine I. Adverse effects of vaccines: Evidence and causality. Washington, DC: The National Academy Press. 2011.
[25] Motala A, Hempel S, Gidengil C, Goetz M, Maglione M, Hall O, et al. Safety of Vaccines Used for Routine Immunization in the United States: An update. PROSPERO 2020 CRD42020180089 2020.
[26] Agency for Healthcare Research and Quality. Safety of Vaccines Used for Routine Immunization in the United States: Research Protocol. April 6, 2020.
[27] Higgins JPT, Green S, (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.
[28] IntHout J, Ioannidis JP, Borm GF. The Hartung-Knapp-Sidik-Jonkman method for random effects meta-analysis is straightforward and considerably outperforms the standard DerSimonian-Laird method. BMC Med Res Methodol. 2014;14:25.
[29] Jackson D, Law M, Rucker G, Schwarzer G. The Hartung-Knapp modification for random-effects meta-analysis: A useful refinement but are there any residual concerns? Stat Med. 2017;36:3923-34.
[30] Santaguida P, Raina P. The Development of the McHarm Quality Assessment Scale for adverse events: Delphi Consensus on important criteria for evaluating harms. McMaster University. 2012.
[31] Methods Guide for Effectiveness and Comparative Effectiveness Reviews. AHRQ Publication No. 10(14)-EHC063-EF. 2011/03/25 ed. Rockville, MD: Agency for Healthcare Research and Quality; January 2014.
[32] Abdelnour A, Silas PE, Lamas MRV, Aragon CFG, Chiu N-C, Chiu C-H, et al. Safety of a quadrivalent meningococcal serogroups A, C, W and Y conjugate vaccine (MenACWY-CRM) administered with routine infant vaccinations: results of an open-label, randomized, phase 3b controlled study in healthy infants. Vaccine. 2014;32:965-72.
[33] Acosta J, Benages C, Díaz MA, Xiberta M, Muñiz F. Preventing pertussis in the early infant: Development and results of a prenatal vaccination program. Acta Medica International. 2016;3:78-81.
[34] Alberer M, Burchard G, Jelinek T, Reisinger E, Beran J, Hlavata LC, et al. Safety and immunogenicity of typhoid fever and yellow fever vaccines when administered concomitantly with quadrivalent meningococcal ACWY glycoconjugate vaccine in healthy adults. Journal of travel medicine. 2015;22:48-56.
[35] Alberer M, Burchard G, Jelinek T, Reisinger EC, Meyer S, Forleo-Neto E, et al. Immunogenicity and safety of concomitant administration of a combined hepatitis A/B vaccine and a quadrivalent meningococcal conjugate vaccine in healthy adults. Journal of travel medicine. 2015;22:105-14.
[36] Amdekar YK, Lalwani SK, Bavdekar A, Balasubramanian S, Chhatwal J, Bhat SR, et al. Immunogenicity and safety of a 13-valent pneumococcal conjugate vaccine in healthy infants

and toddlers given with routine vaccines in India. The Pediatric infectious disease journal. 2013;32:509-16.
[37] Anez G, Hedrick J, Simon MW, Christensen S, Jeanfreau R, Yau E, et al. Immunogenicity and safety of a booster dose of a quadrivalent meningococcal tetanus toxoid-conjugate vaccine (MenACYW-TT) in adolescents and adults: a Phase III randomized study. Hum Vaccin Immunother. 2020:1-7.
[38] Baccarini CI, Simon MW, Brandon D, Christensen S, Jordanov E, Dhingra MS. Safety and Immunogenicity of a Quadrivalent Meningococcal Conjugate Vaccine in Healthy Meningococcal-Naïve Children 2-9 Years of Age: A Phase III, Randomized Study. Pediatric Infectious Disease Journal. 2020:955-60.
[39] Baker MA, Baer B, Kulldorff M, Zichittella L, Reindel R, DeLuccia S, et al. Kawasaki disease and 13-valent pneumococcal conjugate vaccination among young children: A self-controlled risk interval and cohort study with null results. PLoS medicine. 2019;16:e1002844.
[40] Baker MA, Jankosky C, Yih WK, Gruber S, Li L, Cocoros NM, et al. The risk of febrile seizures following influenza and 13-valent pneumococcal conjugate vaccines. Vaccine. 2020;38:2166-71.
[41] Bart S, Cannon K, Herrington D, Mills R, Forleo-Neto E, Lindert K, et al. Immunogenicity and safety of a cell culture-based quadrivalent influenza vaccine in adults: A Phase III, double-blind, multicenter, randomized, non-inferiority study. Human vaccines & immunotherapeutics. 2016;12:2278-88.
[42] Baxter R, Eaton A, Hansen J, Aukes L, Caspard H, Ambrose CS. Safety of quadrivalent live attenuated influenza vaccine in subjects aged 2-49years. Vaccine. 2017;35:1254-8.
[43] Baxter R, Lewis E, Goddard K, Fireman B, Bakshi N, DeStefano F, et al. Acute Demyelinating Events Following Vaccines: A Case-Centered Analysis. Clin Infect Dis. 2016;63:1456-62.
[44] Baxter R, Lewis N, Bohrer P, Harrington T, Aukes L, Klein NP. Sudden-Onset Sensorineural Hearing Loss after Immunization: A Case-Centered Analysis. Otolaryngol Head Neck Surg. 2016;155:81-6.
[45] Becerra-Culqui TA, Getahun D, Chiu V, Sy LS, Tseng HF. Prenatal Tetanus, Diphtheria, Acellular Pertussis Vaccination and Autism Spectrum Disorder. Pediatrics. 2018;142.
[46] Becerra-Culqui TA, Getahun D, Chiu V, Sy LS, Tseng HF. The Association of Prenatal Tetanus, Diphtheria, and Acellular Pertussis (Tdap) Vaccination With Attention-Deficit/Hyperactivity Disorder. American journal of epidemiology. 2020;189:1163-72.
[47] Beran J, Peeters M, Dewe W, Raupachova J, Hobzova L, Devaster JM. Immunogenicity and safety of quadrivalent versus trivalent inactivated influenza vaccine: a randomized, controlled trial in adults. BMC Infectious Diseases. 2013;13:224.
[48] Berenson AB, Hirth JM, Rahman M, Laz TH, Rupp RE, Sarpong KO. Maternal and infant outcomes among women vaccinated against pertussis during pregnancy. Human vaccines & immunotherapeutics. 2016;12:1965-71.
[49] Black S, Klein NP, Shah J, Bedell L, Karsten A, Dull PM. Immunogenicity and tolerability of a quadrivalent meningococcal glycoconjugate vaccine in children 2-10 years of age. Vaccine. 2010;28:657-63.
[50] Block SL, Falloon J, Hirschfield JA, Krilov LR, Dubovsky F, Yi T, et al. Immunogenicity and safety of a quadrivalent live attenuated influenza vaccine in children. Pediatric Infectious Disease Journal. 2012;31:745-51.

[51] Block SL, Klein NP, Sarpong K, Russell S, Fling J, Petrecz M, et al. Lot-to-lot Consistency, Safety, Tolerability and Immunogenicity of an Investigational Hexavalent Vaccine in US Infants. The Pediatric infectious disease journal. 2017;36:202-8.
[52] Block SL, Yi T, Sheldon E, Dubovsky F, Falloon J. A randomized, double-blind noninferiority study of quadrivalent live attenuated influenza vaccine in adults. Vaccine. 2011;29:9391-7.
[53] Bonten MJM, Huijts SM, Bolkenbaas M, Webber C, Patterson S, Gault S, et al. Polysaccharide conjugate vaccine against pneumococcal pneumonia in adults. The New England journal of medicine. 2015;372:1114-25.
[54] Briggs-Steinberg C, Aboudi D, Hodson G, Shah S. Clinical Tolerance of In-Neonatal Intensive Care Unit Administration of Rotavirus Vaccine. Am J Perinatol. 2019.
[55] Burke RM, Tate JE, Dahl RM, Aliabadi N, Parashar UD. Does Rotavirus Vaccination Affect Longer-Term Intussusception Risk in US Infants? J Pediatric Infect Dis Soc. 2020;9:257-60.
[56] Carlin JB, Macartney KK, Lee KJ, Quinn HE, Buttery J, Lopert R, et al. Intussusception risk and disease prevention associated with rotavirus vaccines in Australia's National Immunization Program. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America. 2013;57:1427-34.
[57] Caspard H, Steffey A, Mallory RM, Ambrose CS. Evaluation of the safety of live attenuated influenza vaccine (LAIV) in children and adolescents with asthma and high-risk conditions: a population-based prospective cohort study conducted in England with the Clinical Practice Research Datalink. BMJ Open. 2018;8:e023118.
[58] Chang L-J, Meng Y, Janosczyk H, Landolfi V, Talbot HK, Group QHDS. Safety and immunogenicity of high-dose quadrivalent influenza vaccine in adults >=65years of age: A phase 3 randomized clinical trial. Vaccine. 2019;37:5825-34.
[59] Chang LJ, Hedrick J, Christensen S, Pan J, Jordanov E, Dhingra MS. A Phase II, randomized, immunogenicity and safety study of a quadrivalent meningococcal conjugate vaccine, MenACYW-TT, in healthy adolescents in the United States. Vaccine. 2020a.
[60] Chang LJ, Hedrick J, Christensen S, Pan J, Jordanov E, Dhingra MS. A Phase II, randomized, immunogenicity and safety study of a quadrivalent meningococcal conjugate vaccine, MenACYW-TT, in healthy adolescents in the United States. Vaccine. 2020b.
[61] Chang YC, Chou YJ, Liu JY, Yeh TF, Huang N. Additive benefits of pneumococcal and influenza vaccines among elderly persons aged 75 years or older in Taiwan - A representative population-based comparative study. J Infect. 2012;65:231-8.
[62] Chen LF, Chen HP, Huang YS, Huang KY, Chou P, Lee CC. Pneumococcal pneumonia and the risk of stroke: a population-based follow-up study. PLoS One. 2012;7:e51452.
[63] Chlibek R, Bayas JM, Collins H, de la Pinta MLR, Ledent E, Mols JF, et al. Safety and immunogenicity of an AS01-adjuvanted varicella-zoster virus subunit candidate vaccine against herpes zoster in adults >=50 years of age. The Journal of infectious diseases. 2013;208:1953-61.
[64] Contopoulos-Ioannidis DG, Halpern MS, Maldonado Y. Trends in Hospitalizations for Intussusception in California in Relationship to the Introduction of New Rotavirus Vaccines, 1985-2010. The Pediatric infectious disease journal. 2015;34:712-7.
[65] Cowling BJ, Perera RAPM, Valkenburg SA, Leung NHL, Iuliano AD, Tam YH, et al. Comparative Immunogenicity of Several Enhanced Influenza Vaccine Options for Older Adults: A Randomized, Controlled Trial. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America. 2019.

[66] Cunningham AL, Lal H, Kovac M, Chlibek R, Hwang S-J, Diez-Domingo J, et al. Efficacy of the Herpes Zoster Subunit Vaccine in Adults 70 Years of Age or Older. The New England journal of medicine. 2016;375:1019-32.
[67] Cutland CL, Nolan T, Halperin SA, Kurugol Z, Ahmed K, Perrett KP, et al. Immunogenicity and safety of one or two doses of the quadrivalent meningococcal vaccine MenACWY-TT given alone or with the 13-valent pneumococcal conjugate vaccine in toddlers: A phase III, open-label, randomised study. Vaccine. 2018;36:1908-16.
[68] Dagan R, Patterson S, Juergens C, Greenberg D, Givon-Lavi N, Porat N, et al. Comparative immunogenicity and efficacy of 13-valent and 7-valent pneumococcal conjugate vaccines in reducing nasopharyngeal colonization: a randomized double-blind trial. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America. 2013;57:952-62.
[69] Daley MF, Yih WK, Glanz JM, Hambidge SJ, Narwaney KJ, Yin R, et al. Safety of diphtheria, tetanus, acellular pertussis and inactivated poliovirus (DTaP-IPV) vaccine. Vaccine. 2014;32:3019-24.
[70] Deichmann KA, Ferrera G, Tran C, Thomas S, Eymin C, Baudin M. Immunogenicity and safety of a combined measles, mumps, rubella and varicella live vaccine (ProQuad) administered concomitantly with a booster dose of a hexavalent vaccine in 12-23-month-old infants. Vaccine. 2015;33:2379-86.
[71] DeMeo SD, Raman SR, Hornik CP, Wilson CC, Clark R, Smith PB. Adverse Events After Routine Immunization of Extremely Low-Birth-Weight Infants. JAMA Pediatr. 2015;169:740-5.
[72] DeSilva M, Vazquez-Benitez G, Nordin JD, Lipkind HS, Klein NP, Cheetham TC, et al. Maternal Tdap vaccination and risk of infant morbidity. Vaccine. 2017;35:3655-60.
[73] Dhingra MS, Kundu R, Gupta M, Kanungo S, Ganguly N, Singh MP, et al. Evaluation of safety and immunogenicity of a live attenuated tetravalent (G1-G4) Bovine-Human Reassortant Rotavirus vaccine (BRV-TV) in healthy Indian adults and infants. Vaccine. 2014;32 Suppl 1:A117-23.
[74] Dhingra MS, Peterson J, Hedrick J, Pan J, Neveu D, Jordanov E. Immunogenicity, safety and inter-lot consistency of a meningococcal conjugate vaccine (MenACYW-TT) in adolescents and adults: A Phase III randomized study. Vaccine. 2020;38:5194-201.
[75] Domachowske JB, Pankow-Culot H, Bautista M, Feng Y, Claeys C, Peeters M, et al. A Randomized Trial of Candidate Inactivated Quadrivalent Influenza Vaccine versus Trivalent Influenza Vaccines in Children Aged 3-17 Years. Journal of Infectious Diseases. 2013;207:1878-87.
[76] Donahue JG, Kieke BA, Lewis EM, Weintraub ES, Hanson KE, McClure DL, et al. Near real-time surveillance to assess the safety of the 9-valent human papillomavirus vaccine. Pediatrics. 2019;144.
[77] Duffy J, Weintraub E, Hambidge SJ, Jackson LA, Kharbanda EO, Klein NP, et al. Febrile Seizure Risk After Vaccination in Children 6 to 23 Months. Pediatrics. 2016;138:1-10.
[78] Dunkle LM, Izikson R, Patriarca P, Goldenthal KL, Muse D, Callahan J, et al. Efficacy of Recombinant Influenza Vaccine in Adults 50 Years of Age or Older. The New England journal of medicine. 2017;376:2427-36.
[79] Dunkle LM, Izikson R, Patriarca PA, Goldenthal KL, Muse D, Cox MMJ. Randomized Comparison of Immunogenicity and Safety of Quadrivalent Recombinant Versus Inactivated Influenza Vaccine in Healthy Adults 18-49 Years of Age. The Journal of infectious diseases. 2017;216:1219-26.

[80] Dynavax Technologies C. Safety and Immunogenicity Study of the Hepatitis B Vaccine, HEPLISAV™, Compared to Engerix-B® Vaccine. 2015.
[81] Dynavax Technologies Corporation. Safety and Efficacy of HEPLISAV™ Hepatitis B Virus Vaccine Compared With Engerix-B® Vaccine. 2006.
[82] Eriksson M, Käyhty H, Saha H, Lahdenkari M, Koskinen P, Mäkisalo H, et al. A randomized, controlled trial comparing the immunogenecity and safety of a 23-valent pneumococcal polysaccharide vaccination to a repeated dose 13-valent pneumococcal conjugate vaccination in kidney transplant recipients. Transplant Infectious Disease. 2020;22.
[83] Escolano S, Hill C, Tubert-Bitter P. Intussusception risk after RotaTeq vaccination: evaluation from worldwide spontaneous reporting data using a self-controlled case series approach. Vaccine. 2015;33:1017-20.
[84] Essink B, Fierro C, Rosen J, Figueroa AL, Zhang B, Verhoeven C, et al. Immunogenicity and safety of MF59-adjuvanted quadrivalent influenza vaccine versus standard and alternate B strain MF59-adjuvanted trivalent influenza vaccines in older adults. Vaccine. 2020;38:242-50.
[85] Esteves-Jaramillo A, Koehler T, Jeanfreau R, Neveu D, Jordanov E, Singh Dhingra M. Immunogenicity and safety of a quadrivalent meningococcal tetanus toxoid-conjugate vaccine (MenACYW-TT) in >/=56-year-olds: A Phase III randomized study. Vaccine. 2020;38:4405-11.
[86] Euctr Outside EU/EEA. Safety and Immunogenicity of Meningococcal ACWY Conjugate Versus Polysaccharide Vaccine in Children 2 to 10 Years of Age. 2014.
[87] Fernandes EG, Leshem E, Patel M, Flannery B, Pellini ACG, Veras MA, et al. Hospital-based surveillance of intussusception among infants. Jornal de pediatria. 2016;92:181-7.
[88] Fischer L, Gerstel PF, Poncet A, Siegrist C-A, Laffitte E, Gabay C, et al. Pneumococcal polysaccharide vaccination in adults undergoing immunosuppressive treatment for inflammatory diseases--a longitudinal study. Arthritis Res Ther. 2015;17:151.
[89] Fotso Kamdem A, Vidal C, Pazart L, Leroux F, Pugin A, Savet C, et al. A case-control study of risk factors for intussusception among infants in eastern France after the introduction of the rotavirus vaccine. Vaccine. 2019;37:4587-93.
[90] Frenck RW, Jr., Gurtman A, Rubino J, Smith W, van Cleeff M, Jayawardene D, et al. Randomized, controlled trial of a 13-valent pneumococcal conjugate vaccine administered concomitantly with an influenza vaccine in healthy adults. Clin Vaccine Immunol. 2012;19:1296-303.
[91] Frey SE, Reyes MRA-DL, Reynales H, Bermal NN, Nicolay U, Narasimhan V, et al. Comparison of the safety and immunogenicity of an MF59-adjuvanted with a non-adjuvanted seasonal influenza vaccine in elderly subjects. Vaccine. 2014;32:5027-34.
[92] Garland SM, Cheung T-H, McNeill S, Petersen LK, Romaguera J, Vazquez-Narvaez J, et al. Safety and immunogenicity of a 9-valent HPV vaccine in females 12-26 years of age who previously received the quadrivalent HPV vaccine. Vaccine. 2015;33:6855-64.
[93] Gasparini R, Johnston W, Conversano M, Garscadden A, Alexanderian D, Giglioli N, et al. Immunogenicity and safety of combined tetanus, reduced diphtheria, acellular pertussis vaccine when co-administered with quadrivalent meningococcal conjugate and human papillomavirus vaccines in healthy adolescents. J Vaccines Vaccin. 2014;5:231.
[94] Geier DA, Geier MR. A longitudinal cohort study of childhood MMR vaccination and seizure disorder among American children. Brain and Development. 2020.
[95] Gilca V, Sauvageau C, Panicker G, De Serres G, Ouakki M, Unger ER. Immunogenicity and safety of a mixed vaccination schedule with one dose of nonavalent and one dose of bivalent

HPV vaccine versus two doses of nonavalent vaccine - A randomized clinical trial. Vaccine. 2018;36:7017-24.
[96] Glanz JM, Clarke CL, Xu S, Daley MF, Shoup JA, Schroeder EB, et al. Association between Rotavirus Vaccination and Type 1 Diabetes in Children. JAMA Pediatrics. 2020:E1-E8.
[97] Glover C, Crawford N, Leeb A, Wood N, Macartney K. Active SMS-based surveillance of adverse events following immunisation with influenza and pertussis-containing vaccines in Australian pregnant women using AusVaxSafety. Vaccine. 2020a;38:4892-900.
[98] Greenberg DP, Robertson CA, Landolfi VA, Bhaumik A, Senders SD, Decker MD. Safety and immunogenicity of an inactivated quadrivalent influenza vaccine in children 6 months through 8 years of age. The Pediatric infectious disease journal. 2014;33:630-6.
[99] Greenberg DP, Robertson CA, Noss MJ, Blatter MM, Biedenbender R, Decker MD. Safety and immunogenicity of a quadrivalent inactivated influenza vaccine compared to licensed trivalent inactivated influenza vaccines in adults. Vaccine. 2013;31:770-6.
[100] Greenberg DP, Robertson CA, Talbot HK, Decker MD. Safety and immunogenicity of a quadrivalent influenza vaccine in adults 65 y of age and older. Human vaccines & immunotherapeutics. 2017;13:2058-64.
[101] Greenberg RN, Gurtman A, Frenck RW, Strout C, Jansen KU, Trammel J, et al. Sequential administration of 13-valent pneumococcal conjugate vaccine and 23-valent pneumococcal polysaccharide vaccine in pneumococcal vaccine-naive adults 60-64 years of age. Vaccine. 2014;32:2364-74.
[102] Griffin JB, Yu L, Watson D, Turner N, Walls T, Howe AS, et al. Pertussis Immunisation in Pregnancy Safety (PIPS) Study: A retrospective cohort study of safety outcomes in pregnant women vaccinated with Tdap vaccine. Vaccine. 2018;36:5173-9.
[103] Groome MJ, Tate JE, Arnold M, Chitnis M, Cox S, de Vos C, et al. Evaluation of intussusception after oral monovalent rotavirus vaccination in South Africa. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America. 2019.
[104] Hall C, Abramovitz LM, Bukowinski AT, Ricker AA, Khodr ZG, Gumbs GR, et al. Safety of tetanus, diphtheria, and acellular pertussis vaccination among pregnant active duty U.S. military women. Vaccine. 2020;38:1982-8.
[105] Halperin SA, Donovan C, Marshall GS, Pool V, Decker MD, Johnson DR, et al. Randomized Controlled Trial of the Safety and Immunogenicity of Revaccination With Tetanus-Diphtheria-Acellular Pertussis Vaccine (Tdap) in Adults 10 Years After a Previous Dose. Journal of the Pediatric Infectious Diseases Society. 2019;8:105-14.
[106] Halperin SA, Langley JM, Ye L, MacKinnon-Cameron D, Elsherif M, Allen VM, et al. A Randomized Controlled Trial of the Safety and Immunogenicity of Tetanus, Diphtheria, and Acellular Pertussis Vaccine Immunization During Pregnancy and Subsequent Infant Immune Response. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America. 2018;67:1063-71.
[107] Hansen J, Timbol J, Lewis N, Pool V, Decker MD, Greenberg DP, et al. Safety of DTaP-IPV/Hib vaccine administered routinely to infants and toddlers. Vaccine. 2016;34:4172-9.
[108] Hansen J, Zhang L, Eaton A, Baxter R, Robertson CA, Decker MD, et al. Post-licensure safety surveillance study of routine use of quadrivalent meningococcal diphtheria toxoid conjugate vaccine (MenACWY-D) in infants and children. Vaccine. 2018;36:2133-8.
[109] Hansen J, Zhang L, Klein NP, Robertson CA, Decker MD, Greenberg DP, et al. Post-licensure safety surveillance study of routine use of quadrivalent meningococcal diphtheria toxoid conjugate vaccine. Vaccine. 2017;35:6879-84.

[110] Hartvickson R, Cruz M, Ervin J, Brandon D, Forleo-Neto E, Dagnew AF, et al. Non-inferiority of mammalian cell-derived quadrivalent subunit influenza virus vaccines compared to trivalent subunit influenza virus vaccines in healthy children: a phase III randomized, multicenter, double-blind clinical trial. International journal of infectious diseases : IJID : official publication of the International Society for Infectious Diseases. 2015;41:65-72.
[111] Hattori F, Kawamura Y, Kawada JI, Kojima S, Natsume J, Ito K, et al. Survey of rotavirus-associated severe complications in Aichi Prefecture. Pediatr Int. 2018;60:259-63.
[112] Hawken S, Ducharme R, Rosella LC, Benchimol EI, Langley JM, Wilson K, et al. Assessing the risk of intussusception and rotavirus vaccine safety in Canada. Human vaccines & immunotherapeutics. 2017;13:703-10.
[113] Heyward WL, Kyle M, Blumenau J, Davis M, Reisinger K, Kabongo ML, et al. Immunogenicity and safety of an investigational hepatitis B vaccine with a Toll-like receptor 9 agonist adjuvant (HBsAg-1018) compared to a licensed hepatitis B vaccine in healthy adults 40-70 years of age. Vaccine. 2013;31:5300-5.
[114] Hoffman V, Abu-Elyazeed R, Enger C, Esposito DB, Doherty MC, Quinlan SC, et al. Safety study of live, oral human rotavirus vaccine: A cohort study in United States health insurance plans. Human vaccines & immunotherapeutics. 2018;14:1782-90.
[115] Huang J, Ou HY, Lin J, Karnchanasorn R, Feng W, Samoa R, et al. The impact of hepatitis B vaccination status on the risk of diabetes, implicating diabetes risk reduction by successful vaccination. PLoS ONE. 2015;10.
[116] Huang W-T, Juan Y-C, Liu C-H, Yang Y-Y, Chan KA. Intussusception and Kawasaki disease after rotavirus vaccination in Taiwanese infants. Vaccine. 2020;38:6299-303.
[117] Huh WK, Joura EA, Giuliano AR, Iversen O-E, de Andrade RP, Ault KA, et al. Final efficacy, immunogenicity, and safety analyses of a nine-valent human papillomavirus vaccine in women aged 16-26 years: a randomised, double-blind trial. Lancet (London, England). 2017;390:2143-59.
[118] Hung IFN, Leung AYM, Chu DWS, Leung D, Cheung T, Chan C-K, et al. Prevention of acute myocardial infarction and stroke among elderly persons by dual pneumococcal and influenza vaccination: a prospective cohort study. Clinical Infectious Diseases. 2010;51:1007-16.
[119] Hviid A, Hansen JV, Frisch M, Melbye M. Measles, Mumps, Rubella Vaccination and Autism: A Nationwide Cohort Study. Annals of internal medicine. 2019;170:513-20.
[120] Iwata S, Nakata S, Ukae S, Koizumi Y, Morita Y, Kuroki H, et al. Efficacy and safety of pentavalent rotavirus vaccine in Japan: a randomized, double-blind, placebo-controlled, multicenter trial. Human vaccines & immunotherapeutics. 2013;9:1626-33.
[121] Jackson LA, Gurtman A, van Cleeff M, Jansen KU, Jayawardene D, Devlin C, et al. Immunogenicity and safety of a 13-valent pneumococcal conjugate vaccine compared to a 23-valent pneumococcal polysaccharide vaccine in pneumococcal vaccine-naive adults. Vaccine. 2013;31:3577-84.
[122] Jain A, Marshall J, Buikema A, Bancroft T, Kelly JP, Newschaffer CJ. Autism occurrence by MMR vaccine status among US children with older siblings with and without autism. JAMA. 2015;313:1534-40.
[123] Juergens C, de Villiers PJT, Moodley K, Jayawardene D, Jansen KU, Scott DA, et al. Safety and immunogenicity of 13-valent pneumococcal conjugate vaccine formulations with and without aluminum phosphate and comparison of the formulation of choice with 23-valent pneumococcal polysaccharide vaccine in elderly adults: a randomized open-label trial. Human vaccines & immunotherapeutics. 2014;10:1343-53.

[124] Kantso B, Halkjaer SI, Thomsen OO, Belard E, Gottschalck IB, Jorgensen CS, et al. Immunosuppressive drugs impairs antibody response of the polysaccharide and conjugated pneumococcal vaccines in patients with Crohn's disease. Vaccine. 2015;33:5464-9.
[125] Kharbanda EO, Vazquez-Benitez G, Lipkind HS, Klein NP, Cheetham TC, Naleway AL, et al. Maternal Tdap vaccination: Coverage and acute safety outcomes in the vaccine safety datalink, 2007-2013. Vaccine. 2016;34:968-73.
[126] Kieninger D, Sheldon E, Lin W-Y, Yu C-J, Bayas JM, Gabor JJ, et al. Immunogenicity, reactogenicity and safety of an inactivated quadrivalent influenza vaccine candidate versus inactivated trivalent influenza vaccine: a phase III, randomized trial in adults aged >=18 years. BMC infectious diseases. 2013;13:343.
[127] Kim DS, Shin SH, Lee HJ, Hong YJ, Lee SY, Choi KM, et al. Immunogenicity and safety of 13-valent pneumococcal conjugate vaccine given to korean children receiving routine pediatric vaccines. Pediatric Infectious Disease Journal. 2013;32:266-73.
[128] Kirstein J, Pina M, Pan J, Jordanov E, Dhingra MS. Immunogenicity and safety of a quadrivalent meningococcal tetanus toxoid-conjugate vaccine (MenACYW-TT) in adults 56 years of age and older: a Phase II randomized study. Human Vaccines and Immunotherapeutics. 2020;16:1299-305.
[129] Klein NP, Fireman B, Yih WK, Lewis E, Kulldorff M, Ray P, et al. Measles-mumps-rubella-varicella combination vaccine and the risk of febrile seizures. Pediatrics. 2010;126:e1-8.
[130] Klein NP, Lewis E, Fireman B, Hambidge SJ, Naleway A, Nelson JC, et al. Safety of measles-containing vaccines in 1-year-old children. Pediatrics. 2015;135:e321-9.
[131] Lal H, Cunningham AL, Godeaux O, Chlibek R, Diez-Domingo J, Hwang S-J, et al. Efficacy of an adjuvanted herpes zoster subunit vaccine in older adults. The New England journal of medicine. 2015;372:2087-96.
[132] Langer-Gould A, Qian L, Tartof SY, Brara SM, Jacobsen SJ, Beaber BE, et al. Vaccines and the risk of multiple sclerosis and other central nervous system demyelinating diseases. JAMA neurology. 2014;71:1506-13.
[133] Langley JM, Carmona Martinez A, Chatterjee A, Halperin SA, McNeil S, Reisinger KS, et al. Immunogenicity and safety of an inactivated quadrivalent influenza vaccine candidate: a phase III randomized controlled trial in children. The Journal of infectious diseases. 2013;208:544-53.
[134] Langley JM, Wang L, Aggarwal N, Bueso A, Chandrasekaran V, Cousin L, et al. Immunogenicity and Reactogenicity of an Inactivated Quadrivalent Influenza Vaccine Administered Intramuscularly to Children 6 to 35 Months of Age in 2012-2013: A Randomized, Double-Blind, Controlled, Multicenter, Multicountry, Clinical Trial. Journal of the Pediatric Infectious Diseases Society. 2015;4:242-51.
[135] Layton JB, Butler AM, Li D, Boggess KA, Weber DJ, McGrath LJ, et al. Prenatal Tdap immunization and risk of maternal and newborn adverse events. Vaccine. 2017;35:4072-8.
[136] Layton JB, Butler AM, Panozzo CA, Brookhart MA. Rotavirus vaccination and short-term risk of adverse events in US infants. Paediatr Perinat Epidemiol. 2018;32:448-57.
[137] Lee HJ, Choe YJ, Hong Y-J, Kim K-H, Park SE, Kim Y-K, et al. Immunogenicity and safety of a multicomponent meningococcal serogroup B vaccine in healthy adolescents in Korea--A randomised trial. Vaccine. 2016;34:1180-6.
[138] Lee HJ, Chung M-H, Kim WJ, Hong YJ, Choi KM, Lee J, et al. Immunogenicity and safety of a novel quadrivalent meningococcal conjugate vaccine (MenACWY-CRM) in healthy

873 Korean adolescents and adults. International journal of infectious diseases : IJID : official
874 publication of the International Society for Infectious Diseases. 2014;28:204-10.
875 [139] Leslie DL, Kobre RA, Richmand BJ, Aktan Guloksuz S, Leckman JF. Temporal
876 Association of Certain Neuropsychiatric Disorders Following Vaccination of Children and
877 Adolescents: A Pilot Case-Control Study. Front Psychiatry. 2017;8:3.
878 [140] Li C-Y, Chen L-C, Lin H-Y, Lee M-S, Hung S-K, Lai C-L, et al. Impact of 23-valent
879 pneumococcal polysaccharide vaccination on the frequency of pneumonia-related hospitalization
880 and survival in elderly patients with prostate cancer: A seven-year nationwide matched cohort
881 study. Cancer. 2020.
882 [141] Li R, Stewart B, McNeil MM, Duffy J, Nelson J, Kawai AT, et al. Post licensure
883 surveillance of influenza vaccines in the Vaccine Safety Datalink in the 2013-2014 and 2014-
884 2015 seasons. Pharmacoepidemiology and drug safety. 2016;25:928-34.
885 [142] Li R-c, Huang T, Li Y, Luo D, Tao J, Fu B, et al. Human rotavirus vaccine (RIX4414)
886 efficacy in the first two years of life: a randomized, placebo-controlled trial in China. Human
887 vaccines & immunotherapeutics. 2014;10:11-8.
888 [143] Lombardi F, Belmonti S, Fabbiani M, Morandi M, Rossetti B, Tordini G, et al.
889 Immunogenicity and Safety of the 13-Valent Pneumococcal Conjugate Vaccine versus the 23-
890 Valent Polysaccharide Vaccine in Unvaccinated HIV-Infected Adults: A Pilot, Prospective
891 Controlled Study. PloS one. 2016;11:e0156523.
892 [144] Mallory RM, Nyborg A, Kalyani RN, Yuan Y, Block SL, Dubovsky F. A study to evaluate
893 the immunogenicity and shedding of live attenuated influenza vaccine strains in children 24–<48
894 months of age. Vaccine. 2020;38:1001-8.
895 [145] Mallory RM, Yu J, Kameo S, Tanaka M, Rito K, Itoh Y, et al. The safety and efficacy of
896 quadrivalent live attenuated influenza vaccine in Japanese children aged 2-18 years: Results of
897 two phase 3 studies. Influenza and other respiratory viruses. 2018a;12:438-45.
898 [146] Maréchal C, Lal H, Poder A, Ferguson M, Enweonye I, Heineman TC, et al.
899 Immunogenicity and safety of the adjuvanted recombinant zoster vaccine co-administered with
900 the 23-valent pneumococcal polysaccharide vaccine in adults ≥50 years of age: a randomized
901 trial. Vaccine. 2018;36:4278- 86.
902 [147] Marshall GS, Adams GL, Leonardi ML, Petrecz M, Flores SA, Ngai AL, et al.
903 Immunogenicity, Safety, and Tolerability of a Hexavalent Vaccine in Infants. Pediatrics.
904 2015;136:e323-32.
905 [148] McClure DL, Jacobsen SJ, Klein NP, Naleway AL, Kharbanda EO, Glanz JM, et al.
906 Similar relative risks of seizures following measles containing vaccination in children born
907 preterm compared to full-term without previous seizures or seizure-related disorders. Vaccine.
908 2019;37:76-9.
909 [149] McGeoch LJ, Finn A, Marlow RD. Impact of rotavirus vaccination on intussusception
910 hospital admissions in England. Vaccine. 2020;38:5618-26.
911 [150] Mo Z, Mo Y, Li M, Tao J, Yang X, Kong J, et al. Efficacy and safety of a pentavalent live
912 human-bovine reassortant rotavirus vaccine (RV5) in healthy Chinese infants: A randomized,
913 double-blind, placebo-controlled trial. Vaccine. 2017;35:5897-904.
914 [151] Morgan JL, Baggari SR, McIntire DD, Sheffield JS. Pregnancy outcomes after antepartum
915 tetanus, diphtheria, and acellular pertussis vaccination. Obstetrics and gynecology.
916 2015;125:1433-8.

[152] Munnoch SA, Cashman P, Peel R, Attia J, Hure A, Durrheim DN. Participant-Centered Online Active Surveillance for Adverse Events Following Vaccination in a Large Clinical Trial: Feasibility and Usability Study. Journal of medical Internet research. 2019;21:e14791.
[153] Munoz FM, Bond NH, Maccato M, Pinell P, Hammill HA, Swamy GK, et al. Safety and immunogenicity of tetanus diphtheria and acellular pertussis (Tdap) immunization during pregnancy in mothers and infants: a randomized clinical trial. JAMA. 2014;311:1760-9.
[154] Naleway AL, Mittendorf KF, Irving SA, Henninger ML, Crane B, Smith N, et al. Primary Ovarian Insufficiency and Adolescent Vaccination. Pediatrics. 2018;142.
[155] Nelson JC, Yu O, Dominguez-Islas CP, Cook AJ, Peterson D, Greene SK, et al. Adapting group sequential methods to observational postlicensure vaccine safety surveillance: results of a pentavalent combination DTaP-IPV-Hib vaccine safety study. American Journal of Epidemiology. 2013;177:131-41.
[156] Oberle D, Hoffelner M, Pavel J, Mentzer D, Barth I, Drechsel-Bauerle U, et al. Retrospective multicenter matched case-control study on the risk factors for intussusception in infants less than 1 year of age with a special focus on rotavirus vaccines - the German Intussusception Study. Human vaccines & immunotherapeutics. 2020:1-14.
[157] Ochoa-Gondar O, Vila-Corcoles A, Rodriguez-Blanco T, de Diego-Cabanes C, Hospital-Guardiola I, Jariod-Pamies M, et al. Evaluating the clinical effectiveness of pneumococcal vaccination in preventing myocardial infarction: The CAPAMIS study, three-year follow-up. Vaccine. 2014;32:252-7.
[158] Ofori-Anyinam O, Leroux-Roels G, Drame M, Aerssens A, Maes C, Amanullah A, et al. Immunogenicity and safety of an inactivated quadrivalent influenza vaccine co-administered with a 23-valent pneumococcal polysaccharide vaccine versus separate administration, in adults >=50years of age: Results from a phase III, randomized, non-inferiority trial. Vaccine. 2017;35:6321-8.
[159] Ostergaard L, Lucksinger GH, Absalon J, Beeslaar J, Eiden J, Jansen KU, et al. A phase 3, randomized, active-controlled study to assess the safety and tolerability of meningococcal serogroup B vaccine bivalent rLP2086 in healthy adolescents and young adults. Vaccine. 2016;34:1465-71.
[160] Perez JZV, Aranda JMR, de la O Cavazos M, Osuna MDZ, Davila JP, Elizondo MRB, et al. Randomized clinical trial of the safety and immunogenicity of the Tdap vaccine in pregnant Mexican women. Human Vaccines & Immunotherapeutics. 2017;13:128-35.
[161] Perez-Vilar S, Wernecke M, Arya D, Lo A-C, Lufkin B, Hu M, et al. Surveillance for Guillain-Barre syndrome after influenza vaccination among U.S. Medicare beneficiaries during the 2017-2018 season. Vaccine. 2019;37:3856-65.
[162] Perrett KP, Halperin SA, Nolan T, Martinez Pancorbo C, Tapiero B, Martinon-Torres F, et al. Immunogenicity, transplacental transfer of pertussis antibodies and safety following pertussis immunization during pregnancy: Evidence from a randomized, placebo-controlled trial. Vaccine. 2019.
[163] Petousis-Harris H, Jiang YN, Yu L, Watson D, Walls T, Turner N, et al. A Retrospective Cohort Study of Safety Outcomes in New Zealand Infants Exposed to Tdap Vaccine in Utero. Vaccines. 2019;7:15.
[164] Petrecz M, Ramsey KP, Stek JE, Martin JC, Klopfer SO, Kuter B, et al. Concomitant use of VAQTA® with PedvaxHIB® and Infanrix® in 12 to 17 month old children. Human Vaccines and Immunotherapeutics. 2016;12:503-11.

[165] Pfizer. Study Comparing a 13-valent Pneumococcal Conjugate Vaccine With 23-valent Pneumococcal Polysaccharide Vaccine in Adults. https://ClinicalTrials.gov/show/NCT00427895; 2007.
[166] Puig-Barbera J, Diez-Domingo J, Varea AB, Chavarri GS, Rodrigo JAL, Hoyos SP, et al. Effectiveness of MF59-adjuvanted subunit influenza vaccine in preventing hospitalisations for cardiovascular disease, cerebrovascular disease and pneumonia in the elderly. Vaccine. 2007;25:7313-21.
[167] Richmond PC, Marshall HS, Nissen MD, Jiang Q, Jansen KU, Garces-Sanchez M, et al. Safety, immunogenicity, and tolerability of meningococcal serogroup B bivalent recombinant lipoprotein 2086 vaccine in healthy adolescents: A randomised, single-blind, placebo-controlled, phase 2 trial. The Lancet Infectious Diseases. 2012;12:597-607.
[168] Rivera L, Schwarz TF, Kim KH, Kim YK, Behre U, Cha SH, et al. Immunogenicity and safety of the quadrivalent meningococcal vaccine MenACWY-TT co-administered with a combined diphtheria-tetanus-acellular pertussis vaccine versus their separate administration in adolescents and young adults: A phase III, randomized study. Vaccine. 2018;36:4750-8.
[169] Rodriguez Weber MA, Claeys C, Aranza Doniz C, Feng Y, Innis BL, Jain VK, et al. Immunogenicity and safety of inactivated quadrivalent and trivalent influenza vaccines in children 18-47 months of age. The Pediatric infectious disease journal. 2014;33:1262-9.
[170] Rogers MAM, Basu T, Kim C. Lower Incidence Rate of Type 1 Diabetes after Receipt of the Rotavirus Vaccine in the United States, 2001–2017. Scientific Reports. 2019;9.
[171] Salas A, Pardo-Seco J, Cebey-Lopez M, Martinon-Martinez JM, Gomez-Rial J, Curras-Tuala MJ, et al. Impact of rotavirus vaccination on childhood hospitalizations for seizures: Heterologous or unforeseen direct vaccine effects? Vaccine. 2019;37:3362-8.
[172] Sancovski M, Mesaros N, Feng Y, Ceregido MA, Luyts D, De Barros E. Safety of reduced antigen content diphtheria-tetanus-acellular pertussis vaccine when administered during pregnancy as part of the maternal immunization program in Brazil: a single center, observational, retrospective, cohort study. Human vaccines & immunotherapeutics. 2019;15:2873-81.
[173] Sanofi Pasteur aSC, Sanofi. Safety and Immunogenicity of Fluzone® Quadrivalent, Flublok® Quadrivalent, and Fluzone® High-Dose, Influenza Vaccines, 2018-2019 Formulations. 2018.
[174] Santolaya ME, O'Ryan ML, Valenzuela MT, Prado V, Vergara R, Munoz A, et al. Immunogenicity and tolerability of a multicomponent meningococcal serogroup B (4CMenB) vaccine in healthy adolescents in Chile: a phase 2b/3 randomised, observer-blind, placebo-controlled study. Lancet. 2012;379:617-24.
[175] Schwarz TF, Aggarwal N, Moeckesch B, Schenkenberger I, Claeys C, Douha M, et al. Immunogenicity and Safety of an Adjuvanted Herpes Zoster Subunit Vaccine Coadministered With Seasonal Influenza Vaccine in Adults Aged 50 Years or Older. The Journal of infectious diseases. 2017;216:1352-61.
[176] Schwarz TF, Flamaing J, Rumke HC, Penzes J, Juergens C, Wenz A, et al. A randomized, double-blind trial to evaluate immunogenicity and safety of 13-valent pneumococcal conjugate vaccine given concomitantly with trivalent influenza vaccine in adults aged >=65 years. Vaccine. 2011;29:5195-202.
[177] Senders S, Bhuyan P, Jiang Q, Absalon J, Eiden JJ, Jones TR, et al. Immunogenicity, Tolerability and Safety in Adolescents of Bivalent rLP2086, a Meningococcal Serogroup B Vaccine, Coadministered with Quadrivalent Human Papilloma Virus Vaccine. The Pediatric infectious disease journal. 2016;35:548-54.

[178] Seo YB, Choi WS, Lee J, Song JY, Cheong HJ, Kim WJ. Comparison of immunogenicity and safety of an influenza vaccine administered concomitantly with a 13-valent pneumococcal conjugate vaccine or 23-valent polysaccharide pneumococcal vaccine in the elderly. Clinical and Experimental Vaccine Research. 2017;6:38-44.
[179] Shakib JH, Korgenski K, Sheng X, Varner MW, Pavia AT, Byington CL. Tetanus, diphtheria, acellular pertussis vaccine during pregnancy: pregnancy and infant health outcomes. The Journal of pediatrics. 2013;163:1422-4.
[180] Shimada K, Morinaga H, Kiyanagi T, Miyazaki T, Nishitani-Yokoyama M, Okai I, et al. Safety and Efficacy of Simultaneous Inoculations of Pneumococcal and Influenza Vaccines in Patients with Coronary Artery Disease. J Atheroscler Thromb. 2020.
[181] Shiramoto M, Hanada R, Juergens C, Shoji Y, Yoshida M, Ballan B, et al. Immunogenicity and safety of the 13-valent pneumococcal conjugate vaccine compared to the 23-valent pneumococcal polysaccharide vaccine in elderly Japanese adults. Human vaccines & immunotherapeutics. 2015;11:2198-206.
[182] Siriwardena AN, Asghar Z, Coupland CC. Influenza and pneumococcal vaccination and risk of stroke or transient ischaemic attack-matched case control study. Vaccine. 2014;32:1354-61.
[183] Song JY, Cheong HJ, Hyun HJ, Seo YB, Lee J, Wie S-H, et al. Immunogenicity and safety of a 13-valent pneumococcal conjugate vaccine and an MF59-adjuvanted influenza vaccine after concomitant vaccination in 60-year-old adults. Vaccine. 2017;35:313-20.
[184] Song JY, Cheong HJ, Noh JY, Choi MJ, Yoon JG, Lee SN, et al. Immunogenicity and safety of a tetanus-diphtheria vaccine and a 13-valent pneumococcal conjugate vaccine after concomitant vaccination in >= 50-year-old adults. BMC infectious diseases. 2018;18:628.
[185] Song JY, Cheong HJ, Tsai TF, Chang H-A, Choi MJ, Jeon JH, et al. Immunogenicity and safety of concomitant MF59-adjuvanted influenza vaccine and 23-valent pneumococcal polysaccharide vaccine administration in older adults. Vaccine. 2015a;33:4647-52.
[186] Song JY, Cheong HJ, Tsai TF, Chang HA, Choi MJ, Jeon JH, et al. Immunogenicity and safety of concomitant MF59-adjuvanted influenza vaccine and 23-valent pneumococcal polysaccharide vaccine administration in older adults. Vaccine. 2015b;33:4647-52.
[187] Stockwell MS, Broder K, LaRussa P, Lewis P, Fernandez N, Sharma D, et al. Risk of fever after pediatric trivalent inactivated influenza vaccine and 13-valent pneumococcal conjugate vaccine. JAMA pediatrics. 2014;168:211-9.
[188] Stockwell MS, Broder KR, Lewis P, Jakob K, Iqbal S, Fernandez N, et al. Assessing Fever Frequency After Pediatric Live Attenuated Versus Inactivated Influenza Vaccination. Journal of the Pediatric Infectious Diseases Society. 2017;6:e7-e14.
[189] Stowe J, Andrews N, Ladhani S, Miller E. The risk of intussusception following monovalent rotavirus vaccination in England: A self-controlled case-series evaluation. Vaccine. 2016;34:3684-9.
[190] Strezova A, Lal H, Enweonye I, Campora L, Beukelaers P, Segall N, et al. The adjuvanted recombinant zoster vaccine co-administered with a tetanus, diphtheria and pertussis vaccine in adults aged >=50 years: A randomized trial. Vaccine. 2019;37:5877-85.
[191] Svensson T, Kattstrom M, Hammarlund Y, Roth D, Andersson PO, Svensson M, et al. Pneumococcal conjugate vaccine triggers a better immune response than pneumococcal polysaccharide vaccine in patients with chronic lymphocytic leukemia A randomized study by the Swedish CLL group. Vaccine. 2018;36:3701-7.

[192] Tapiero B, Halperin SA, Dionne M, Meekison W, Diaz-Mitoma F, Zickler P, et al. Safety and immunogenicity of a hexavalent vaccine administered at 2, 4 and 6 months of age with or without a heptavalent pneumococcal conjugate vaccine: a randomized, open-label study. Pediatric Infectious Disease Journal. 2013;32:54-61.
[193] Tate JE, Mwenda JM, Armah G, Jani B, Omore R, Ademe A, et al. Evaluation of Intussusception after Monovalent Rotavirus Vaccination in Africa. N Engl J Med. 2018;378:1521-8.
[194] Tate JE, Yen C, Steiner CA, Cortese MM, Parashar UD. Intussusception Rates Before and After the Introduction of Rotavirus Vaccine. Pediatrics. 2016;138.
[195] Thompson AR, Klein NP, Downey HJ, Patterson S, Sundaraiyer V, Watson W, et al. Coadministration of 13-valent pneumococcal conjugate and quadrivalent inactivated influenza vaccines in adults previously immunized with polysaccharide pneumococcal vaccine 23: a randomized clinical trial. Human vaccines & immunotherapeutics. 2019;15:444-51.
[196] Timmermann CAG, Osuna CE, Steuerwald U, Weihe P, Poulsen LK, Grandjean P. Asthma and allergy in children with and without prior measles, mumps, and rubella vaccination. Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology. 2015;26:742-9.
[197] Tinoco JC, Pavia-Ruz N, Cruz-Valdez A, Aranza Doniz C, Chandrasekaran V, Dewe W, et al. Immunogenicity, reactogenicity, and safety of inactivated quadrivalent influenza vaccine candidate versus inactivated trivalent influenza vaccine in healthy adults aged >=18 years: a phase III, randomized trial. Vaccine. 2014;32:1480-7.
[198] Togashi T, Okada K, Yamaji M, Thompson A, Gurtman A, Cutler M, et al. Immunogenicity and Safety of a 13-Valent Pneumococcal Conjugate Vaccine Given With DTaP Vaccine in Healthy Infants in Japan. The Pediatric infectious disease journal. 2015;34:1096-104.
[199] Treanor JT, Albano FR, Sawlwin DC, Graves Jones A, Airey J, Formica N, et al. Immunogenicity and safety of a quadrivalent inactivated influenza vaccine compared with two trivalent inactivated influenza vaccines containing alternate B strains in adults: A phase 3, randomized noninferiority study. Vaccine. 2017;35:1856-64.
[200] Tregnaghi M, Lopez P, Stamboulian D, Grana G, Odrljin T, Bedell L, et al. Immunogenicity and safety of a quadrivalent meningococcal polysaccharide CRM conjugate vaccine in infants and toddlers. International journal of infectious diseases : IJID : official publication of the International Society for Infectious Diseases. 2014;26:22-30.
[201] Tseng H-F, Sy LS, Ackerson BK, Hechter RC, Tartof SY, Haag M, et al. Safety of Quadrivalent Meningococcal Conjugate Vaccine in 11- to 21-Year-Olds. Pediatrics. 2017;139.
[202] Tseng HF, Sy LS, Liu ILA, Qian L, Marcy SM, Weintraub E, et al. Postlicensure surveillance for pre-specified adverse events following the 13-valent pneumococcal conjugate vaccine in children. Vaccine. 2013;31:2578-83.
[203] Tseng HF, Sy LS, Qian L, Liu I-LA, Mercado C, Lewin B, et al. Pneumococcal Conjugate Vaccine Safety in Elderly Adults. Open forum infectious diseases. 2018;5:ofy100.
[204] Uhlig U, Kostev K, Schuster V, Koletzko S, Uhlig HH. Impact of rotavirus vaccination in Germany: rotavirus surveillance, hospitalization, side effects and comparison of vaccines. The Pediatric infectious disease journal. 2014;33:e299-304.
[205] Uno Y, Uchiyama T, Kurosawa M, Aleksic B, Ozaki N. Early exposure to the combined measles-mumps-rubella vaccine and thimerosal-containing vaccines and risk of autism spectrum disorder. Vaccine. 2015;33:2511-6.

[206] Vaarala O, Jokinen J, Lahdenkari M, Leino T. Rotavirus Vaccination and the Risk of Celiac Disease or Type 1 Diabetes in Finnish Children at Early Life. The Pediatric infectious disease journal. 2017;36:674-5.
[207] Van Damme P, Meijer CJLM, Kieninger D, Schuyleman A, Thomas S, Luxembourg A, et al. A phase III clinical study to compare the immunogenicity and safety of the 9-valent and quadrivalent HPV vaccines in men. Vaccine. 2016;34:4205-12.
[208] Vandecasteele SJ, De Bacquer D, Caluwe R, Ombelet S, Van Vlem B. Immunogenicity and safety of the 13-valent Pneumococcal Conjugate vaccine in 23-valent pneumococcal polysaccharide vaccine-naive and pre-immunized patients under treatment with chronic haemodialysis: a longitudinal quasi-experimental phase IV study. Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases. 2018;24:65-71.
[209] Vesikari T, Brodszki N, van Damme P, Diez-Domingo J, Icardi G, Petersen LK, et al. A Randomized, Double-Blind, Phase III Study of the Immunogenicity and Safety of a 9-Valent Human Papillomavirus L1 Virus-Like Particle Vaccine (V503) Versus Gardasil in 9-15-Year-Old Girls. The Pediatric infectious disease journal. 2015;34:992-8.
[210] Vesikari T, Karvonen A, Prymula R, Schuster V, Tejedor J, Cohen R, et al. Efficacy of human rotavirus vaccine against rotavirus gastroenteritis during the first 2 years of life in European infants: randomised, double-blind controlled study. Lancet. 2007;370:1757-63.
[211] Vesikari T, Wysocki J, Beeslaar J, Eiden J, Jiang Q, Jansen KU, et al. Immunogenicity, Safety, and Tolerability of Bivalent rLP2086 Meningococcal Group B Vaccine Administered Concomitantly With Diphtheria, Tetanus, and Acellular Pertussis and Inactivated Poliomyelitis Vaccines to Healthy Adolescents. Journal of the Pediatric Infectious Diseases Society. 2016;5:180-7.
[212] Vila-Corcoles A, Ochoa-Gondar O, de Diego C, Satue E, Aragón M, Vila-Rovira A, et al. Evaluating clinical effectiveness of 13-valent pneumococcal conjugate vaccination against pneumonia among middle-aged and older adults in Catalonia: results from the EPIVAC cohort study. BMC Infectious Diseases. 2018;18:196.
[213] Villa M, Black S, Groth N, Rothman KJ, Apolone G, Weiss NS, et al. Safety of MF59-adjuvanted influenza vaccination in the elderly: results of a comparative study of MF59-adjuvanted vaccine versus nonadjuvanted influenza vaccine in northern Italy. American journal of epidemiology. 2013;178:1139-45.
[214] Walter EB, Klein NP, Wodi AP, Rountree W, Todd CA, Wiesner A, et al. Fever after influenza, diphtheria-tetanus-acellular pertussis, and pneumococcal vaccinations. Pediatrics. 2020;145.
[215] Wang L, Chandrasekaran V, Domachowske JB, Li P, Innis BL, Jain VK. Immunogenicity and Safety of an Inactivated Quadrivalent Influenza Vaccine in US Children 6-35 Months of Age During 2013-2014: Results From A Phase II Randomized Trial. Journal of the Pediatric Infectious Diseases Society. 2016;5:170-9.
[216] Wang SV, Abdurrob A, Spoendlin J, Lewis E, Newcomer SR, Fireman B, et al. Methods for addressing "innocent bystanders" when evaluating safety of concomitant vaccines. Pharmacoepidemiology and drug safety. 2018;27:405-12.
[217] Yih WK, Lieu TA, Kulldorff M, Martin D, McMahill-Walraven CN, Platt R, et al. Intussusception risk after rotavirus vaccination in U.S. infants. The New England journal of medicine. 2014;370:503-12.

[218] Yung CF, Chan SP, Soh S, Tan A, Thoon KC. Intussusception and Monovalent Rotavirus Vaccination in Singapore: Self-Controlled Case Series and Risk-Benefit Study. J Pediatr. 2015;167:163-8.e1.
[219] Yung CF, Ma X, Cheung YB, Oh BK, Soh S, Thoon KC. Kawasaki Disease following administration of 13-valent pneumococcal conjugate vaccine in young children. Scientific reports. 2019;9:14705.
[220] Zahid M, Singla I, Good CB, Stone RA, Kim S, Fine MJ, et al. Associations between Pneumococcal Vaccination and Adverse Outcomes in Patients with Suspected Acute Coronary Syndrome. Adv Infect Dis. 2012:122-34.
[221] Erratum: v72P10 Meningococcal B Adolescent Vaccine Study group. Immunogenicity and tolerability of a multicomponent meningococcal serogroup B (4CMenB) vaccine in healthy adolescents in Chile: a phase 2b/3 randomised, observer-blind, placebo-controlled study (The Lancet (2012) 379 (617-624)). Lancet. 2015;385:1728.
[222] AstraZeneca. A Phase3 Study to Evaluate the Efficacy and Safety of MEDI3250 in Healthy Japanese Children Age 7 Years Through 18 Years. 2015.
[223] AZ Sint-Jan AV, Pfizer. 13 Valent Pneumococcal Conjugate Vaccine in Chronic Dialysis Patients. 2013.
[224] Baker MA, Lieu TA, Li LL, Hua W, Qiang YD, Kawai AT, et al. A Vaccine Study Design Selection Framework for the Postlicensure Rapid Immunization Safety Monitoring Program. American Journal of Epidemiology. 2015;181:608-18.
[225] Baxter R, Lewis E, Fireman B, DeStefano F, Gee J, Klein NP. Case-centered Analysis of Optic Neuritis After Vaccines. Clin Infect Dis. 2016;63:79-81.
[226] Clinical Trials Registry-India. CTRI/2012/07/002820: A study to evaluate safety of Rotavirus vaccine in Healthy Adult Volunteers followed by Safety, Tolerability and Immunogenicity evaluation in healthy infants. July 31, 2013.
[227] Colindres R, Wascotte V, Brecx A, Clarke C, Hervé C, Kim JH, et al. Post hoc analysis of reactogenicity trends between dose 1 and dose 2 of the adjuvanted recombinant zoster vaccine in two parallel randomized trials. Human Vaccines and Immunotherapeutics. 2020.
[228] Columbia University, Centers for Disease Control Prevention. FeverText: Assessing Fever Rates After Vaccination During the 2011-12 Influenza Season Using Text Messaging. 2011.
[229] Columbia University, Centers for Disease Control Prevention. Assessing Fever Rates in Children Ages 24 to 59 Months After Live Attenuated Influenza Vaccine (LAIV) or Inactivated Influenza Vaccines (IIV) Using Text Messaging for U.S. Influenza Vaccines in 2012-2013 & 2013-2014. 2013.
[230] Cowling BJ, Thompson MG, Ng TWY, Fang VJ, Perera RAPM, Leung NHL, et al. Comparative Reactogenicity of Enhanced Influenza Vaccines in Older Adults. The Journal of infectious diseases. 2020;222:1383-91.
[231] DeSilva M, Vazquez-Benitez G, Nordin JD, Lipkind HS, Romitti PA, DeStefano F, et al. Tdap Vaccination During Pregnancy and Microcephaly and Other Structural Birth Defects in Offspring. JAMA. 2016;316:1823-5.
[232] Dezure A, Marechal C, Lal H, Poder A, Ferguson M, Enweonye I, et al. Immunogenicity and safety of an adjuvanted herpes zoster subunit vaccine candidate when coadministered with a 23-valent pneumococcal polysaccharide vaccine in adults 50 years of age or older: a phase iii, randomized clinical trial. Journal of the american pharmacists association. 2018;58:e72- e3.
[233] Dynavax Technologies Corporation. Safety and Immunogenicity Study of the Hepatitis B Virus (HBV) Vaccine, HEPLISAV Compared to Engerix-B Vaccine. 2010.

[234] Euctr BE. Evaluation of the immunogenicity and safety of GlaxoSmithKline (GSK) Biologicals' Quadrivalent Influenza Vaccine Influsplit™ Tetra (Fluarix™ Tetra) (GSK2321138A) when co-administered with Pneumovax™ 23 in adults 50 years of age and older. http://wwwwhoint/trialsearch/Trial2aspx?TrialID=EUCTR2014-001118-24-BE. 2014.
[235] Euctr CZ. Immunogenicity and safety study of Boostrix in pregnant women. http://wwwwhoint/trialsearch/Trial2aspx?TrialID=EUCTR2014-001119-38-CZ. 2015.
[236] Everett D. Hepatitis B Vaccine (Recombinant), Adjuvanted (Heplisav-B): Review of Safety. Vaccines and Related Biological Products Advisory Committee Meeting: U.S. Food & Drug Administration; July 28, 2017.
[237] Garland SM, Pitisuttithum P, Ngan HYS, Cho CH, Lee CY, Chen CA, et al. Efficacy, Immunogenicity, and Safety of a 9-Valent Human Papillomavirus Vaccine: Subgroup Analysis of Participants From Asian Countries. The Journal of infectious diseases. 2018;218:95-108.
[238] Giuliano AR, Joura EA, Garland SM, Huh WK, Iversen O-E, Kjaer SK, et al. Nine-valent HPV vaccine efficacy against related diseases and definitive therapy: comparison with historic placebo population. Gynecologic oncology. 2019;154:110-7.
[239] GlaxoSmithKline. To Test 2 Doses of GSK Biologicals' Oral Live Attenuated Human Rotavirus (HRV) Vaccine in Healthy Infants in Co-administration With Specific Childhood Vaccines. https://ClinicalTrials.gov/show/NCT00140686; 2004.
[240] GlaxoSmithKline. Immunogenicity and Safety Study of a GSK Influenza Vaccine Candidate in Adults. 2008.
[241] GlaxoSmithKline. Immunogenicity and Safety Study of GSK Biologicals' Herpes Zoster Vaccine With Various Formulations in Adults >= 50 Years. 2009.
[242] GlaxoSmithKline. Immunogenicity and Safety Study of a GlaxoSmithKline Biologicals' Candidate Influenza Vaccine in Healthy Children. https://ClinicalTrials.gov/show/NCT00985790; 2009.
[243] GlaxoSmithKline. Safety and Immunogenicity Study of GSK Biologicals' Seasonal Influenza Candidate Vaccine (GSK2321138A). 2010.
[244] GlaxoSmithKline. A Study to Evaluate the Safety and Immunogenicity of GSK Biologicals' Seasonal Influenza Vaccine in Children. 2010.
[245] GlaxoSmithKline. A Study to Evaluate the Safety and Immunogenicity of GSK Biologicals' Seasonal Influenza Vaccine in Adults. 2010.
[246] GlaxoSmithKline. Study to Assess the Efficacy, Immunogenicity and Safety of Liquid Human Rotavirus Vaccine, in Healthy Chinese Infants. 2010.
[247] GlaxoSmithKline. Study to Evaluate Efficacy, Safety and Immunogenicity of GSK Biologicals' Herpes Zoster (HZ) Vaccine GSK1437173A in Adults Aged 50 Years and Older. 2010.
[248] GlaxoSmithKline. Immunogenicity and Safety Study of GSK Biologicals' Influenza Vaccine When Administered in Children. https://ClinicalTrials.gov/show/NCT01196988; 2010.
[249] GlaxoSmithKline. Study to Evaluate Immunogenicity and Safety of GlaxoSmithKline (GSK) Biologicals' Quadrivalent Influenza Vaccine GSK2282512A When Administered to Children 6 to 35 Months of Age. 2012.
[250] GlaxoSmithKline. Immunogenicity and Safety Study of GlaxoSmithKline (GSK) Biologicals' Meningococcal Conjugate Vaccine (GSK134612) When Co-administered With Boostrix® in Subjects Between 11 and 25 Years of Age. 2014.

[251] GlaxoSmithKline. Study to Evaluate Immunogenicity and Safety of GlaxoSmithKline (GSK) Biologicals' Quadrivalent Influenza Vaccine GSK2282512A When Administered to Children From 6 to 35 Months of Age. 2014.
[252] GlaxoSmithKline. Study to Evaluate the Immunogenicity and Safety of GlaxoSmithKline (GSK) Biologicals' Herpes Zoster Vaccine GSK1437173A When Co-administered With GSK Biologicals' Seasonal Influenza Vaccine GSK2321138A in Adults Aged 50 Years and Older. 2014.
[253] GlaxoSmithKline. Study to Assess the Immunogenicity and Safety of GlaxoSmithKline (GSK) Biologicals' Herpes Zoster Subunit (HZ/su) Vaccine (GSK1437173A) When Co-administered With GSK Biologicals' Diphtheria, Tetanus and Pertussis Vaccine (Boostrix®) in Adults Aged 50 Years and Older. 2015.
[254] GlaxoSmithKline. Evaluation of the Immunogenicity and Safety of GlaxoSmithKline (GSK) Biologicals' Quadrivalent Influenza Vaccine Influsplit™ Tetra (Fluarix™ Tetra) (GSK2321138A) When Co-administered With Pneumovax™ 23 in Adults 50 Years of Age and Older. 2015.
[255] GlaxoSmithKline. Study to Evaluate Immunogenicity and Safety Study of GSK Biologicals' Herpes Zoster (HZ) Vaccine GSK1437173A When Co-administered With Pneumovax 23™ in Adults Aged 50 Years and Older. 2015.
[256] GlaxoSmithKline. A Post-marketing, Observational, Retrospective Study to Assess the Safety of RefortrixTM (Tdap) When Administered During Pregnancy in a Maternal Immunization Program in Brazil. 2017.
[257] GlaxoSmithKline. Immunogenicity and Safety Study of GlaxoSmithKline (GSK) Biologicals' Boostrix™ Vaccine in Pregnant Women. 2017.
[258] GlaxoSmithKline. Study to Evaluate Efficacy, Safety and Immunogenicity of GlaxoSmithKline (GSK) Biologicals' Herpes Zoster (HZ) Vaccine GSK1437173A in Adults Aged 70 Years and Older. 2017.
[259] GlaxoSmithKline. Immunogenicity and Safety Study of Infanrix Hexa in Healthy Infants Born to Mothers Vaccinated With Boostrix™ During Pregnancy or Immediately Post-delivery. 2018.
[260] GlaxoSmithKline. Evaluation of Immunogenicity and Safety of a Booster Dose of Infanrix Hexa™ in Healthy Infants Born to Mothers Vaccinated With Boostrix™ During Pregnancy or Immediately Post-delivery. 2019.
[261] Guevara A, Cabello R, Woelber L, Moreira ED, Jr., Joura E, Reich O, et al. Antibody persistence and evidence of immune memory at 5years following administration of the 9-valent HPV vaccine. Vaccine. 2017;35:5050-7.
[262] Guevara AM, Suarez E, Victoria A, Ngan HYS, Hirschberg AL, Fedrizzi E, et al. Maternal transfer of anti HPV 6 and 11 antibodies upon immunization with the 9-valent HPV vaccine. Human Vaccines and Immunotherapeutics. 2019;15:141-5.
[263] Hambidge SJ, Newcomer SR, Narwaney KJ, Glanz JM, Daley MF, Xu S, et al. Timely versus delayed early childhood vaccination and seizures. Pediatrics. 2014;133:e1492-9.
[264] Hedrick J, Christensen S, Chang LJ, Pan J, Jordanov E, Dhinghra M. Study of the immunogenicity and safety of an investigational quadrivalent meningococcal conjugate vaccine (MENACYW-TT) when co-administered with other vaccines in healthy adolescents. Open forum infectious diseases. 2018;5:S570- .

[265] Hedrick J, Simon MW, Christensen S, Anez G, Pan J, Jordanov E, et al. Immunogenicity and safety of a booster dose of a quadrivalent meningococcal conjugate vaccine (menacyw-TT) in adolescents and adults. Open forum infectious diseases. 2019;6:S959- .
[266] Helsinki University Central Hospital, Tampere University Hospital. Immunogenicity of 13-valent Pneumococcal Conjugate Vaccine Compared to the Pneumococcal Polysaccharide Vaccine in Adult Kidney and Liver Transplant Patients. https://ClinicalTrials.gov/show/NCT01781871; 2013.
[267] Hospital Universitario Dr. Jose E. Gonzalez. Immunogenicity and Safety of an Acellular DPT Vaccine During Pregnancy. 2011.
[268] Hyer R, McGuire DK, Xing B, Jackson S, Janssen R. Safety of a two-dose investigational hepatitis B vaccine, HBsAg-1018, using a toll-like receptor 9 agonist adjuvant in adults. Vaccine. 2018;36:2604-11.
[269] Hyer RN, Janssen RS. Immunogenicity and safety of a 2-dose hepatitis B vaccine, HBsAg/CpG 1018, in persons with diabetes mellitus aged 60-70years. Vaccine. 2019;37:5854-61.
[270] Jeanfreau R, Esteves-Jaramillo A, Neveu D, Jordanov E, Dhingra MS. Immunogenicity and safety of a quadrivalent meningococcal conjugate vaccine (MenACYW-TT) administered in individuals 56 years of age and older. Open forum infectious diseases. 2018;5.
[271] Joura EA, Giuliano AR, Iversen O-E, Bouchard C, Mao C, Mehlsen J, et al. A 9-valent HPV vaccine against infection and intraepithelial neoplasia in women. The New England journal of medicine. 2015;372:711-23.
[272] Kantso B, Halkjaer S, Thomsen O, Belard E, Gottschalck IB, Jorgensen CS, et al. Specific antibody response to two pneumococcal vaccines in crohn's disease patients treated with immunosuppressive drugs alone or in combination with biological therapy. Gastroenterology. 2015;148:S176- S7.
[273] Karolinska University Hospital. Pneumococcal Vaccine in Untreated CLL Patients. 2013.
[274] Kharbanda EO, Vazquez-Benitez G, Lipkind HS, Klein NP, Cheetham TC, Naleway A, et al. Evaluation of the association of maternal pertussis vaccination with obstetric events and birth outcomes. JAMA. 2014;312:1897-904.
[275] Korea University Guro Hospital. Influenza Vaccine and Pneumococcal Vaccine. https://ClinicalTrials.gov/show/NCT02582047; 2012.
[276] Korea University Guro Hospital. Immunogenicity and Safety of a Tetanus-diphtheria Vaccine and a 13-valent Pneumococcal Conjugate Vaccine. 2013.
[277] Korea University Guro Hospital, Novartis. MF59-adjuvanted Influenza Vaccine and 23-valent Pneumococcal Polysaccharide Vaccine. 2013.
[278] Korea University Guro Hospital, Pfizer. Immunogenicity and Safety of PCV13 and Fluad in Adults Aged ≥60 Years. 2014.
[279] Laval University. Immunogenicity and Safety of Gardasil-9 and Cervarix. 2015.
[280] Li R-C, Huang T, Li Y, Wang L-H, Tao J, Fu B, et al. Immunogenicity and reactogenicity of the human rotavirus vaccine, RIX4414 oral suspension, when co-administered with routine childhood vaccines in Chinese infants. Human vaccines & immunotherapeutics. 2016;12:785-93.
[281] Li R-C, Li Y-P, Mo Z-J, Luo D, Huang T, Kong J-L, et al. Reactogenicity and safety of a liquid human rotavirus vaccine (RIX4414) in healthy adults, children and infants in China: randomized, double-blind, placebo-controlled Phase I studies. Human vaccines & immunotherapeutics. 2013;9:1638-42.

[282] Lopez-Fauqued M, Campora L, Delannois F, El Idrissi M, Oostvogels L, De Looze FJ, et al. Safety profile of the adjuvanted recombinant zoster vaccine: Pooled analysis of two large randomised phase 3 trials. Vaccine. 2019;37:2482-93.
[283] Luxembourg A, Brown D, Bouchard C, Giuliano AR, Iversen OE, Joura EA, et al. Phase II studies to select the formulation of a multivalent HPV L1 virus-like particle (VLP) vaccine. Human Vaccines and Immunotherapeutics. 2015;11:1313-22.
[284] Mayrand MH, Bautista O, Moeller E, Ritter M, Luxembourg A. End of study efficacy, immunogenicity and safety of a novel 9-valent HPV l1 virus-like particle vaccine in 16-26 year old women. International journal of gynecology and obstetrics (varpagings). 2015;131:E270.
[285] McElhaney JE, Lal H, Cunningham AL, Levin MJ, Chlibek R, Diez-Domingo J, et al. Efficacy, immunogenicity and safety of an investigational subunit adjuvanted herpes zoster vaccine in adults aged 60 years and older: results from the zoe-50 and zoe-70 efficacy studies. Open forum infectious diseases. 2016;3.
[286] MedImmune LLC. A Study to Evaluate the Immunogenicity of Quadrivalent LAIV in Adults 18 to 49 Years of Age. https://ClinicalTrials.gov/show/NCT00860067; 2009.
[287] MedImmune LLC. A Study to Evaluate the Immunogenicity of Quadrivalent Live Attenuated Influenza Vaccine (LAIV) in Children. https://ClinicalTrials.gov/show/NCT01091246; 2010.
[288] MedImmune LLC. Postmarketing Safety Study of Q/LAIV in Subjects 2 Through 49 Years of Age. https://ClinicalTrials.gov/show/NCT01985997; 2014.
[289] MedImmune LLC, AstraZeneca. Evaluate the Shedding and Immunogencity of Different Formulations of FluMist in Children 24 to <48 Months of Age. 2017.
[290] Merck Sharp, Dohme Corp. Efficacy, Safety, and Immunogenicity of V260 in Healthy Chinese Infants (V260-024). 2015.
[291] Merck Sharp, Dohme Corp. Concomitant Use of Hepatitis A Vaccine, Inactivated With Haemophilus b Conjugate Vaccine (Meningococcal Protein Conjugate) and Diphtheria and Tetanus Toxoids and Acellular Pertussis Vaccine Adsorbed Given to Healthy Children 15 Months of Age (V251-068). 2006.
[292] Merck Sharp, Dohme Corp. Broad Spectrum HPV (Human Papillomavirus) Vaccine Study in 16-to 26-Year-Old Women (V503-001). 2007.
[293] Merck Sharp, Dohme Corp. Immunogenicity and Safety Study of Proquad® and Infanrix® Hexa When Administered Concomitantly (V221-035). 2007.
[294] Merck Sharp, Dohme Corp. Phase III Placebo-controlled Study of V260 (RotaTeq™) in Japanese Healthy Infants (V260-029)(COMPLETED). https://ClinicalTrials.gov/show/NCT00718237; 2008.
[295] Merck Sharp, Dohme Corp. A Study of V503, a 9-valent Human Papillomavirus (9vHPV) Vaccine in Females 12-26 Years of Age Who Have Previously Received GARDASIL™ (V503-006). 2010.
[296] Merck Sharp, Dohme Corp. Immunogenicity and Tolerability of V503 Versus GARDASIL (V503-009). 2011.
[297] Merck Sharp, Dohme Corp. A Study to Compare Immune Response of V503 to Gardasil in 16- to 26-year-old Men (V503-020). 2014.
[298] Merck Sharp, Dohme Corp., MCM Vaccines B.V. A Study of V419 Given Concomitantly With Prevnar 13™ and RotaTeq™ (V419-006). 2011.

[299] Merck Sharp, Dohme Corp., MCM Vaccines B.V. Safety, Tolerability, and Immunogenicity of V419 Given Concomitantly With Prevnar 13™ and RotaTeq™ (V419-005). 2011.
[300] Munoz FM, Bond NH, Maccato M. Safety and Immunogenicity of Tetanus Diphtheria and Acellular Pertussis (Tdap) Immunization During Pregnancy in Mothers and Infants: A Randomized Clinical Trial (vol 311, pg 1760, 2014). Jama-Journal of the American Medical Association. 2017;317:442-.
[301] National Institute of Allergy Infectious Diseases. Pertussis Vaccine in Healthy Pregnant Women. 2009.
[302] Novartis, Novartis Vaccines. Study to Evaluate the Safety of Novartis MenACWY Conjugate Vaccine When Administered With Routine Infant Vaccinations to Healthy Infants. 2008.
[303] Novartis, Novartis Vaccines. A Phase 3b, Randomized, Open-Label Study to Evaluate the Safety and Immunogenicity of Select Travel Vaccines When Administered Concomitantly With MenACWY in Adults. 2011.
[304] Novartis, Novartis Vaccines. A Phase 4, Placebo-Controlled, Randomized Study to Evaluate the Immunogenicity and Safety of HPV and Tdap When Administered With MenACWY in Adolescents. 2011.
[305] Novartis Vaccines, GlaxoSmithKline, Novartis. Study to Evaluate the Safety and Immunogenicity of Combined Hepatitis A/B Vaccine With MenACWY-CRM Conjugate Vaccine. 2011.
[306] Novartis Vaccines, Novartis. Safety and Immunogenicity of Meningococcal ACWY Conjugate Versus Polysaccharide Vaccine in Children 2 to 10 Years of Age. 2006.
[307] Novartis Vaccines, Novartis. A Study to Evaluate Safety and Immune Response of Novartis Meningococcal ACWY Vaccine In Infants. 2007.
[308] Novartis Vaccines, Novartis. Safety, Tolerability and Immunogenicity of Novartis Meningococcal B Recombinant Vaccine Administered to Healthy Adolescents According to Different Vaccination Schedules. 2008.
[309] Novartis Vaccines, Novartis. A Multi-center, Observer-blind, Placebo-controlled, Randomized Study to Evaluate the Immunogenicity and Safety of MenACWY in Adolescents and Adults in Korea. 2010.
[310] Novartis Vaccines, Novartis. Safety and Immunogenicity of MF59C.1 Adjuvanted Trivalent Subunit Influenza Vaccine in Elderly Subjects. 2010.
[311] Novartis Vaccines, Novartis. Safety and Immunogenicity of Three Influenza Vaccines Adults Ages 18 and Older. 2014.
[312] Novartis Vaccines, Novartis. Safety and Immunogenicity of Three Influenza Vaccines in Children Aged 4 Years Old to Less Than 18 Years Old. 2014.
[313] Novartis Vaccines, Novartis. Safety and Immunogenicity Study of Two Doses of Novartis Meningococcal Serogroup B Recombinant Vaccine in Adolescents Aged 11-17 Years. 2014.
[314] Novartis Vaccines and Diagnostics S.r.l. A phase III, double-blind, randomized, placebo-controlled, multi-country and multi-center study to assess the efficacy and safety of two doses of GSK Biologicals' oral live attenuated human rotavirus (HRV) vaccine in healthy infants.\. 2014.
[315] Oostvogels L, Heineman TC, Johnson RW, Levin MJ, McElhaney JE, Van den Steen P, et al. Medical conditions at enrollment do not impact efficacy and safety of the adjuvanted recombinant zoster vaccine: a pooled post-hoc analysis of two parallel randomized trials. Human vaccines & immunotherapeutics. 2019;15:2865-72.

[316] Perrett K, Garcia IC, Halperin S, Nolan T, Virta M, Stranak Z, et al. Pertussis vaccination during pregnancy: a multicountry randomised placebo-controlled trial assessing immunogenicity and safety of dTpa in mothers and their infants. Bjog. 2019;126:173- 4.
[317] Perrett KP, Halperin SA, Nolan T, Martinez AC, Martinon-Torres F, Garcia-Sicilia J, et al. Impact of tetanus-diphtheria-acellular pertussis immunization during pregnancy on subsequent infant immunization seroresponses: follow-up from a large randomized placebo-controlled trial. Vaccine. 2020;38:2105-14.
[318] Petersen LK, Restrepo J, Moreira ED, Iversen OE, Pitisuttithum P, Van Damme P, et al. Impact of baseline covariates on the immunogenicity of the 9-valent HPV vaccine – A combined analysis of five phase III clinical trials. Papillomavirus Research. 2017;3:105-15.
[319] Peterson J, Hedrick J, Pan J, Neveu D, Jordanov E, Dhingra MS. Immunogenicity and safety of a quadrivalent meningococcal conjugate vaccine (menacyw-TT) administered in adults 18-55 years of age. Open forum infectious diseases. 2019;6:S957- .
[320] Peterson J, Hedrick J, Pan J, Neveu D, Jordanov E, Dhingra MS. Immunogenicity and safety of a quadrivalent meningococcal conjugate vaccine (menacyw-TT) administered in adolescents 10-17 years of age. Open forum infectious diseases. 2019;6:S958- .
[321] Pfizer. Study Evaluating the Impact of a 13-valent Pneumococcal Conjugate Vaccine on Nasopharyngeal Colonization. 2007.
[322] Pfizer. Study Evaluating 13 Valent Pneumococcal Conjugate Vaccine With Trivalent Inactivated Influenza Vaccine. https://ClinicalTrials.gov/show/NCT00521586; 2007.
[323] Pfizer. Study Evaluating Safety and Immunogenicity of 13-Valent Pneumococcal Conjugate Vaccine With Influenza Vaccine in Adults. https://ClinicalTrials.gov/show/NCT00492557; 2007.
[324] Pfizer. Study Evaluating the Effiacy of a 13-Valent Pneumococcal Conjugate Vaccine (13vPnC) in Adults. 2008.
[325] Pfizer. A Study Evaluating Safety And Immunogenicity Of Meningococcal B Rlp2086 Vaccine In Adolescents. 2009.
[326] Pfizer. Trial Evaluating a 13-valent Pneumococcal Conjugate Vaccine Given With Diphtheria, Tetanus, and Acellular Pertussis Vaccine (DTaP) in Healthy Japanese Infants. 2010.
[327] Pfizer. A Trial to Assess the Safety, Tolerability and Immunogenicity of Repevax and rLP2086 Vaccine When Given Together in Healthy Subjects Aged >=11 to <19 Years. 2011.
[328] Pfizer. A Global Phase 3 Safety Study of 120 mcg rLP2086 Vaccine in Adolescents and Young Adults Aged 10 to 25 Years. 2012.
[329] Pfizer. Immunogenicity and Safety Study of 1 and 2 Doses of GlaxoSmithKline (GSK) Biologicals' Meningococcal Vaccine MenACWY-TT (GSK134612) in Toddlers, Persistence up to 5 Years After Vaccination and Co-administration With Pfizer's Prevenar 13™Vaccine. 2013.
[330] Pfizer. Concomitant Administration of 13-valent Pneumococcal Conjugate Vaccine (13vPnC) With Influenza Vaccine in 23-valent Pneumococcal Polysaccharide (23vPS) Pre-vaccinated Adults. 2015.
[331] Pfizer Winawoso. Study Evaluating A 13-Valent Pneumococcal Conjugate Vaccine Administered To Infants In Korea. https://ClinicalTrials.gov/show/NCT00689351; 2008.
[332] Protein Sciences Corporation. Protective Efficacy of Flublok® Quadrivalent Versus Licensed Inactivated Influenza Vaccine in Adults ≥50 Years of Age. 2015.
[333] Protein Sciences Corporation, Syneos Health, Department of Health Human Services. Safety and Immunogenicity of Flublok Quadrivalent vs IIV4 in Adults 18-49 Years of Age. 2015.

[334] Rowhani-Rahbar A, Fireman B, Lewis E, Nordin J, Naleway A, Jacobsen SJ, et al. Effect of age on the risk of Fever and seizures following immunization with measles-containing vaccines in children. JAMA pediatrics. 2013;167:1111-7.
[335] Ruiz-Sternberg AM, Moreira ED, Jr., Restrepo JA, Lazcano-Ponce E, Cabello R, Silva A, et al. Efficacy, immunogenicity, and safety of a 9-valent human papillomavirus vaccine in Latin American girls, boys, and young women. Papillomavirus research (Amsterdam, Netherlands). 2018;5:63-74.
[336] Sanofi Pasteur a Sanofi Company. Immune Lot Consistency, Immunogenicity, and Safety of an Investigational Quadrivalent Meningococcal Conjugate Vaccine.
[337] Sanofi Pasteur a Sanofi Company, Merck Sharp, Dohme Corp., Sanofi. Study of PR5I, a Pediatric Combination Vaccine With Enhanced Hepatitis B Component Given Concomitantly With Prevnar®. 2006.
[338] Sanofi Pasteur a Sanofi Company, Sanofi. 36-Month Post-marketing Surveillance and Analysis of Menactra Vaccine in 2-10 Year Olds. 2005.
[339] Sanofi Pasteur a Sanofi Company, Sanofi. Descriptive, Post-marketing, Surveillance Safety Study of Menactra Vaccine. 2005.
[340] Sanofi Pasteur a Sanofi Company, Sanofi. Database Surveillance Safety Study of PENTACEL® Vaccine. 2008.
[341] Sanofi Pasteur a Sanofi Company, Sanofi. A Study of Influenza Virus Vaccines in Children and Adults. 2009.
[342] Sanofi Pasteur a Sanofi Company, Sanofi. Study of Quadrivalent Influenza Vaccine Among Children. 2010.
[343] Sanofi Pasteur a Sanofi Company, Sanofi. Study of Quadrivalent Influenza Vaccine Among Adults. https://ClinicalTrials.gov/show/NCT01218646; 2010.
[344] Sanofi Pasteur a Sanofi Company, Sanofi. Post-licensure Safety Surveillance Study of Menactra Vaccine When Administered As a 2-dose Schedule to Children. 2011.
[345] Sanofi Pasteur a Sanofi Company, Sanofi. Safety and Immunogenicity in Adults of Revaccination With Adacel® Vaccine 10 Years After a Previous Dose. 2011.
[346] Sanofi Pasteur a Sanofi Company, Sanofi. Immunogenicity and Safety of an Investigational Quadrivalent Meningococcal Conjugate Vaccine in Healthy Adolescents. 2014.
[347] Sanofi Pasteur a Sanofi Company, Sanofi. Immunogenicity and Safety of a Booster Dose of an Investigational Quadrivalent Meningococcal Conjugate Vaccine. 2016.
[348] Sanofi Pasteur a Sanofi Company, Sanofi. Immunogenicity and Safety of an Investigational Quadrivalent Meningococcal Conjugate Vaccine in Adults 56 Years and Older. 2016.
[349] Sanofi Pasteur a Sanofi Company, Sanofi. Safety and Immunogenicity of High-Dose Quadrivalent Influenza Vaccine in Participants ≥65 Years in the US. 2017.
[350] Sanofi Pasteur aSC, Sanofi. Study of a Quadrivalent Meningococcal Conjugate Vaccine in Subjects Aged 56 and Older. https://ClinicalTrials.gov/show/NCT01732627; 2012.
[351] Sanofi Pasteur aSC, Sanofi. Study of an Investigational Quadrivalent Meningococcal Conjugate Vaccine Administered in Children Aged 2 to 9 Years. https://ClinicalTrials.gov/show/NCT03077438; 2017.
[352] Scott Halperin, IWK Health Centre, Sanofi Pasteur a Sanofi Company, Dalhousie University. Pertussis Maternal Immunization Study. 2007.
[353] Scott J. Statistical Analysis of Heplisav-B AMI Risk. Vaccines and Related Biological Products Advisory Committee Meeting: U.S. Food & Drug Administration; July 28, 2017.

[354] Seqirus. Phase 3 Safety and Immunogenicity Study of aQIV in Elderly Adults.
[355] Seqirus. A Study to Evaluate the Immunogenicity and Safety of bioCSL Quadrivalent Influenza Vaccine (QIV) in Adults Aged 18 Years and Above. 2014.
[356] Simon MW, Brandon D, Christensen S, Baccarini C, Jordanov E, Dhingra MS. Safety and immunogenicity of a quadrivalent meningococcal conjugate vaccine (menacyw-TT) administered in healthy meningococcal vaccine-naÃ¯ve children (2-9 years). Open forum infectious diseases. 2019;6:S958- S9.
[357] Statens Serum Institut. Pneumococcal vaccination of Crohn patients - A randomized, non-blinded phase 4 clinical trial with the purpose of investigating the immune response against two different pneumococcal vaccines in patients with Crohn's disease. 2013.
[358] Statens Serum Institut, Hvidovre University Hospital, Herlev Hospital. Pneumococcal Vaccination of Crohn Patients. 2013.
[359] The University of Hong Kong, Centers for Disease Control Prevention. Immunogenicity of Alternative Annual Influenza Vaccination Strategies in Older Adults in Hong Kong. 2017.
[360] U.S Food and Drug Administration. Clinical Review, November 9, 2017-Heplisav-B: BLA Clinical Review Memorandum. 2017.
[361] University D, Centers for Disease Control Prevention, Kaiser Permanente. Fever After Simultaneous Versus Sequential Vaccination in Young Children. 2018.
[362] University of Newcastle, National Health and Medical Research Council. Pneumococcal polysaccharide vaccine versus Normal Saline for primary prevention of heart attacks and strokes in at-risk Australians aged 55-60. 2015.
[363] University of Siena, Ministry of Education Universities, Research Italy. Serological Response to Antipneumococcal Vaccination and Impact on Streptococcus Pneumoniae Nasal Carriage in HIV Adults. 2011.
[364] van Werkhoven CH, Bonten MJM. The Community-Acquired Pneumonia immunization Trial in Adults (CAPiTA): what is the future of pneumococcal conjugate vaccination in elderly? Future microbiology. 2015;10:1405-13.
[365] Vila-Corcoles A, Ochoa-Gondar O, Rodriguez-Blanco T, de Diego-Cabanes C, Satue-Gracia E, Vila-Rovira A, et al. Evaluating clinical effectiveness of pneumococcal vaccination in preventing stroke: the CAPAMIS Study, 3-year follow-up. Journal of stroke and cerebrovascular diseases : the official journal of National Stroke Association. 2014;23:1577-84.
[366] Vila-Corcoles A, Ochoa-Gondar O, Rodriguez-Blanco T, Gutierrez-Perez A, Vila-Rovira A, Gomez F, et al. Clinical effectiveness of pneumococcal vaccination against acute myocardial infarction and stroke in people over 60 years: the CAPAMIS study, one-year follow-up. BMC Public Health. 2012;12:222.
[367] Wyeth is now a wholly owned subsidiary of Pfizer. Study to Evaluate a 13-valent Pneumococcal Conjugate Vaccine in Elderly Subjects. https://ClinicalTrials.gov/show/NCT00269672; 2005.
[368] Wyeth is now a wholly owned subsidiary of Pfizer. Study Evaluating the Safety, Tolerability and Immunogenicity of 13vPnC as a 2-Dose Regimen or With 23vPS. https://ClinicalTrials.gov/show/NCT00574548; 2007.
[369] Yih K, Lieu T, Kulldorff M, Martin D, McMahill-Walraven C, Platt R, et al. Intussusception Risk After Rotavirus Vaccination in U.S. Infants. Available at: http://www.mini-sentinel.org/work_products/PRISM/Mini-Sentinel_PRISM_Rotavirus-and-intussusception-Report.pdf. Mini-Sentinel Coordinating Center; June 2013.

[370] Andrews N, Stowe J, Miller E, Svanstrom H, Johansen K, Bonhoeffer J, et al. A collaborative approach to investigating the risk of thrombocytopenic purpura after measles-mumps-rubella vaccination in England and Denmark. Vaccine. 2012;30:3042-6.
[371] Andrews N, Stowe J, Wise L, Miller E. Post-licensure comparison of the safety profile of diphtheria/tetanus/whole cell pertussis/haemophilus influenza type b vaccine and a 5-in-1 diphtheria/tetanus/acellular pertussis/haemophilus influenza type b/polio vaccine in the United Kingdom. Vaccine. 2010;28:7215-20.
[372] Armah GE, Sow SO, Breiman RF, Dallas MJ, Tapia MD, Feikin DR, et al. Efficacy of pentavalent rotavirus vaccine against severe rotavirus gastroenteritis in infants in developing countries in sub-Saharan Africa: a randomised, double-blind, placebo-controlled trial. Lancet. 2010;376:606-14.
[373] Barbosa CM, Terreri MT, Rosario PO, de Moraes-Pinto MI, Silva CA, Hilario MO. Immune response and tolerability of varicella vaccine in children and adolescents with systemic lupus erythematosus previously exposed to varicella-zoster virus. Clin Exp Rheumatol. 2012;30:791-8.
[374] Barrett PN, Berezuk G, Fritsch S, Aichinger G, Hart MK, El-Amin W, et al. Efficacy, safety, and immunogenicity of a Vero-cell-culture-derived trivalent influenza vaccine: a multicentre, double-blind, randomised, placebo-controlled trial. Lancet. 2011;377:751-9.
[375] Baxter R, Bakshi N, Fireman B, Lewis E, Ray P, Vellozzi C, et al. Lack of association of guillain-barre syndrome with vaccinations. Clin Infect Dis. 2013;57:197-204.
[376] Baxter R, Toback SL, Sifakis F, Hansen J, Bartlett J, Aukes L, et al. A postmarketing evaluation of the safety of Ann Arbor strain live attenuated influenza vaccine in adults 18-49 years of age. Vaccine. 2012;30:3053-60.
[377] Baxter R, Toback SL, Sifakis F, Hansen J, Bartlett J, Aukes L, et al. A postmarketing evaluation of the safety of Ann Arbor strain live attenuated influenza vaccine in children 5 through 17 years of age. Vaccine. 2012;30:2989-98.
[378] Baxter R, Tran TN, Hansen J, Emery M, Fireman B, Bartlett J, et al. Safety of ZostavaxTM--a cohort study in a managed care organization. Vaccine. 2012;30:6636-41.
[379] Benchimol EI, Hawken S, Kwong JC, Wilson K. Safety and utilization of influenza immunization in children with inflammatory bowel disease. Pediatrics. 2013;131:e1811-20.
[380] Bernsen RMD, de Jongste JC, Koes BW, Aardoom HA, van der Wouden JC. Diphtheria tetanus pertussis poliomyelitis vaccination and reported atopic disorders in 8-12-year-old children. Vaccine. 2006;24:2035-42.
[381] Bertuola F, Morando C, Menniti-Ippolito F, Da Cas R, Capuano A, Perilongo G, et al. Association between drug and vaccine use and acute immune thrombocytopenia in childhood: a case-control study in Italy. Drug Safety. 2010;33:65-72.
[382] Bhatla N, Suri V, Basu P, Shastri S, Datta SK, Bi D, et al. Immunogenicity and safety of human papillomavirus-16/18 AS04-adjuvanted cervical cancer vaccine in healthy Indian women.[Erratum appears in J Obstet Gynaecol Res. 2010 Apr;36(2):466]. J Obstet Gynaecol Res. 2010;36:123-32.
[383] Block SL, Brown DR, Chatterjee A, Gold MA, Sings HL, Meibohm A, et al. Clinical trial and post-licensure safety profile of a prophylactic human papillomavirus (Types 6, 11, 16, and 18) L1 virus-like particle vaccine. Pediatric Infectious Disease Journal. 2010;29:95-101.
[384] Block SL, Vesikari T, Goveia MG, Rivers SB, Adeyi BA, Dallas MJ, et al. Efficacy, immunogenicity, and safety of a pentavalent human-bovine (WC3) reassortant rotavirus vaccine at the end of shelf life. Pediatrics. 2007;119:11-8.

[385] Buttery JP, Danchin MH, Lee KJ, Carlin JB, McIntyre PB, Elliott EJ, et al. Intussusception following rotavirus vaccine administration: post-marketing surveillance in the National Immunization Program in Australia. Vaccine. 2011;29:3061-6.
[386] Capeding MRZ, Nohynek H, Pascual LG, Kayhty H, Sombrero LT, Eskola J, et al. The immunogenicity of three Haemophilus influenzae type B conjugate vaccines after a primary vaccination series in Philippine infants. American Journal of Tropical Medicine and Hygiene. 1996;55:516-20.
[387] Chang C-C, Chang M-H, Lin T-Y, Lee H-C, Hsieh W-S, Lee P-I. Experience of pentavalent human-bovine reassortant rotavirus vaccine among healthy infants in Taiwan. Journal of the Formosan Medical Association. 2009;108:280-5.
[388] Chao C, Klein NP, Velicer CM, Sy LS, Slezak JM, Takhar H, et al. Surveillance of autoimmune conditions following routine use of quadrivalent human papillomavirus vaccine. J Intern Med. 2012;271:193-203.
[389] Chen RT, Glasser JW, Rhodes PH, Davis RL, Barlow WE, Thompson RS, et al. Vaccine safety datalink project: A new tool for improving vaccine safety monitoring in the United States. Pediatrics. 1997;99:765-73.
[390] Clark LR, Myers ER, Huh W, Joura EA, Paavonen J, Perez G, et al. Clinical trial experience with prophylactic human papillomavirus 6/11/16/18 vaccine in young black women. Journal of Adolescent Health. 2013;52:322-9.
[391] Crawford NW, Cheng A, Andrews N, Charles PG, Clothier HJ, Day B, et al. Guillain-Barre syndrome following pandemic (H1N1) 2009 influenza A immunisation in Victoria: a self-controlled case series. Med J Aust. 2012;197:574-8.
[392] De Carvalho N, Teixeira J, Roteli-Martins CM, Naud P, De Borba P, Zahaf T, et al. Sustained efficacy and immunogenicity of the HPV-16/18 AS04-adjuvanted vaccine up to 7.3 years in young adult women. Vaccine. 2010;28:6247-55.
[393] Dennehy PH, Brady RC, Halperin SA, Ward RL, Alvey JC, Fischer FH, Jr., et al. Comparative evaluation of safety and immunogenicity of two dosages of an oral live attenuated human rotavirus vaccine. Pediatric Infectious Disease Journal. 2005;24:481-8.
[394] Dodds L, Macdonald N, Scott J, Spencer A, Allen VM, McNeil S. The association between influenza vaccine in pregnancy and adverse neonatal outcomes. J Obstet Gynaecol Can. 2012;34:714-20.
[395] Duderstadt SK, Rose CE, Jr., Real TM, Sabatier JF, Stewart B, Ma G, et al. Vaccination and risk of type 1 diabetes mellitus in active component U.S. Military, 2002-2008. Vaccine. 2012;30:813-9.
[396] Eder L, Law T, Chandran V, Shanmugarajah S, Shen H, Rosen CF, et al. Association between environmental factors and onset of psoriatic arthritis in patients with psoriasis. Arthritis Care and Research. 2011;63:1091-7.
[397] Englund JA, Walter E, Black S, Blatter M, Nyberg J, Ruben FL, et al. Safety and immunogenicity of trivalent inactivated influenza vaccine in infants: a randomized double-blind placebo-controlled study. Pediatric Infectious Disease Journal. 2010;29:105-10.
[398] Eurich DT, Johnstone JJ, Minhas-Sandhu JK, Marrie TJ, Majumdar SR. Pneumococcal vaccination and risk of acute coronary syndromes in patients with pneumonia: population-based cohort study. Heart. 2012;98:1072-7.
[399] Farez MF, Ysrraelit MC, Fiol M, Correale J. H1N1 vaccination does not increase risk of relapse in multiple sclerosis: a self-controlled case-series study. Multiple Sclerosis. 2012;18:254-6.

[400] Fell DB, Sprague AE, Liu N, Yasseen AS, 3rd, Wen S-W, Smith G, et al. H1N1 influenza vaccination during pregnancy and fetal and neonatal outcomes. American Journal of Public Health. 2012;102:e33-40.
[401] Frey S, Vesikari T, Szymczakiewicz-Multanowska A, Lattanzi M, Izu A, Groth N, et al. Clinical efficacy of cell culture-derived and egg-derived inactivated subunit influenza vaccines in healthy adults. Clinical Infectious Diseases. 2010;51:997-1004.
[402] Gallagher CM, Goodman MS. Hepatitis B Vaccination of Male Neonates and Autism Diagnosis, NHIS 1997-2002. Journal of Toxicology and Environmental Health-Part a-Current Issues. 2010;73:1665-77.
[403] Garbe E, Andersohn F, Bronder E, Salama A, Klimpel A, Thomae M, et al. Drug-induced immune thrombocytopaenia: results from the Berlin Case-Control Surveillance Study. European Journal of Clinical Pharmacology. 2012;68:821-32.
[404] Gee J, Naleway A, Shui I, Baggs J, Yin R, Li R, et al. Monitoring the safety of quadrivalent human papillomavirus vaccine: findings from the Vaccine Safety Datalink. Vaccine. 2011;29:8279-84.
[405] Gilbertson DT, Guo H, Arneson TJ, Collins AJ. The association of pneumococcal vaccination with hospitalization and mortality in hemodialysis patients. Nephrology Dialysis Transplantation. 2011;26:2934-9.
[406] Giuliano AR, Palefsky JM, Goldstone S, Moreira ED, Jr., Penny ME, Aranda C, et al. Efficacy of quadrivalent HPV vaccine against HPV Infection and disease in males.[Erratum appears in N Engl J Med. 2011 Apr 14;364(15):1481]. New England Journal of Medicine. 2011;364:401-11.
[407] Glanz JM, Newcomer SR, Hambidge SJ, Daley MF, Narwaney KJ, Xu S, et al. Safety of trivalent inactivated influenza vaccine in children aged 24 to 59 months in the vaccine safety datalink. Archives of Pediatrics & Adolescent Medicine. 2011;165:749-55.
[408] Gold M, Dugdale S, Woodman RJ, McCaul KA. Use of the Australian Childhood Immunisation Register for vaccine safety data linkage. Vaccine. 2010;28:4308-11.
[409] Gotoh K, Ito Y, Suzuki E, Kaneko K, Kiuchi T, Ando H, et al. Effectiveness and safety of inactivated influenza vaccination in pediatric liver transplant recipients over three influenza seasons. Pediatric Transplantation. 2011;15:112-6.
[410] Greene SK, Rett M, Weintraub ES, Li L, Yin R, Amato AA, et al. Risk of Confirmed Guillain-Barre Syndrome Following Receipt of Monovalent Inactivated Influenza A (H1N1) and Seasonal Influenza Vaccines in the Vaccine Safety Datalink Project, 2009-2010. American Journal of Epidemiology. 2012;175:1100-9.
[411] Greenhawt MJ, Spergel JM, Rank MA, Green TD, Masnoor D, Sharma H, et al. Safe administration of the seasonal trivalent influenza vaccine to children with severe egg allergy. Ann Allergy Asthma Immunol. 2012;109:426-30.
[412] Grimaldi-Bensouda L, Alperovitch A, Besson G, Vial C, Cuisset J-M, Papeix C, et al. Guillain-Barre syndrome, influenzalike illnesses, and influenza vaccination during seasons with and without circulating A/H1N1 viruses. American Journal of Epidemiology. 2011;174:326-35.
[413] Groves FD, Gridley G, Wacholder S, Shu XO, Robison LL, Neglia JP, et al. Infant vaccinations and risk of childhood acute lymphoblastic leukaemia in the USA. Br J Cancer. 1999;81:175-8.
[414] Gruber C, Warner J, Hill D, Bauchau V, Group ES. Early atopic disease and early childhood immunization--is there a link? Allergy. 2008;63:1464-72.

[415] Gwini SM, Coupland CAC, Siriwardena AN. The effect of influenza vaccination on risk of acute myocardial infarction: self-controlled case-series study. Vaccine. 2011;29:1145-9.
[416] Halasa N, Englund JA, Nachman S, Weinberg GA, Huber VC, Allison K, et al. Safety of live attenuated influenza vaccine in mild to moderately immunocompromised children with cancer. Vaccine. 2011;29:4110-5.
[417] Hambidge SJ, Ross C, Glanz J, McClure D, Daley MF, Xu S, et al. Trivalent inactivated influenza vaccine is not associated with sickle cell crises in children. Pediatrics. 2012;129:e54-9.
[418] Hambidge SJ, Ross C, McClure D, Glanz J, team VSD. Trivalent inactivated influenza vaccine is not associated with sickle cell hospitalizations in adults from a large cohort. Vaccine. 2011;29:8179-81.
[419] Hedlund J, Christenson B, Lundbergh P, Ortqvist A. Effects of a large-scale intervention with influenza and 23-valent pneumococcal vaccines in elderly people: A 1-year follow-up. Vaccine. 2003;21:3906-11.
[420] Hummel M, Fuchtenbusch M, Schenker M, Ziegler AG. No major association of breast-feeding, vaccinations, and childhood viral diseases with early islet autoimmunity in the German BABYDIAB study. Diabetes Care. 2000;23:969-74.
[421] Hurst FP, Lee JJ, Jindal RM, Agodoa LY, Abbott KC. Outcomes associated with influenza vaccination in the first year after kidney transplantation. Clin J Am Soc Nephrol. 2011;6:1192-7.
[422] Huu TN, Toan NT, Tuan HM, Viet HL, Le Thanh Binh P, Yu TW, et al. Safety and reactogenicity of primary vaccination with the 10-valent pneumococcal non-typeable Haemophilus influenzae protein D conjugate vaccine in Vietnamese infants: a randomised, controlled trial. BMC Infectious Diseases. 2013;13:95.
[423] Iorio A, Basileo M, Marcucci M, Guercini F, Camilloni B, Paccamiccio E, et al. Influenza vaccination and vitamin K antagonist treatment: a placebo-controlled, randomized, double-blind crossover study. Archives of Internal Medicine. 2010;170:609-16.
[424] Irving SA, Kieke BA, Donahue JG, Mascola MA, Baggs J, DeStefano F, et al. Trivalent inactivated influenza vaccine and spontaneous abortion. Obstet Gynecol. 2013;121:159-65.
[425] Isai A, Durand J, Le Meur S, Hidalgo-Simon A, Kurz X. Autoimmune disorders after immunisation with Influenza A/H1N1 vaccines with and without adjuvant: EudraVigilance data and literature review. Vaccine. 2012;30:7123-9.
[426] Italian Multicenter Study Group for D, Vaccine Safety in C. Effectiveness and safety of the A-H1N1 vaccine in children: a hospital-based case-control study. BMJ Open. 2011;1:e000167.
[427] Jackson LA, Gaglani MJ, Keyserling HL, Balser J, Bouveret N, Fries L, et al. Safety, efficacy, and immunogenicity of an inactivated influenza vaccine in healthy adults: a randomized, placebo-controlled trial over two influenza seasons. BMC Infectious Diseases. 2010;10:71.
[428] Johnstone J, Loeb M, Teo KK, Gao P, Dyal L, Liu L, et al. Influenza vaccination and major adverse vascular events in high risk patients. Circulation. 2012;21.
[429] Kang S, Kim KH, Kim YT, Kim YT, Kim JH, Song YS, et al. Safety and immunogenicity of a vaccine targeting human papillomavirus types 6, 11, 16 and 18: a randomized, placebo-controlled trial in 176 Korean subjects. Int J Gynecol Cancer. 2008;18:1013-9.
[430] Kawamura N, Tokoeda Y, Oshima M, Okahata H, Tsutsumi H, Van Doorn LJ, et al. Efficacy, safety and immunogenicity of RIX4414 in Japanese infants during the first two years of life. Vaccine. 2011;29:6335-41.
[431] Kelly H, Carcione D, Dowse GK, Effler P. The vaccine-attributable risk for febrile convulsions following influenza vaccine. Pediatric Infectious Disease Journal. 2012;31:792.

[432] Kerdpanich A, Chokephaibulkit K, Watanaveeradej V, Vanprapar N, Simasathien S, Phavichitr N, et al. Immunogenicity of a live-attenuated human rotavirus RIX4414 vaccine with or without buffering agent. Human Vaccines. 2010;6:254-62.
[433] Khalil M, Al-Mazrou Y, Findlow H, Chadha H, Bosch Castells V, Johnson DR, et al. Safety and immunogenicity of a meningococcal quadrivalent conjugate vaccine in five- to eight-year-old Saudi Arabian children previously vaccinated with two doses of a meningococcal quadrivalent polysaccharide vaccine. Clin Vaccine Immunol. 2012;19:1561-6.
[434] Khatun S, Akram Hussain SM, Chowdhury S, Ferdous J, Hossain F, Begum SR, et al. Safety and immunogenicity profile of human papillomavirus-16/18 AS04 adjuvant cervical cancer vaccine: a randomized controlled trial in healthy adolescent girls of Bangladesh. Japanese Journal of Clinical Oncology. 2012;42:36-41.
[435] Kim DS, Lee TJ, Kang JH, Kim J-H, Lee JH, Ma SH, et al. Immunogenicity and safety of a pentavalent human-bovine (WC3) reassortant rotavirus vaccine in healthy infants in Korea. Pediatric Infectious Disease Journal. 2008;27:177-8.
[436] Kim JS, Bae CW, Lee KY, Park MS, Choi YY, Kim KN, et al. Immunogenicity, reactogenicity and safety of a human rotavirus vaccine (RIX4414) in Korean infants: A randomized, double-blind, placebo-controlled, phase IV study. Human Vaccines and Immunotherapeutics. 2012;8:806-12.
[437] Kim SC, Song YS, Kim Y-T, Kim YT, Ryu K-S, Gunapalaiah B, et al. Human papillomavirus 16/18 AS04-adjuvanted cervical cancer vaccine: immunogenicity and safety in 15-25 years old healthy Korean women. Journal of Gynecologic Oncology. 2011;22:67-75.
[438] Klein NP, Aukes L, Lee J, Fireman B, Shapira SK, Slade B, et al. Evaluation of immunization rates and safety among children with inborn errors of metabolism. Pediatrics. 2011;127:e1139-e46.
[439] Klein NP, Hansen J, Chao C, Velicer C, Emery M, Slezak J, et al. Safety of quadrivalent human papillomavirus vaccine administered routinely to females. Archives of Pediatrics & Adolescent Medicine. 2012;166:1140-8.
[440] Klein NP, Massolo ML, Greene J, Dekker CL, Black S, Escobar GJ. Risk factors for developing apnea after immunization in the neonatal intensive care unit. Pediatrics. 2008;121:463-9.
[441] Klein NP, Reisinger KS, Johnston W, Odrljin T, Gill CJ, Bedell L, et al. Safety and immunogenicity of a novel quadrivalent meningococcal CRM-conjugate vaccine given concomitantly with routine vaccinations in infants.[Erratum appears in Pediatr Infect Dis J. 2012 Oct;31(10):1105]. Pediatric Infectious Disease Journal. 2012;31:64-71.
[442] Langley JM, Aoki F, Ward BJ, McGeer A, Angel JB, Stiver G, et al. A nasally administered trivalent inactivated influenza vaccine is well tolerated, stimulates both mucosal and systemic immunity, and potentially protects against influenza illness. Vaccine. 2011;29:1921-8.
[443] Lee GM, Greene SK, Weintraub ES, Baggs J, Kulldorff M, Fireman BH, et al. H1N1 and seasonal influenza vaccine safety in the vaccine safety datalink project. American Journal of Preventive Medicine. 2011;41:121-8.
[444] Lee S, Park WB, Shin K-H, Ahn DH, Yoon SH, Cho J-Y, et al. Immunogenicity and safety of a single intramuscular dose of a diphtheria-tetanus toxoid (Td) vaccine (GC1107) in Korean adults. Vaccine. 2011;29:7638-43.

[445] Levin MJ, Moscicki A-B, Song L-Y, Fenton T, Meyer WA, 3rd, Read JS, et al. Safety and immunogenicity of a quadrivalent human papillomavirus (types 6, 11, 16, and 18) vaccine in HIV-infected children 7 to 12 years old. J Acquir Immune Defic Syndr. 2010;55:197-204.
[446] Li R, Li Y, Radley D, Liu Y, Huang T, Sings HL, et al. Safety and immunogenicity of a vaccine targeting human papillomavirus types 6, 11, 16 and 18: a randomized, double-blind, placebo-controlled trial in Chinese males and females. Vaccine. 2012;30:4284-91.
[447] Lin TH, Lin SY, Lin CH, Lin RI, Lin HC, Chiu TH, et al. AdimFlu-S((registered trademark)) influenza A (H1N1) vaccine during pregnancy: The Taiwanese Pharmacovigilance Survey. Vaccine. 2012;30:2671-5.
[448] Ma X, Does MB, Metayer C, Russo C, Wong A, Buffler PA. Vaccination history and risk of childhood leukaemia. Int J Epidemiol. 2005;34:1100-9.
[449] Macaladad N, Marcano T, Guzman M, Moya J, Jurado F, Thompson M, et al. Safety and immunogenicity of a zoster vaccine in varicella-zoster virus seronegative and low-seropositive healthy adults. Vaccine. 2007;25:2139-44.
[450] MacArthur AC, McBride ML, Spinelli JJ, Tamaro S, Gallagher RP, Theriault GP. Risk of childhood leukemia associated with vaccination, infection, and medication use in childhood: the Cross-Canada Childhood Leukemia Study. Am J Epidemiol. 2008;167:598-606.
[451] Madhi SA, Cunliffe NA, Steele D, Witte D, Kirsten M, Louw C, et al. Effect of human rotavirus vaccine on severe diarrhea in African infants. New England Journal of Medicine. 2010;362:289-98.
[452] Madhi SA, Dittmer S, Kuwanda L, Venter M, Cassim H, Lazarus E, et al. Efficacy and immunogenicity of influenza vaccine in HIV-infected children: a randomized, double-blind, placebo controlled trial. Aids. 2013;27:369-79.
[453] Madhi SA, Maskew M, Koen A, Kuwanda L, Besselaar TG, Naidoo D, et al. Trivalent inactivated influenza vaccine in African adults infected with human immunodeficient virus: double blind, randomized clinical trial of efficacy, immunogenicity, and safety. Clinical Infectious Diseases. 2011;52:128-37.
[454] Mallory RM, Malkin E, Ambrose CS, Bellamy T, Shi L, Yi T, et al. Safety and Immunogenicity Following Administration of a Live, Attenuated Monovalent 2009 H1N1 Influenza Vaccine to Children and Adults in Two Randomized Controlled Trials. PLoS ONE [Electronic Resource]. 2010;5:e13755.
[455] Matheson MC, Haydn Walters E, Burgess JA, Jenkins MA, Giles GG, Hopper JL, et al. Childhood immunization and atopic disease into middle-age--a prospective cohort study. Pediatr Allergy Immunol. 2010;21:301-6.
[456] Mills R, Tyring SK, Levin MJ, Parrino J, Li X, Coll KE, et al. Safety, tolerability, and immunogenicity of zoster vaccine in subjects with a history of herpes zoster. Vaccine. 2010;28:4204-9.
[457] Mommers M, Weishoff-Houben M, Swaen GM, Creemers H, Freund H, Dott W, et al. Infant immunization and the occurrence of atopic disease in Dutch and German children: a nested case-control study. Pediatr Pulmonol. 2004;38:329-34.
[458] Moreira Jr ED, Palefsky JM, Giuliano AR, Goldstone S, Aranda C, Jessen H, et al. Safety and reactogenicity of a quadrivalent human papillomavirus (types 6, 11, 16, 18) L1 viral-like-particle vaccine in older adolescents and young adults. Human Vaccines. 2011;7:768-75.
[459] Morgan TM, Schlegel C, Edwards KM, Welch-Burke T, Zhu Y, Sparks R, et al. Vaccines are not associated with metabolic events in children with urea cycle disorders. Pediatrics. 2011;127:e1147-e53.

[460] Mullooly JP, Schuler R, Barrett M, Maher JE. Vaccines, antibiotics, and atopy. Pharmacoepidemiology and Drug Safety. 2007;16:275-88.
[461] Mullooly JP, Schuler R, Mesa J, Drew L, DeStefano F, team VSD. Wheezing lower respiratory disease and vaccination of premature infants. Vaccine. 2011;29:7611-7.
[462] Murray AV, Reisinger KS, Kerzner B, Stek JE, Sausser TA, Xu J, et al. Safety and tolerability of zoster vaccine in adults >=60 years old. Human Vaccines. 2011;7:1130-6.
[463] Nakajima K, Dharmage SC, Carlin JB, Wharton CL, Jenkins MA, Giles GG, et al. Is childhood immunisation associated with atopic disease from age 7 to 32 years? Thorax. 2007;62:270-5.
[464] Narang A, Bose A, Pandit AN, Dutta P, Kang G, Bhattacharya SK, et al. Immunogenicity, reactogenicity and safety of human rotavirus vaccine (RIX4414) in Indian infants. Human Vaccines. 2009;5:414-9.
[465] Ngan HYS, Cheung ANY, Tam KF, Chan KKL, Tang HW, Bi D, et al. Human papillomavirus-16/18 AS04-adjuvanted cervical cancer vaccine: Immunogenicity and safety in healthy Chinese women from Hong Kong. Hong Kong Medical Journal. 2010;16:171-9.
[466] Nordin JD, Kharbanda EO, Benitez GV, Nichol K, Lipkind H, Naleway A, et al. Maternal safety of trivalent inactivated influenza vaccine in pregnant women. Obstet Gynecol. 2013;121:519-25.
[467] O'Leary ST, Glanz JM, McClure DL, Akhtar A, Daley MF, Nakasato C, et al. The risk of immune thrombocytopenic purpura after vaccination in children and adolescents. Pediatrics. 2012;129:248-55.
[468] Omenaca F, Sarlangue J, Szenborn L, Nogueira M, Suryakiran PV, Smolenov IV, et al. Safety, reactogenicity and immunogenicity of the human rotavirus vaccine in preterm European Infants: a randomized phase IIIb study. Pediatric Infectious Disease Journal. 2012;31:487-93.
[469] Omer SB, Goodman D, Steinhoff MC, Rochat R, Klugman KP, Stoll BJ, et al. Maternal influenza immunization and reduced likelihood of prematurity and small for gestational age births: a retrospective cohort study. PLoS Med. 2011;8:e1000441.
[470] Pagaoa MA, Okcu MF, Bondy ML, Scheurer ME. Associations between vaccination and childhood cancers in Texas regions. Journal of Pediatrics. 2011;158:996-1002.
[471] Pahud BA, Rowhani-Rahbar A, Glaser C, Gavali S, Salibay CJ, Fireman B, et al. Lack of association between childhood immunizations and encephalitis in California, 1998-2008. Vaccine. 2012;30:247-53.
[472] Patel MM, Lopez-Collada VR, Bulhoes MM, De Oliveira LH, Bautista Marquez A, Flannery B, et al. Intussusception risk and health benefits of rotavirus vaccination in Mexico and Brazil. New England Journal of Medicine. 2011;364:2283-92.
[473] Phua KB, Lim FS, Lau YL, Nelson EAS, Huang LM, Quak SH, et al. Rotavirus vaccine RIX4414 efficacy sustained during the third year of life: A randomized clinical trial in an Asian population. Vaccine. 2012;30:4552-7.
[474] Phua KB, Quak SH, Lee BW, Emmanuel SC, Goh P, Han HH, et al. Evaluation of RIX4414, A live, attenuated rotavirus vaccine, in a randomized, double-blind, placebo-controlled phase 2 trial involving 2464 Singaporean infants. Journal of Infectious Diseases. 2005;192:S6-S16.
[475] Ray P, Black S, Shinefield H, Dillon A, Carpenter D, Lewis E, et al. Risk of rheumatoid arthritis following vaccination with tetanus, influenza and hepatitis B vaccines among persons 15-59 years of age. Vaccine. 2011;29:6592-7.

[476] Richards JL, Hansen C, Bredfeldt C, Bednarczyk RA, Steinhoff MC, Adjaye-Gbewonyo D, et al. Neonatal outcomes after antenatal influenza immunization during the 2009 H1N1 influenza pandemic: impact on preterm birth, birth weight, and small for gestational age birth. Clinical Infectious Diseases. 2013;56:1216-22.
[477] Roteli-Martins CM, Naud P, De Borba P, Teixeira JC, De Carvalho NS, Zahaf T, et al. Sustained immunogenicity and efficacy of the HPV-16/18 AS04-adjuvanted vaccine: Up to 8.4 years of follow-up. Human Vaccines and Immunotherapeutics. 2012;8:381-8.
[478] Rowhani-Rahbar A, Klein NP, Lewis N, Fireman B, Ray P, Rasgon B, et al. Immunization and bell's palsy in children: A case-centered analysis. American Journal of Epidemiology. 2012;175:878-85.
[479] Ruiz-Palacios GM, Perez-Schael I, Velazquez FR, Abate H, Breuer T, Clemens SC, et al. Safety and efficacy of an attenuated vaccine against severe rotavirus gastroenteritis. New England Journal of Medicine. 2006;354:11-22.
[480] Santosham M, Wolff M, Reid R, Hohenboken M, Bateman M, Goepp J, et al. The efficacy in Navajo infants of a conjugate vaccine consisting of Haemophilus influenzae type b polysaccharide and Neisseria meningitidis outer-membrane protein complex. New England Journal of Medicine. 1991;324:1767-72.
[481] Schmader KE, Levin MJ, Gnann JW, Jr., McNeil SA, Vesikari T, Betts RF, et al. Efficacy, safety, and tolerability of herpes zoster vaccine in persons aged 50-59 years. Clinical Infectious Diseases. 2012;54:922-8.
[482] Schwarz TF, Huang LM, Medina DM, Valencia A, Lin TY, Behre U, et al. Four-year follow-up of the immunogenicity and safety of the HPV-16/18 AS04-adjuvanted vaccine when administered to adolescent girls aged 10-14 years. Journal of Adolescent Health. 2012;50:187-94.
[483] Shui IM, Baggs J, Patel M, Parashar UD, Rett M, Belongia EA, et al. Risk of intussusception following administration of a pentavalent rotavirus vaccine in US infants. JAMA. 2012;307:598-604.
[484] Siberry GK, Williams PL, Lujan-Zilbermann J, Warshaw MG, Spector SA, Decker MD, et al. Phase I/II, open-label trial of safety and immunogenicity of meningococcal (groups A, C, Y, and W-135) polysaccharide diphtheria toxoid conjugate vaccine in human immunodeficiency virus-infected adolescents. Pediatric Infectious Disease Journal. 2010;29:391-6.
[485] Simberkoff MS, Arbeit RD, Johnson GR, Oxman MN, Boardman KD, Williams HM, et al. Safety of herpes zoster vaccine in the shingles prevention study: a randomized trial. Annals of Internal Medicine. 2010;152:545-54.
[486] Siriwardena AN, Gwini SM, Coupland CAC. Influenza vaccination, pneumococcal vaccination and risk of acute myocardial infarction: matched case-control study. CMAJ Canadian Medical Association Journal. 2010;182:1617-23.
[487] Sow PS, Watson-Jones D, Kiviat N, Changalucha J, Mbaye KD, Brown J, et al. Safety and immunogenicity of human papillomavirus-16/18 AS04-adjuvanted vaccine: a randomized trial in 10-25-year-old HIV-Seronegative African girls and young women. Journal of Infectious Diseases. 2013;207:1753-63.
[488] Steele AD, Reynders J, Scholtz F, Bos P, de Beer MC, Tumbo J, et al. Comparison of 2 different regimens for reactogenicity, safety, and immunogenicity of the live attenuated oral rotavirus vaccine RIX4414 coadministered with oral polio vaccine in South African infants. Journal of Infectious Diseases. 2010;202 Suppl:S93-100.

[489] Stowe J, Andrews N, Bryan P, Seabroke S, Miller E. Risk of convulsions in children after monovalent H1N1 (2009) and trivalent influenza vaccines: A database study. Vaccine. 2011;29:9467-72.
[490] Sun YL, Christensen J, Hviid A, Li J, Vedsted P, Olsen J, et al. Risk of Febrile Seizures and Epilepsy After Vaccination With Diphtheria, Tetanus, Acellular Pertussis, Inactivated Poliovirus, and Haemophilus Influenzae Type b. Jama-Journal of the American Medical Association. 2012;307:823-31.
[491] Talaat KR, Greenberg ME, Lai MH, Hartel GF, Wichems CH, Rockman S, et al. A single dose of unadjuvanted novel 2009 H1N1 vaccine is immunogenic and well tolerated in young and elderly adults. Journal of Infectious Diseases. 2010;202:1327-37.
[492] Tanner K, Roy N, Merrill RM, Sauder C, Houtz DR, Smith ME. Case-control study of risk factors for spasmodic dysphonia: A comparison with other voice disorders. Laryngoscope. 2012;122:1082-92.
[493] Thomson JA, Widjaja C, Darmaputra AAP, Lowe A, Matheson MC, Bennett CM, et al. Early childhood infections and immunisation and the development of allergic disease in particular asthma in a high-risk cohort: A prospective study of allergy-prone children from birth to six years. Pediatr Allergy Immunol. 2010;21:1076-85.
[494] Ting SCH, Crooks SW, South G. The effect of influenza vaccination on the incidence of chronic obstructive pulmonary disease exacerbations in the immediate postvaccination period. J Epidemiol Community Health. 2011;65:157-9.
[495] Tokars JI, Lewis P, DeStefano F, Wise M, Viray M, Morgan O, et al. The risk of Guillain-Barre syndrome associated with influenza A (H1N1) 2009 monovalent vaccine and 2009-2010 seasonal influenza vaccines: results from self-controlled analyses. Pharmacoepidemiol Drug Saf. 2012;21:546-52.
[496] Treanor JJ, El Sahly H, King J, Graham I, Izikson R, Kohberger R, et al. Protective efficacy of a trivalent recombinant hemagglutinin protein vaccine (FluBlok[REGISTERED]) against influenza in healthy adults: a randomized, placebo-controlled trial. Vaccine. 2011;29:7733-9.
[497] Tse A, Tseng HF, Greene SK, Vellozzi C, Lee GM, Group VSDRCAIW. Signal identification and evaluation for risk of febrile seizures in children following trivalent inactivated influenza vaccine in the Vaccine Safety Datalink Project, 2010-2011. Vaccine. 2012;30:2024-31.
[498] Tseng HF, Liu A, Sy L, Marcy SM, Fireman B, Weintraub E, et al. Safety of zoster vaccine in adults from a large managed-care cohort: a Vaccine Safety Datalink study. J Intern Med. 2012;271:510-20.
[499] Tseng HF, Slezak JM, Quinn VP, Sy LS, Van den Eeden SK, Jacobsen SJ. Pneumococcal vaccination and risk of acute myocardial infarction and stroke in men. JAMA. 2010;303:1699-706.
[500] Uno Y, Uchiyama T, Kurosawa M, Aleksic B, Ozaki N. The combined measles, mumps, and rubella vaccines and the total number of vaccines are not associated with development of autism spectrum disorder: The first case-control study in Asia. Vaccine. 2012;30:4292-8.
[501] Velazquez FR, Colindres RE, Grajales C, Hernandez MT, Mercadillo MG, Torres FJ, et al. Postmarketing surveillance of intussusception following mass introduction of the attenuated human rotavirus vaccine in Mexico. Pediatric Infectious Disease Journal. 2012;31:736-44.
[502] Velentgas P, Amato AA, Bohn RL, Chan KA, Cochrane T, Funch DP, et al. Risk of Guillain-Barre syndrome after meningococcal conjugate vaccination. Pharmacoepidemiol Drug Saf. 2012;21:1350-8.

[503] Vermeulen JN, Lange JMA, Tyring SK, Peters PH, Nunez M, Poland G, et al. Safety, tolerability, and immunogenicity after 1 and 2 doses of zoster vaccine in healthy adults >=60 years of age. Vaccine. 2012;30:904-10.
[504] Vesikari T, Clark HF, Offit PA, Dallas MJ, DiStefano DJ, Goveia MG, et al. Effects of the potency and composition of the multivalent human-bovine (WC3) reassortant rotavirus vaccine on efficacy, safety and immunogenicity in healthy infants. Vaccine. 2006;24:4821-9.
[505] Vesikari T, Karvonen A, Bouckenooghe A, Suryakiran PV, Smolenov I, Han HH. Immunogenicity, reactogenicity and safety of the human rotavirus vaccine RIX4414 oral suspension (liquid formulation) in Finnish infants. Vaccine. 2011;29:2079-84.
[506] Vesikari T, Karvonen A, Korhonen T, Espo M, Lebacq E, Forster J, et al. Safety and immunogenicity of RIX4414 live attenuated human rotavirus vaccine in adults, toddlers and previously uninfected infants. Vaccine. 2004;22:2836-42.
[507] Vesikari T, Karvonen A, Puustinen L, Zeng S-Q, Szakal ED, Delem A, et al. Efficacy of RIX4414 live attenuated human rotavirus vaccine in Finnish infants. Pediatric Infectious Disease Journal. 2004;23:937-43.
[508] Vesikari T, Matson DO, Dennehy P, Van Damme P, Santosham M, Rodriguez Z, et al. Safety and efficacy of a pentavalent human-bovine (WC3) reassortant rotavirus vaccine. New England Journal of Medicine. 2006;354:23-33.
[509] Wang IK, Lin CL, Lin PC, Liang CC, Liu YL, Chang CT, et al. Effectiveness of influenza vaccination in patients with end-stage renal disease receiving hemodialysis: a population-based study. PLoS ONE [Electronic Resource]. 2013;8:e58317.
[510] Weinberg A, Levin MJ, Macgregor RR. Safety and immunogenicity of a live attenuated varicella vaccine in VZV-seropositive HIV-infected adults. Human Vaccines. 2010;6:318-21.
[511] Whitehouse AJO, Maybery M, Wray JA, Hickey M. No association between early gastrointestinal problems and autistic-like traits in the general population. Dev Med Child Neurol. 2011;53:457-62.
[512] Wilson K, Hawken S, Kwong JC, Deeks S, Crowcroft NS, Van Walraven C, et al. Adverse events following 12 and 18 month vaccinations: a population-based, self-controlled case series analysis. PLoS ONE [Electronic Resource]. 2011;6:e27897.
[513] Wise ME, Viray M, Sejvar JJ, Lewis P, Baughman AL, Connor W, et al. Guillain-Barre Syndrome During the 2009-2010 H1N1 Influenza Vaccination Campaign: Population-based Surveillance Among 45 Million Americans. American Journal of Epidemiology. 2012;175:1110-9.
[514] Xu R, Luo Y, Chambers C. Assessing the effect of vaccine on spontaneous abortion using time-dependent covariates Cox models. Pharmacoepidemiology and Drug Safety. 2012.
[515] Yih WK, Lee GM, Lieu TA, Ball R, Kulldorff M, Rett M, et al. Surveillance for Adverse Events Following Receipt of Pandemic 2009 H1N1 Vaccine in the Post-Licensure Rapid Immunization Safety Monitoring (PRISM) System, 2009-2010. American Journal of Epidemiology. 2012;175:1120-8.
[516] Yu O, Bohlke K, Hanson CA, Delaney K, Rees TG, Zavitkovsky A, et al. Hepatitis B vaccine and risk of autoimmune thyroid disease: A Vaccine Safety Datalink study. Pharmacoepidemiology and Drug Safety. 2007;16:736-45.
[517] Zaman K, Sack DA, Yunus M, Arifeen SE, Podder G, Azim T, et al. Successful co-administration of a human rotavirus and oral poliovirus vaccines in Bangladeshi infants in a 2-dose schedule at 12 and 16 weeks of age. Vaccine. 2009;27:1333-9.

[518] Zhang J, Xie F, Delzell E, Chen L, Winthrop KL, Lewis JD, et al. Association between vaccination for herpes zoster and risk of herpes zoster infection among older patients with selected immune-mediated diseases. JAMA. 2012;308:43-9.
[519] Christie CDC, Duncan ND, Thame KA, Onorato MT, Smith HD, Malcolm LG, et al. Pentavalent rotavirus vaccine in developing countries: safety and health care resource utilization. Pediatrics. 2010;126:e1499-506.
[520] EU/EEA EO. Study to evaluate the efficacy, safety and immunogenicity of two or three doses of GSK Biologicals' oral live attenuated human rotavirus (HRV) vaccine given concomitantly with routine EPI vaccinations in healthy infants. http://wwwwhoint/trialsearch/Trial2aspx?TrialID=EUCTR2015-001485-26-Outside-EU/EEA. 2015.
[521] EU/EEA EO. A phase III study to assess the efficacy, immunogenicity and safety of GSK Biologicals' human rotavirus (HRV) vaccine given concomitantly with routine expanded program on immunisation (EPI) vaccinations including oral poliovirus vaccine (OPV) in healthy infants across 6 countries in Latin America. http://wwwwhoint/trialsearch/Trial2aspx?TrialID=EUCTR2015-001540-10-Outside-EU/EEA. 2015.
[522] Euctr Outside EU/EEA. A phase II study to assess the safety and immunogenicity of GlaxoSmithKline Biologicals' rotavirus vaccine, RIX4414 when administered to HIV infected infants in South Africa. http://wwwwhoint/trialsearch/Trial2aspx?TrialID=EUCTR2015-001484-39-Outside-EU/EEA. 2015.
[523] Euctr Outside EU/EEA. Study to evaluate the efficacy and safety of two doses of GSK Biologicals' oral live attenuated human rotavirus (HRV) vaccine in healthy infants. 2015.
[524] Euctr Outside EU/EEA. Study to evaluate immunogenicity, reactogenicity and safety of Rotarix™ vaccine in Korean infants. http://wwwwhoint/trialsearch/Trial2aspx?TrialID=EUCTR2015-001545-81-Outside-EU/EEA. 2015.
[525] Euctr Outside EU/EEA. Study to evaluate the efficacy and safety of two doses of GSK Biologicals' oral live attenuated human rotavirus (HRV) vaccine in healthy infants. http://wwwwhoint/trialsearch/Trial2aspx?TrialID=EUCTR2015-001541-92-Outside-EU/EEA. 2015.
[526] GlaxoSmithKline. A Study to Test 2 Doses of GSK Biologicals' Oral Live Attenuated Human Rotavirus (HRV) Vaccine in Healthy Infants. https://ClinicalTrials.gov/show/NCT00197210; 2003.
[527] GlaxoSmithKline. Study of 2 Doses of HRV Vaccine Given Concomitantly With Routine EPI Vaccinations Including OPV in Healthy Infants. 2003.
[528] GlaxoSmithKline. Year 3 Extension for Efficacy Follow-up in Subjects Vaccinated in Studies Rota-028, 029 or 030 (NCT00197210). https://ClinicalTrials.gov/show/NCT00329745; 2007.
[529] GlaxoSmithKline Biologicals. **A phase III, double-blind, randomized, placebo-controlled, multi-country and multi-center study to assess the efficacy and safety of two doses of GSK Biologicals' oral live attenuated human rotavirus (HRV) vaccine in healthy infants** 2015.
[530] GlaxoSmithKline Biologicals. Study to evaluate immunogenicity, reactogenicity and safety of Rotarix™ vaccine in Korean infants.

http://wwwwhoint/trialsearch/Trial2aspx?TrialID=EUCTR2015-001545-81-Outside-EU/EEA. 2015.
[531] Goveia MG, Rodriguez ZM, Dallas MJ, Itzler RF, Boslego JW, Heaton PM, et al. Safety and efficacy of the pentavalent human-bovine (WC3) reassortant rotavirus vaccine in healthy premature infants. Pediatric Infectious Disease Journal. 2007;26:1099-104.
[532] Grant LR, Watt JP, Weatherholtz RC, Moulton LH, Reid R, Santosham M, et al. Efficacy of a pentavalent human-bovine reassortant rotavirus vaccine against rotavirus gastroenteritis among American Indian children. Pediatric Infectious Disease Journal. 2012;31:184-8.
[533] Guillermo M, al. e. RIX4414 (Rotarix™) has demonstrated efficacy during the first 2 years of life in infants from 11 Latin American countries. 10th International Rotavirus Symposium (IRS). Bangkok, Thailand2012.
[534] Hemming-Harlo M, Lahdeaho M-L, Maki M, Vesikari T. Rotavirus Vaccination Does Not Increase Type 1 Diabetes and May Decrease Celiac Disease in Children and Adolescents. The Pediatric infectious disease journal. 2019;38:539-41.
[535] Laserson KF, Nyakundi D, Feikin DR, Nyambane G, Cook E, Oyieko J, et al. Safety of the pentavalent rotavirus vaccine (PRV), RotaTeq([REGISTERED]), in Kenya, including among HIV-infected and HIV-exposed infants. Vaccine. 2012;30 Suppl 1:A61-70.
[536] Lau YL, Nelson EA, Poon KH, Chan PK, Chiu S, Sung R, et al. Efficacy, safety and immunogenicity of a human rotavirus vaccine (RIX4414) in Hong Kong children up to three years of age: a randomized, controlled trial. Vaccine. 2013;31:2253-9.
[537] Lee KJ, Carlin JB. Fractional polynomial adjustment for time-varying covariates in a self-controlled case series analysis. Stat Med. 2014;33:105-16.
[538] Oxman MN, Levin MJ, Johnson GR, Schmader KE, Straus SE, Gelb LD, et al. A vaccine to prevent herpes zoster and postherpetic neuralgia in older adults. N Engl J Med. 2005;352:2271-84.
[539] Phua KB, al. e. Human rotavirus vaccine RIX4414 (Rotarix™) is highly efficacious in infants from Asia during the first two years of life. 13th International Congress on Infectious Diseases (ICID). Kuala Lumpur, Malaysia2008.
[540] Phua KB, al. e. Human rotavirus vaccine RIX4414 (Rotarix™) is highly efficacious in Asian infants during the first three years of life. 13th Asian Pacific Congress of Pediatrics (APCP). Shanghai, China2009.
[541] Phua KB, al. e. Human rotavirus vaccine RIX4414 is highly efficacious in Asian infants during the third year of life. 27th Annual Meeting of the European Society for Paediatric Infectious Diseases (ESPID). Brussels, Belgium2009.
[542] Phua KB, al. e. Efficacy of rotavirus vaccine RIX4414 during the first 3 years of life: a randomised, double-blind, placebo-controlled study in infants from Hong Kong, Singapore and Taiwan. 10th International Rotavirus Symposium (IRS). Bangkok, Thailand2012.
[543] Phua KB, Lim FS, Lau YL, Nelson EAS, Huang LM, Quak SH, et al. Safety and efficacy of human rotavirus vaccine during the first 2 years of life in Asian infants: randomised, double-blind, controlled study. Vaccine. 2009;27:5936-41.
[544] Popmihajlov Z, Pang L, Brown E, Joshi A, Su SC, Kaplan SS, et al. A post hoc analysis utilizing the FDA toxicity grading scale to assess injection site adverse events following immunization with the live attenuated Zoster Vaccine (ZVL). Human Vaccines and Immunotherapeutics. 2018;14:2916-20.
[545] Rodriguez ZM, Goveia MG, Stek JE, Dallas MJ, Boslego JW, DiNubile MJ, et al. Concomitant use of an oral live pentavalent human-bovine reassortant rotavirus vaccine with

licensed parenteral pediatric vaccines in the United States. Pediatric Infectious Disease Journal. 2007;26:221-7.
[546] Sow SO, Tapia M, Haidara FC, Ciarlet M, Diallo F, Kodio M, et al. Efficacy of the oral pentavalent rotavirus vaccine in Mali. Vaccine. 2012;30 Suppl 1:A71-8.
[547] Tregnaghi MW, Abate HJ, Valencia A, Lopez P, Da Silveira TR, Rivera L, et al. Human rotavirus vaccine is highly efficacious when coadministered with routine expanded program of immunization vaccines including oral poliovirus vaccine in Latin America. Pediatric Infectious Disease Journal. 2011;30:e103-e8.
[548] Zaman K, Dang DA, Victor JC, Shin S, Yunus M, Dallas MJ, et al. Efficacy of pentavalent rotavirus vaccine against severe rotavirus gastroenteritis in infants in developing countries in Asia: a randomised, double-blind, placebo-controlled trial. Lancet. 2010;376:615-23.
[549] Zaman K, Yunus M, El Arifeen S, Azim T, Faruque ASG, Huq E, et al. Methodology and lessons-learned from the efficacy clinical trial of the pentavalent rotavirus vaccine in Bangladesh. Vaccine. 2012;30 Suppl 1:A94-100.
[550] Dudley MZ, Halsey NA, Omer SB, Orenstein WA, O'Leary ST, Limaye RJ, et al. The state of vaccine safety science: systematic reviews of the evidence. Lancet Infect Dis. 2020;20:e80-e9.
[551] Lu H-L, Ding Y, Goyal H, Xu H-G. Association Between Rotavirus Vaccination and Risk of Intussusception Among Neonates and Infants: A Systematic Review and Meta-analysis. JAMA network open. 2019;2:e1912458.
[552] Health Resources and Services Administration. Vaccine Injury Table.
[553] Hartung DM, Zarin DA, Guise JM, McDonagh M, Paynter R, Helfand M. Reporting discrepancies between the ClinicalTrials.gov results database and peer-reviewed publications. Ann Intern Med. 2014;160:477-83.
[554] Tang E, Ravaud P, Riveros C, Perrodeau E, Dechartres A. Comparison of serious adverse events posted at ClinicalTrials.gov and published in corresponding journal articles. BMC Med. 2015;13:189.
[555] HealthIT.gov. Interoperability Proving Ground.

2122 **Table 1. Included vaccines, populations, and recent changes (within five years)**

Vaccine (abbreviation; brand name)	Populations recommended for routine use	Recent changes to formulation, age indication, or dosing (within last five years)
9-valent human papillomavirus (HPV9; Gardasil 9®)	Adults, children	Gardasil 9 approval expanded to include use in women and men 27 through 45 years of age in 2018. Gardasil 9 approved as a two-dose series if first dose initiated 9-14 years of age (otherwise three-dose series as before) in 2016. Catch-up HPV vaccination recommended for all persons through age 26 years in 2019.
13-valent pneumococcal conjugate (PCV13; Prevnar 13®)	Adults, children	Age indications were expanded from younger than 18 years and older than 50 years to include adults aged 18-49 years in 2016.
23-valent pneumococcal polysaccharide (PPSV23; Pneumovax®)	Adults, children	None
Diphtheria, tetanus, and acellular pertussis (DTaP; Daptacel®, Infanrix®)	Children	None
Haemophilus influenzae type b (Hib; ActHIB®, Hiberix®, PedvaxHIB®)	Children	Hiberix approved in 2016 as a three-dose primary series at ages 2, 4, and 6 months (initially approved only as a booster dose for ages 15 months through 4 years).
Hepatitis A (HepA; Havrix®, Vaqta®)	Adults, children	None
Hepatitis B (HepB; Engerix-B®, Recombivax HB®, HEPLISAV-B®)	Adults, children, pregnant women (except for HEPLISAV-B, which is not recommended for children and pregnant women)	HEPLISAV-B approved in 2017.
Hepatitis A-Hepatitis B (HepA-HepB; Twinrix®)	Adults	None
Inactivated poliovirus (IPV; IPOL®)	Children	None
Influenza, inactivated (IIV; Afluria Quadrivalent®, Fluarix Quadrivalent®, Flucelvax Quadrivalent®, Flulaval Quadrivalent®, Fluzone High Dose Quadrivalent®, Fluzone Quadrivalent®)	Adults, children, pregnant women (except for Fluzone High Dose Quadrivalent, which is for adults aged 65 years and older)	Afluria Quadrivalent and Flucelvax Quadrivalent approved in 2016. Fluzone High Dose Quadrivalent approved in 2019. Flulaval Quadrivalent expanded use to 6 months of age and older in 2016. Afluria Quadrivalent and Fluarix Quadrivalent expanded use to 6 months of age and older in 2018. Fluzone Quadrivalent dose for children aged 6 through 35 months was updated to be either 0.25 mL or 0.5 mL in 2018.
Influenza, inactivated, adjuvanted (aIIV; Fluad®, Fluad Quadrivalent®)	Adults aged 65 years and older	Fluad approved in 2015; Fluad Quadrivalent approved in 2020. Changes to influenza strains for vaccine made annually.
Influenza, recombinant (RIV; Flublok Quadrivalent®)	Adults, pregnant women	Flublok Quadrivalent approved in 2017. Changes to influenza strains for vaccine made annually.
Influenza, live attenuated (LAIV; FluMist Quadrivalent®)	Adults (through 49 years of age), children	Changes to influenza vaccine strains made annually.
Measles, mumps, rubella (MMR; M-M-R II®)	Adults, children	None
Serogroup A, C, W, and Y meningococcal (MenACWY-D, Menactra®; Men-ACWY-CRM,	Adults, children	MenQuadFi (MenACWY-TT) was approved in 2020.

Vaccine (abbreviation; brand name)	Populations recommended for routine use	Recent changes to formulation, age indication, or dosing (within last five years)
Menveo®; MenACWY-TT, MenQuadFi®)		
Serogroup B meningococcal (MenB-FHbp, Trumenba®; MenB-4C, Bexsero®)	Adults, children	None.
Rotavirus (RV; Rotarix®, RotaTeq®)	Children	None
Tetanus, diphtheria, and acellular pertussis (Tdap; Adacel®, Boostrix®)	Children, adults, pregnant women	Adacel approved for repeat dose in people 10 through 64 years of age in 2019. ACIP recommendation updated to allow for use of Tdap or Td as decennial booster, wound prophylaxis, and catch up vaccination in 2020.
Tetanus, diphtheria (Td; TDVAX®, Tenivac®)	Adults	None
Varicella (VAR; Varivax®)	Children, adults	None
Zoster recombinant (RZV; Shingrix®)	Adults	Shingrix was approved in 2017. (Use of live zoster vaccine [Zostavax] was discontinued in November 2020.)

Abbreviations: aIIV—Adjuvanted inactivated influenza vaccine; DTaP— Diphtheria and tetanus toxoids and acellular pertussis vaccine; HepA—Hepatitis A vaccine; HepB—Hepatitis B vaccine; HepA-HepB—Hepatitis A and Hepatitis B vaccines; Hib— *Haemophilus influenzae* type b vaccine; HPV9—9-valent human papillomavirus vaccine; IIV—Inactivated influenza vaccine; IPV—Inactivated poliovirus vaccine; LAIV—Live attenuated influenza vaccine; MenACWY—Serogroups A, C, W, and Y meningococcal vaccine; MenB—Serogroup B meningococcal vaccine; MMR—Measles, mumps, and rubella vaccine; MMR-V—Measles, mumps, rubella, and varicella vaccine; PCV13—13-valent pneumococcal conjugate vaccine; PPSV23—23-valent pneumococcal polysaccharide vaccine; RIV—Recombinant influenza vaccine; RV—Rotavirus vaccine; RZV—Recombinant zoster vaccine; Td—Tetanus and diphtheria toxoids; Tdap—Tetanus and diphtheria toxoids and acellular pertussis vaccine; VAR—Varicella vaccine
Note: Combination vaccines that incorporate existing vaccines (e.g., DTaP-IPV/Hib) were also assessed, and are summarized in the Appendix.

2135 **Table 2. Strength of Evidence (SoE) for safety of vaccines in children**

Vaccine (abbreviation; brand name[s])	Synthesis of SoE* and findings for vaccines currently in use in children
9-valent human papillomavirus (HPV9; Gardasil 9®)	Low: No evidence of increased risk of autoimmune disease, birth defects, death, reproductive system events, seizures, spontaneous abortion
13-valent pneumococcal conjugate (PCV13; Prevnar 13®)	Low: Increased risk of febrile seizures Moderate: No evidence of increased risk of death Low: No evidence of increased risk of asthma, cardiovascular events, intussusception, meningitis, reproductive system events, seizures
23-valent pneumococcal polysaccharide (PPSV23; Pneumovax®)	Insufficient evidence to draw conclusions about key adverse events
Diphtheria, tetanus, and acellular pertussis (DTaP; Daptacel®, Infanrix®)	Moderate: No evidence of increased risk of type 1 diabetes mellitus Low: No evidence of increased risk of asthma or death
Tetanus, diphtheria, and acellular pertussis (Tdap; Adacel®, Boostrix®)	Low: No evidence of increased risk of cardiovascular events, death
Haemophilus influenzae type b (Hib; PedvaxHIB®, ActHIB®, Hiberix®)	Moderate**:** No evidence of increased risk of serious adverse events in short term
Hepatitis A (HepA; Havrix®, Vaqta®)	Moderate: Increased risk of idiopathic thrombocytopenic purpura
Hepatitis B (HepB; Engerix-B®, Recombivax HB®)	Moderate: No evidence of increased risk of multiple sclerosis
Inactivated poliovirus (IPV; IPOL®)	Insufficient evidence to draw conclusions about key adverse events
Influenza, inactivated (IIV; Afluria Quadrivalent®, Fluarix Quadrivalent®, Flulaval Quadrivalent®, Fluzone Quadrivalent®, Flucelvax Quadrivalent®)	Moderate: No evidence of increased risk of death Low: No evidence of increased risk of anaphylaxis or systemic allergic reaction, asthma, autoimmune disease, cardiovascular events, febrile seizures, seizures
Influenza, live attenuated (LAIV; FluMist Quadrivalent®)	Low: No evidence of increased risk of death or seizures
Measles, mumps, and rubella (MMR; M-M-R II®)	High: No evidence of increased risk of autism High: Increased risk of anaphylaxis in children with allergies; increased risk of febrile seizures Moderate: Increased risk of idiopathic thrombocytopenic purpura Low: No evidence of increased risk for asthma

Vaccine (abbreviation; brand name[s])	Synthesis of SoE* and findings for vaccines currently in use in children
Meningococcal, A, C, W, and Y (MenACWY; MenACWY-D [Menactra®], MenACWY-CRM [Menveo®], MenACWY-TT [MenQuadFi®])	Moderate: No evidence of increased risk of cardiovascular events, diabetes, febrile seizures, intussusception, idiopathic thrombocytopenic purpura, Kawasaki disease, seizures Moderate: Increased risk of anaphylaxis in children with allergies Low: No evidence of increased risk of acute disseminated encephalomyelitis, anaphylaxis or systemic allergic reaction, asthma, autoimmune disease, death, encephalitis/encephalopathy, meningitis, multiple sclerosis, reproductive system events, transverse myelitis
Meningococcal B (MenB; MenB-4C [Bexsero®], MenB-FHbp [Trumenba®])	Moderate: No evidence of increased risk of anaphylaxis or systemic allergic reaction, reproductive system events Low: No evidence of increased risk of asthma, death, seizures
Rotavirus (RV; Rotarix®, RotaTeq®)	High: No evidence of increased risk of diabetes Moderate: No evidence of increased risk of intussusception (moderate SoE for increased risk from prior report was not confirmed when combining all available trials, though some observational studies showed increased risk). No evidence of increased risk of asthma, autoimmune disease, death, encephalitis/encephalopathy, febrile seizures, idiopathic thrombocytopenic purpura, seizures, stroke Low: No evidence of increased risk of anaphylaxis or systemic allergic reaction, autoimmune thyroiditis (Hashimoto's disease), Kawasaki disease, meningitis, reproductive system events
Varicella (VAR; Varivax®)	High: Increased risk of anaphylaxis Moderate: Increased risk of idiopathic thrombocytopenic purpura among children aged 11 to 17 years

*Please see Appendix Table 5a for a description of the SoE and findings from the prior 2014 report (including adverse events not examined as key adverse events in the update), the update, and the synthesis across the report and update (including for combination vaccines).

Table 3. Strength of Evidence (SoE) for safety of vaccines in adults

Vaccine (abbreviation; brand name[s])	Synthesis of SoE and findings for vaccines currently in use in adults
9-valent human papillomavirus (HPV9; Gardasil 9®)	Insufficient evidence to draw conclusions; see Table 2 for studies that combined children and adults
13-valent pneumococcal conjugate (PCV13; Prevnar 13®)	Moderate: No evidence of increased risk of cardiovascular events, herpes zoster, myocardial infarction, reproductive system events, stroke Low: No evidence of increased risk of acute disseminated encephalomyelitis, anaphylaxis or systemic allergic reaction, asthma, autoimmune disease, death, encephalitis/encephalopathy, herpes zoster, idiopathic thrombocytopenic purpura, meningitis, seizures
23-valent pneumococcal polysaccharide (PPSV23; Pneumovax®)	High: No evidence of increased risk of cardiovascular or cerebrovascular events in adults aged 65 years and older Moderate: No evidence of increased risk of death
Hepatitis A (HepA; Havrix®, Vaqta®)	Insufficient evidence to draw conclusions about key adverse events

Vaccine (abbreviation; brand name[s])	Synthesis of SoE and findings for vaccines currently in use in adults
Hepatitis B (HepB; Engerix-B®, Recombivax HB®, HEPLISAV-B®)	Moderate: No evidence of increased risk of multiple sclerosis (for hepatitis B vaccines except HEPLISAV-B, for which there was insufficient evidence) Moderate: No evidence of increased risk of diabetes (across all hepatitis B vaccines) Moderate: Increased risk of anaphylaxis in patients allergic to yeast (for hepatitis B vaccines except HEPLISAV-B, for which there were no studies) Low: No evidence of increased risk of asthma, autoimmune disease, cardiovascular events, death, herpes zoster, reproductive system events; stroke for HEPLISAV-B
Influenza, inactivated (IIV; Afluria Quadrivalent®, Flucelvax Quadrivalent®, Fluarix Quadrivalent®, Flulaval Quadrivalent®, Fluzone High Dose Quadrivalent®, Fluzone Quadrivalent®)	Low: No evidence of increased risk of asthma, cardiovascular events, death, myocardial infarction, reproductive system events, seizures, stroke
Influenza, inactivated, adjuvanted (aIIV; Fluad®, Fluad Quadrivalent®)	Moderate: No evidence of increased risk of stroke Low: No evidence of increased risk of asthma, autoimmune disease, cardiovascular events, death, encephalitis/encephalopathy, Guillain-Barré syndrome, idiopathic thrombocytopenic purpura, myocardial infarction, seizures
Influenza, recombinant (RIV; Flublok Quadrivalent®)	Low: No evidence of increased risk of cardiovascular events, death, encephalitis/encephalopathy, myocardial infarction, reproductive system events, stroke
Influenza, live attenuated (LAIV; FluMist Quadrivalent®)	Insufficient evidence to draw conclusions about key adverse events
Measles, mumps, and rubella (MMR; M-M-R II®)	Moderate: No evidence of increased risk of type 1 diabetes mellitus
Meningococcal A, C, W, and Y (MenACWY; MenACWY-D [Menactra®], MenACWY-CRM [Menveo®], MenACWY-TT [MenQuadFi®])	Moderate: No evidence of increased risk of death Low: No evidence of increased risk of cardiovascular events, myocardial infarction, stroke
Meningococcal B (MenB; MenB-4C [Bexsero®],MenB-FHbp [Trumenba®])	Insufficient evidence to draw conclusions about key adverse events; see Table 2 for studies that combined children and adults
Tetanus, diphtheria, and acellular pertussis (Tdap; Adacel®, Boostrix®) and tetanus and diphtheria (Td; TDVAX, Tenivac®)	High: Increased risk of anaphylaxis
Varicella (VAR; Varivax®)	Insufficient evidence to draw conclusions about key adverse events
Zoster recombinant (RZV; Shingrix®)	High: No evidence of increased risk of herpes zoster Moderate: No evidence of increased risk of amyotrophic lateral sclerosis, anaphylaxis or systemic allergic reaction, asthma, cardiovascular events, death, diabetes, encephalitis/encephalopathy, Guillain-Barré syndrome, idiopathic

Vaccine (abbreviation; brand name[s])	Synthesis of SoE and findings for vaccines currently in use in adults
	thrombocytopenic purpura, meningitis, myocardial infarction, reproductive system events, seizures, stroke Low: No evidence of increased risk of acute disseminated encephalomyelitis, angioedema, ataxia, autoimmune disease, autoimmune thyroiditis (Hashimoto's disease)

*Please see Appendix Table 6a for a description of the SoE and findings from the prior 2014 report (including adverse events not examined as key adverse events in the update), the update, and the synthesis across the report and update (including for combination vaccines).

Table 4. Strength of Evidence (SoE) for safety of vaccines in pregnant women

Vaccine (abbreviation; brand name[s])	Synthesis of SOE* and findings for vaccines currently in use in pregnant women
Hepatitis B (HepB; Engerix-B®, Recombivax HB®)	Insufficient evidence to draw conclusions about key adverse events
Influenza, inactivated (IIV; Afluria Quadrivalent®, Flucelvax Quadrivalent®, Fluarix Quadrivalent®, Flulaval Quadrivalent®, Fluzone Quadrivalent®)	Insufficient evidence to draw conclusions about key adverse events
Influenza, recombinant (RIV; Flublok Quadrivalent®)	Insufficient evidence to draw conclusions about key adverse events
Tetanus, diphtheria, and acellular pertussis (Tdap; Adacel®, Boostrix®)	Moderate: No evidence of increased risk of maternal cardiovascular events, maternal death, maternal diabetes, eclampsia/pre-eclampsia, preterm labor, maternal reproductive system events, stillbirth, cardiovascular events in infants, death in infants, encephalitis/encephalopathy in infants, seizures in infants Low: No evidence of increased risk of maternal encephalitis/encephalopathy, autism in infants, birth defects in infants, febrile seizures in infants

*Please see Appendix Table 7a for a description of the SoE and findings from the prior 2014 report, the update, and the synthesis across the report and update.