

UC Irvine

UC Irvine Previously Published Works

Title

Superior Global Cognition in Oldest-Old Is Associated with Resistance to Neurodegenerative Pathologies: Results from The 90+ Study.

Permalink

<https://escholarship.org/uc/item/26r0803m>

Journal

Journal of Alzheimer's Disease, 93(2)

ISSN

1387-2877

Authors

Biswas, Roshni
Kawas, Claudia
Montine, Thomas J
et al.

Publication Date

2023

DOI

10.3233/jad-221062

Peer reviewed

Superior Global Cognition in Oldest-Old Is Associated with Resistance to Neurodegenerative Pathologies: Results from The 90+ Study

Roshni Biswas^{a,*}, Claudia Kawas^{a,b}, Thomas J. Montine^c, Syed A. Bukhari^c, Luohua Jiang^d and Maria M. Corrada^{a,d,*}

^a*Department of Neurology, University of California, Irvine, CA, USA*

^b*Department of Neurobiology and Behavior, University of California, Irvine, CA, USA*

^c*Department of Pathology, Stanford University, Palo Alto, CA, USA*

^d*Department of Epidemiology and Biostatistics, University of California, Irvine, CA, USA*

Handling Associate Editor: Erin Abner

Accepted 8 March 2023

Pre-press 10 April 2023

Abstract.

Background: Some oldest-old individuals can maintain superior cognition despite advanced age. Little is known about the neuropathological changes in the brains of oldest-old superior cognitive performers.

Objective: Our objective was to examine the associations between Alzheimer's disease (AD) and non-AD neuropathologic features in relation to superior cognitive performance in oldest-old individuals.

Methods: We analyzed brain autopsy data from 102 participants with normal cognition from *The 90+ Study*. Superior global cognitive performers (SGCP) were defined as having Mini-Mental State Examination (MMSE) score ≥ 28 in the last visit 12 to 2 months before death. To examine the associations between individual and multiple comorbid neuropathologic features with SGCP status we used multiple logistic regression models adjusting for age, sex, and education.

Results: Alzheimer's disease neuropathological change (ADNC) and low levels of vascular pathologic change were not associated with superior cognition. In contrast, participants with limbic (OR = 8.37; 95% CI: 1.48–47.44) and neocortical (OR = 10.80; 95% CI: 1.03–113.82) Lewy body disease (LBD), or with hippocampal sclerosis (HS) (OR = 5.28; 95% CI: 1.10–25.47) were more likely to be non-SGCP. High total burden of multiple comorbid neuropathologic features was associated with a lower likelihood of being SGCP.

Conclusion: Oldest-old superior cognitive performers were resilient to ADNC and low levels of vascular pathologic change and were resistant to non-AD neurodegenerative changes and multiple comorbid neuropathologic features. Understanding the factors underlying the ability of superior cognitive performers to resist these changes might provide useful insights on maintenance of superior cognition despite advanced age.

Keywords: Alzheimer's disease, cognitive aging, neurodegenerative disease, oldest-old, successful aging, vascular

*Correspondence to: Roshni Biswas, University of California, Irvine, Hewitt Hall Room 1515, Irvine, CA 92697, USA. Tel.: +1 949 824 9176; E-mail: roshni.biswas@uci.edu. and Maria M. Corrada, University of California, Irvine, Hewitt Hall Room 1513, Irvine, CA 92697, USA. Tel.: +1 949 824 9109; E-mail: mcorrada@uci.edu.

INTRODUCTION

Cognitive impairment is considered an almost unavoidable consequence of aging [1, 2], and several autopsy studies have found associations between

cognitive impairment and age-related neuropathologic features in the brain [3–7]. However, few studies have examined neuropathologic features in relation to superior cognitive performance. The Northwestern University SuperAging Study defined “SuperAgers” as individuals having similar memory to younger individuals and found significantly lower frequency of AD neuropathologic change (ADNC), i.e., neurofibrillary tangles and amyloid plaques, in the anterior cingulate cortex of SuperAgers [8–12] when compared to age-matched controls. In another study of 10 SuperAgers, they observed sparse to frequent tangles in the hippocampus and no neocortical tangles in 90% (9/10) of SuperAgers [11]. While there is some information on the association between superior cognition in advanced age and ADNC, not much is known about the associations of superior cognition with non-AD neuropathologic features or with multiple comorbid neuropathologic features. Examining brain autopsy data in oldest-old superior cognitive performers could reveal valuable insights on whether their high level of performance results from cognitive resilience despite neuropathologic changes or from resistance to developing such changes [13, 14]. This could inform prevention strategies targeted towards specific neuropathologic features that reduce the likelihood of superior cognitive performance.

Previous autopsy studies on *The 90+ Study* participants reported that individual and multiple comorbid neuropathologic features were common in oldest-old individuals and were associated with increased likelihood and severity of cognitive impairment [6, 14, 15]. In this study, we explored the superior performance end of the cognitive spectrum using brain autopsy data from *The 90+ Study*—an ongoing, longitudinal, community-based study of aging and dementia on individuals aged 90 years and older, the oldest-old [15, 16]. Our objectives were: 1) to examine the associations between individual AD and non-AD neuropathologic features with superior cognitive performance in the oldest-old, and 2) to examine the associations between multiple comorbid neuropathologic features with superior cognitive performance in the oldest-old.

METHODS

Participants

The initial participants of *The 90+ Study* were survivors of the Leisure World Cohort Study (LWCS) [17], an epidemiologic study on the members of a

retirement community in Orange County, California. *The 90+ Study* commenced in 2003 when LWCS participants aged 90 or older on January 1, 2003, were invited to join. A similar invitation was extended on January 1, 2008, and every year thereafter to those turning 90 years old. More recently, volunteers aged 90 years and older who were residents of Orange County, California, and lived within a two hour drive of the study location, were recruited in the study through an open recruitment [18]. While the LWCS participants were recruited regardless of cognitive diagnosis, the volunteers had no or mild dementia. Currently, approximately two thirds of *The 90+ Study* participants are from the original LWCS and one third are volunteers from open recruitment.

All 1,335 participants (966 from LWCS and 369 volunteers) who agreed to in-person examination were invited to be part of the autopsy program. As of February 2022, 623 participants (308 from LWCS and 315 volunteers) had enrolled in the autopsy program. Of the 623 participants, 481 have died, 452 (94%) came to autopsy, and we have completed the autopsy evaluation and final cognitive diagnosis for 407 participants. Only the 117 participants with normal cognitive diagnosis at the time of death were considered for this analysis (Fig. 1).

Standard protocol approvals, registrations, and patient consents

All participants or their designated informants provided consent to participate in the study and the study was approved by the University of California, Irvine (UCI) Institutional Review Board (IRB).

Neuropathological assessment

The UCI pathology team procured the brain specimens, which were fixed in formalin and then sent to the Department of Pathology at Stanford University for inspection, dissection, and (immuno)histopathologic evaluation according to current consensus criteria while blinded to cognitive diagnosis and any other participant information. The neuropathologic features were scored as follows: 1) ADNC (0 = not 1 = low; 2 = intermediate; 3 = high) was based on the National Institute on Aging-Alzheimer’s Association (NIA-AA) “ABC” score, which incorporates Thal Phase for amyloid- β (A β) plaques, Braak staging for neurofibrillary tangles, and Consortium to Establish a Registry for AD (CERAD) staging for neuritic plaques [19, 20];




Fig. 1. Participant flowchart: *The 90+ Study*. CIND, Cognitive impairment, no dementia; MMSE, Mini-Mental State Examination; non-SGCP, non-Superior Global Cognitive Performers; SGCP, Superior Global Cognitive Performers.

134 2) Cerebral amyloid angiopathy (CAA) (0 = none;
 135 1 = mild; 2 = moderate; 3 = severe) [21]; 3) Microin-
 136 farcts (0 = none; 1 = 1; 2 = 2; 3 = 3+) [19, 20]; 4)
 137 Atherosclerosis (0 = none; 1 = mild when present at
 138 branch points in the circle of Willis; 2 = moderate
 139 when present at branch points and elsewhere in
 140 the circle of Willis; 3 = severe when present on the
 141 cerebral convexity); 5) Arteriolosclerosis (0 = none;
 142 1 = mild for partial replacement of vascular smooth
 143 muscle cells; 2 = moderate for full replacement of
 144 vascular smooth muscle cells; 3 = severe for con-
 145 centric thickening of the sclerotic vessel wall);
 146 6) Lewy body disease (LBD) (0 = none/olfactory;
 147 1 = brainstem-predominant; 2 = limbic [transitional];
 148 3 = neocortical [diffuse]) (2 individuals with LBD
 149 only in the amygdala were included in the 0
 150 LBD category) [22]; 7) Hippocampal sclerosis (HS)

(0 = absent; 1 = present in either right or left or
 151 both) [19, 20]; 8) Limbic-predominant age-related
 152 TDP-43 encephalopathy neuropathological change
 153 (LATE-NC) (0 = none; 1 = amygdala only; 2 = plus
 154 hippocampus; 3 = plus middle frontal gyrus) [23]; 9)
 155 Age-related tau astroglialopathy (ARTAG) (0 = none;
 156 1 = occasional; 2 = numerous) [24].
 157

158 *Assessment of comorbid pathology indices*

159 To examine the comorbid neuropathologic features
 160 at different severity levels, we followed the meth-
 161 ods from a previous study on the Nun Study and the
 162 Honolulu-Asia Aging Study (HAAS) [7] to compute
 163 summary comorbid neuropathology indices. Each
 164 specific neuropathologic feature was given an index
 165 value of 0 for absent/negligible levels, 0.4 for inter-

mediate, or 1 to denote severe levels of pathologic change. The index values for the specific neuropathologic features were then added into an index. As done by the previous study, we chose 0.4 as the intermediate level, to distinguish the influence of a single severe pathology (comorbid pathology index 1.0) from 2 or 3 intermediate pathologies (comorbid pathology indices 0.8 and 1.2, respectively). Specific neuropathologic feature index values were assigned as follows: 1) AD index value was 0 for low/not, 0.4 for intermediate, and 1.0 for high ADNC; 2) CAA index value was 0 for was low, 0.4 for mild/moderate, and 1.0 for severe CAA; 3) Microinfarcts index value was 0 for no, 0.4 for 1-2, and 1.0 for 3 or more microinfarcts; 4) Atherosclerosis index value was 0 for no, 0.4 for mild/moderate, and 1.0 for severe atherosclerosis; 5) Arteriolosclerosis index value was 0 for no, 0.4 for mild, and 1.0 for moderate/severe arteriolosclerosis; 6) LBD index value was 0 for none/olfactory LBD, 0.4 for brainstem-predominant/limbic LBD, and 1.0 for neocortical (diffuse) LBD; 7) HS index value was 0 for absent, 0.4 for unilateral, and 1.0 for bilateral HS; (\8) LATE-NC index value was 0 for no, 0.4 for amygdala, and 1.0 for hippocampal/cortical LATE-NC; (\9) ARTAG index value was 0 for no, 0.4 for occasional, and 1.0 for numerous ARTAG. Additionally, to derive binary severe vascular index values, all vascular neuropathologies, i.e., CAA, microinfarcts, atherosclerosis, and arteriosclerosis, were further recoded as- absent (=0) when the specific vascular pathology index value were 0 or 0.4, and present (=1) when the index value was 1.0.

Four different comorbid neuropathology indices were calculated as follows: 1) Total neuropathology index was the sum of all (i.e., ADNC, CAA, microinfarcts, atherosclerosis, arteriosclerosis, LBD, HS, LATE-NC, and ARTAG) neuropathology index values; 2) Vascular index was the sum of CAA, microinfarcts, atherosclerosis, and arteriosclerosis index values; 3) Severe vascular index was the sum of severe vascular index values for CAA, microinfarcts, atherosclerosis, and arteriosclerosis; 4) Neurodegenerative index was the sum of index values for LBD, HS, LATE-NC, and ARTAG.

Cognitive diagnosis

Full clinical evaluations of participants carried out every six months consisted of a neurologic examination, physical examination, full neuropsychological battery [25], review of medical history, examination of medication containers, and interviews with infor-

ants. A final cognitive diagnosis was assigned after death in a multidisciplinary consensus conference using all available information from the longitudinal evaluations, brain imaging when clinically available, and medical records. All cognitive diagnosis assignments were done blinded to pathologic evaluation results. Participants were diagnosed as having normal cognition, cognitive impairment but no dementia (CIND), or dementia. A dementia diagnosis was made according to *Diagnostic and Statistical Manual of Mental Disorders* 4th Edition diagnostic criteria [26]. CIND was assigned when cognitive or functional impairments were present but did not meet criteria for dementia.

Superior global cognitive performer definition

The neuropsychological battery included the Mini-Mental State Examination (MMSE) to assess global cognition [27]. For this work, we assessed maintenance of superior global cognition until the time of death. Superior global cognitive performers (SGCP) were defined as individuals who had: 1) normal diagnosis in the consensus conference and, 2) an MMSE score of 28 or above in the last visit 12 to 2 months before death. Non- superior global cognitive performers (non-SGCP) were individuals who also had normal diagnosis in the consensus conference but had an MMSE score below 28. This cut-off was based on a previous study on *The 90+ Study* participants reporting a mean MMSE score of 28 in cognitively normal participants [28]. To limit the possibility of classification error due to poor cognitive performance resulting from terminal illness, we did not consider MMSE scores in the last 2 months of life.

Statistical analysis

We describe participant characteristics and proportion of participants overall and by SGCP status. We calculated the different comorbid indices, examined their distribution by SGCP status, and then categorized the indices into smaller groups according to their distribution, such that each category had sufficient number of SGCP and non-SGCP participants. The total neuropathology index was categorized as <1.8 (reference group), 1.8-<3, 3-<4, and 4+. We did not categorize the vascular index as there was no clear relationship pattern with SGCP status. The severe vascular index was categorized as 0 (reference group), 1, 2, and 3. The neurodegenerative index was categorized as 0 (reference group), 0.4-<1, and 1+. To

Table 1
 Characteristics of autopsied cognitively normal participants: *The 90+ Study*

Characteristics	All participants (N = 102)	Superior global cognitive performers (SGCP) (N = 71)	Non-Superior global cognitive performers (Non SGCP) (N = 31)
Sex			
Male	41 (40.2%)	27 (38.0%)	14 (45.2%)
Female	61 (59.8%)	44 (62.0%)	17 (54.8%)
Age of death			
Mean (SD)	97.6 (3.3)	97.4 (3.4)	98.0 (2.9)
Age of visit for MMSE			
Mean (SD)	97.1 (3.3)	96.9 (3.4)	97.5 (2.9)
Education			
Less than college	44 (43.1%)	31 (43.7%)	13 (41.9%)
College or higher	58 (56.9%)	40 (56.3%)	18 (58.1%)
Leisure World Cohort participant			
No	51 (50.0%)	34 (47.9%)	17 (54.8%)
Yes	51 (50.0%)	37 (52.1%)	14 (45.2%)
Brain weight (grams)			
Mean (SD)	1,160.9 (110.1)	1,161.4 (111.7)	1,159.7 (108.4)
N (Missing)	96 (6)	65 (6)	31 (0)

MMSE, Mini-Mental State Examination.

examine the associations with individual and multiple comorbid neuropathologic features, we calculated odds ratios (OR) and 95% confidence intervals (CI) using multiple logistic regression models adjusting for age at visit (last visit when MMSE was done), sex, and college education (yes/no). For all analyses, our outcome of interest was SGCP status, and we modeled the probability of being a non-SGCP. Therefore, in our results, an OR above 1 would indicate a higher likelihood of being a non-SGCP which also indicates a lower likelihood of being a SGCP. Conversely, an OR below 1 would indicate a lower likelihood of being a non-SGCP which also indicates a higher likelihood of being a SGCP.

Secondary analyses

To assess the robustness of our analysis, we did additional secondary analyses. First, since some of the neuropathologic features had low prevalence in our study population, which could lead to sparse data bias, we conducted exact logistic regressions for all neuropathologic features in relation to superior cognition. Second, we tried two alternative MMSE cut-off points (27 and 29) to define superior cognitive performers. To examine the associations of superior cognition using these alternative cut-offs with individual neuropathologic features, we repeated the multiple logistic regression analyses adjusting for age at visit, sex, and college education to calculate OR and 95% CI. All analyses were performed using SAS 9.4 (SAS Institute; Cary, NC, US).

RESULTS

Out of 117 autopsied participants with normal cognitive diagnosis, 102 had at least one MMSE score between 12 to 2 months before death (Fig. 1). Overall, the average age of our study participants at the time of their last visit was 97.1 years. Most participants were women (59.8%) and college-educated (56.9%). Among these participants, 71 (69.6%) were classified as SGCP while 31 (30.4%) were non-SGCPs. There was no significant difference in age, sex, education, LWC participation, and brain weight between SGCPs and non-SGCPs (Table 1).

Distribution of individual and multiple comorbid pathologies by SGCP status

The distribution of neuropathologic features by SGCP status are shown in Table 2. ADNC was not different between SGCP and non-SGCP participants with most participants in both groups having low and intermediate ADNC (66.2% of SGCPs versus 61.3% of non-SGCPs). When separately examining the distribution of the A score, Braak tangle stage, and C scores we found no difference in the SGCP and non-SGCP. Among vascular neuropathologic features, CAA was observed in about half of the participants in both groups. Severe CAA was less common in SGCPs (5.6%) compared to non-SGCPs (16.1%). Microinfarcts were slightly more common in SGCPs (22.5%) compared to non-SGCPs (19.4%). Mild or moderate atherosclerosis was present in more

Table 2
Associations between individual neuropathological features and superior global cognitive performance in the oldest-old

Pathologies	All Participants (N = 102)	Superior global cognitive performers (SGCP) (N = 71)	Non-Superior global cognitive performers (Non SGCP) (N = 31)	OR (95% CI) ^a
AD neuropathologic change (ADNC)				
Not	13 (12.7%)	9 (12.7%)	4 (12.9%)	1.00 (Reference)
Low	28 (27.5%)	17 (23.9%)	11 (35.5%)	1.24 (0.29–5.28)
Intermediate	38 (37.3%)	30 (42.3%)	8 (25.8%)	0.53 (0.12–2.25)
High	23 (22.5%)	15 (21.1%)	8 (25.8%)	1.09 (0.23–5.06)
NIA-AA A Score (Thal Phase for Aβ plaques)				
Phase 0	13 (12.7%)	9 (12.7%)	4 (12.9%)	1.00 (Reference)
Phase 1 or 2	30 (29.4%)	20 (28.2%)	10 (32.3%)	0.90 (0.20–3.92)
Phase 3	18 (17.6%)	14 (19.7%)	4 (12.9%)	0.59 (0.11–3.09)
Phase 4 or 5	41 (40.2%)	28 (39.4%)	13 (41.9%)	0.92 (0.23–3.68)
Braak Tangle Stage				
0,1,2	8 (7.8%)	5 (7.0%)	3 (9.7%)	1.00 (Reference)
3	20 (19.6%)	14 (19.7%)	6 (19.4%)	0.64 (0.11–3.69)
4	49 (48.0%)	35 (49.3%)	14 (45.2%)	0.66 (0.14–3.20)
5	24 (23.5%)	16 (22.5%)	8 (25.8%)	0.79 (0.14–4.44)
6	1 (1.0%)	1 (1.4%)	0 (0.0%)	–
NIA-AA C Score (CERAD neuritic plaques)				
None	32 (31.4%)	21 (29.6%)	11 (35.5%)	1.00 (Reference)
Sparse	14 (13.7%)	10 (14.1%)	4 (12.9%)	0.70 (0.17–2.81)
Moderate	10 (9.8%)	8 (11.3%)	2 (6.5%)	0.41 (0.07–2.39)
Frequent	46 (45.1%)	32 (45.1%)	14 (45.2%)	0.83 (0.31–2.26)
Cerebral Amyloid Angiopathy (CAA)				
None	49 (48.0%)	33 (46.5%)	16 (51.6%)	1.00 (Reference)
Mild	10 (9.8%)	7 (9.9%)	3 (9.7%)	0.89 (0.20–4.08)
Moderate	34 (33.3%)	27 (38.0%)	7 (22.6%)	0.44 (0.15–1.31)
Severe	9 (8.8%)	4 (5.6%)	5 (16.1%)	2.77 (0.62–12.36)
Microinfarcts				
0	80 (78.4%)	55 (77.5%)	25 (80.6%)	1.00 (Reference)
1	18 (17.6%)	14 (19.7%)	4 (12.9%)	0.59 (0.17–2.02)
2	3 (2.9%)	1 (1.4%)	2 (6.5%)	2.28 (0.30–17.52) ^b
3+	1 (1.0%)	1 (1.4%)	0 (0%)	–
Atherosclerosis				
None	24 (24.5%)	18 (26.5%)	6 (20.0%)	1.00 (Reference)
Mild	45 (45.9%)	32 (47.1%)	13 (43.3%)	1.21 (0.38–3.86)
Moderate	26 (26.5%)	17 (25.0%)	9 (30.0%)	1.71 (0.49–5.96)
Severe	3 (3.1%)	1 (1.5%)	2 (6.7%)	5.72 (0.42–78.78)
Arteriolosclerosis				
None	12 (11.8%)	9 (12.7%)	3 (9.7%)	1.00 (Reference)
Mild	30 (29.4%)	22 (31.0%)	8 (25.8%)	1.04 (0.22–5.06)
Moderate	60 (58.8%)	40 (56.3%)	20 (64.5%)	1.37 (0.31–6.02)
Severe	0 (0%)	0 (0%)	0 (0%)	NA
Lewy Body Disease (LBD)				
None	84 (82.4%)	64 (90.1%)	20 (64.5%)	1.00 (Reference)
Brainstem	7 (6.9%)	4 (5.6%)	3 (9.7%)	2.36 (0.47–11.73)
Limbic	7 (6.9%)	2 (2.8%)	5 (16.1%)	8.37 (1.47–47.44)
Neocortical	4 (3.9%)	1 (1.4%)	3 (9.7%)	10.80 (1.03–113.82)
Hippocampal sclerosis (HS)				
Absent	94 (92.2%)	68 (95.8%)	26 (83.9%)	1.00 (Reference)
Present	8 (7.8%)	3 (4.2%)	5 (16.1%)	5.28 (1.10–25.47)
LATE-NC				
None	78 (76.5%)	57 (80.3%)	21 (67.7%)	1.00 (Reference)
Amygdala	7 (6.9%)	5 (7.0%)	2 (6.5%)	1.26 (0.21–7.40)
Hippocampus	15 (14.7%)	9 (12.7%)	6 (19.4%)	2.49 (0.83–7.46) ^c
Cortex	2 (2.0%)	0 (0%)	2 (6.5%)	–

(Continued)

Table 2
(Continued)

Pathologies	All Participants (N = 102)	Superior global cognitive performers (SGCP) (N = 71)	Non-Superior global cognitive performers (Non SGCP) (N = 31)	OR (95% CI) ^a
ARTAG				
None	40 (39.2%)	29 (40.8%)	11 (35.5%)	1.00 (Reference)
Occasional	54 (52.9%)	38 (53.5%)	16 (51.6%)	1.05 (0.42–2.64)
Numerous	8 (7.8%)	4 (5.6%)	4 (12.9%)	2.22 (0.44–11.10)

^aMultiple logistic regression adjusting for age at visit for last MMSE, sex, and education used to calculate Odds ratio (OR) and 95% Confidence Interval (CI). Values in bold are statistically significant. ^bOR calculated for 2+ microinfarcts. 1 SGCP with 3+ microinfarct was included in this analysis. ^cOR calculated for hippocampal/cortical LATE-NC. 2 persons non-SGCPs with LATE-NC in cortex were included in this analysis. AD, Alzheimer’s disease; ARTAG, Age-related tau astrogliaopathy; LATE-NC, Limbic-predominant age-related TDP-43 encephalopathy neuropathological change; NIA-AA, National Institute on Aging-Alzheimer’s Association.




Fig. 2. Distribution of the comorbid neuropathology index scores by superior global cognitive performance in the oldest-old. Numbers inside bars denote frequency.

322 than 70% of participants in both the SGCPs and non-
 323 SGCPs while mild or moderate arteriolosclerosis was
 324 present in above 85% participants in both groups.
 325 The severe vascular pathologic changes were infre-
 326 quent in both groups and none of the participants
 327 in either group had severe arteriolosclerosis. The
 328 non-AD neurodegenerative neuropathologic features
 329 were less frequent in SGCPs than in non-SGCPs. HS
 330 was observed in 4.2% of SGCPs compared to 16.1%
 331 of non-SGCPs. LBD was observed in 9.9% of SGCPs
 332 compared to 35.5% of non-SGCPs. LATE-NC was
 333 observed in 19.3% of SGCPs compared to 32.3%
 334 non-SGCPs. HS and LATE-NC co-existed in 4 par-

335 ticipants, all of whom were non-SGCPs (not shown
 336 in table). Occasional ARTAG was observed in half
 337 of the participants in both groups, while numerous
 338 ARTAG were observed in 5.6% of SGCPs versus
 339 12.9% non-SGCPs (Table 2).

340 To examine comorbid pathology indices by SGCP
 341 status, we first evaluated their distribution (Fig. 2).
 342 The total neuropathology index scores ranged from 0
 343 to 4.8 with SGCPs having lower index scores while
 344 non-SGCPs tended to be on the higher index score
 345 range, as shown by the red stacked bars in Fig. 2A.
 346 Overall, most participants had scores between 1.8 to
 347 2.8 (Fig. 2A). The vascular index scores ranged from

348 0 to 2.4 but did not show a clear pattern of distribution
 349 based on SGCP status (Fig. 2B). The severe vascular
 350 index scores ranged from 0 to 3 with SGCPs having
 351 lower scores (Fig. 2C). The neurodegenerative index
 352 scores ranged from 0 to 2.4 with most SGCPs having
 353 lower scores. However, the only participant with a
 354 score of 2.4 in the neurodegenerative pathology index
 355 was a person categorized as SGCP (Fig. 2D).

356 Association of ADNC and total neuropathology 357 index with superior cognitive performance

358 We used logistic regression models adjusting for
 359 age, sex, and education to examine the associations
 360 of ADNC, A score, Braak tangle stage, and C score,
 361 as well as total pathology index with SGCP status.
 362 There was no association between ADNC severity
 363 level and SGCP status. The ORs ranged from 0.53
 364 to 1.24, and there was no clear pattern of associa-
 365 tion with different levels of ADNC severity (Table 2).
 366 Similarly, there was no association between the indi-
 367 vidual ADNC pathologies (A score, Braak stage and
 368 C score) with SGCP status (Table 2).

369 There was a significantly higher likelihood of being
 370 a non-SGCP with increasing total neuropathology
 371 index score. Compared to the reference category
 372 (0-<1.8 score), for the total neuropathology index cat-
 373 egory 1.8-<3 the OR was 5.87 (95% CI:1.23,28.05),
 374 for 3-<4 category the OR was 3.75 (95% CI: 0.62,
 375 22.82), and for 4+ category the OR was 18.31 (95%
 376 CI: 2.34,143.19) (Fig. 3A).

377 Association between individual and multiple 378 vascular neuropathologic features and superior 379 cognitive performance

380 Logistic regression models adjusting for age,
 381 sex, and education were also used to examine the
 382 associations of individual and comorbid vascular neu-
 383 ropathologic features with SGCP status. There were
 384 trends that participants with severe vascular patho-
 385 logic changes were more likely to be non-SGCP
 386 (Table 2). For example, the ORs for severe CAA com-
 387 pared to no CAA was 2.77 (95% CI:0.62–12.36),
 388 for 2+ microinfarcts compared to no microinfarcts
 389 2.28 (95%CI:0.30–17.52), for severe atherosclerosis
 390 compared to no atherosclerosis was 5.72 (95%CI:
 391 0.41–78.78) and no participant had severe arterio-
 392 sclerosis. However, none of these ORs reached
 393 statistical significance.

394 Unlike the vascular index, which showed no clear
 395 relationship pattern to SGCP status, for the severe




Fig. 3. Association between comorbid neuropathology indices and superior global cognitive performance in the oldest-old. A) Association with total neuropathology (ADNC, CAA, microinfarcts, atherosclerosis, arteriosclerosis, LBD, HS, LATE-NC, and ARTAG) index scores. B) Association with severe vascular (CAA, microinfarcts, atherosclerosis, and arteriosclerosis) index scores. C) Association with neurodegenerative (LBD, HS, LATE-NC, and ARTAG) index scores. ADNC, Alzheimer's disease neuropathological change; ARTAG, Age-related tau astroglial pathology; CAA, cerebral amyloid angiopathy; HS, hippocampal sclerosis; LATE-NC, limbic-predominant age-related TDP-43 encephalopathy neuropathological change; LBD, Lewy body disease; Non-SGCP, Non-Superior Global Cognitive Performers; SGCP, Superior Global Cognitive Performers.

396 vascular index, increasing scores tend to be associ-
 397 ated with increased likelihood of being a non-SGCP,
 398 although the ORs were not statistically significant
 399 (Fig. 3B).

400 Association between individual and multiple 401 neurodegenerative neuropathologic features and 402 superior cognitive performance

403 As above, logistic regression models adjusting
 404 for age, sex, and education were used to examine

405 the associations of individual and comorbid neu- 455
406 rodegenerative neuropathologic features with SGCP 456
407 status. Compared to participants with no LBD, those 457
408 having LBD were more likely to be non-SGCP, 458
409 and the association was significant for limbic LBD 459
410 (OR = 8.36; 95%CI: 1.47–47.44) and neocortical LBD 460
411 (OR = 10.80; 95%CI: 1.02–113.82). Participants with 461
412 HS were more likely to be non-SGCP compared 462
413 to those without (OR = 5.28; 95%CI: 1.10–25.47). 463
414 Compared to participants with no LATE-NC, for all 464
415 levels of LATE-NC there was a non-significant but 465
416 increased likelihood of being a non-SGCP. Compared 466
417 to participants without ARTAG, those with numer- 467
418 ous ARTAG had increased, although non-significant, 468
419 likelihood of being a non-SGCP (Table 2). 469

420 There was a significantly higher likelihood of being 470
421 a non-SGCP with increasing neurodegenerative index 471
422 score, with OR = 6.93 (95%CI = 1.66–28.83) in par- 472
423 ticipants with 1+ scores compared to participants with 473
424 score 0 (Fig. 3C). 474

425 *Secondary analyses*

426 Given the sparse data for some of the neuropathologic 475
427 features, we conducted exact logistic regression 476
428 to confirm the robustness of our findings. For most 477
429 neuropathologic features, the magnitude of the unad- 478
430 justed ORs from the exact logistic regression analyses 479
431 were very similar to the maximum likelihood esti- 480
432 mates (MLE) of ORs from unadjusted and adjusted 481
433 logistic regressions (Supplementary Table 1). The 482
434 only exceptions were neocortical LBD and HS pres- 483
435 ence, for which the 95% CI were wider and the ORs 484
436 slightly attenuated and thus non-significant (Supple- 485
437 mentary Table 1). 486

438 We also conducted secondary analyses chang- 487
439 ing the MMSE cut-off for superior performance 488
440 to 27 and 29 and examined the associations with 489
441 neuropathologic features. With the lower cut-off 490
442 of 27, more participants (87 out of 102) were 491
443 categorized as superior performers and fewer (15 492
444 participants) were non-superior performers (Supple- 493
445 mentary Table 2). The results for the association 494
446 of individual neuropathologic features with supe- 495
447 rior cognition remained mostly unchanged with the 496
448 following few differences (Supplementary Table 2). 497
449 For ADNC, the ORs increased in magnitude rang- 498
450 ing from 0.9 to 6.6 (versus 0.53 to 1.24 with the 499
451 original cut-off of 28) but remained non-significant 500
452 with wider 95% CIs and no clear pattern of asso- 501
453 ciation. All levels of CAA were associated with 502
454 non-significant increased likelihood of being a non-

superior performer. For limbic LBD the OR was 455
attenuated (OR = 3.62) and not associated with being 456
non-superior performer. The OR for HS was higher 457
(OR = 7.08) and remained significantly associated 458
with being non-superior performer (Supplementary 459
Table 2). 460

461 With the higher MMSE cut-off of 29, fewer indi- 462
463 viduals (49 out of 102) were superior cognitive 464
465 performers and more (53 participants) non-superior 466
467 performers. Even with this cut-off, the results for the 468
469 associations continued to remain mostly unchanged 470
471 with the following few differences. Mild (OR = 0.12) 472
473 and moderate (OR = 0.17) CAA were associated with 474
475 significantly decreased odds of being a non-superior 476
477 performer, whereas with the original cut-off of 28 478
479 these associations were non-significant. For limbic 480
481 (OR = 5.25) and neocortical (OR = 3.11) LBD, the 482
483 ORs were attenuated and not significantly associ- 484
485 ated with being non-superior performer. For HS, 486
487 the OR was attenuated (OR = 2.27) and not signif- 488
489 icantly associated with increased odds of being a 489
490 non-superior performer (Supplementary Table 3). 491

492 Switching the MMSE cut-off values to lower and 493
494 higher levels resulted in attenuation of some of the 494
495 ORs, but the direction of association remained con- 495
496 sistent. The secondary analyses did not have much of 496
497 an impact on our overall results and conclusions and 497
498 suggest the robustness of our findings. 498

483 *Comparison of autopsied and non-autopsied* 484 484 *participants*

485 A substantial proportion of *The 90+ Study* par- 486
487 ticipants invited to join the autopsy program did 487
488 not consent to autopsy. Therefore, we compared the 488
489 autopsied and non-autopsied individuals to explore 489
490 differences between the two groups. Compared to 490
491 those who did not consent to autopsy, the autopsy 491
492 group included fewer women (66.9% versus 74.3%), 492
493 were slightly younger (93.0 versus 93.8 years), had 493
494 higher education (51.9% college or higher versus 494
495 38.1%), and fewer were LWC participants (49.4% 495
versus 92.4%) (Supplementary Table 4). 496

496 **DISCUSSION**

497 Our study aimed to evaluate the neuropathologi- 497
498 cal findings in superior global cognitive performers 498
499 that distinguish them from non-superior performing 499
500 peers using autopsy data from 102 cognitively nor- 500
501 mal participants of *The 90+ Study*. Neither ADNC 501
502 nor vascular neuropathologic features were associ-

ated with superior cognitive performance. However, among the few cognitively normal participants with severe vascular pathologic changes, we observed a suggestion of higher likelihood of not having superior cognition, i.e., lower likelihood of superior cognition. Participants with LBD and HS were more likely to be non-SGCP, i.e., less likely to have superior cognition. There was a trend that presence of LATE-NC was related to higher likelihood of being non-SGCP, i.e., a lower likelihood of superior cognitive performance, although the results were not statistically significant. Presence of higher levels of multiple comorbid neuropathologic features (as indicated by the comorbid neuropathology indices) were associated with increased likelihood of being non-SGCP, i.e., a lower likelihood of superior cognitive performance.

ADNC was very common, found in 87% of our study participants but was not associated with superior cognitive performance. This finding is expected given that ADNC is almost ubiquitous in this oldest-old cohort [14], and has been commonly reported in individuals without dementia in a recent meta-analysis on 17 population based autopsy studies with age of death 65 years and above [29]. Furthermore, a previous analysis on eight participants from *The 90+ Study* reported the presence of plaques and tangles—the hallmarks of ADNC—in the brains of “SuperAgers” [30]. However, previous autopsy studies in the Northwestern University SuperAging Study cohort reported that compared to age-matched controls, fewer neurofibrillary tangles and amyloid plaques were observed in the anterior cingulate and entorhinal cortex of “SuperAgers”, defined as 80 and older individuals having similar memory to participants 20 to 30 years younger [8, 12]. Methodological differences including difference in superior cognitive performer definition, brain regions evaluated, and larger sample size could all have contributed to the incongruence with our study results.

We also found that maintenance of superior cognitive performance in this oldest-old cohort was associated with lower levels of comorbid neuropathologic features in the brain. While no study has analyzed multiple comorbid neuropathologic features in cognitively superior individuals, previous reports from *The 90+ Study* [6, 31] and other cohorts [7, 32, 33] have shown the association between mixed neuropathologic features and increased likelihood and severity of cognitive impairment. Results from the Nun Study and HAAS reported that comorbid neuropathologic features were common and the cumulative burden was the most relevant determinant

of cognitive impairment [7]. A large study on clinic-based population from the NIH-funded Alzheimer Disease Research Centers on individuals with and without dementia with mean age 81 years reported that mixed neuropathologic features were common and high comorbid neuropathology was associated with lower MMSE scores [5]. Results from the Nun Study and HAAS also reported that cognitive resilience, despite high levels of neurofibrillary tangles, was strongly associated with minimal comorbid neuropathologic features [7]. Another study from the Religious Orders Study (ROS), and the Rush Memory and Aging Project (MAP) on older adults with a mean age of 87 years reported comorbid ADNC, vascular and non-AD neurodegenerative neuropathologic features as well as their increasing severity were related to faster rate of cognitive decline in individuals without dementia [32, 34]. Results from Adult Changes in Thought (ACT) study comparing vascular neuropathologic features in ADNC resistant groups to their age-matched AD dementia groups reported that resistant individuals had less arteriolosclerosis, microinfarcts and CAA, and maintained high cognitive scores over time [35]. Given the commonality of ADNC in our study participants, it is possible that the added presence of vascular and neurodegenerative neuropathologic features lead to worsening of cognitive performance in cognitively normal oldest-old.

Among vascular neuropathologic features, CAA and microinfarcts were less common while lower levels of atherosclerosis and arteriolosclerosis were relatively common in all participants. However, very few cognitively normal participants had severe levels of any vascular neuropathology, and none had severe arteriolosclerosis. No previous autopsy study has explored vascular neuropathologic features in relation to superior cognitive performance. However, vascular neuropathologies have reportedly been associated with cognitive impairment. CAA has been associated with an increased likelihood of dementia and a faster rate of global cognitive decline [36]. Previous studies in our cohort [4] as well as other cohorts [37–40] reported that microinfarcts independently and in presence of ADNC were associated with increased odds of dementia and cognitive impairment. A previous autopsy study on Brazilian population aged >50 years found an association between high-grade atherosclerosis and dementia [41]. A study on participants of the Baltimore Longitudinal Study of Aging with mean age 87.6 years observed high prevalence of atherosclerosis and found it to be an independent risk factor for dementia

607 [42]. In our study, there were trends that partici- 659
608 pants with severe vascular pathologic changes were 660
609 less likely to be superior cognitive performers, but 661
610 none of the associations were statistically significant. 662
611 Given the high mortality in the presence of vascular 663
612 disease, it is possible that individuals with severe vas- 664
613 cular pathologic changes do not survive to become 665
614 oldest-old [43–45]. As a result, there could be low 666
615 prevalence of severe vascular pathologies in cogni- 667
616 tively normal participants and consequently, limiting 668
617 the power of our data to determine whether absence of 669
618 severe vascular pathologic changes is indeed related 670
619 to maintaining superior cognition. 671

620 With multiple comorbid severe vascular pathologic 672
621 changes as determined by the severe vascular index, 673
622 we observed a trend showing decreased likelihood 674
623 of superior cognitive performance. A previous study 675
624 in the ROS-MAP participants reported mixed vascu- 676
625 lar disease associated with faster cognitive decline, 677
626 especially in the presence of atherosclerosis and 678
627 arteriolosclerosis [46]. Our findings are consistent 679
628 with available literature and allude to severe vascular 680
629 pathologic changes independently and in combina- 681
630 tion having a more deleterious effect on cognition, 682
631 while at lower levels their effect is not as prominent. 683
632 However, given the small sample size of individuals 684
633 with severe vascular pathologic changes, the inter- 685
634 pretation of our results warrants caution. 686

635 Among neurodegenerative neuropathologic fea- 687
636 tures, LBD and HS were rare in these cognitively 688
637 normal oldest old and were associated with a lower 689
638 likelihood of superior cognition. About 20% of our 690
639 participants had LATE-NC and showed trends of 691
640 association with a lower likelihood of superior cog- 692
641 nition. Despite the comorbidity with HS [47], only 693
642 4 of our 102 participants had HS with LATE-NC 694
643 and none had superior cognition. In the autopsy study 695
644 on Northwestern SuperAging study participants, the 696
645 authors found a single case of HS and 2 cases of 697
646 LBD out of 10 SuperAger cases studied [11]. Our 698
647 results are consistent in reporting a low prevalence of 699
648 these neuropathologic features in superior cognitive 700
649 performers. 701

650 We further observed that high levels of total 702
651 burden of non-AD neurodegenerative neuropatholo- 703
652 gic features was less likely in superior cognitive 704
653 performers. Despite limited evidence on the associ- 705
654 ation with superior cognition, in published literature 706
655 there is a known association of both individual and 707
656 comorbid non-AD neurodegenerative neuropatholo- 708
657 gies with global cognitive impairment and dementia 709
658 [39, 48–50]. Previous studies have reported that LBD

independently and in presence of ADNC were asso- 659
ciated with increased odds of dementia [5, 39] and 660
cognitive decline [34]. Studies from our cohort [14] 661
and other cohorts [23, 48, 49] have reported associa- 662
tion of independent and comorbid LATE-NC and HS 663
with global cognitive impairment. Our findings add 664
to the evidence by reporting an association between 665
superior cognitive performance in oldest old and 666
absence of independent and comorbid non-AD neu- 667
rodegenerative neuropathologic features. 668

669 Resistance is inferred when observed levels of 670
dementia-related neuropathologic features is lower 671
than expected based on individual characteristics 672
while cognitive resilience is inferred from a higher 673
level of cognitive functioning than expected despite 674
high neuropathologic burden [13, 14]. Our results 675
indicate that superior cognitive performance in 676
oldest-old can be explained by concepts of both 677
resilience and resistance. In cognitively normal 678
oldest-old, the superior global cognitive perform- 679
ers were resilient to ADNC, i.e., they maintained 680
superior cognition despite having ADNC. On the 681
other hand, other neurodegenerative neuropathologic 682
features, particularly LBD and HS, were very rare 683
indicating resistance, i.e., their ability to escape the 684
development of these changes in the brains. For 685
vascular neuropathologic features, the results were 686
mixed showing resilience at lower levels and poten- 687
tial resistance to higher levels of vascular pathologic 688
change needed to maintain superior cognitive perfor- 689
mance. 690

691 One of the major strengths of our study is the 692
sample size given the advanced age of our popula- 693
tion. Our results are derived by analyzing autopsy 694
data from 102 cognitively normal individuals who 695
died at a mean age of 97 years. Second, we used 696
cognitive test scores between 12 to 2 months before 697
death. In published literature there is no consensus 698
definition for superior cognitive performance. Most 699
studies define superior cognitive performance at a 700
specific age range relative to either age-matched peers 701
or younger cohorts. However, individuals who are 702
superior performers at a particular age might decline 703
progressively over time and this decline would be 704
faster in oldest-old. By relying on MMSE perfor- 705
mance in the last year of life, we could ensure that 706
the participants identified as SGCPs maintained high 707
performance throughout life. By excluding test scores 708
from the last two months before death, we could 709
avoid misclassifying a SGCP who had lower scores 710
due to terminal illness. Third, unlike other studies 711
that identify superior performers in memory, we used 712

MMSE which is a measure of global cognition. This is important as superior memory performance does not always indicate superior performance in other domains [51, 52]. By evaluating global cognition, we aimed to identify individuals with superior performance in memory as well as other cognitive domains.

Our study had some limitations. First, despite the relatively large sample size, very few participants had severe levels of vascular and neurodegenerative neuropathologic features which could lead to sparse data bias. One hallmark sign of sparse data bias in multivariate analysis is that, as more variables are added to the regression model the coefficient estimates get further away from the null [53]. However, this was not the case in our study as the adjusted and unadjusted ORs were very similar. In addition, the ORs from the exact logistic regression, which can be used to address sparse data bias when sample sizes are small, showed very similar associations to the ORs from MLE logistic regressions suggesting the robustness of our findings. Nonetheless, we should be cautious in drawing inference on the associations of vascular and neurodegenerative neuropathologic features with superior cognition, and future work should examine these associations in other datasets of oldest-old individuals. Second, MMSE test scores are related to education level and our participants were highly educated [54, 55]. We found about 70% cognitively normal participants were SGCPs. It is possible that in some cases the high MMSE scores are due to better test taking abilities and not necessarily superior cognition levels. Nonetheless, since all participants were cognitively normal at the time of death, we expect most participants to have relatively high MMSE scores. Moreover, in our secondary analyses we examined all associations using alternative lower and higher cut-off values of MMSE, and yet got very similar results. These additional findings suggest that our findings are robust. Third, *The 90+ Study* participants are mostly White college-educated women. However, given that among individuals aged 85 and older there are two females for every male, and around 81% are Non-Hispanic White [56], our results are somewhat generalizable to oldest-old US population in terms of sex and race distribution. Nonetheless, we acknowledge that the lack of ethnoracial and educational diversity in our cohort precludes us from generalizing the findings from this study to a more diverse population. Fourth, since we did not examine cognitive trajectories and only assessed cross-sectional superior cognition at the time of death, we did not test preserved cognition over time. As a result, indi-

viduals starting with MMSE scores below the cut-off and maintaining the same over time would not qualify as SGCP despite no cognitive decline. Future work should analyze individuals with stable cognitive trajectories over time and examine the association with neuropathologic features. Last, *The 90+ Study* participants who consented to the autopsy study had significant demographic differences from the non-autopsied participants. This could make our findings less reflective of the general population.

To our knowledge, this is the first study that examines associations between individual AD and non-AD neuropathologic features as well as multiple comorbid neuropathologic features with superior cognitive performance. We report new evidence about the neuropathologic features in the brains of oldest-old individuals who maintain superior cognition till the time of death. Our findings suggests that while AD and low levels of vascular pathologic changes are common in oldest old superior cognitive performers, they remain resistant to severe vascular and non-AD neurodegenerative changes, in particular, LBD and HS. They also have lesser total burden of comorbid neuropathologic features. Therefore, preventive measures targeted towards increasing resistance and resilience to non-AD comorbid neuropathologies in the brain might be useful in maintenance of superior cognition despite advanced age.

ACKNOWLEDGMENTS

We thank the participants and staff of *The 90+ Study*.

FUNDING

This research was funded by grants from the NIH (R01AG21055, UF1 AG057707, and P30AG066519). The study sponsors had no role in the study design; in the collection, analysis, and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

CONFLICT OF INTEREST

Roshni Biswas, Claudia Kawas, Syed A Bukhari, and Maria Corrada declare no conflicts of interest.

From 2019-2022, Luohua Jiang served as a paid advisor on the Scientific Advisory Board of Birkeland Current, a small business that was funded by National Institute on Aging to develop a medical device that

monitors impacts of cognitive decline on Activities of Daily Living.

Thomas Montine receives royalties from Up To Date.

DATA AVAILABILITY

The data supporting the findings of this study are available from the corresponding author upon reasonable request.

SUPPLEMENTARY MATERIAL

The supplementary material is available in the electronic version of this article: <https://dx.doi.org/10.3233/JAD-221062>.

REFERENCES

- [1] Lipnicki DM, Crawford JD, Dutta R, Thalamuthu A, Kochan NA, Andrews G, Lima-Costa MF, Castro-Costa E, Brayne C, Matthews FE (2017) Age-related cognitive decline and associations with sex, education and apolipoprotein E genotype across ethnocultural groups and geographic regions: A collaborative cohort study. *PLoS Med* **14**, e1002261.
- [2] Salthouse TA (2009) When does age-related cognitive decline begin? *Neurobiol Aging* **30**, 507-514.
- [3] Boyle PA, Yu L, Leurgans SE, Wilson RS, Brookmeyer R, Schneider JA, Bennett DA (2019) Attributable risk of Alzheimer's dementia attributed to age-related neuropathologies. *Ann Neurol* **85**, 114-124.
- [4] Corrada MM, Sonnen JA, Kim RC, Kawas CH (2016) Microinfarcts are common and strongly related to dementia in the oldest-old: The 90+ study. *Alzheimers Dement* **12**, 900-908.
- [5] Godrich D, Martin ER, Schellenberg G, Pericak-Vance MA, Cuccaro M, Scott WK, Kukull W, Montine T, Beecham GW (2022) Neuropathological lesions and their contribution to dementia and cognitive impairment in a heterogeneous clinical population. *Alzheimers Dement* **18**, 2403-2412.
- [6] Kawas CH, Kim RC, Sonnen JA, Bullain SS, Trieu T, Corrada MM (2015) Multiple pathologies are common and related to dementia in the oldest-old: The 90+ Study. *Neurology* **85**, 535-542.
- [7] White LR, Edland SD, Hemmy LS, Montine KS, Zarow C, Sonnen JA, Uyehara-Lock JH, Gelber RP, Ross GW, Petrovitch H, Masaki KH, Lim KO, Launer LJ, Montine TJ (2016) Neuropathologic comorbidity and cognitive impairment in the Nun and Honolulu-Asia Aging Studies. *Neurology* **86**, 1000-1008.
- [8] Gefen T, Peterson M, Papastefan ST, Martersteck A, Whitney K, Rademaker A, Bigio EH, Weintraub S, Rogalski E, Mesulam M-M (2015) Morphometric and histologic substrates of cingulate integrity in elders with exceptional memory capacity. *J Neurosci* **35**, 1781-1791.
- [9] Gefen T, Papastefan ST, Rezvanian A, Bigio EH, Weintraub S, Rogalski E, Mesulam M-M, Geula C (2018) Von Economo neurons of the anterior cingulate across the lifespan and in Alzheimer's disease. *Cortex* **99**, 69-77.
- [10] Gefen T, Kawles A, Makowski-Woidan B, Engelmeyer J, Ayala I, Abbassian P, Zhang H, Weintraub S, Flanagan ME, Mao Q, Bigio EH, Rogalski E, Mesulam MM, Geula C (2021) Paucity of entorhinal cortex pathology of the Alzheimer's type in SuperAgers with superior memory performance. *Cereb Cortex* **31**, 3177-3183.
- [11] Rogalski E, Gefen T, Mao Q, Connelly M, Weintraub S, Geula C, Bigio EH, Mesulam M-M (2019) Cognitive trajectories and spectrum of neuropathology in SuperAgers: The first 10 cases. *Hippocampus* **29**, 458-467.
- [12] Rogalski EJ, Gefen T, Shi J, Samimi M, Bigio E, Weintraub S, Geula C, Mesulam M-M (2013) Youthful memory capacity in old brains: Anatomic and genetic clues from the Northwestern SuperAging Project. *J Cogn Neurosci* **25**, 29-36.
- [13] Montine TJ, Cholerton BA, Corrada MM, Edland SD, Flanagan ME, Hemmy LS, Kawas CH, White LR (2019) Concepts for brain aging: Resistance, resilience, reserve, and compensation. *Alzheimers Res Ther* **11**, 22.
- [14] Montine TJ, Corrada MM, Kawas C, Bukhari S, White L, Tian L, Cholerton B (2022) Association of cognition and dementia with neuropathologic changes of Alzheimer disease and other conditions in the oldest-old. *Neurology* **99**, e1067-e1078.
- [15] Corrada MM, Berlau DJ, Kawas CH (2012) A population-based clinicopathological study in the oldest-old: The 90+ study. *Curr Alzheimer Res* **9**, 709-717.
- [16] Kawas CH (2008) The oldest old and the 90+ Study. *Alzheimers Dement* **4**, S56-S59.
- [17] Paganini-Hill A, Ross RK, Henderson BE (1986) Prevalence of chronic disease and health practices in a retirement community. *J Chronic Dis* **39**, 699-707.
- [18] Melikyan ZA, Greenia DE, Corrada MM, Hester MM, Kawas CH, Grill JD (2019) Recruiting the oldest-old for clinical research. *Alzheimer Dis Assoc Disord* **33**, 160-162.
- [19] Hyman BT, Phelps CH, Beach TG, Bigio EH, Cairns NJ, Carrillo MC, Dickson DW, Duyckaerts C, Frosch MP, Masliah E, Mirra SS, Nelson PT, Schneider JA, Thal DR, Thies B, Trojanowski JQ, Vinters HV, Montine TJ (2012) National Institute on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease. *Alzheimers Dement* **8**, 1-13.
- [20] Montine TJ, Phelps CH, Beach TG, Bigio EH, Cairns NJ, Dickson DW, Duyckaerts C, Frosch MP, Masliah E, Mirra SS, Nelson PT, Schneider JA, Thal DR, Trojanowski JQ, Vinters HV, Hyman BT (2012) National Institute on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease: A practical approach. *Acta Neuropathol (Berl)* **123**, 1-11.
- [21] Vonsattel JPG, Myers RH, Tessa Hedley-Whyte E, Ropper AH, Bird ED, Richardson Jr EP (1991) Cerebral amyloid angiopathy without and with cerebral hemorrhages: A comparative histological study. *Ann Neurol* **30**, 637-649.
- [22] McKeith IG, Boeve BF, Dickson DW, Halliday G, Taylor J-P, Weintraub D, Aarsland D, Galvin J, Attems J, Ballard CG, Bayston A, Beach TG, Blanc F, Bohnen N, Bonanni L, Bras J, Brundin P, Burn D, Chen-Plotkin A, Duda JE, El-Agnaf O, Feldman H, Ferman TJ, ffytche D, Fujishiro H, Galasko D, Goldman JG, Gomperts SN, Graff-Radford NR, Honig LS, Iranzo A, Kantarci K, Kaufer D, Kukull W, Lee VMY, Leverenz JB, Lewis S, Lippa C, Lunde A, Masellis M, Masliah E, McLean P, Mollenhauer B, Montine TJ, Moreno E, Mori E, Murray M, O'Brien JT, Orimo S, Postuma RB, Ramaswamy S, Ross OA, Salmon DP, Singleton A, Taylor A, Thomas A, Tiraboschi P, Toledo JB,

- 927 Trojanowski JQ, Tsuang D, Walker Z, Yamada M, Kosaka K
928 (2017) Diagnosis and management of dementia with Lewy
929 bodies. *Neurology* **89**, 88.
- 930 [23] Nelson PT, Dickson DW, Trojanowski JQ, Jack CR, Boyle
931 PA, Arfanakis K, Rademakers R, Alafuzoff I, Attems J,
932 Brayne C, Coyle-Gilchrist ITS, Chui HC, Fardo DW, Flana-
933 gan ME, Halliday G, Hokkanen SRK, Hunter S, Jicha GA,
934 Katsumata Y, Kawas CH, Keene CD, Kovacs GG, Kukull
935 WA, Levey AI, Makkinejad N, Montine TJ, Murayama S,
936 Murray ME, Nag S, Rissman RA, Seeley WW, Sperling
937 RA, White III CL, Yu L, Schneider JA (2019) Limbic-
938 predominant age-related TDP-43 encephalopathy (LATE):
939 Consensus working group report. *Brain* **142**, 1503-1527.
- 940 [24] Kovacs GG, Ferrer I, Grinberg LT, Alafuzoff I, Attems J,
941 Budka H, Cairns NJ, Crary JF, Duyckaerts C, Ghetti B,
942 Halliday GM, Ironside JW, Love S, Mackenzie IR, Munoz
943 DG, Murray ME, Nelson PT, Takahashi H, Trojanowski
944 JQ, Ansorge O, Arzberger T, Baborie A, Beach TG, Bien-
945 niek KF, Bigio EH, Bodi I, Dugger BN, Feany M, Gelpi E,
946 Gentleman SM, Giaccone G, Hatanpaa KJ, Heale R, Hof
947 PR, Hofer M, Hortobágyi T, Jellinger K, Jicha GA, Ince
948 P, Kofler J, Kövari E, Kril JJ, Mann DM, Matej R, McKee
949 AC, McLean C, Milenkovic I, Montine TJ, Murayama S,
950 Lee EB, Rahimi J, Rodriguez RD, Rozemüller A, Schnei-
951 der JA, Schultz C, Seeley W, Seilhean D, Smith C, Tagliavini
952 F, Takao M, Thal DR, Toledo JB, Tolnay M, Troncoso JC,
953 Vinters HV, Weis S, Wharton SB, White CL, Wisniewski T,
954 Woulfe JM, Yamada M, Dickson DW (2016) Aging-related
955 tau astroglialopathy (ARTAG): Harmonized evaluation strat-
956 egy. *Acta Neuropathol (Berl)* **131**, 87-102.
- 957 [25] Whittle C, Corrada MM, Dick M, Ziegler R, Kahle-
958 Wroblewski K, Paganini-Hill A, Kawas C (2007) Neuropsy-
959 chological data in nondemented oldest old: The 90+ Study.
960 *J Clin Exp Neuropsychol* **29**, 290-299.
- 961 [26] GUZE SB (1995) Diagnostic and Statistical Manual of Men-
962 tal Disorders, 4th ed. (DSM-IV). *Am J Psychiatry* **152**,
963 1228-1228.
- 964 [27] Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental
965 state": A practical method for grading the cognitive state of
966 patients for the clinician. *J Psychiatr Res* **12**, 189-198.
- 967 [28] Melikyan ZA, Corrada MM, Dick MB, Whittle C, Paganini-
968 Hill A, Kawas CH (2019) Neuropsychological test norms
969 in cognitively intact oldest-old. *J Int Neuropsychol Soc* **25**,
970 530-545.
- 971 [29] Azarpazhooh MR, Avan A, Cipriano LE, Munoz DG, Erfan-
972 ian M, Amiri A, Stranges S, Hachinski V (2020) A
973 third of community-dwelling elderly with intermediate and
974 high level of Alzheimer's neuropathologic changes are not
975 demented: A meta-analysis. *Ageing Res Rev* **58**, 101002.
- 976 [30] Rezvanian A, Ohm DT, Kukreja L, Gefen TD, Weintraub
977 S, Rogalski E, Kim R, Aguirre C, Corrada M, Mesulam M-
978 M, Kawas C, Geula C (2016) The oldest-old with preserved
979 cognition and the full range of Alzheimer pathology. *Society
980 for Neuroscience*.
- 981 [31] Robinson JL, Corrada MM, Kovacs GG, Dominique M,
982 Caswell C, Xie SX, Lee VM-Y, Kawas CH, Trojanowski
983 JQ (2018) Non-Alzheimer's contributions to dementia and
984 cognitive resilience in The 90+ Study. *Acta Neuropathol
985 (Berl)* **136**, 377-388.
- 986 [32] Boyle PA, Wang T, Yu L, Wilson RS, Dawe R, Arfanakis K,
987 Schneider JA, Bennett DA (2021) To what degree is late life
988 cognitive decline driven by age-related neuropathologies?
989 *Brain* **144**, 2166-2175.
- 990 [33] J. Montine T, A. Sonnen J, S. Montine K, K. Crane P, B.
991 Larson E (2012) Adult Changes in Thought Study: Demen-
tia is an Individually varying convergent syndrome with
prevalent clinically silent diseases that may be modified by
some commonly used therapeutics. *Curr Alzheimer Res* **9**,
718-723.
- [34] Boyle P, Yu L, Wilson R, Schneider J, Bennett D (2013)
Relation of neuropathology with cognitive decline among
older persons without dementia. *Front Aging Neurosci* **5**,
50.
- [35] Latimer CS, Burke BT, Liachko NF, Currey HN, Kilgore
MD, Gibbons LE, Henriksen J, Darvas M, Domoto-
Reilly K, Jayadev S, Grabowski TJ, Crane PK, Larson
EB, Kraemer BC, Bird TD, Keene CD (2019) Resis-
tance and resilience to Alzheimer's disease pathology are
associated with reduced cortical pTau and absence of
limbic-predominant age-related TDP-43 encephalopathy in
a community-based cohort. *Acta Neuropathol Commun* **7**,
91.
- [36] Boyle PA, Yu L, Nag S, Leurgans S, Wilson RS, Ben-
nett DA, Schneider JA (2015) Cerebral amyloid angiopathy
and cognitive outcomes in community-based older persons.
Neurology **85**, 1930.
- [37] Arvanitakis Z, Leurgans SE, Barnes LL, Bennett DA,
Schneider JA (2011) Microinfarct pathology, dementia, and
cognitive systems. *Stroke* **42**, 722-727.
- [38] Brundel M, de Bresser J, van Dillen JJ, Kappelle LJ, Biessels
GJ (2012) Cerebral microinfarcts: A systematic review of
neuropathological studies. *J Cereb Blood Flow Metab* **32**,
425-436.
- [39] Sonnen JA, Larson EB, Crane PK, Haneuse S, Li G,
Schellenberg GD, Craft S, Leverenz JB, Montine TJ
(2007) Pathological correlates of dementia in a longitudinal,
population-based sample of aging. *Ann Neurol* **62**, 406-413.
- [40] Troncoso JC, Zonderman AB, Resnick SM, Crain B, Plet-
nikova O, O'Brien RJ (2008) Effect of infarcts on dementia
in the Baltimore longitudinal study of aging. *Ann Neurol* **64**,
168-176.
- [41] Suemoto CK, Nitrini R, Grinberg LT, Ferretti RE, Far-
fel JM, Leite RE, Menezes PR, Fregni F, Jacob-Filho W,
Pasqualucci CA; Brazilian Aging Brain Study Group (2011)
Atherosclerosis and dementia. *Stroke* **42**, 3614-3615.
- [42] Dolan H, Crain B, Troncoso J, Resnick SM, Zonderman AB,
Obrien RJ (2010) Atherosclerosis, dementia, and Alzheimer
disease in the Baltimore Longitudinal Study of aging cohort.
Ann Neurol **68**, 231-240.
- [43] Hilal S, Doolabi A, Vrooman H, Ikram MK, Ikram MA,
Vernooij MW (2021) Clinical relevance of cortical cerebral
microinfarcts on 1.5T magnetic resonance imaging in the
late-adult population. *Stroke* **52**, 922-930.
- [44] van Oijen M, Jan de Jong F, Witteman JCM, Hofman A,
Koudstaal PJ, Breteler MMB (2007) Atherosclerosis and
risk for dementia. *Ann Neurol* **61**, 403-410.
- [45] Yilmaz P, Ikram MA, Ikram MK, Niessen WJ, Viswanathan
A, Charidimou A, Vernooij MW (2019) Application of an
imaging-based sum score for cerebral amyloid angiopa-
thy to the general population: Risk of major neurological
diseases and mortality. *Front Neurol* **10**, 1276.
- [46] Lamar M, Leurgans S, Kapasi A, Barnes LL, Boyle PA,
Bennett DA, Arfanakis K, Schneider JA (2022) Complex
profiles of cerebrovascular disease pathologies in the aging
brain and their relationship with cognitive decline. *Stroke*
53, 218-227.
- [47] Nelson PT, Schmitt FA, Lin Y, Abner EL, Jicha GA, Patel
E, Thomason PC, Neltner JH, Smith CD, Santacruz KS,
Sonnen JA, Poon LW, Gearing M, Green RC, Woodard JL,
Van Eldik LJ, Kryscio RJ (2011) Hippocampal sclerosis in

- 1057 advanced age: Clinical and pathological features. *Brain J*
1058 *Neurol* **134**, 1506-1518.
- 1059 [48] Flanagan ME, Cholerton B, Latimer CS, Hemmy LS,
1060 Edland SD, Montine KS, White LR, Montine TJ (2018)
1061 TDP-43 neuropathologic associations in the Nun Study and
1062 the Honolulu-Asia Aging Study. *J Alzheimers Dis* **66**, 1549-
1063 1558.
- 1064 [49] Nag S, Yu L, Capuano AW, Wilson RS, Leurgans SE, Ben-
1065 nett DA, Schneider JA (2015) Hippocampal sclerosis and
1066 TDP-43 pathology in aging and Alzheimer disease. *Ann*
1067 *Neurol* **77**, 942-952.
- 1068 [50] Nag S, Barnes LL, Yu L, Buchman AS, Bennett DA, Schnei-
1069 der JA, Wilson RS (2021) Association of Lewy bodies with
1070 age-related clinical characteristics in Black and White de-
1071 cedents. *Neurology* **97**, e825.
- 1072 [51] Corrada MM, Melikyan ZA, Dominguez EN, Ho C, Sajjadi
1073 SA, Kawas CH (2021) Super-agers in memory are not nec-
1074 essarily super-agers in other cognitive domains. *Alzheimers*
Dement **17**, e055708.
- [52] Yu J, Collinson SL, Liew TM, Ng T-P, Mahendran R, Kua
1075 E-H, Feng L (2020) Super-cognition in aging: Cognitive
1076 profiles and associated lifestyle factors. *Appl Neuropsychol*
1077 *Adult* **27**, 497-503.
- [53] Greenland S, Mansournia MA, Altman DG (2016) Sparse
1079 data bias: A problem hiding in plain sight. *BMJ* **352**, i1981.
1080
- [54] Black SA, Espino DV, Mahurin R, Lichtenstein MJ, Hazuda
1081 HP, Fabrizio D, Ray LA, Markides KS (1999) The influence
1082 of noncognitive factors on the Mini-Mental State Exam-
1083 ination in older Mexican-Americans: Findings from the
1084 Hispanic EPESE. *J Clin Epidemiol* **52**, 1095-1102.
1085
- [55] Crum RM, Anthony JC, Bassett SS, Folstein MF (1993)
1086 Population-based norms for the Mini-Mental State Exam-
1087 ination by age and educational level. *JAMA* **269**, 2386-2391.
1088
- [56] Roberts AW, Ogunwole SU, Blakeslee L, Rabe MA (2018)
1089 The population years and older in the United States: 2016.
1090 American Community Survey Reports. United States Cen-
1091 sus Bureau. Accessed on September 23, 2022.
1092