

UCLA

UCLA Previously Published Works

Title

Novel Agents for the Pharmacological Treatment of Alcohol Use Disorder.

Permalink

<https://escholarship.org/uc/item/26z37645>

Journal

Drugs, 82(3)

ISSN

0012-6667

Authors

Burnette, Elizabeth M

Nieto, Steven J

Grodin, Erica N

et al.

Publication Date

2022-02-01

DOI

10.1007/s40265-021-01670-3

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial License, available at <https://creativecommons.org/licenses/by-nc/4.0/>

Peer reviewed



Novel Agents for the Pharmacological Treatment of Alcohol Use Disorder

Elizabeth M. Burnette^{1,2} · Steven J. Nieto¹ · Erica N. Grodin¹ · Lindsay R. Meredith¹ · Brian Hurley³ · Karen Miotto⁴ · Artha J. Gillis⁴ · Lara A. Ray^{1,4}

Accepted: 21 December 2021 / Published online: 8 February 2022
© The Author(s) 2022

Abstract

Alcohol use disorder (AUD) is a highly prevalent but severely under-treated disorder, with only three widely-approved pharmacotherapies. Given that AUD is a very heterogeneous disorder, it is unlikely that one single medication will be effective for all individuals with an AUD. As such, there is a need to develop new, more effective, and diverse pharmacological treatment options for AUD with the hopes of increasing utilization and improving care. In this qualitative literature review, we discuss the efficacy, mechanism of action, and tolerability of approved, repurposed, and novel pharmacotherapies for the treatment of AUD with a clinical perspective. Pharmacotherapies discussed include: disulfiram, acamprosate, naltrexone, nalmefene, topiramate, gabapentin, varenicline, baclofen, sodium oxybate, aripiprazole, ondansetron, mifepristone, ibudilast, suvorexant, prazosin, doxazosin, *N*-acetylcysteine, GET73, ASP8062, ABT-436, PF-5190457, and cannabidiol. Overall, many repurposed and novel agents discussed in this review demonstrate clinical effectiveness and promise for the future of AUD treatment. Importantly, these medications also offer potential improvements towards the advancement of precision medicine and personalized treatment for the heterogeneous AUD population. However, there remains a great need to improve access to treatment, increase the menu of approved pharmacological treatments, and de-stigmatize and increase treatment-seeking for AUD.

Key Points

Alcohol use disorder is severely under-treated, and the development of new and more effective pharmacotherapies is necessary.

The repurposed and novel agents discussed in this qualitative literature review offer promise for the future of AUD treatment, including advancements toward precision medicine for the heterogeneous AUD population.

1 Introduction

Alcohol use disorder (AUD) is a highly prevalent, chronic, relapsing condition characterized by an impaired ability to stop or control alcohol use despite clinically significant impairment, distress, or other adverse consequences [1, 2]. It is the most common substance use disorder: in 2016, 100.4 million people globally were estimated to have an AUD [3]. This disorder represents a significant public health concern. The WHO estimates that alcohol consumption is responsible for 5.9% of all deaths (7.6% in men, 4.0% in women) and 5.1% of global disease burden [4]. Alcohol use and misuse is thought to contribute to over 200 related diseases and health conditions globally, including cardiovascular disease, cancer, liver cirrhosis, and injuries [4]. AUD is also often comorbid with other substance use disorders, major depressive disorder, bipolar disorder, and other psychiatric disorders [2].

In the USA alone, AUD is estimated to contribute to about 88,000 deaths each year [5]. An estimated 44.6 million adults per year in the USA suffer from AUD, and 93.4 million (approximately 33%) adults in the USA will meet or have met AUD criteria at some point in their life [2].

✉ Lara A. Ray
lararay@psych.ucla.edu

¹ Department of Psychology, University of California, Los Angeles, 1285 Franz Hall, Box 951563, Los Angeles, CA 90095-1563, USA

² Neuroscience Interdepartmental Program, University of California, Los Angeles, Los Angeles, CA, USA

³ Department of Public Health, Los Angeles County, CA, USA

⁴ Department of Psychiatry and Biobehavioral Sciences, University of California, Los Angeles, Los Angeles, CA, USA

Furthermore, the economic burden of AUD is estimated to be approximately \$250 billion across the USA [6].

Although AUD is an important public health concern, the disorder remains severely under treated with only 7% of adults with AUD in the USA [7] and less than 10% in Europe [8] receiving pharmacotherapy and/or psychotherapy treatment. Furthermore, the lag between the age at which AUD onsets and the age of first AUD treatment has been estimated to be eight years on average [7]. Pharmacotherapies have seen limited use in the treatment of AUD, partially due to lack of addiction treatment training for medical professionals, lack of awareness regarding medication options (e.g., limited marketing), reluctance to prescribe and take medications, perceived low efficacy of medication, and stigma surrounding treatment. Indeed, only three pharmacotherapies are approved by the US Food and Drug Administration (FDA) for use in AUD treatment. These medications are disulfiram (Antabuse), acamprosate, and naltrexone (formulated for oral administration or extended-release injection) [9–11]. The European Medicines Agency (EMA) similarly recognizes only these same three medications, as well as nalmefene, as established pharmacotherapies for AUD [12]. These medications are only modestly effective [13] and are under-utilized in treatment. A study conducted in 2019 found that only 1.6% of adults in the USA with past-year AUD received evidence-based AUD pharmacotherapies [14].

Treatment outcomes for AUD differ widely across patients and medications. While abstinence may be desirable, it is infrequently obtained, such that only 16% of individuals in treatment for AUD achieve abstinence [15]. Furthermore, evidence does not support abstinence as the only approach in the treatment of AUD [16]. Not all individuals with AUD consider abstinence to be a goal of their recovery; only 2–6% of goals set in patient-driven treatment center on attaining alcohol abstinence [17]. Non-abstinent recovery, including reductions in drinking and heavy drinking in particular (e.g., controlled drinking / harm reduction), has been recognized to have health and societal value and has gained traction as a treatment target [18–20]. Indeed, non-abstinent AUD recovery has been shown to be sustainable for up to 10 years following treatment [21]. Despite growing recognition of the benefits of harm reduction, however, the most commonly prescribed pharmacotherapy to treat AUD remains disulfiram, a medication advised strictly for abstinence [22]. Furthermore, the heterogeneity of AUD suggests that it will be unlikely that one single medication will be effective for all individuals with an AUD. Therefore, there is a pressing need for the development of novel, diverse, and effective pharmacological treatment options for AUD with the hopes of increasing utilization and improving care. The focus of the current review is to summarize pharmacotherapies for AUD with a clinical perspective. Specifically, this review provides a brief overview of currently approved medications

and identifies new and repurposed agents “on the horizon” for which evidence indicates a potentially effective application toward AUD treatment.

2 Pharmacological Treatments for AUD

The following sections examine pharmacotherapies approved or in-development for AUD. Medications were selected for this qualitative review by considering gaps in existing review articles and the expertise of all authors; information was gathered via qualitative PubMed literature searches. This review is organized into sections based on drug approval status, namely (a) approved medications, (b) repurposed (i.e., off-label) medications, and (c) novel agents. Within each medication section, we examine the basic mechanism of action, evaluate preclinical research testing the efficacy of the medication in mitigating alcohol-related behaviors in animal models, and review clinical findings from human laboratory studies and randomized controlled trials (RCTs) where available. We also examine the tolerability and potential personalized applications of each drug by identifying populations in which the drug may be particularly effective, and indicate treatment targets (drinking reduction, achievement, or maintenance of abstinence). This information is summarized in Table 1. We conclude by discussing future directions for the development of pharmacological treatments and precision medicine for AUD.

2.1 Approved Agents

This section briefly describes medications currently approved by agencies in many countries including the US FDA and EMA for the treatment of AUD: disulfiram, acamprosate, and naltrexone (oral and extended-release), as well as nalmefene, which is EMA-approved. This section also reviews baclofen and sodium oxybate, which are medications approved for AUD treatment by agencies in European countries but not by the FDA or EMA. As many of these pharmacotherapies have been extensively discussed in the literature, our review of these medications primarily focuses on clinical trials and recent meta-analyses.

2.1.1 Disulfiram

Disulfiram, the first medication FDA-approved for AUD treatment (in 1951), is an aldehyde dehydrogenase inhibitor that acts by blocking the metabolism of alcohol, increasing acetaldehyde concentration. Acetaldehyde, a toxic metabolite of ethanol, produces an alcohol-induced aversive response, characterized by nausea, vomiting, sweating, flushing, and heart palpitations [23]. Unsurprisingly, these unpleasant effects give disulfiram a relatively poor adherence

Table 1 Pharmacotherapies approved and under development for the treatment of AUD

Agent	Mechanism of action	Approval status	Preclinical results	Clinical results	Target outcome
Disulfiram	Aldehyde dehydrogenase inhibitor; dopamine beta-hydroxylase inhibitor	FDA- and EMA-approved for AUD	Mixed results: prevents heavy but not moderate drinking [297]; chronic alcohol induces disulfiram tolerance [298]	Mixed results; drinking outcomes improved with supervised administration [29–33]	Achieve and maintain abstinence
Acamprosate	Modulates glutamatergic activity	FDA- and EMA-approved for AUD	Reduced ethanol intake, withdrawal symptoms, place preference [299–302]	Mixed results; Reduced risk of relapse, increased cumulative abstinence duration [36, 44, 49]. Negative studies show no benefit over placebo [37–42]	Achieve and maintain abstinence
Naltrexone	Opioid receptor antagonist	FDA- and EMA-approved for AUD	Reduced binge drinking, ethanol preference, motor impairment and sedation [303–305]	Reduced risk of relapse, binge drinking, and craving [52–59]; modest effect sizes [62–64]	Achieve and maintain abstinence and reduce drinking
Nalmefene	Opioid receptor antagonist and partial agonist	EMA-approved for AUD; FDA-approved for opioid overdose	Reduced alcohol seeking, relapse and binge-like drinking, neuroinflammation [306–308]	Reduced binge drinking and total alcohol consumption [71–73, 75]	Reduce drinking
Baclofen	GABA _B agonist	Approved for AUD in France	Reduced ethanol self-administration and motivational properties [309–312]	Mixed results; increased rates of abstinence, time to first relapse, possibly reduced heavy-drinking days [84–86, 91, 92]	Achieve and maintain abstinence
Sodium oxybate	Modulates GABA activity	Approved for AUD in Italy and Austria	Reduced ethanol self-administration, withdrawal symptoms, and relapse-like drinking [247, 313]	Effective treatment of alcohol withdrawal syndrome, increased abstinence [99, 100]	Reduce withdrawal symptoms and achieve abstinence
Topiramate	Inhibits glutamatergic activity and increases GABA activity	Repurposed	Reduced ethanol intake in rodent models [113–115]	Reduced drinks per day and percent heavy-drinking days and increased percent days abstinent [112, 116–119]	Achieve abstinence and reduce drinking
Gabapentin	Modulates GABA activity	Repurposed	Mixed results: reduced or increased ethanol intake [123–125]	Reduced percent heavy-drinking days, alcohol consumption, and abstinence rates and increased time to relapse, with outcomes improved among those with alcohol withdrawal symptoms [126–135]	Achieve abstinence and reduce drinking
Varenicline	Nicotinic acetylcholine receptor agonist	Repurposed	Reduced ethanol seeking, intake, and binge-like consumption [138–140]	Reduced percent heavy-drinking days and drinks per day with outcomes improved for those who smoke cigarettes [142–144, 146–149]	Reduce drinking
Aripiprazole	Dopamine receptor partial agonist and 5-HT _{2A} receptor antagonist	Repurposed	Reduce ethanol-induced place preference and ethanol consumption [151–153]	Mixed results with outcomes improved for more impulsive individuals; reduced heavy-drinking days and increased days abstinent at rates comparable to naltrexone [154–160]	Reduce drinking
Ondansetron	5-HT ₃ receptor antagonist	Repurposed	Blocked sensitization to locomotor stimulant effects, reduced voluntary ethanol intake [167–169]	Reduced drinks per day and increased days abstinent in patients with early-onset AUD [170–174]	Achieve abstinence and reduce drinking

Table 1 (continued)

Agent	Mechanism of action	Approval status	Preclinical results	Clinical results	Target outcome
Mifepristone	Glucocorticoid receptor antagonist	Repurposed	Reduced ethanol-seeking in rodents [177, 178] Mixed results in primates; reduced chronic voluntary alcohol use but did not prevent relapse [179, 180]	Reduced alcohol craving and consumption [178, 181, 182]	Reduce drinking
lbutilast	Selective PDE inhibitor and M1F inhibitor	Repurposed	Reduced ethanol intake [188–191]	Reduced craving, odds of heavy drinking, and neural cue-reactivity [192, 193]	Reduce drinking
Prazosin and Doxazosin	Adrenergic α -1 antagonists	Repurposed	Reduced relapse, ethanol intake, and stress-induced ethanol seeking [201–205]	Mixed results; reduced craving and alcohol consumption. Outcomes improved among those with alcohol withdrawal symptoms [206–211]	Reduce drinking
N-Acetyl-cysteine	Modulates synaptic glutamatergic activity	Repurposed	Reduced ethanol intake and relapse [215–220]	Increased abstinence rates and reduced drinks per week among those with cannabis use disorder [222, 223]	Reduce drinking
Suvorexant	Dual orexin antagonist	Repurposed	Reduced ethanol consumption and relapse [229–238]	–	Unknown
ABT-436	Highly selective vasopressin type 1B receptor antagonist	Novel agent	Attenuated ethanol reinstatement and reduced ethanol intake [241, 242]	Increased days abstinent but did not reduce heavy drinking or craving [244]	Achieve and maintain abstinence
GET73	Gamma-hydroxybutyrate analogue; negative allosteric modulator at mGluR5	Novel agent	Reduced ethanol intake and suppressed relapse [246]	–	Unknown
ASP8062	Positive allosteric modulator of GABA _B receptor	Novel agent	Reduced ethanol self-administration [259]	–	Unknown
PF-5190457	Ghrelin receptor inverse agonist	Novel agent	Reduced ethanol intake and preference [266–270]	Reduced alcohol craving and cue-reactivity [272]	Unknown
Cannabidiol	Diverse pharmacological effects	Novel agent	Reduced ethanol administration and relapse-like behavior [279]	–	Unknown

5-HT₃ serotonin receptor 3, 5-HT_{2A} serotonin receptor 2A, AUD alcohol use disorder, EMA European Medicines Agency, FDA United States Food and Drug Administration, GABA gamma-aminobutyric acid, mGluR5 metabotropic glutamate receptor 5, MIF macrophage migration inhibitory factor, PDE phosphodiesterase

rate [24, 25]. Disulfiram may also act on dopamine systems, as its major metabolite inhibits the enzyme dopamine beta-hydroxylase (DBH), which aids in metabolizing dopamine into noradrenaline [22, 26, 27]. Serum DBH activity is associated with withdrawal symptoms, and disulfiram has been shown to reduce serum DBH levels [28].

A meta-analysis of 22 RCTs found that disulfiram saw increased success rates compared to placebo in open-label studies only—blinded trials yielded no statistical significance between disulfiram and placebo [29]. However, while these findings do not appear to support the use of disulfiram for treating AUD, this outcome may be due to placebo effects. Research showed that placebo-treated individuals showed decreases in cue-reactivity to alcohol stimuli in a sample of 38 participants [30], and experiencing an acetaldehyde reaction did not necessarily improve treatment response in a sample of 46 participants [31]. Instead, a patient simply being aware of a potential adverse reaction appears to be enough to influence drinking behavior [32].

Disulfiram ingestion under supervision (to ensure adherence) saw significantly better success rates compared to non-supervised treatment, suggesting that supervised administration of disulfiram may still have a place in treating individuals struggling to attain sobriety [33], and unsupervised disulfiram may also be helpful for individuals highly motivated for abstinence. However, adherence management issues limit the utility of disulfiram in the treatment of AUD, and the disulfiram-ethanol interaction can sometimes present as a medical emergency. Therefore, disulfiram is only recommended in the maintenance of abstinence; using this medication to reduce drinking is not advised [34].

2.1.2 Acamprosate

While the specific mechanisms through which acamprosate works to treat AUD remain under investigation, it is thought to act on the glutamatergic system as an *N*-methyl-D-aspartic acid (NMDA) receptor partial co-agonist [35]. This may reduce neuronal hyperexcitability, a phenomenon that occurs in acute withdrawal and protracted abstinence from alcohol.

A meta-analysis of 27 RCTs with a total of 7519 participants found that acamprosate treatment reduced risk of abstinent patients returning to any drinking, but did not reduce rates of binge drinking [36]. A number of trials have also found that acamprosate did not show a significant benefit over placebo [37–41]. In particular, a large scale trial (COMBINE), which compared acamprosate, naltrexone, and behavioral therapies, both individually and combined with each other, against placebo ($N = 1383$), found that acamprosate had no significant effect on drinking in comparison to placebo, either alone or in combination with naltrexone and/or behavioral intervention [42]. Placebo effects in this trial may explain some of these negative outcomes, as might

differences in trial design and patient characteristics: COMBINE required 4 days of pre-trial abstinence while European trials with positive outcomes were typically conducted in inpatient populations requiring complete detoxification [43]. Overall, however, a Cochrane meta-analysis of 24 RCTs with 6915 participants found that acamprosate significantly reduced the risk of any drinking and increased cumulative duration of abstinence [44].

Acamprosate is generally well tolerated [34] and may also have neuroprotective effects [45, 46]. As chronic alcohol abuse is associated with neuronal changes related to NMDA receptors, this neuroprotection may be particularly important in AUD treatment [47, 48]. Acamprosate is recommended for the achievement and maintenance of complete abstinence, rather than for the reduction of drinking or prevention of relapse in the event of drinking [49]. It is FDA-approved for abstinence maintenance in AUD patients who are abstinent when beginning treatment.

2.1.3 Naltrexone (Oral and Extended-Release)

Naltrexone, the best-studied of these three commonly approved medications, was originally approved to treat opioid use disorder. Naltrexone is an antagonist of the mu opioid receptor (with additional affinity for kappa and delta opioid receptors). By attenuating alcohol-induced opioid-dergic activity in the mesolimbic dopamine system, opioid antagonists like naltrexone, modulate the rewarding effects of alcohol (mediated by dopamine), thereby reducing alcohol consumption [50, 51].

In 1994, the FDA approved oral naltrexone to treat alcohol dependence after two independent 12-week trials, which included 97 and 70 participants, respectively, found that naltrexone significantly decreased drinking days and relapse rates [52, 53]. Additional recent trials demonstrate that naltrexone reduces the rewarding effects of alcohol [54, 55], alcohol craving [40, 56], drinks per drinking day [57], and relapse rates [58, 59], strengthening the initial findings.

Extended-release injectable naltrexone (i.e., Vivitrol), administered once monthly by a medical professional, may be beneficial for individuals who are more sensitive to naltrexone's adverse side effects or have difficulty adhering to oral medication [60]. A six-month multisite trial of 380 mg injectable naltrexone in 624 patients with alcohol dependence found significant reductions in heavy-drinking days versus placebo [61]. However, evidence for naltrexone remains mixed. Another RCT reported no significant differences between naltrexone and placebo ($N = 183$), and that the clinical utility of naltrexone was limited by its adverse effects [62]. Additionally, clinical trials of naltrexone often yield modest effect sizes [63]. A meta-analysis of 53 naltrexone RCTs with a total of 9140 participants found naltrexone

to significantly reduce both the risk of return to any drinking and return to binge drinking [36]; however, both of these associations were modest in magnitude [64].

Oral and injectable naltrexone show similar decreases in likelihood of binge drinking [65] and both are generally well tolerated, with fairly mild side effects [66]. Importantly, however, naltrexone does block the therapeutic effects of opioid analgesics and can precipitate opioid withdrawal in patients who have developed physical dependence to opioids [34]; therefore, individuals who are prescribed naltrexone for AUD must be monitored for opioid use and withdrawal. Of note, naltrexone is contraindicated for patients with acute hepatitis or liver failure, and should be “carefully considered” in patients with acute liver disease [67], potentially limiting its use in the alcohol-associated liver disease (AALD) population. In summary, naltrexone appears to have a moderate effect on the reduction of alcohol use.

2.1.4 Nalmefene

Nalmefene works in a similar manner to naltrexone as an antagonist at the mu and delta opioid receptor, but is also a kappa opioid receptor partial agonist [68]. Via its kappa agonist activity, particularly centered in dopaminergic nucleus accumbens circuitry, nalmefene may reduce motivation for self-administration and withdrawal-induced alcohol consumption [69, 70].

Nalmefene was approved by the EMA in 2013 for the reduction of alcohol consumption among patients with AUD. Approval followed findings from three multicenter 6-month clinical trials enrolling 604, 667, and 718 individuals, respectively, in which participants took the medication or placebo on an as-needed basis [71–73]. In these trials, nalmefene decreased total alcohol consumption and heavy-drinking days. However, the drug’s approval based on these studies has received criticism due to limited evidence of efficacy, especially as these trials were conducted only against placebo rather than an active medication comparison (e.g., naltrexone) [34, 74]. A more recent meta-analysis of the efficacy of nalmefene, which included five RCTs ($N = 2567$), found that participants treated with nalmefene had 1.65 fewer heavy-drinking days per month than participants treated with placebo after 6 months [75].

Studies indicate that nalmefene is associated with more adverse events and study dropouts compared to placebo, and the most frequently reported of these are dizziness, nausea, vomiting, insomnia, and headache [36, 76]. Overall, while nalmefene may reduce heavy-drinking rates, its effects on other outcomes [77] remain small-to-moderate, and study withdrawals related to adverse events are common in nalmefene trials. However, similar to naltrexone, this medication may appeal to patients with goals of reducing alcohol

consumption and those reluctant to engage in abstinence-based treatment.

2.1.5 Baclofen

Baclofen acts on the γ -amino butyric acid (GABA) system as a GABA_B agonist. It was approved for treatment of AUD in France in 2018 [78] and has been used off-label for AUD for over a decade in other countries, especially other European countries and Australia [79, 80].

Clinical trials of baclofen have produced mixed results [65–67]. A recent meta-analysis of 12 RCTs ($N = 703$) showed that baclofen in comparison to placebo was associated with higher abstinence rates [81–83]; however, it did not increase abstinent days or decrease craving, heavy drinking, depression, or anxiety [84]. Another meta-analysis of 13 RCTs ($N = 1492$) indicated that baclofen (again vs placebo) was associated with longer time to relapse and a larger percentage of abstinent patients [85]. Furthermore, greater alcohol use at baseline was correlated with a greater treatment effect [85]. In contrast, however, another recent meta-analysis of 12 RCTs with 1128 total participants found no significant differences between baclofen and placebo in primary (i.e., percent abstinent days, percent heavy-drinking days, return to any drinking, and study dropout) or secondary outcomes of interest (i.e., anxiety and craving); however, baclofen did increase depression and adverse effects including sedation and vertigo [86].

Baclofen’s significant adverse side effects include drowsiness, sedation, headache, vertigo, confusion, perspiration, muscle stiffness and/or abnormal movements, slurred speech, and numbness [86–88]. Tolerance to baclofen may also develop with chronic administration [89, 90]. Additionally, dose and sex may moderate baclofen tolerability and response; escalation of dosage in response to developing tolerance can increase sedative side effects, which affect women significantly more than men at the same dose [91]. Importantly, cessation or reduction in dose can precipitate potentially life-threatening withdrawal syndrome [90].

The variability in baclofen’s effectiveness seen across studies may be partially explained by high baclofen pharmacokinetic variability seen among individuals with AUD. This heterogeneity is an important factor to take into account when considering baclofen as an AUD treatment [92]. Of note, there is also some evidence that baclofen might be particularly useful in treatment of AUD among individuals with liver disease [93, 94].

In summary, baclofen seems to effectively promote abstinence; however, it shows mixed results regarding its clinical effects on non-abstinence outcomes (e.g., reduction in heavy drinking), significant adverse side effects, and inter-individual variability in response. As such, baclofen as a treatment

of AUD—as well as its optimal dosage—continues to be debated [95].

2.1.6 Sodium Oxybate

Sodium oxybate (SMO), the sodium salt of gamma-hydroxybutyrate (GHB), has been utilized as a medication for a number of disorders. SMO is approved in Italy and Austria for the treatment of AUD [96]. SMO acts on the GABA system both directly as a GABA_B partial agonist and indirectly through GHB-derived GABA [97, 98].

A Cochrane meta-analysis of 13 RCTs with a total of 649 participants found that SMO (50 mg) was effective compared to placebo in the treatment of alcohol withdrawal syndrome (AWS) and in preventing relapses in previously detoxified participants, and that SMO was more effective than naltrexone and disulfiram in maintaining abstinence [99]. Another recent meta-analysis found that SMO, compared to placebo, increased abstinence rates by up to 34% in a sample of 711 participants with very high drinking risk levels [100]. However, use of GHB as a recreational drug raises concern regarding the abuse potential of SMO. Indeed, in another clinical study with a sample of 48 participants, craving for SMO was observed in 10% of participants with psychiatric comorbidities [101], indicating that patients may be at risk of developing craving for and abusing SMO.

SMO may be particularly effective in combination with other pharmacotherapies. It was shown to reduce relapses when administered with naltrexone and escitalopram [102]. A study of 52 subjects with chronic, treatment-resistant alcohol dependence found that participants stayed in treatment significantly longer when SMO was co-administered with disulfiram than with SMO alone [103]. Additionally, in a case study of seven partial- and non-responders to SMO treatment, co-administration with nalmefene was effective in promoting abstinence, reducing heavy-drinking episodes, and importantly, suppressing craving for SMO [104].

SMO craving and abuse potential may limit its use; however, at therapeutic doses, SMO abuse seems to be a relatively infrequent phenomenon among patients without psychiatric comorbidities or poly-drug use [96, 100, 105]. Safety data show that serious adverse events with SMO treatment are rare, with the most common adverse effects being transitory dizziness and vertigo [96]. Overall, the effectiveness of SMO lies largely in reducing withdrawal symptoms and increasing abstinence rates. Its therapeutic use remains controversial due to its potential for craving and abuse, and further study on co-administration with other pharmacotherapies appears promising.

2.2 Repurposed Agents

The following sections address medications that have been approved by the FDA for applications other than AUD but are in development to be repurposed (i.e., used off-label) to treat AUD.

2.2.1 Topiramate

Despite that the anti-convulsant medication topiramate has not been approved by the FDA or EMA for AUD treatment, the US Department of Veterans Affairs recommends topiramate as a treatment for AUD, and it has been suggested by the American Psychiatric Association (APA) for the pharmacological treatment of patients with AUD [106, 107]. Currently, the specific mechanisms of action of topiramate remain under investigation. However, the drug is thought to inhibit glutamate α -amino 3-hydroxy-5-methyl-4-isoxazole-propionic acid (AMPA) and kainate receptors [108–110] and L-type calcium channels [111], as well as enhance the inhibitory activity of GABA [109]. These effects, taken together, work to attenuate dopaminergic activity mesolimbic reward circuits, thereby reducing both alcohol craving and withdrawal symptoms [112].

Preclinically, topiramate was seen to decrease ethanol consumption in rodent models [113–115]. Clinical studies have found that topiramate reduced craving and alcohol drinking in a sample of 150 participants over 12 weeks [112], and decreased drinking days, heavy-drinking days, and number of drinks per day in a sample of 85 participants in a 14-week trial [116]. A large, multisite 14-week RCT that enrolled 371 participants with alcohol dependence found reductions in heavy-drinking days and improvements in various self-reported drinking-related outcomes [117]. A meta-analysis including seven RCTs of topiramate for AUD with a total of 1125 participants found that topiramate significantly increased number of days abstinence and decreased heavy-drinking days compared to placebo [118]. Of interest, this meta-analysis also directly compared effect sizes for topiramate and naltrexone, finding that topiramate was significantly more effective than naltrexone in reducing binge drinking, but not in improving abstinence [118]. Another recent month-long study of 94 patients that directly compared topiramate and baclofen demonstrated that baclofen was overall more effective (i.e., 61% of participants receiving baclofen achieved abstinence vs 38% with topiramate) and better tolerated (i.e., 33% of participants receiving topiramate dropped out of the study vs 24% with baclofen) than topiramate [119].

The potential of reduced tolerability and significant side effects such as pruritis, paresthesia, anorexia, dysgeusia, dizziness, nervousness, and cognitive impairment, including difficulty with concentration and attention, are concerns for

the use of topiramate in the treatment of AUD [50]. These cognitive impairments were found to be dose-associated and include impairments in working memory and verbal fluency, as well as mental slowing [116]. Topiramate's side effects, especially those affecting memory and cognition, present reasons for concern and warrant long-term trials to further investigate these effects. Slow dose titration and ongoing patient monitoring is also recommended.

It is worth noting a potential strength of topiramate: the possibility of starting treatment while patients are still actively drinking, allowing topiramate to serve as a harm-reduction or abstinence-initiation treatment, rather than only being used to prevent relapse in already detoxified patients [120]. Topiramate has already been demonstrably effective in clinical applications across a number of drinking outcomes [121], but longer-term trials of topiramate are recommended to optimize dose and duration, as well as further exploring side effects.

2.2.2 Gabapentin

Another anti-convulsant medication, gabapentin, is also suggested by the APA for use in AUD treatment with the goal of reducing alcohol consumption or achieving abstinence [107]. It is thought to modulate GABA activity by indirectly interacting with voltage-gated calcium channels [122]. Preclinical studies of gabapentin's effects on ethanol consumption have shown mixed results, with some studies finding that gabapentin reduced ethanol intake in alcohol-dependent rodents [123] and others finding that it increased self-administration and binge-like drinking [124, 125].

Gabapentin has shown promising results in human laboratory studies and clinical trials. Gabapentin reduced percent heavy-drinking days and drinks per day, as well as increased percentage abstinent days in a 28-day trial with 60 participants [126]. In another 10-day study with 21 participants, gabapentin significantly delayed return to heavy drinking [127]. In a proof-of-concept one-week human laboratory trial ($N = 33$), participants treated with gabapentin showed significant decreases in alcohol craving in comparison to placebo [128], and a follow-up 12-week RCT with 150 participants found that gabapentin significantly increased abstinence rates and percentage of no heavy-drinking days [129]. Gabapentin may also be particularly effective in combination with other medications. For instance, a combination of gabapentin and flumazenil over 39 days in 60 participants increased percentage of abstinent days and delayed time to first heavy-drinking day compared to placebo. This trial also showed an interaction effect with pre-study alcohol withdrawal, such that those with more severe withdrawal symptoms at baseline benefitted more from the treatment during the trial [130]. In combination with naltrexone, gabapentin was also shown to be more effective (i.e., decreased

heavy-drinking days, fewer drinks per drinking day, and delayed time to heavy drinking) compared with naltrexone alone in a 6-week trial with 150 participants [131].

A Cochrane review of 25 studies ($N = 2641$) found anti-convulsants, including gabapentin, to significantly reduce both heavy-drinking days and drinks per drinking day in comparison to placebo. Anticonvulsants were also associated with longer time-to-relapse and fewer heavy-drinking days compared to naltrexone [132]. A recent meta-analysis of 7 RCTs with a total of 751 participants found that gabapentin compared to placebo only significantly decreased percent heavy-drinking days, although trend-level effect estimates were shown for other alcohol-related outcomes (i.e., relapse, drinks per day, complete abstinence, percent days abstinent, and concentration of gamma-glutamyl transferase (GGT)) [133]. In a 16-week RCT with 96 participants, gabapentin improved the percentage of individuals with no heavy-drinking days and increased total abstinence rates over placebo, especially among participants with more severe pre-study withdrawal [134]. However, another recent 6-month multisite RCT of extended-release gabapentin conducted in 346 participants with at least moderate AUD found no effects over placebo for any clinical outcomes including abstinence rate, percent heavy-drinking days, drinks per drinking day, or alcohol craving [135]. While these findings may be related to potential pharmacokinetic issues relating to the specific formulation of extended-release gabapentin used in the trial, it is also possible that gabapentin may simply be more effective in patients with more clinically relevant and severe symptoms of alcohol withdrawal [130, 134].

The most common adverse events associated with gabapentin treatment are somnolence, dizziness, peripheral edema, and ataxia or gait disorder [136]. Notably, gabapentin does carry the potential for misuse and abuse, particularly in individuals with opioid use disorders [137]; therefore, recommendation of gabapentin to individuals with comorbid substance use disorder should be carefully considered. In summary, gabapentin shows some clinical efficacy, especially in populations with more severe withdrawal symptomatology, but more extensive investigation is recommended.

2.2.3 Varenicline

Varenicline is a nicotinic acetylcholine receptor agonist (partial at $\alpha 4\beta 2$; full at $\alpha 7$) that is used for the treatment of nicotine dependence. In rodent models, varenicline was shown to reduce ethanol seeking, intake, and binge-like consumption [138–140]. Additionally, the combination of varenicline and naltrexone decreased alcohol drinking more effectively than either drug on its own [141].

Clinically, varenicline was found to significantly reduce weekly percent heavy-drinking days, drinks per day, drinks per drinking day, and alcohol craving compared to placebo

in a 13-week multisite RCT with 200 participants [142]. However, a 12-week RCT ($N = 160$) found no effect of varenicline over placebo on heavy-drinking days [143], and a recent 6-week human laboratory study in 47 participants found that varenicline did not significantly attenuate cue-induced alcohol craving relative to placebo [144].

Varenicline appears to be especially relevant to heavy-drinking smokers. Approximately 20%–25% of current smokers are estimated to also be heavy drinkers [145]. A human laboratory study with 20 heavy-drinking smokers on varenicline for 7 days found that varenicline significantly reduced the number of drinks consumed and increased the likelihood of complete abstinence in the human laboratory paradigm [146]. Moderator analyses of data collected in the above multisite trial [142] indicated that varenicline was more efficacious in reducing drinking among smokers who also reduced their cigarette smoking [147] and among individuals with lower severity of AUD, such that that varenicline significantly reduced percent heavy-drinking days and drinks per drinking day in low-severity individuals, while the most severe group showed no differences between varenicline and placebo on drinking outcomes [148].

Findings also indicate that varenicline may be more effective as an AUD treatment in men, as a 16-week RCT with 131 participants found that varenicline combined with medication management decreased percent heavy-drinking days only in men, but improved smoking abstinence in both men and women [149]. Varenicline is well tolerated, suggesting that it may serve as a promising option for AUD treatment, especially in the case of individuals co-using alcohol and nicotine.

2.2.4 Aripiprazole

Aripiprazole is an antipsychotic drug that acts as a partial agonist at the dopamine D2 and 5-HT1A receptors and as an antagonist at the 5-HT2A receptor [150]. Preclinically, aripiprazole has been shown to reduce ethanol-induced place preference and decrease drinking behaviors in mice [151], as well as reducing alcohol consumption in alcohol-preferring rats [152, 153].

A human laboratory study in 18 healthy participants indicated that aripiprazole affected outcomes related to alcohol consumption (i.e., subjective response to alcohol), including reducing the euphoric effects of alcohol and increasing sedative effects [154]. In a sample of 30 participants with AUD, aripiprazole compared to placebo, was found to attenuate cue-induced neural activation in the ventral striatum, a region associated with reward [155]. Another recent clinical laboratory study in which 99 participants with AUD took aripiprazole over eight days found that aripiprazole reduced the number of drinks consumed in a bar lab setting, especially among individuals

with low self-control, and prolonged the latency to drink in individuals with high impulsivity [156].

Results from clinical trials of aripiprazole are mixed. A 16-week RCT directly comparing aripiprazole against naltrexone ($N = 75$) found that the two medications were overall comparable in reducing heavy-drinking days and increasing days of abstinence, although patients treated with naltrexone had reduced craving compared to aripiprazole [157]. A 35-day open-label trial assessing the combination of aripiprazole and topiramate in 13 heavy-drinking participants found significant reductions in alcohol use. Additionally, this trial provided no evidence that the side effects of the two medications are additive, indicating that the combination is generally safe and well tolerated [158]. However, another recent five-week study assessing topiramate, aripiprazole, and their combination with 90 participants found that effects on drinking reduction were due to topiramate, while no significant findings were seen for aripiprazole for any outcomes [159]. Another multicenter RCT enrolling 295 participants over 12 weeks found no significant difference between the aripiprazole and placebo groups in percentage of participants completely abstinent from alcohol throughout the study or time to first drinking day; however, the average number of drinks per drinking day was significantly lower for aripiprazole [160]. The aripiprazole group yielded significantly higher discontinuation rates and earlier discontinuation, largely due to adverse events associated with side effects, especially when dosage exceeded 15 mg/day. The most common side effects cited in cases of discontinuation were fatigue, insomnia, restlessness, anxiety, and deficits in attention. Of note, long-term use of antipsychotics like aripiprazole may be associated with more severe adverse effects such as tardive dyskinesia, with risk factors including older age and female sex [161, 162].

In summary, these findings suggest that aripiprazole may be more effective at lower doses and in more impulsive individuals, although larger confirmatory studies are needed to pursue these personalized medicine approaches.

2.2.5 Ondansetron

Ondansetron is a 5-HT3 antagonist that is used to treat nausea and vomiting. Although the specific mechanism of action remains under investigation, ondansetron may address serotonergic dysfunction common in early-onset AUD [163, 164], and may reduce alcohol craving via 5-HT3 projections to dopaminergic connections in the midbrain. Preclinically, 5-HT3 antagonism has been shown to block acquisition and maintenance of ethanol self-administration [165] and reduce ethanol-associated dopamine concentration in the nucleus accumbens [166]. Ondansetron was also found to block the development and expression of sensitization to the

locomotor stimulant effects of ethanol [167] and reduce voluntary ethanol intake, preference, and withdrawal seizures in rodent models [168, 169].

Clinically, ondansetron may be particularly effective in combination with naltrexone. In an 8-week RCT in 20 participants with early-onset AUD, ondansetron and naltrexone (in comparison with placebo) significantly reduced drinks per drinking day and trended towards an increase in percentage of days abstinent [170]. Another combination study in 90 participants after 7 days on ondansetron and naltrexone found that the combination decreased craving for alcohol and ventral striatal activation to alcohol cues [171].

Ondansetron may also be suitable for individuals with biological predisposition to early-onset AUD. In an 11-week RCT of 271 participants, ondansetron was shown to reduce self-reported drinking such that patients with early-onset AUD who received ondansetron reported fewer drinks per day and more days of abstinence compared to placebo [172]. Ondansetron, combined with cognitive behavioral therapy in an 8-week open-label trial ($N = 40$) comparing effects in early-onset versus late-onset AUD, found that drinks per drinking day and alcohol-related problems (i.e., Drinker Inventory of Consequences; DrInC) were significantly decreased among those with early-onset AUD compared to those with late-onset AUD [173]. Furthermore, in an 11-week study with 253 participants, ondansetron at 4 $\mu\text{g/kg}$ reduced overall craving significantly in participants with early-onset AUD, while a lower dose (1 $\mu\text{g/kg}$) reduced craving in participants with late-onset AUD. These reductions in craving were also associated with reduced drinking (i.e., drinks per drinking day) and an increased percentage of abstinent days [174].

Ondansetron is well tolerated with relatively mild side effects including diarrhea, constipation, and headache [175]. Overall, these studies suggest a potential role for ondansetron as an AUD treatment, especially in participants with early-onset AUD and possibly in combination with naltrexone.

2.2.6 Mifepristone

Mifepristone (RU-486), a glucocorticoid receptor antagonist, works on the stress system by regulating the amygdala [176]. Preclinically, both systemic and central amygdala injections of mifepristone were shown to suppress yohimbine stress-induced reinstatement of alcohol seeking, indicating that the central amygdala plays an important role in mifepristone's effects on ethanol-seeking [177, 178]. Recent preclinical research in primates has shown mixed results, such that mifepristone was shown to decrease chronic voluntary alcohol consumption in rhesus macaques at doses of 30 and 56 mg/kg per day [179], but had no effect on alcohol-seeking or self-administration in baboons at slightly

lower doses of 10–30 mg/kg per day [180]. Of note, in the former study, cessation of mifepristone treatment resulted in a rapid return to baseline intake levels, and mifepristone was not effective in preventing a relapse during early abstinence.

In a human laboratory study with 56 alcohol-dependent participants, mifepristone (taken for one week) was effective in reducing alcohol craving and consumption relative to placebo, improved liver-function markers, and was overall well tolerated [178]. A two-week Phase 4 RCT examining the effects of mifepristone on cognition in AUD was recently conducted, but recruitment challenges rendered the results inconclusive [181, 182]. Of note, in this trial, participants who received mifepristone had higher Beck Depression Inventory scores compared to placebo at 4 weeks post-randomization despite similar scores at baseline between the two groups, indicating a greater severity of depression symptoms caused by the medication. Additional clinical research on mifepristone as a treatment for AUD and its potential side effects is warranted.

2.2.7 Ibudilast

Ibudilast is an inhibitor of phosphodiesterases (PDE)-3, -4, -10, and -11 [183] and macrophage migration inhibitory factor (MIF) [184]. Ibudilast has been shown to dose-dependently suppress pro-inflammatory cytokines, such as interleukins IL-1 β , IL-6, and tumor necrosis factor alpha (TNF- α), and to increase the anti-inflammatory cytokine IL-10 and neurotrophic factors [185]. As increases in inflammation are seen in AUD [186], the effects of ibudilast in treating AUD are thought to be driven by its anti-inflammatory and pro-neurotrophic qualities [187].

Preclinically, ibudilast was demonstrated to reduce alcohol intake in two rat models, and decreased drinking selectively in alcohol-dependent mice in comparison to non-dependent mice [188]. These preclinical findings align with prior rodent studies in which pharmacological inhibition of PDE also reduced alcohol intake [189–191].

In a 7-day human laboratory crossover trial ($N = 24$), ibudilast was well tolerated and decreased tonic craving in comparison to placebo. Additionally, ibudilast improved mood during exposure to alcohol and stress cues, and reduced the mood-altering and stimulant effects of alcohol among participants with more severe depressive symptoms [192]. Another recent 2-week RCT enrolling 52 participants found that ibudilast also significantly decreased the odds of heavy drinking during the trial by 45% compared to placebo and attenuated neural response to alcohol cues in the ventral striatum [193, 194].

Ibudilast appears to be well tolerated [195], with common adverse side effects including gastrointestinal symptoms (nausea, vomiting, abdominal pain, and diarrhea),

headaches, and depression [192, 196]. In the aforementioned 2-week RCT, no significant differences in side effects were seen between groups [193]. In summary, early findings from clinical studies of ibudilast for AUD treatment appear promising and warrant further clinical investigation.

2.2.8 Prazosin and Doxazosin

Prazosin and doxazosin are alpha-1 adrenergic receptor antagonists with similar chemical structures that can readily cross the blood-brain barrier and block noradrenergic excitation of the mesolimbic dopaminergic system [197, 198]. Both medications are used for the treatment of hypertension and benign prostatic hyperplasia. While these medications show good safety and tolerability, doxazosin appears to have a better clinical profile, such as improved absorption profile and a longer half-life, leading to less frequent dosing. Adrenergic receptors regulate sympathetic nervous system activity through activation of the neurotransmitter norepinephrine [199]. Stress physiology is disrupted with chronic alcohol use, particularly during early alcohol abstinence. In early abstinence, individuals with AUD experience more emotional dysregulation, stress, and alcohol cravings, all of which can increase the risk of relapse. Thus, alpha-1 blockers like prazosin and doxazosin may help to normalize these stress system changes seen in AUD [200].

Preclinical work found that prazosin reduced ethanol-related operant responding in acute withdrawal for dependent, but not non-dependent rodents [201]. In addition, prazosin treatment prevented yohimbine stress-induced reinstatement of ethanol seeking [202] and attenuated ethanol intake during relapse in alcohol-preferring rats [203]. Doxazosin decreased voluntary alcohol intake in alcohol-preferring rats in a dose-dependent manner, and this effect was likely not due to general motor impairment [204]. Further, doxazosin significantly reduced voluntary ethanol intake in a rodent model of AUD and stress exposure [205].

In humans, an early 6-week pilot study with 24 randomized participants with AUD revealed that prazosin treatment was associated with fewer drinking days per week and fewer drinks per week than placebo [206]. More recently, 92 participants with AUD completed a 12-week double-blind study with prazosin and medication management [207]. Results from intent-to-treat analyses showed that prazosin participants had greater reductions in heavy drinking and rates of drinking over time, although these effects were modest. A 10-week RCT comprising 41 individuals with AUD showed no significant effect of doxazosin over placebo on quantity of alcohol consumption (i.e., drinks per week, heavy-drinking days) [208]. However, further examination of clinical moderators revealed that doxazosin significantly reduced drinks per week and heavy-drinking days among individuals with higher family history of AUD and higher

standing diastolic blood pressure [209]. Similarly, moderator analyses from a 12-week RCT of prazosin ($N = 100$) found that one's degree of alcohol withdrawal symptoms predicted clinical response, such that participants with high levels of withdrawal symptoms benefited the most from treatment (e.g., reduced craving and heavy-drinking days; improved mood symptoms) [210]. A small meta-analysis of 6 studies with a total of 319 participants tested the effectiveness of drugs acting on adrenergic receptors for AUD and found a significant treatment effect of prazosin and doxazosin on alcohol consumption but not abstinence [211]. In summary, prazosin and doxazosin show some early promise as a treatment for AUD and may be particularly beneficial as a harm-reduction approach and for individuals with significant alcohol withdrawal symptoms or family history of AUD as well as comorbid post-traumatic stress disorder (PTSD).

2.2.9 N-Acetylcysteine

Alcohol use has been shown to alter the glutamate system. Chronic and binge drinking inhibit glutamate levels and transmission through blockade of NMDA receptors, subsequently leading to elevated glutamate levels during alcohol withdrawal [212–214]. *N*-Acetylcysteine (NAC) is a cysteine prodrug that works to restore glutamatergic tone in reward circuitry by improving the expression and function of the cysteine-glutamate exchanger and normalizing glial glutamate transporters [215, 216].

Preclinically, NAC has been shown to reduce withdrawal effects [217], block the development of behavioral sensitization to alcohol [218], and attenuate biological adaptations induced by alcohol cessation (i.e., blocked the reduction of transcription factor Δ FosB in the nucleus accumbens; reduced corticosterone and leptin levels) [218–220]. Additionally, NAC reduced ethanol-seeking and self-administration in rodents [215, 216]. NAC and aspirin co-administration also reduced ethanol intake and relapse binge drinking in ethanol-preferring rats [221].

In a recent meta-analysis of seven RCTs with a total of 245 participants, NAC compared to placebo was shown to reduce craving symptoms across a number of substance use disorders [222]. A secondary analysis of a 12-week, multisite RCT of NAC to treat cannabis use disorder in 277 participants found that NAC increased odds of between-visit abstinence, and reduced alcohol consumption (i.e., drinks per week and drinking days per week) by 30% [223]. However, a recent 5-day human laboratory study ($N = 9$) found that NAC did not attenuate alcohol self-administration [212]. Additional clinical trials will also examine the effectiveness of NAC in adolescent and adult samples of AUD ([224], NCT03707951). These trials include samples with comorbid psychopathology (i.e., PTSD) and will use neuroimaging

methods to examine the neural circuitry underlying NAC's modulation of relevant metabolites and neural reactivity to alcohol cues.

Orally-administered NAC appears to be well tolerated, with the most common adverse effects being nausea and diarrhea [212, 222]. Of note, NAC is currently being tested in adolescent AUD, a unique prospect as no other AUD pharmacotherapies are yet approved for adolescents [225]. In summary, NAC represents a promising potential treatment for AUD and merits further exploration.

2.2.10 Suvorexant

Suvorexant is a dual orexin antagonist that is used for the treatment of insomnia. The orexin/hypocretin system is well known for its role in sleep-wake regulation but has more recently been implicated in AUD [226]. Orexins are neuropeptides that are densely localized in the lateral hypothalamus. Orexins A and B bind to the G-protein coupled orexin-1 and orexin-2 receptors. Orexin A has equal affinity for both receptors, whereas orexin B has selectivity for orexin-2 receptors. The dense orexin projections from the lateral hypothalamus to the ventral tegmental area provide neurobiological support that orexins may influence responses to rewarding stimuli, including alcohol [227, 228].

Animal models demonstrate that orexin-1 receptor antagonists reduce alcohol drinking in alcohol-dependent mice and alcohol-preferring rats [229–233]. Orexin-2 antagonists also reduce alcohol drinking and reinstatement/relapse in mice and rats [234–236]. Similarly, dual orexin antagonists reduce alcohol consumption in alcohol-preferring rats [237, 238]. Given the effectiveness of orexin antagonists in reducing alcohol drinking at the preclinical level of analysis, suvorexant has garnered interest as a drug that can be repurposed to treat AUD [239]. While no animal or human laboratory study has directly examined the efficacy of suvorexant on alcohol-related behaviors, two ongoing Phase 2 RCTs will assess suvorexant's potential as a treatment for AUD (NCT04229095) and comorbid AUD + insomnia (NCT03897062). The sedative effects of suvorexant are of notable concern, particularly if individuals engage in alcohol drinking during treatment [240]. Thus, the level of patient monitoring throughout the treatment phase and finding the optimal time of dosing to negate the additive sedative effects are important factors to consider to fully evaluate its therapeutic potential.

2.3 Novel Agents

The following sections discuss novel agents that have not yet been approved for any use. These medications are early in development and the majority do not yet have available

clinical findings on their application toward the treatment of AUD.

2.3.1 ABT-436

ABT-436 is an orally active, highly selective vasopressin type 1B (V1B) receptor antagonist. V1B antagonists attenuate basal hypothalamic-pituitary-adrenal (HPA) axis activity and have shown favorable effects in rat models of alcohol dependence, including attenuating reinstatement of alcohol self-administration [241], and diminishing alcohol intake by alcohol-preferring and alcohol-dependent rats [241, 242].

ABT-436 has been shown to attenuate basal HPA axis activity in humans [243]. A 12-week RCT enrolling 150 participants found ABT-436 to be associated with an increased percentage of days of abstinence compared to placebo, as well as significantly reduced cigarette use [244]. However, no differences were found on heavy-drinking days or alcohol craving. A subgroup analysis also showed that ABT-436 appeared to have greater efficacy among participants with high baseline stress levels. This study was the first Phase 2 clinical trial that tested a V1B receptor antagonist for AUD.

The finding that ABT-436 reduced both alcohol drinking and smoking indicate a potential use for V1B antagonists to treat co-use of alcohol and nicotine. ABT-436 is generally well tolerated in humans, with the most common side effect being diarrhea. Furthermore, results indicate that patients with high levels of stress may specifically benefit from medications targeting the vasopressin receptor.

2.3.2 GET73

N-[(4-Trifluoromethyl) benzyl] 4-methoxybutyramide (GET73) is a GHB analogue that has shown promising in vitro and in vivo preclinical results as a potential agent for the treatment of AUD. The addition of GET73 to cultures of rat hippocampal neurons rescued negative ethanol-induced effects, reductions in cell viability, and increases in reactive oxygen species production, providing evidence for a neuroprotective role of GET73 as an AUD treatment [245]. In vivo, GET73 treatment at low, non-sedative doses (5–50 mg/kg) reduced alcohol intake and suppressed relapse in alcohol-preferring rats, as well as exerting anxiolytic effects [246]. These effects are similar to those seen with GHB administration [247]; however, GET73 was shown not to bind to either high- or low-affinity GHB binding sites in rat cortical membranes [246]. Recent studies indicate that GET73 may act as a negative allosteric modulator at the metabotropic glutamate subtype 5 receptor (mGluR5) [248]; however, the complete mechanism of action of GET73 remains unclear.

A translational study examining GET73-alcohol interactions found that neither 30 nor 100 mg/kg GET73

administered in rats (equivalent to doses employed in humans) potentiated alcohol-induced intoxication. Additionally, GET73 administered both in the presence and absence of alcohol was well tolerated in two samples of 14 and 11 participants, with no severe adverse events and no difference in adverse events [249]. A Phase I clinical trial in two samples of 48 and 32 participants found that both single doses (up to 600 mg) and repeated ascending doses (up to 450 mg twice /day) of GET-73 were safe and well tolerated [250]. Another inpatient human laboratory study conducted by the same group confirmed the safety and tolerability of GET73 such that no serious or severe adverse events occurred when GET73 was co-administered with alcohol. Co-administration also did not affect the pharmacokinetics of either GET73 or alcohol. However, GET73 also had no effect on alcohol cue-induced craving or self-administration, warranting additional research [251]. A clinical trial of the effects of GET73 on magnetic resonance spectroscopy (MRS) measures of glutamate and GABA levels in individuals with AUD was recently completed (NCT03418623); however, the results have not yet been posted, and another trial, which includes a free-drinking bar lab component, is ongoing (NCT04831684).

2.3.3 ASP8062

In comparison to the drawbacks presented by GABA_B agonists like baclofen and SMO, positive allosteric modulators (PAMs) at this receptor present a potential alternative [252]. By binding to a different, non-competitive allosteric site on the GABA_B receptor, PAMs allow endogenous GABA, binding at its original orthosteric site, to retain its potency and efficacy, reducing the risk of tolerance development and side effects [253, 254]. A number of PAMs have been studied as potential pharmacotherapies for AUD, and have demonstrated reductions in alcohol-associated behaviors and ethanol self-administration in preclinical models [252, 255–258]. When directly contrasted against baclofen, a GABA_B PAM had a better profile, with dose-dependent reduction of relapse-like alcohol drinking and with no signs of sedation [257].

One such novel GABA_B PAM, ASP8062, appears particularly promising as it has been shown to significantly increase the affinity and efficacy of endogenous GABA binding in human and rat GABA_B receptors in vitro and with oral administration in an in vivo rodent model of fibromyalgia, demonstrating the ability of oral formulated ASP8062 to cross the blood-brain-barrier [259]. ASP8062 has recently progressed to clinical development. Two Phase I clinical trials with a combined total of 112 participants evaluated single ascending doses and multiple ascending doses of ASP8062, respectively [260]. These studies found that ASP8062 was

well tolerated in humans, with no evidence of drug effects on safety, cognition, drug withdrawal, or suicidal ideation. One additional clinical trial, assessing the safety and efficacy of ASP8062 for alcohol use disorder (NCT05096117), is currently underway. Overall, ASP8062, and GABA_B PAMs in general, appear to be well tolerated in humans and decrease alcohol self-administration in animals. These agents present a potential pathway to better utilize the GABAergic system and reduce the side effects seen with GABA_B agonists.

2.3.4 PF-5190457

Ghrelin, a peptide produced by endocrine cells primarily in the stomach [261], is thought to regulate growth hormone secretion, food intake, and glucose homeostasis [262]. Ghrelin is also thought to play a role in AUD. Ghrelin signaling is required for stimulation of the reward system by alcohol, and higher ghrelin levels are associated with higher self-reported measures of alcohol craving [263]. In human laboratory studies, intravenous ghrelin administration has been shown to increase the urge to drink, increase alcohol self-administration, and modulate brain activity in regions involved in reward processing and stress regulation [264, 265].

Preclinical studies with ghrelin receptor (GHS-R1a) antagonists have shown reductions in alcohol conditioned place preference, alcohol intake and preference, and alcohol-elicited nucleus accumbens dopamine release in rodents [266–270].

PF-5190457 is a ghrelin receptor inverse agonist that inhibits GHS-R1a constitutive activity as well as blocking its activation by ghrelin [271]. In a preliminary clinical study in 12 heavy-drinking individuals, PF-5190457, compared to placebo, reduced alcohol craving and cue-reactivity to alcohol. Additionally, when administered in combination with alcohol, PF-5190457 was safe and well-tolerated with no drug-alcohol interactions [272]. This was the first clinical study of a GHS-R1a inverse agonist in a sample of heavy alcohol drinkers.

PF-5190457 may increase somnolence, heart rate, and lower blood glucose concentrations [273], although clinical results indicate that these side effects were not exacerbated by alcohol co-administration and in general PF-5190457 is well tolerated. In summary, preclinical and early clinical evidence support additional research toward investigating PF-5190457 as a pharmacological approach to treat AUD.

2.3.5 Cannabidiol

Cannabidiol (CBD), one of the main compounds found in *Cannabis sativa*, has shown promise as a novel therapeutic to treat AUD. CBD is nonintoxicating and has diverse pharmacological effects throughout the central nervous system, including functioning as a negative allosteric modulator of

CB₁ and CB₂ receptors [274, 275], and blocking anandamide uptake and inhibiting enzymatic hydrolysis [276]. CBD may also interact with non-endocannabinoid signaling systems, including the serotonergic system [277] and the opioidergic system [278], among others.

Preclinical studies have shown that CBD reduces alcohol administration, decreases motivation for alcohol, reduces relapse-like behavior, and improves withdrawal symptoms in animals exposed to chronic alcohol [279]. Evidence in healthy individuals demonstrates that CBD is well tolerated, does not interact with the subjective effects of alcohol, and has no abuse liability [280]. Two recent studies investigated signals for potential efficacy of CBD as a treatment for heroin use disorder [281] and cannabis use disorder [282]. Regarding heroin use disorder, acute CBD reduced cue-induced craving for heroin and reduced anxiety in a sample of 42 abstinent individuals, which was maintained one-week following the last CBD exposure. The cannabis use disorder study found that CBD was more efficacious at reducing cannabis use than placebo in a sample of 48 subjects. In both clinical samples, CBD was well tolerated and not associated with serious adverse events. CBD is currently being evaluated as a potential treatment for AUD, AUD with comorbid PTSD, and alcohol withdrawal in AUD in three clinical trials (NCT03252756, NCT03248167, NCT04205682). In brief, preclinical evidence and clinical evidence in other substance use disorders indicate the promise of CBD as a novel therapeutic for AUD.

3 Discussion

This qualitative literature review discusses the efficacy, mechanism of action, and tolerability of approved, repurposed, and novel pharmacotherapies for the treatment of AUD. This information is summarized in Table 1. As of 2018, the APA recommends acamprosate and naltrexone for the treatment of AUD and suggests gabapentin and topiramate for patients with the goal of reducing alcohol consumption or achieving abstinence, while disulfiram is suggested for achieving and maintaining abstinence only [107]. Similarly, while not included in the APA's recommendations, aripiprazole and mifepristone are associated with drinking reduction, while baclofen shows association with abstinence and mixed results with drinking reduction. Additional repurposed medications show clinical effectiveness for the treatment of AUD. Some of these appear to have particular promise in specific cases, such as varenicline's use for nicotine and alcohol co-users [147], baclofen for individuals with liver disease [93], and aripiprazole for more impulsive individuals [156]. Novel agents such as GET73 [245] and ASP8062 [260] have also reduced alcohol intake in preclinical studies. In summary, while currently approved

medications are somewhat effective, there remains a crucial need to develop new and improved pharmacotherapies for AUD. Novel and repurposed agents show significant promise as treatments that may improve upon currently approved pharmacotherapies.

Medication development has been identified as a critical priority for AUD research [283]. While considerable progress has been made in this field, there are a number of areas which require our attention to realize the benefit of AUD pharmacotherapy. First, despite the prevalence of AUD, the rate of seeking treatment for AUD remains very low [284]. In order for anyone to benefit from the advances in medication development reviewed herein, the treatment gap must be closed. This will require engagement at multiple levels, from prevention to public education about AUD and the available treatments. Researchers and clinicians can help in these efforts by reducing stigma surrounding AUD and other substance use disorders by choosing appropriate language to describe these disorders and the people who are affected by them [285].

A related issue is the need to improve access to FDA-approved pharmacotherapies for AUD. A recent analysis of the 2019 National Survey on Drug Use and Health found that only 1.6% of people with a past-year AUD received an evidence-based medication to treat their AUD [14]. Medication use was associated with living in a large metropolitan area, use of the emergency department, and receiving mental health care, indicating that these services and residing in an urban environment appear to increase access to evidence-based medications. There is also a great need to improve the education of physicians and clinicians on the availability of evidence-based treatments for AUD. Ongoing efforts in this area are underway by the American Society of Addiction Medicine and the American Academy of Addiction Psychiatry, as well as by the National Institute on Alcohol Abuse and Alcoholism.

To further improve access to treatments and increase treatment-seeking, there is a need to increase the menu of approved pharmacological treatments towards AUD, especially those that have shown promise internationally. Currently, the FDA only accepts two primary outcomes for Phase 3 trials: abstinence and no heavy-drinking days. These outcomes do not always mirror the goals of patients with AUD for their recovery, which may be better reflected by a harm reduction endpoint. Recent work has found that harm reduction endpoints, including reductions in WHO-based drinking levels [286], are associated with improvements in physical health and quality of life [287] and can be used as efficacy outcomes in clinical trials [288]. The acceptance of these outcomes as clinical trial endpoints could have a substantial impact on the medication development field and ultimately result in a larger pharmacotherapy toolbox for clinicians.

Another area of development is the move towards personalized treatment, also referred to as precision medication. AUD is a highly heterogeneous disorder, and it is unlikely that any medication will work for all individuals with an AUD. As such, there have been efforts to use precision medicine approaches to tailor pharmacotherapies to individuals with different presentations of AUD. Studies have taken several approaches towards personalized treatments, such as pharmacogenetics, sex differences, family history, severity of alcohol withdrawal, drinking phenotypes, and biobehavioral markers [283, 289–293]. However, even these efforts may be overly simplistic given the complex nature of AUD. It is likely that personalized treatment approaches will need to account for multiple factors to truly tailor treatments to individual patients [294]. Conversely, the public health significance of the improved efficacy of AUD pharmacotherapy with clinically accessible phenotypes argue for wider dissemination and implementation of precision treatment recommendations identified to date.

A final issue to consider is the need to develop treatments for individuals with AUD and comorbid psychiatric disorders and for individuals with AUD and AALD. AUD often co-occurs with other psychiatric disorders, including other substance use disorders, personality disorders, major depressive disorder, anxiety disorders, and PTSD [295]. The development of integrated treatments, including combined behavioral and pharmacological interventions, which simultaneously address AUD and other co-occurring disorders, is difficult due to the complexity of treating multiple disorders and the limited understanding of the underlying mechanisms. However, this is a necessary area of research given the high rates of comorbidity in the AUD population. Treatment options for individuals with AALD are limited; of the FDA-approved medications, only acamprosate can be used without concerns of hepatotoxicity [296]. Of the non-FDA medications that may prove useful in this population, only baclofen has been evaluated in an RCT [94]. There is a clear need to develop additional treatments for this population.

4 Conclusions

In conclusion, the three widely approved medications for AUD (i.e., disulfiram, acamprosate, and naltrexone) are both modestly effective and underutilized. The heterogeneity inherent in AUD also presents the issue that no one medication is likely to be effective for all individuals with AUD. Therefore, it is necessary to develop more effective, novel, and diverse pharmacotherapy options and to improve access to treatment. The repurposed and novel medications reviewed in this article demonstrate promise for the future of

AUD treatment. Additionally, the future directions indicated above offer potential improvements toward the advancement of precision medicine and personalized treatment for the heterogeneous AUD population.

Declarations

Funding This work was supported in part by the National Institute on Alcohol Abuse and Alcoholism (K24AA025704 to LAR; F31AA028976 to EMB; F32AA029288 to SJN).

Conflict of interest The authors declare they have no conflict of interest.

Author contributions This article was conceptualized by LAR, ENG, and EMB. Literature review and original draft preparation was conducted by EMB, ENG, SJN, and LRM. All authors reviewed and commented on previous versions of the manuscript, and all authors read and approved the final manuscript. All authors agree to be accountable for the work presented in the manuscript.

Ethics approval Not applicable.

Consent to participate Not applicable.

Consent for publication Not applicable.

Code availability Not applicable.

Availability of data and material Not applicable.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License, which permits any non-commercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc/4.0/>.

References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders [Internet]. 5th ed. American Psychiatric Association; 2013 [cited 2021 Mar 15]. <https://doi.org/10.1176/appi.books.9780890425596>.
2. Grant BF, Goldstein RB, Saha TD, Chou SP, Jung J, Zhang H, et al. Epidemiology of DSM-5 alcohol use disorder: results from the national epidemiologic survey on alcohol and related conditions III. *JAMA Psychiat*. 2015;72(8):757–66.
3. Degenhardt L, Charlson F, Ferrari A, Santomauro D, Erskine H, Mantilla-Herrera A, et al. The global burden of disease attributable to alcohol and drug use in 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Psychiatry*. 2018;5(12):987–1012.

4. World Health Organization. Global status report on alcohol and health 2014. World Health Organization; 2014.
5. Stahre M, Roeber J, Kanny D, Brewer RD, Zhang X. Contribution of excessive alcohol consumption to deaths and years of potential life lost in the United States. *Prev Chronic Dis*. 2014;26(11):E109.
6. Sacks JJ, Gonzales KR, Bouchery EE, Tomedi LE, Brewer RD. 2010 national and state costs of excessive alcohol consumption. *Am J Prev Med*. 2015;49(5):e73–9.
7. Hasin DS, Stinson FS, Ogburn E, Grant BF. Prevalence, correlates, disability, and comorbidity of DSM-IV alcohol abuse and dependence in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry*. 2007;64(7):830–42.
8. Rehm J, Shield KD, Gmel G, Rehm MX, Frick U. Modeling the impact of alcohol dependence on mortality burden and the effect of available treatment interventions in the European Union. *Eur Neuropsychopharmacol*. 2013;23(2):89–97.
9. Petrakis IL. A rational approach to the pharmacotherapy of alcohol dependence. *J Clin Psychopharmacol*. 2006;26(Suppl 1):S3–12.
10. Pettinati HM, Rabinowitz AR. Choosing the right medication for the treatment of alcoholism. *Curr Psychiatry Rep*. 2006;8(5):383–8.
11. Zindel LR, Kranzler HR. Pharmacotherapy of alcohol use disorders: seventy-five years of progress. *J Stud Alcohol Drugs Suppl*. 2014;75(Suppl 17):79–88.
12. Wallhed Finn S, Lundin A, Sjöqvist H, Danielsson A-K. Pharmacotherapy for alcohol use disorders—unequal provision across sociodemographic factors and co-morbid conditions. A cohort study of the total population in Sweden. *Drug Alcohol Depend*. 2021;227:108964.
13. Ray LA, Bujarski S, Grodin E, Hartwell E, Green R, Venegas A, et al. State-of-the-art behavioral and pharmacological treatments for alcohol use disorder. *Am J Drug Alcohol Abuse*. 2019;45(2):124–40.
14. Han B, Jones CM, Einstein EB, Powell PA, Compton WM. Use of medications for alcohol use disorder in the US: results from the 2019 national survey on drug use and health. *JAMA Psychiatry* [Internet]. 2021 [cited 2021 Jun 16]. <https://jamanetwork.com/journals/jamapsychiatry/fullarticle/2781290>.
15. Fan AZ, Chou SP, Zhang H, Jung J, Grant BF. Prevalence and correlates of past-year recovery from DSM-5 alcohol use disorder: results from national epidemiologic survey on alcohol and related conditions-III. *Alcohol Clin Exp Res*. 2019;43(11):2406–20.
16. Henssler J, Müller M, Carreira H, Bschor T, Heinz A, Baethge C. Controlled drinking—non-abstinent versus abstinent treatment goals in alcohol use disorder: a systematic review, meta-analysis and meta-regression. *Addiction*. 2021;116(8):1973–87.
17. Fentress TSP, Wald S, Brah A, Leemon G, Reyes R, Alkhamees F, et al. Dual study describing patient-driven harm reduction goal-setting among people experiencing homelessness and alcohol use disorder. *Exp Clin Psychopharmacol*. 2021;29(3):261–71.
18. Charlet K, Heinz A. Harm reduction—a systematic review on effects of alcohol reduction on physical and mental symptoms. *Addict Biol*. 2017;22(5):1119–59.
19. Marenmami I, Cibin M, Pani PP, Rossi A, Turchetti G. Harm Reduction as “Continuum Care” in alcohol abuse disorder. *Int J Environ Res Public Health*. 2015;12(11):14828–41.
20. Witkiewitz K, Morris J, Tucker JA. Commentary on Henssler et al.: the public health case for promoting and valuing drinking reductions in the treatment of alcohol use disorder. *Addiction*. 2021;116(8):1988–9.
21. Witkiewitz K, Wilson AD, Roos CR, Swan JE, Votaw VR, Stein ER, et al. Can individuals with alcohol use disorder sustain non-abstinent recovery? Non-abstinent outcomes 10 years after alcohol use disorder treatment. *J Addict Med*. 2021;15(4):303–10.
22. Mohapatra S, Rath NR. Disulfiram induced psychosis. *Clin Psychopharmacol Neurosci*. 2017;15(1):68–9.
23. Vallari RC, Pietruszko R. Human aldehyde dehydrogenase: mechanism of inhibition of disulfiram. *Science*. 1982;216(4546):637–9.
24. Fuller RK, Branchey L, Brightwell DR, Derman RM, Emrick CD, Iber FL, et al. Disulfiram treatment of alcoholism. A veterans administration cooperative study. *JAMA*. 1986;256(11):1449–55.
25. Garbutt JC, West SL, Carey TS, Lohr KN, Crews FT. Pharmacological treatment of alcohol dependence: a review of the evidence. *JAMA*. 1999;281(14):1318–25.
26. Kosten TR, Wu G, Huang W, Harding MJ, Hamon SC, Lapalain J, et al. Pharmacogenetic randomized trial for cocaine abuse: disulfiram and dopamine β -hydroxylase. *Biol Psychiatry*. 2013;73(3):219–24.
27. Devoto P, Flore G, Saba P, Cadeddu R, Gessa GL. Disulfiram stimulates dopamine release from noradrenergic terminals and potentiates cocaine-induced dopamine release in the prefrontal cortex. *Psychopharmacology*. 2012;219(4):1153–64.
28. Bagdy G, Arató M. Serum dopamine-beta-hydroxylase activity and alcohol withdrawal symptoms. *Drug Alcohol Depend*. 1987;19(1):45–50.
29. Skinner MD, Lahmek P, Pham H, Aubin H-J. Disulfiram efficacy in the treatment of alcohol dependence: a meta-analysis. *PLoS ONE*. 2014;9(2):e87366.
30. Skinner MD, Coudert M, Berlin I, Passeri E, Michel L, Aubin H-J. Effect of the threat of a disulfiram-ethanol reaction on cue reactivity in alcoholics. *Drug Alcohol Depend*. 2010;112(3):239–46.
31. Mutschler J, Dirican G, Funke S, Obermann C, Grosshans M, Mann K, et al. Experienced acetaldehyde reaction does not improve treatment response in outpatients treated with supervised disulfiram. *Clin Neuropharmacol*. 2011;34(4):161–5.
32. Mutschler J, Grosshans M, Soyka M, Rösner S. Current findings and mechanisms of action of disulfiram in the treatment of alcohol dependence. *Pharmacopsychiatry*. 2016;49(4):137–41.
33. Fuller RK, Gordis E. Does disulfiram have a role in alcoholism treatment today? *Addiction*. 2004;99(1):21–4.
34. Kranzler HR, Soyka M. Diagnosis and pharmacotherapy of alcohol use disorder: a review. *JAMA*. 2018;320(8):815–24.
35. Mason BJ, Heyser CJ. Acamprosate: a prototypic neuromodulator in the treatment of alcohol dependence. *CNS Neurol Disord Drug Targets*. 2010;9(1):23–32.
36. Jonas DE, Amick HR, Feltner C, Bobashev G, Thomas K, Wines R, et al. Pharmacotherapy for adults with alcohol use disorders in outpatient settings: a systematic review and meta-analysis. *JAMA*. 2014;311(18):1889–900.
37. Paille FM, Guelfi JD, Perkins AC, Royer RJ, Steru L, Parot P. Double-blind randomized multicentre trial of acamprosate in maintaining abstinence from alcohol. *Alcohol Alcohol*. 1995;30(2):239–47.
38. Chick J, Howlett H, Morgan MY, Ritson B. United Kingdom Multicentre Acamprosate Study (UKMAS): a 6-month prospective study of acamprosate versus placebo in preventing relapse after withdrawal from alcohol. *Alcohol Alcohol*. 2000;35(2):176–87.
39. Kampman KM, Pettinati HM, Lynch KG, Xie H, Dackis C, Oslin DW, et al. Initiating acamprosate within-detoxification versus post-detoxification in the treatment of alcohol dependence. *Addict Behav*. 2009;34(6–7):581–6.
40. Richardson K, Baillie A, Reid S, Morley K, Teesson M, Sanbale C, et al. Do acamprosate or naltrexone have an effect

- on daily drinking by reducing craving for alcohol? *Addiction*. 2008;103(6):953–9.
41. Mason BJ, Goodman AM, Chabac S, Leher P. Effect of oral acamprosate on abstinence in patients with alcohol dependence in a double-blind, placebo-controlled trial: the role of patient motivation. *J Psychiatr Res*. 2006;40(5):383–93.
 42. Anton RF, O'Malley SS, Ciraulo DA, Cisler RA, Couper D, Donovan DM, et al. Combined pharmacotherapies and behavioral interventions for alcohol dependence: the COMBINE study: a randomized controlled trial. *JAMA*. 2006;295(17):2003–17.
 43. Swift RM, Aston ER. Pharmacotherapy for alcohol use disorder: current and emerging therapies. *Harv Rev Psychiatry*. 2015;23(2):122–33.
 44. Rösner S, Hackl-Herrwerth A, Leucht S, Leher P, Vecchi S, Soyka M. Acamprosate for alcohol dependence. *Cochrane Database Syst Rev*. 2010;(9):CD004332.
 45. Littleton JM. Acamprosate in alcohol dependence: implications of a unique mechanism of action. *J Addict Med*. 2007;1(3):115–25.
 46. Koob GF, Mason BJ, De Witte P, Littleton J, Higgins GR. Potential neuroprotective effects of acamprosate. *Alcohol Clin Exp Res*. 2002;26(4):586–92.
 47. Davidson M, Shanley B, Wilce P. Increased NMDA-induced excitability during ethanol withdrawal: a behavioural and histological study. *Brain Res*. 1995;674(1):91–6.
 48. Grant KA, Valverius P, Hudspeth M, Tabakoff B. Ethanol withdrawal seizures and the NMDA receptor complex. *Eur J Pharmacol*. 1990;176(3):289–96.
 49. Rösner S, Leucht S, Leher P, Soyka M. Acamprosate supports abstinence, naltrexone prevents excessive drinking: evidence from a meta-analysis with unreported outcomes. *J Psychopharmacol*. 2008;22(1):11–23.
 50. Mitchell JM, O'Neil JP, Janabi M, Marks SM, Jagust WJ, Fields HL. Alcohol consumption induces endogenous opioid release in the human orbitofrontal cortex and nucleus accumbens. *Sci Transl Med*. 2012;4(116):116ra6.
 51. Nestler EJ. Is there a common molecular pathway for addiction? *Nat Neurosci*. 2005;8(11):1445–9.
 52. O'Malley SS, Jaffe AJ, Chang G, Schottenfeld RS, Meyer RE, Rounsaville B. Naltrexone and coping skills therapy for alcohol dependence. A controlled study. *Arch Gen Psychiatry*. 1992;49(11):881–7.
 53. Volpicelli JR, Alterman AI, Hayashida M, O'Brien CP. Naltrexone in the treatment of alcohol dependence. *Arch Gen Psychiatry*. 1992;49(11):876–80.
 54. Drobos DJ, Anton RF, Thomas SE, Voronin K. Effects of naltrexone and nalmefene on subjective response to alcohol among non-treatment-seeking alcoholics and social drinkers. *Alcohol Clin Exp Res*. 2004;28(9):1362–70.
 55. Ray LA, Hutchison KE. Effects of naltrexone on alcohol sensitivity and genetic moderators of medication response: a double-blind placebo-controlled study. *Arch Gen Psychiatry*. 2007;64(9):1069–77.
 56. O'Malley SS, Krishnan-Sarin S, Farren C, Sinha R, Kreek MJ. Naltrexone decreases craving and alcohol self-administration in alcohol-dependent subjects and activates the hypothalamo-pituitary-adrenocortical axis. *Psychopharmacology*. 2002;160(1):19–29.
 57. O'Malley SS, Corbin WR, Leeman RF, DeMartini KS, Fucito LM, Ikomi J, et al. Reduction of alcohol drinking in young adults by naltrexone: a double-blind, placebo-controlled, randomized clinical trial of efficacy and safety. *J Clin Psychiatry*. 2015;76(2):e207–213.
 58. Morris PL, Hopwood M, Whelan G, Gardiner J, Drummond E. Naltrexone for alcohol dependence: a randomized controlled trial. *Addiction*. 2001;96(11):1565–73.
 59. Bouza C, Angeles M, Magro A, Muñoz A, Amate JM. Efficacy and safety of naltrexone and acamprosate in the treatment of alcohol dependence: a systematic review. *Addiction*. 2004;99(7):811–28.
 60. Johnson BA. Naltrexone long-acting formulation in the treatment of alcohol dependence. *Ther Clin Risk Manag*. 2007;3(5):741–9.
 61. Garbutt JC, Kranzler HR, O'Malley SS, Gastfriend DR, Pettinati HM, Silverman BL, et al. Efficacy and tolerability of long-acting injectable naltrexone for alcohol dependence: a randomized controlled trial. *JAMA*. 2005;293(13):1617–25.
 62. Kranzler HR, Modesto-Lowe V, Van Kirk J. Naltrexone vs. nefazodone for treatment of alcohol dependence. A placebo-controlled trial. *Neuropsychopharmacology*. 2000;22(5):493–503.
 63. Streeton C, Whelan G. Naltrexone, a relapse prevention maintenance treatment of alcohol dependence: a meta-analysis of randomized controlled trials. *Alcohol Alcohol*. 2001;36(6):544–52.
 64. Mark TL, Kassed CA, Vandivort-Warren R, Levit KR, Kranzler HR. Alcohol and opioid dependence medications: prescription trends, overall and by physician specialty. *Drug Alcohol Depend*. 2009;99(1–3):345–9.
 65. Busch AC, Denduluri M, Glass J, Hetzel S, Guhani SP, Gassman M, et al. PredischARGE injectable versus oral naltrexone to improve postdischarge treatment engagement among hospitalized veterans with alcohol use disorder: a randomized pilot proof-of-concept study. *Alcohol Clin Exp Res*. 2017;41(7):1352–60.
 66. Rösner S, Hackl-Herrwerth A, Leucht S, Vecchi S, Srisuranant M, Soyka M. Opioid antagonists for alcohol dependence. *Cochrane Database Syst Rev*. 2010;(12):CD001867.
 67. McDonough M. Naltrexone and liver disease. *Aust Prescrip*. 2015;38(5):151.
 68. Bart G, Schluger JH, Borg L, Ho A, Bidlack JM, Kreek MJ. Nalmefene induced elevation in serum prolactin in normal human volunteers: partial kappa opioid agonist activity? *Neuropsychopharmacology*. 2005;30(12):2254–62.
 69. Rose JH, Karkhanis AN, Steiniger-Brach B, Jones SR. Distinct effects of nalmefene on dopamine uptake rates and kappa opioid receptor activity in the nucleus accumbens following chronic intermittent ethanol exposure. *Int J Mol Sci*. 2016;17(8):1216.
 70. Nealey KA, Smith AW, Davis SM, Smith DG, Walker BM. κ -opioid receptors are implicated in the increased potency of intra-accumbens nalmefene in ethanol-dependent rats. *Neuropharmacology*. 2011;61:35–42.
 71. Mann K, Bladström A, Torup L, Gual A, van den Brink W. Extending the treatment options in alcohol dependence: a randomized controlled study of as-needed nalmefene. *Biol Psychiatry*. 2013;73(8):706–13.
 72. van den Brink W, Aubin H-J, Bladström A, Torup L, Gual A, Mann K. Efficacy of as-needed nalmefene in alcohol-dependent patients with at least a high drinking risk level: results from a subgroup analysis of two randomized controlled 6-month studies. *Alcohol Alcohol*. 2013;48(5):570–8.
 73. Gual A, He Y, Torup L, van den Brink W, Mann K, ESENSE 2 Study Group. A randomised, double-blind, placebo-controlled, efficacy study of nalmefene, as-needed use, in patients with alcohol dependence. *Eur Neuropsychopharmacol*. 2013;23(11):1432–42.
 74. Naudet F, Fitzgerald N, Braillon A. Nalmefene for alcohol dependence: a NICE decision? *Lancet Psychiatry*. 2016;3(12):1104–5.
 75. Palpacuer C, Laviolle B, Boussageon R, Reymann JM, Bellissant E, Naudet F. Risks and benefits of nalmefene in the treatment of adult alcohol dependence: a systematic literature review and meta-analysis of published and unpublished double-blind randomized controlled trials. *PLoS Med*. 2015;12(12):e1001924.
 76. Johansen KGV, Tarp S, Astrup A, Lund H, Pagsberg AK, Christensen R. Harms associated with taking nalmefene for

- substance use and impulse control disorders: a systematic review and meta-analysis of randomised controlled trials. *PLoS ONE*. 2017;12(8):e0183821.
77. Palpacuer C, Duprez R, Huneau A, Locher C, Boussageon R, Laviolle B, et al. Pharmacologically controlled drinking in the treatment of alcohol dependence or alcohol use disorders: a systematic review with direct and network meta-analyses on nalmefene, naltrexone, acamprosate, baclofen and topiramate. *Addiction*. 2018;113(2):220–37.
 78. Naudet F, Braillon A. Baclofen and alcohol in France. *Lancet Psychiatry*. 2018;5(12):961–2.
 79. Agabio R, Sinclair JM, Addolorato G, Aubin H-J, Beraha EM, Caputo F, et al. Baclofen for the treatment of alcohol use disorder: the Cagliari statement. *Lancet Psychiatry*. 2018;5(12):957–60.
 80. Leggio L, Garbutt JC, Addolorato G. Effectiveness and safety of baclofen in the treatment of alcohol dependent patients. *CNS Neurol Disord Drug Targets*. 2010;9(1):33–44.
 81. Garbutt JC, Kampov-Polevoy AB, Gallop R, Kalka-Juhl L, Flannery BA. Efficacy and safety of baclofen for alcohol dependence: a randomized, double-blind, placebo-controlled trial. *Alcohol Clin Exp Res*. 2010;34(11):1849–57.
 82. Addolorato G, Caputo F, Capristo E, Domenicali M, Bernardi M, Janiri L, et al. Baclofen efficacy in reducing alcohol craving and intake: a preliminary double-blind randomized controlled study. *Alcohol Alcohol*. 2002;37(5):504–8.
 83. Morley KC, Baillie A, Leung S, Addolorato G, Leggio L, Haber PS. Baclofen for the treatment of alcohol dependence and possible role of comorbid anxiety. *Alcohol Alcohol*. 2014;49(6):654–60.
 84. Rose AK, Jones A. Baclofen: its effectiveness in reducing harmful drinking, craving, and negative mood. A meta-analysis. *Addiction*. 2018;113(8):1396–406.
 85. Pierce M, Sutherland A, Beraha EM, Morley K, van den Brink W. Efficacy, tolerability, and safety of low-dose and high-dose baclofen in the treatment of alcohol dependence: a systematic review and meta-analysis. *Eur Neuropsychopharmacol*. 2018;28(7):795–806.
 86. Minozzi S, Saulle R, Rösner S. Baclofen for alcohol use disorder. *Cochrane Database Syst Rev*. 2018;11:CD012557.
 87. Hauser P, Fuller B, Ho SB, Thuras P, Kern S, Dieperink E. The safety and efficacy of baclofen to reduce alcohol use in veterans with chronic hepatitis C: a randomized controlled trial. *Addiction*. 2017;112(7):1173–83.
 88. Reynaud M, Aubin H-J, Trinquet F, Zakine B, Dano C, Dematteis M, et al. A randomized, placebo-controlled study of high-dose baclofen in alcohol-dependent patients—the ALPADIR study. *Alcohol Alcohol*. 2017;52(4):439–46.
 89. Chartier M, Tannous S, Benturquia N, Labat L, Reis R, Risède P, et al. Baclofen-induced neuro-respiratory toxicity in the rat: contribution of tolerance and characterization of withdrawal syndrome. *Toxicol Sci*. 2018;164(1):153–65.
 90. Santos C, Olmedo RE. Sedative-hypnotic drug withdrawal syndrome: recognition and treatment. *Emerg Med Pract*. 2017;19(3):1–20.
 91. Garbutt JC, Kampov-Polevoy AB, Pedersen C, et al. Efficacy and tolerability of baclofen in a U.S. community population with alcohol use disorder: a dose-response, randomized, controlled trial. *Neuropsychopharmacol*. 2021;46:2250–6.
 92. Farokhnia M, Deschaine SL, Sadighi A, Farinelli LA, Lee MR, Akhlaghi F, et al. A deeper insight into how GABA-B receptor agonism via baclofen may affect alcohol seeking and consumption: lessons learned from a human laboratory investigation. *Mol Psychiatry*. 2021;26(2):545–55.
 93. Morley KC, Baillie A, Fraser I, Furneaux-Bate A, Dore G, Roberts M, et al. Baclofen in the treatment of alcohol dependence with or without liver disease: multisite, randomised, double-blind, placebo-controlled trial. *Br J Psychiatry*. 2018;212(6):362–9.
 94. Addolorato G, Leggio L, Ferrulli A, Cardone S, Vonghia L, Mirijello A, et al. Effectiveness and safety of baclofen for maintenance of alcohol abstinence in alcohol-dependent patients with liver cirrhosis: randomised, double-blind controlled study. *Lancet*. 2007;370(9603):1915–22.
 95. Leggio L, Litten RZ. The GABA-B receptor agonist baclofen helps patients with alcohol use disorder: why these findings matter. *Neuropsychopharmacol*. 2021;46:2228–9.
 96. Addolorato G, Lesch O-M, Maremmani I, Walter H, Nava F, Raffaillac Q, et al. Post-marketing and clinical safety experience with sodium oxybate for the treatment of alcohol withdrawal syndrome and maintenance of abstinence in alcohol-dependent subjects. *Expert Opin Drug Saf*. 2020;19(2):159–66.
 97. Skala K, Caputo F, Mirijello A, Vassallo G, Antonelli M, Ferrulli A, et al. Sodium oxybate in the treatment of alcohol dependence: from the alcohol withdrawal syndrome to the alcohol relapse prevention. *Expert Opin Pharmacother*. 2014;15(2):245–57.
 98. Busardò FP, Kyriakou C, Napoletano S, Marinelli E, Zaami S. Clinical applications of sodium oxybate (GHB): from narcolepsy to alcohol withdrawal syndrome. *Eur Rev Med Pharmacol Sci*. 2015;19(23):4654–63.
 99. Leone MA, Vigna-Taglianti F, Avanzi G, Brambilla R, Faggiano F. Gamma-hydroxybutyrate (GHB) for treatment of alcohol withdrawal and prevention of relapses. *Cochrane Database Syst Rev*. 2010;(2):CD006266.
 100. van den Brink W, Addolorato G, Aubin H-J, Benyamina A, Caputo F, Dematteis M, et al. Efficacy and safety of sodium oxybate in alcohol-dependent patients with a very high drinking risk level. *Addict Biol*. 2018;23(4):969–86.
 101. Caputo F, Francini S, Brambilla R, Vigna-Taglianti F, Stoppo M, Del Re A, et al. Sodium oxybate in maintaining alcohol abstinence in alcoholic patients with and without psychiatric comorbidity. *Eur Neuropsychopharmacol*. 2011;21(6):450–6.
 102. Mannucci C, Pichini S, Spagnolo EV, Calapai F, Gangemi S, Navarra M, et al. Sodium oxybate therapy for alcohol withdrawal syndrome and keeping of alcohol abstinence. *Curr Drug Metab*. 2018;19(13):1056–64.
 103. Maremmani AGI, Pani PP, Rovai L, Pacini M, Dell’Osso L, Maremmani I. Long-term γ -hydroxybutyric acid (GHB) and disulfiram combination therapy in GHB treatment-resistant chronic alcoholics. *Int J Environ Res Public Health*. 2011;8(7):2816–27.
 104. Caputo F, Maremmani AGI, Addolorato G, Domenicali M, Zoli G, D’Amore A, et al. Sodium oxybate plus nalmefene for the treatment of alcohol use disorder: a case series. *J Psychopharmacol*. 2016;30(4):402–9.
 105. Caputo F, Francini S, Stoppo M, Lorenzini F, Vignoli T, Del Re A, et al. Incidence of craving for and abuse of gamma-hydroxybutyric acid (GHB) in different populations of treated alcoholics: an open comparative study. *J Psychopharmacol*. 2009;23(8):883–90.
 106. Department of Defense. VA/DoD clinical practice guideline for the management of substance use disorders. 2015.
 107. Reus VI, Fochtmann LJ, Bukstein O, Eyler AE, Hilty DM, Horvitz-Lennon M, et al. The American psychiatric association practice guideline for the pharmacological treatment of patients with alcohol use disorder. *Am J Psychiatry*. 2018;175(1):86–90.
 108. Shalaby HN, El-Tanbouly DM, Zaki HF. Topiramate mitigates 3-nitropropionic acid-induced striatal neurotoxicity via modulation of AMPA receptors. *Food Chem Toxicol*. 2018;118:227–34.
 109. Huang Y-L, Peng M, Zhu G. Topiramate in the treatment of antipsychotic-induced hyperprolactinemia. *Med Hypotheses*. 2020;138:109607.

110. Kranzler HR, Covault J, Feinn R, Armeli S, Tennen H, Arias AJ, et al. Topiramate treatment for heavy drinkers: moderation by a GRIK1 polymorphism. *Am J Psychiatry*. 2014;171(4):445–52.
111. Zhang X, Velumian AA, Jones OT, Carlen PL. Modulation of high-voltage-activated calcium channels in dentate granule cells by topiramate. *Epilepsia*. 2000;41(S1):52–60.
112. Johnson BA, Ait-Daoud N, Bowden CL, DiClemente CC, Roache JD, Lawson K, et al. Oral topiramate for treatment of alcohol dependence: a randomised controlled trial. *Lancet*. 2003;361(9370):1677–85.
113. Hargreaves GA, McGregor IS. Topiramate moderately reduces the motivation to consume alcohol and has a marked antidepressant effect in rats. *Alcohol Clin Exp Res*. 2007;31(11):1900–7.
114. Knapp CM, Mercado M, Markley TL, Crosby S, Ciraulo DA, Kornetsky C. Zonisamide decreases ethanol intake in rats and mice. *Pharmacol Biochem Behav*. 2007;87(1):65–72.
115. Breslin FJ, Johnson BA, Lynch WJ. Effect of topiramate treatment on ethanol consumption in rats. *Psychopharmacology*. 2010;207(4):529–34.
116. Knapp CM, Ciraulo DA, Sarid-Segal O, Richardson MA, Devine E, Streeter CC, et al. Zonisamide, topiramate, and levetiracetam: efficacy and neuropsychological effects in alcohol use disorders. *J Clin Psychopharmacol*. 2015;35(1):34–42.
117. Johnson BA, Rosenthal N, Capece JA, Wiegand F, Mao L, Beyers K, et al. Topiramate for treating alcohol dependence: a randomized controlled trial. *JAMA*. 2007;298(14):1641–51.
118. Blodgett JC, Re ACD, Maisel NC, Finney JW. A meta-analysis of topiramate's effects for individuals with alcohol use disorders. *Alcohol Clin Exp Res*. 2014;38(6):1481–8.
119. Jose NA, Yadav P, Kapoor A, Mahla VP. Comparison between baclofen and topiramate in alcohol dependence: a prospective study. *Ind Psychiatry J*. 2019;28(1):44–50.
120. Kenna GA, Lomastro TL, Schiesl A, Leggio L, Swift RM. Review of topiramate: an antiepileptic for the treatment of alcohol dependence. *Curr Drug Abuse Rev*. 2009;2(2):135–42.
121. Olmsted CL, Kockler DR. Topiramate for alcohol dependence. *Ann Pharmacother*. 2008;42(10):1475–80.
122. Sills GJ. The mechanisms of action of gabapentin and pregabalin. *Curr Opin Pharmacol*. 2006;6(1):108–13.
123. Roberto M, Gilpin NW, O'Dell LE, Cruz MT, Morse AC, Siggins GR, et al. Cellular and behavioral interactions of gabapentin with alcohol dependence. *J Neurosci*. 2008;28(22):5762–71.
124. Besheer J, Frisbee S, Randall PA, Jaramillo AA, Masciello M. Gabapentin potentiates sensitivity to the interoceptive effects of alcohol and increases alcohol self-administration in rats. *Neuropharmacology*. 2016;101:216–24.
125. Ozburn AR, Metten P, Potretzke S, Townsley KG, Blednov YA, Crabbe JC. Effects of pharmacologically targeting neuroimmune pathways on alcohol drinking in mice selectively bred to drink to intoxication. *Alcohol Clin Exp Res*. 2020;44(2):553–66.
126. Furiere FA, Nakamura-Palacios EM. Gabapentin reduces alcohol consumption and craving: a randomized, double-blind, placebo-controlled trial. *J Clin Psychiatry*. 2007;68(11):1691–700.
127. Brower KJ, Myra Kim H, Strobbe S, Karam-Hage MA, Consens F, Zucker RA. A randomized double-blind pilot trial of gabapentin versus placebo to treat alcohol dependence and comorbid insomnia. *Alcohol Clin Exp Res*. 2008;32(8):1429–38.
128. Mason BJ, Light JM, Williams LD, Drobos DJ. Proof-of-concept human laboratory study for protracted abstinence in alcohol dependence: effects of gabapentin. *Addict Biol*. 2009;14(1):73–83.
129. Mason BJ, Quello S, Goodell V, Shadan F, Kyle M, Begovic A. Gabapentin treatment for alcohol dependence: a randomized clinical trial. *JAMA Intern Med*. 2014;174(1):70–7.
130. Anton RF, Myrick H, Baros AM, Latham PK, Randall PK, Wright TM, et al. Efficacy of a combination of flumazenil and gabapentin in the treatment of alcohol dependence: relationship to alcohol withdrawal symptoms. *J Clin Psychopharmacol*. 2009;29(4):334–42.
131. Anton RF, Myrick H, Wright TM, Latham PK, Baros AM, Waid LR, et al. Gabapentin combined with naltrexone for the treatment of alcohol dependence. *Am J Psychiatry*. 2011;168(7):709–17.
132. Pani PP, Trogu E, Pacini M, Maremmanni I. Anticonvulsants for alcohol dependence. *Cochrane Database Syst Rev*. 2014;13(2):CD008544.
133. Kranzler HR, Feinn R, Morris P, Hartwell EE. A meta-analysis of the efficacy of gabapentin for treating alcohol use disorder. *Addiction*. 2019;114(9):1547–55.
134. Anton RF, Latham P, Voronin K, Book S, Hoffman M, Prisciandaro J, et al. Efficacy of gabapentin for the treatment of alcohol use disorder in patients with alcohol withdrawal symptoms: a randomized clinical trial. *JAMA Intern Med*. 2020;180(5):728–36.
135. Falk DE, Ryan ML, Fertig JB, Devine EG, Cruz R, Brown ES, et al. Gabapentin enacarbil extended-release for alcohol use disorder: a randomized, double-blind, placebo-controlled, multisite trial assessing efficacy and safety. *Alcohol Clin Exp Res*. 2019;43(1):158–69.
136. Wiffen PJ, Derry S, Bell RF, Rice AS, Tölle TR, Phillips T, et al. Gabapentin for chronic neuropathic pain in adults. *Cochrane Database Syst Rev*. 2017;6:CD007938.
137. Smith RV, Havens JR, Walsh SL. Gabapentin misuse, abuse and diversion: a systematic review. *Addiction*. 2016;111(7):1160–74.
138. Steensland P, Simms JA, Holgate J, Richards JK, Bartlett SE. Varenicline, an $\alpha 4 \beta 2$ nicotinic acetylcholine receptor partial agonist, selectively decreases ethanol consumption and seeking. *Proc Natl Acad Sci USA*. 2007;104(30):12518–23.
139. Patkar OL, Belmer A, Tarren JR, Holgate JY, Bartlett SE. The effect of varenicline on binge-like ethanol consumption in mice is $\beta 4$ nicotinic acetylcholine receptor-independent. *Neurosci Lett*. 2016;28(633):235–9.
140. Kamens HM, Silva C, Peck C, Miller CN. Varenicline modulates ethanol and saccharin consumption in adolescent male and female C57BL/6J mice. *Brain Res Bull*. 2018;138:20–5.
141. Froehlich JC, Fischer SM, Dilley JE, Nicholson ER, Smith TN, Filosa NJ, et al. Combining varenicline (Chantix) with naltrexone decreases alcohol drinking more effectively than does either drug alone in a rodent model of alcoholism. *Alcohol Clin Exp Res*. 2016;40(9):1961–70.
142. Litten RZ, Ryan ML, Fertig JB, Falk DE, Johnson B, Dunn KE, et al. A double-blind, placebo-controlled trial assessing the efficacy of varenicline tartrate for alcohol dependence. *J Addict Med*. 2013;7(4):277–86.
143. de Bejczy A, Löf E, Walther L, Guterstam J, Hammarberg A, Asanovska G, et al. Varenicline for treatment of alcohol dependence: a randomized, placebo-controlled trial. *Alcohol Clin Exp Res*. 2015;39(11):2189–99.
144. Miranda R, O'Malley SS, Treloar Padovano H, Wu R, Falk DE, Ryan ML, et al. Effects of alcohol cue reactivity on subsequent treatment outcomes among treatment-seeking individuals with alcohol use disorder: a multisite randomized, double-blind, placebo-controlled clinical trial of varenicline. *Alcohol Clin Exp Res*. 2020;44(7):1431–43.
145. Anthony JC, Echeagaray-Wagner F. Epidemiologic analysis of alcohol and tobacco use. *Alcohol Res Health*. 2000;24(4):201–8.
146. McKee SA, Harrison ELR, O'Malley SS, Krishnan-Sarin S, Shi J, Tetrault JM, et al. Varenicline reduces alcohol self-administration in heavy-drinking smokers. *Biol Psychiatry*. 2009;66(2):185–90.
147. Falk DE, Castle I-JP, Ryan M, Fertig J, Litten RZ. Moderators of varenicline treatment effects in a double-blind,

- placebo-controlled trial for alcohol dependence: an exploratory analysis. *J Addict Med*. 2015;9(4):296–303.
148. Donato S, Green R, Ray LA. Alcohol use disorder severity moderates clinical response to varenicline. *Alcohol Clin Exp Res*. 2021;45(9):1877–87.
 149. O'Malley SS, Zweben A, Fucito LM, Wu R, Piepmeyer ME, Ockert DM, et al. Effect of varenicline combined with medical management on alcohol use disorder with comorbid cigarette smoking. *JAMA Psychiat*. 2018;75(2):129–38.
 150. Aripiprazole PE. Aripiprazole. *Am J Hosp Palliat Care*. 2017;34(2):180–5.
 151. Shibasaki M, Kurokawa K, Mizuno K, Ohkuma S. Effect of aripiprazole on anxiety associated with ethanol physical dependence and on ethanol-induced place preference. *J Pharmacol Sci*. 2012;118(2):215–24.
 152. Nirogi R, Kandikere V, Jayarajan P, Bhyrapuneni G, Saralaya R, Muddana N, et al. Aripiprazole in an animal model of chronic alcohol consumption and dopamine D₂ receptor occupancy in rats. *Am J Drug Alcohol Abuse*. 2013;39(2):72–9.
 153. Ingman K, Kupila J, Hyytiä P, Korpi ER. Effects of aripiprazole on alcohol intake in an animal model of high-alcohol drinking. *Alcohol Alcohol*. 2006;41(4):391–8.
 154. Kranzler HR, Covault J, Pierucci-Lagha A, Chan G, Douglas K, Arias AJ, et al. Effects of aripiprazole on subjective and physiological responses to alcohol. *Alcohol Clin Exp Res*. 2008;32(4):573–9.
 155. Myrick H, Li X, Randall PK, Henderson S, Voronin K, Anton RF. The effect of aripiprazole on cue-induced brain activation and drinking parameters in alcoholics. *J Clin Psychopharmacol*. 2010;30(4):365–72.
 156. Anton RF, Schacht JP, Voronin KE, Randall PK. Aripiprazole suppression of drinking in a clinical laboratory paradigm: influence of impulsivity and self-control. *Alcohol Clin Exp Res*. 2017;41(7):1370–80.
 157. Martinotti G, Di Nicola M, Di Giannantonio M, Janiri L. Aripiprazole in the treatment of patients with alcohol dependence: a double-blind, comparison trial vs. naltrexone. *J Psychopharmacol*. 2009;23(2):123–9.
 158. Kenna GA, Leggio L, Swift RM. A safety and tolerability laboratory study of the combination of aripiprazole and topiramate in volunteers who drink alcohol. *Hum Psychopharmacol*. 2009;24(6):465–72.
 159. Haass-Koffler CL, Goodyear K, Zywiak WH, Leggio L, Kenna GA, Swift RM. Comparing and combining topiramate and aripiprazole on alcohol-related outcomes in a human laboratory study. *Alcohol Alcohol*. 2018;53(3):268–76.
 160. Anton RF, Kranzler H, Breder C, Marcus RN, Carson WH, Han J. A randomized, multicenter, double-blind, placebo-controlled study of the efficacy and safety of aripiprazole for the treatment of alcohol dependence. *J Clin Psychopharmacol*. 2008;28(1):5–12.
 161. Heitzmann E, Javelot H, Weiner L, Michel B. A case of aripiprazole-induced tardive dyskinesia with dramatic evolution. *Case Rep Psychiatry*. 2016;2016:7031245.
 162. Macaluso M, Flynn A, Preskorn S. Determining whether a definitive causal relationship exists between aripiprazole and tardive dyskinesia and/or dystonia in patients with major depressive disorder, part 4: case report data. *J Psychiatr Pract*. 2016;22(3):203–20.
 163. Johnson BA, Ait-Daoud N. Neuropharmacological treatments for alcoholism: scientific basis and clinical findings. *Psychopharmacology*. 2000;149(4):327–44.
 164. Johnson BA. Update on neuropharmacological treatments for alcoholism: scientific basis and clinical findings. *Biochem Pharmacol*. 2008;75(1):34–56.
 165. Rodd ZA, Bell RL, Oster SM, Toalston JE, Pommer TJ, McBride WJ, et al. Serotonin-3 receptors in the posterior ventral tegmental area regulate ethanol self-administration of alcohol-preferring (P) rats. *Alcohol*. 2010;44(3):245–55.
 166. Campbell AD, McBride WJ. Serotonin-3 receptor and ethanol-stimulated dopamine release in the nucleus accumbens. *Pharmacol Biochem Behav*. 1995;51(4):835–42.
 167. Umathe SN, Bhutada PS, Raut VS, Jain NS, Mundhada YR. The 5-HT₃ receptor antagonist, ondansetron, blocks the development and expression of ethanol-induced locomotor sensitization in mice. *Behav Pharmacol*. 2009;20(1):78–83.
 168. Tomkins DM, Le AD, Sellers EM. Effect of the 5-HT₃ antagonist ondansetron on voluntary ethanol intake in rats and mice maintained on a limited access procedure. *Psychopharmacology*. 1995;117(4):479–85.
 169. Kostowski W, Bisaga A, Jankowska E, Krzaścik P. Studies on the effects of certain 5-HT₃ receptor antagonists on ethanol preference and withdrawal seizures in the rat. *Pol J Pharmacol*. 1994;46(3):133–7.
 170. Johnson BA, Ait-Daoud N, Prihoda TJ. Combining ondansetron and naltrexone effectively treats biologically predisposed alcoholics: from hypotheses to preliminary clinical evidence. *Alcohol Clin Exp Res*. 2000;24(5):737–42.
 171. Myrick H, Anton RF, Li X, Henderson S, Randall PK, Voronin K. Effect of naltrexone and ondansetron on alcohol cue-induced activation of the ventral striatum in alcohol-dependent people. *Arch Gen Psychiatry*. 2008;65(4):466–75.
 172. Johnson BA, Roache JD, Javors MA, DiClemente CC, Cloninger CR, Prihoda TJ, et al. Ondansetron for reduction of drinking among biologically predisposed alcoholic patients: a randomized controlled trial. *JAMA*. 2000;284(8):963–71.
 173. Kranzler HR, Pierucci-Lagha A, Feinn R, Hernandez-Avila C. Effects of ondansetron in early- versus late-onset alcoholics: a prospective, open-label study. *Alcohol Clin Exp Res*. 2003;27(7):1150–5.
 174. Johnson BA, Roache JD, Ait-Daoud N, Zanca NA, Velazquez M. Ondansetron reduces the craving of biologically predisposed alcoholics. *Psychopharmacology*. 2002;160(4):408–13.
 175. Roila F, Tonato M, Basurto C, Bracarda S, Sassi M, Lupattelli M, et al. Ondansetron. *Eur J Cancer*. 1993;29A(Suppl 1):S16–21.
 176. Shen WW. Anticraving therapy for alcohol use disorder: a clinical review. *Neuropsychopharmacol Rep*. 2018;38(3):105–16.
 177. Simms JA, Haass-Koffler CL, Bito-Onon J, Li R, Bartlett SE. Mifepristone in the central nucleus of the amygdala reduces yohimbine stress-induced reinstatement of ethanol-seeking. *Neuropsychopharmacology*. 2012;37(4):906–18.
 178. Vendruscolo LF, Estey D, Goodell V, Macshane LG, Logrip ML, Schlosburg JE, et al. Glucocorticoid receptor antagonism decreases alcohol seeking in alcohol-dependent individuals. *J Clin Invest*. 2015;125(8):3193–7.
 179. Jimenez VA, Walter NAR, Shnitko TA, Newman N, Diem K, Vanderhoof L, et al. Mifepristone decreases chronic voluntary ethanol consumption in rhesus macaques. *J Pharmacol Exp Ther*. 2020;375(2):258–67.
 180. Holtyn AF, Weerts EM. Evaluation of mifepristone effects on alcohol-seeking and self-administration in baboons. *Exp Clin Psychopharmacol*. 2019;27(3):227–35.
 181. Donoghue K, Rose A, Coulton S, Milward J, Reed K, Drummond C, et al. Double-blind, 12 month follow-up, placebo-controlled trial of mifepristone on cognition in alcoholics: the MIFCOG trial protocol. *BMC Psychiatry*. 2016;24(16):40.
 182. Donoghue K, Rose A, Coulton S, Coleman R, Milward J, Philips T, et al. Double-blind, placebo-controlled trial of mifepristone on cognition and depression in alcohol dependence. *Trials*. 2020;21(1):796.

183. Gibson LCD, Hastings SF, McPhee I, Clayton RA, Darroch CE, Mackenzie A, et al. The inhibitory profile of Ibudilast against the human phosphodiesterase enzyme family. *Eur J Pharmacol.* 2006;538(1–3):39–42.
184. Cho Y, Crichlow GV, Vermeire JJ, Leng L, Du X, Hodsdon ME, et al. Allosteric inhibition of macrophage migration inhibitory factor revealed by ibudilast. *Proc Natl Acad Sci USA.* 2010;107(25):11313–8.
185. Mizuno T, Kurotani T, Komatsu Y, Kawanokuchi J, Kato H, Mitsuma N, et al. Neuroprotective role of phosphodiesterase inhibitor ibudilast on neuronal cell death induced by activated microglia. *Neuropharmacology.* 2004;46(3):404–11.
186. Mayfield J, Harris RA. The neuroimmune basis of excessive alcohol consumption. *Neuropsychopharmacology.* 2017;42(1):376.
187. Johnson K, Matsuda K, Iwaki Y. Ibudilast for the treatment of drug addiction and other neurological conditions. *Clin Investig.* 2014;4(3):269–79.
188. Bell RL, Lopez MF, Cui C, Egli M, Johnson KW, Franklin KM, et al. Ibudilast reduces alcohol drinking in multiple animal models of alcohol dependence. *Addict Biol.* 2015;20(1):38–42.
189. Blednov YA, Benavidez JM, Black M, Harris RA. Inhibition of phosphodiesterase 4 reduces ethanol intake and preference in C57BL/6J mice. *Front Neurosci.* 2014;8:129.
190. Logrip ML, Vendruscolo LF, Schlosburg JE, Koob GF, Zorrilla EP. Phosphodiesterase 10A regulates alcohol and saccharin self-administration in rats. *Neuropsychopharmacology.* 2014;39(7):1722–31.
191. Wen R-T, Zhang M, Qin W-J, Liu Q, Wang W-P, Lawrence AJ, et al. The phosphodiesterase-4 (PDE4) inhibitor rolipram decreases ethanol seeking and consumption in alcohol-preferring Fawn-Hooded rats. *Alcohol Clin Exp Res.* 2012;36(12):2157–67.
192. Ray LA, Bujarski S, Shoptaw S, Roche DJ, Heinzerling K, Miotto K. Development of the neuroimmune modulator ibudilast for the treatment of alcoholism: a randomized, placebo-controlled, human laboratory trial. *Neuropsychopharmacology.* 2017;42(9):1776–88.
193. Grodin EN, Bujarski S, Towns B, Burnette E, Nieto S, Lim A, et al. Ibudilast, a neuroimmune modulator, reduces heavy drinking and alcohol cue-elicited neural activation: a randomized trial. *Transl Psychiatry.* 2021;11(1):355.
194. Burnette EM, Ray LA, Irwin MR, Grodin EN. Ibudilast attenuates alcohol cue-elicited frontostriatal functional connectivity in alcohol use disorder. *Alcohol Clin Exp Res.* 2021;45(10):2017–28.
195. Rolan P, Gibbons JA, He L, Chang E, Jones D, Gross MI, et al. Ibudilast in healthy volunteers: safety, tolerability and pharmacokinetics with single and multiple doses. *Br J Clin Pharmacol.* 2008;66(6):792–801.
196. Fox RJ, Coffey CS, Conwit R, Cudkowicz ME, Gleason T, Goodman A, et al. Phase 2 trial of ibudilast in progressive multiple sclerosis. *N Engl J Med.* 2018;379(9):846–55.
197. Sommermeier H, Frielingsdorf J, Knorr A. Effects of prazosin on the dopaminergic neurotransmission in rat brain. *Eur J Pharmacol.* 1995;276(3):267–70.
198. Leggio L, Kenna GA. Doxazosin for alcoholism. *Alcohol Clin Exp Res.* 2013;37(2):191–3.
199. Perez DM. α 1-Adrenergic receptors in neurotransmission, synaptic plasticity, and cognition. *Front Pharmacol.* 2020;11:581098.
200. Haass-Koffler CL, Swift RM, Leggio L. Noradrenergic targets for the treatment of alcohol use disorder. *Psychopharmacology.* 2018;235(6):1625–34.
201. Walker BM, Rasmussen DD, Raskind MA, Koob GF. α 1-noradrenergic receptor antagonism blocks dependence-induced increases in responding for ethanol. *Alcohol.* 2008;42(2):91–7.
202. Funk D, Coen K, Tamadon S, Li Z, Loughlin A, Lê AD. Effects of prazosin and doxazosin on yohimbine-induced reinstatement of alcohol seeking in rats. *Psychopharmacology.* 2016;233(11):2197–207.
203. Froehlich JC, Hausauer B, Fischer S, Wise B, Rasmussen DD. Prazosin reduces alcohol intake in an animal model of alcohol relapse. *Alcohol Clin Exp Res.* 2015;39(8):1538–46.
204. O'Neil ML, Beckwith LE, Kincaid CL, Rasmussen DD. The α 1-adrenergic receptor antagonist, doxazosin, reduces alcohol drinking in alcohol-preferring (P) rats. *Alcohol Clin Exp Res.* 2013;37(2):202–12.
205. Lopez MF, Reasons SE, Carper BA, Nolen TL, Williams RL, Becker HC. Evaluation of the effect of doxazosin and zonisamide on voluntary ethanol intake in mice that experienced chronic intermittent ethanol exposure and stress. *Alcohol.* 2020;89:37–42.
206. Simpson TL, Saxon AJ, Meredith CW, Malte CA, McBride B, Ferguson LC, et al. A pilot trial of the α -1 adrenergic antagonist, prazosin, for alcohol dependence. *Alcohol Clin Exp Res.* 2009;33(2):255–63.
207. Simpson TL, Saxon AJ, Stappenbeck C, Malte CA, Lyons R, Tell D, et al. Double-blind randomized clinical trial of the α -1 noradrenergic antagonist prazosin for alcohol use disorder. *Am J Psychiatry.* 2018;175(12):1216–24.
208. Kenna GA, Haass-Koffler CL, Zywiak WH, Edwards SM, Brickley MB, Swift RM, et al. Role of the α 1 blocker doxazosin in alcoholism: a proof-of-concept randomized controlled trial. *Addict Biol.* 2016;21(4):904–14.
209. Haass-Koffler CL, Goodyear K, Zywiak WH, Magill M, Eltinge SE, Wallace PM, et al. Higher pretreatment blood pressure is associated with greater alcohol drinking reduction in alcohol-dependent individuals treated with doxazosin. *Drug Alcohol Depend.* 2017;1(177):23–8.
210. Sinha R, Wemm S, Fogelman N, Milivojevic V, Morgan PM, Angarita GA, et al. Moderation of Prazosin's efficacy by alcohol withdrawal symptoms. *American Journal of Psychiatry [Internet].* 2020 [cited 2021 Sep 16]. <https://doi.org/10.1176/appi.ajp.2020.20050609>.
211. Vanderkam P, Solinas M, Ingrand I, Doux N, Ebrahimighavani S, Jaafari N, et al. Effectiveness of drugs acting on adrenergic receptors in the treatment for tobacco or alcohol use disorders: systematic review and meta-analysis. *Addiction.* 2021;116(5):1011–20.
212. Stoops WW, Strickland JC, Hays LR, Rayapati AO, Lile JA, Rush CR. Influence of n-acetylcysteine maintenance on the pharmacodynamic effects of oral ethanol. *Pharmacol Biochem Behav.* 2020;198:173037.
213. Chen X, Michaelis ML, Michaelis EK. Effects of chronic ethanol treatment on the expression of calcium transport carriers and NMDA/glutamate receptor proteins in brain synaptic membranes. *J Neurochem.* 1997;69(4):1559–69.
214. Dahchour A, De Witte P. Excitatory and inhibitory amino acid changes during repeated episodes of ethanol withdrawal: an in vivo microdialysis study. *Eur J Pharmacol.* 2003;459(2–3):171–8.
215. Lebourgeois S, González-Marín MC, Jeanblanc J, Naassila M, Vilpoux C. Effect of N-acetylcysteine on motivation, seeking and relapse to ethanol self-administration. *Addict Biol.* 2018;23(2):643–52.
216. Lebourgeois S, González-Marín MC, Antol J, Naassila M, Vilpoux C. Evaluation of N-acetylcysteine on ethanol self-administration in ethanol-dependent rats. *Neuropharmacology.* 2019;15(150):112–20.
217. Mocelin R, Marcon M, da Rosa Araujo AS, Herrmann AP, Piatto A. Withdrawal effects following repeated ethanol exposure are prevented by N-acetylcysteine in zebrafish. *Prog Neuropsychopharmacol Biol Psychiatry.* 2019;13(93):161–70.
218. Morais-Silva G, Alves GC, Marin MT. N-Acetylcysteine treatment blocks the development of ethanol-induced behavioural

- sensitization and related Δ FosB alterations. *Neuropharmacology*. 2016;110(Pt A):135–42.
219. Schneider R, Santos CF, Clarimundo V, Dalmaz C, Elisabetsky E, Gomez R. *N*-Acetylcysteine prevents behavioral and biochemical changes induced by alcohol cessation in rats. *Alcohol*. 2015;49(3):259–63.
 220. Schneider R, Bandiera S, Souza DG, Bellaver B, Caletti G, Quincozes-Santos A, et al. *N*-Acetylcysteine prevents alcohol related neuroinflammation in rats. *Neurochem Res*. 2017;42(8):2135–41.
 221. Israel Y, Quintanilla ME, Ezquer F, Morales P, Santapau D, Berrios-Cárcano P, Ezquer M, Olivares B, Herrera-Marschitz M. Aspirin and *N*-acetylcysteine co-administration markedly inhibit chronic ethanol intake and block relapse binge drinking: Role of neuroinflammation-oxidative stress self-perpetuation. *Addict Biol*. 2021;26(1):e12853.
 222. Duailibi MS, Cordeiro Q, Brietzke E, Ribeiro M, LaRowe S, Berk M, et al. *N*-Acetylcysteine in the treatment of craving in substance use disorders: Systematic review and meta-analysis. *Am J Addict*. 2017;26(7):660–6.
 223. Squeglia LM, Tomko RL, Baker NL, McClure EA, Book GA, Gray KM. The effect of *N*-acetylcysteine on alcohol use during a cannabis cessation trial. *Drug Alcohol Depend*. 2018;01(185):17–22.
 224. Back SE, Gray K, Ana ES, Jones JL, Jarnecke AM, Joseph JE, et al. *N*-Acetylcysteine for the treatment of comorbid alcohol use disorder and posttraumatic stress disorder: design and methodology of a randomized clinical trial. *Contemp Clin Trials*. 2020;91:105961.
 225. Hammond CJ. The role of pharmacotherapy in the treatment of adolescent substance use disorders. *Child Adolesc Psychiatr Clin N Am*. 2016;25(4):685–711.
 226. Walker LC, Ch'ng SS, Lawrence AJ. Role of lateral hypothalamic orexin (hypocretin) neurons in alcohol use and abuse: recent advances. *Curr Pharmacol Rep*. 2016;2(6):241–52.
 227. Korotkova TM, Sergeeva OA, Eriksson KS, Haas HL, Brown RE. Excitation of ventral tegmental area dopaminergic and nondopaminergic neurons by orexins/hypocretins. *J Neurosci*. 2003;23(1):7–11.
 228. Nakamura T, Uramura K, Nambu T, Yada T, Goto K, Yanagisawa M, et al. Orexin-induced hyperlocomotion and stereotypy are mediated by the dopaminergic system. *Brain Res*. 2000;873(1):181–7.
 229. Lei K, Kwok C, Darevsky D, Wegner SA, Yu J, Nakayama L, et al. Nucleus accumbens shell orexin-1 receptors are critical mediators of binge intake in excessive-drinking individuals. *Front Neurosci*. 2019;13:88.
 230. Moorman DE, James MH, Kilroy EA, Aston-Jones G. Orexin/hypocretin-1 receptor antagonism reduces ethanol self-administration and reinstatement selectively in highly-motivated rats. *Brain Res*. 2017;1654(Pt A):34–42.
 231. Brown RM, Kim AK, Khoo SY-S, Kim JH, Jupp B, Lawrence AJ. Orexin-1 receptor signalling in the prelimbic cortex and ventral tegmental area regulates cue-induced reinstatement of ethanol-seeking in iP rats. *Addict Biol*. 2016;21(3):603–12.
 232. Ubaldi M, Giordano A, Severi I, Li H, Kallupi M, de Guglielmo G, et al. Activation of hypocretin-1/orexin-a neurons projecting to the bed nucleus of the stria terminalis and paraventricular nucleus is critical for reinstatement of alcohol seeking by neuropeptide S. *Biol Psychiatry*. 2016;79(6):452–62.
 233. Lawrence AJ, Cowen MS, Yang H-J, Chen F, Oldfield B. The orexin system regulates alcohol-seeking in rats. *Br J Pharmacol*. 2006;148(6):752–9.
 234. Shoblock JR, Welty N, Aluisio L, Fraser I, Motley ST, Morton K, et al. Selective blockade of the orexin-2 receptor attenuates ethanol self-administration, place preference, and reinstatement. *Psychopharmacology*. 2011;215(1):191–203.
 235. Brown RM, Khoo SY-S, Lawrence AJ. Central orexin (hypocretin) 2 receptor antagonism reduces ethanol self-administration, but not cue-conditioned ethanol-seeking, in ethanol-preferring rats. *Int J Neuropsychopharmacol*. 2013;16(9):2067–79.
 236. Barson JR, Ho HT, Leibowitz SF. Anterior thalamic paraventricular nucleus is involved in intermittent access ethanol drinking: role of orexin receptor 2. *Addict Biol*. 2015;20(3):469–81.
 237. Anderson RI, Becker HC, Adams BL, Jesudason CD, Rorick-Kehn LM. Orexin-1 and orexin-2 receptor antagonists reduce ethanol self-administration in high-drinking rodent models. *Front Neurosci*. 2014;8:33.
 238. Srinivasan S, Simms JA, Nielsen CK, Lieske SP, Bito-Onon JJ, Yi H, et al. The dual orexin/hypocretin receptor antagonist, almorexant, in the ventral tegmental area attenuates ethanol self-administration. *PLoS ONE*. 2012;7(9):e44726.
 239. Campbell EJ, Norman A, Bonomo Y, Lawrence AJ. Suvorexant to treat alcohol use disorder and comorbid insomnia: plan for a phase II trial. *Brain Res*. 2020;1(1728):146597.
 240. Sun H, Yee KL, Gill S, Liu W, Li X, Panebianco D, et al. Psychomotor effects, pharmacokinetics and safety of the orexin receptor antagonist suvorexant administered in combination with alcohol in healthy subjects. *J Psychopharmacol*. 2015;29(11):1159–69.
 241. Zhou Y, Colombo G, Carai MAM, Ho A, Gessa GL, Kreek MJ. Involvement of arginine vasopressin and V1b receptor in alcohol drinking in Sardinian alcohol-preferring rats. *Alcohol Clin Exp Res*. 2011;35(10):1876–83.
 242. Edwards S, Guerrero M, Ghoneim OM, Roberts E, Koob GF. Evidence that vasopressin V1b receptors mediate the transition to excessive drinking in ethanol-dependent rats. *Addict Biol*. 2012;17(1):76–85.
 243. Katz DA, Locke C, Liu W, Zhang J, Achari R, Wesnes KA, et al. Single-dose interaction study of the arginine vasopressin type 1B receptor antagonist ABT-436 and alcohol in moderate alcohol drinkers. *Alcohol Clin Exp Res*. 2016;40(4):838–45.
 244. Ryan ML, Falk DE, Fertig JB, Rendenbach-Mueller B, Katz DA, Tracy KA, et al. A phase 2, double-blind, placebo-controlled randomized trial assessing the efficacy of ABT-436, a novel V1b receptor antagonist, for alcohol dependence. *Neuropsychopharmacology*. 2017;42(5):1012–23.
 245. Tomasini MC, Borelli AC, Beggiato S, Tanganelli S, Loche A, Cacciaglia R, et al. GET73 prevents ethanol-induced neurotoxicity in primary cultures of rat hippocampal neurons. *Alcohol Alcohol*. 2016;51(2):128–35.
 246. Loche A, Simonetti F, Lobina C, Carai MAM, Colombo G, Castelli MP, et al. Anti-alcohol and anxiolytic properties of a new chemical entity, GET73. *Front Psychiatry*. 2012;3:8.
 247. Colombo G, Carai MAM, Gessa GL. Suppression by γ -hydroxybutyric acid of “alcohol deprivation effect” in rats: pre-clinical evidence of its anti-relapse properties. *Front Psychiatry*. 2012;3:95.
 248. Beggiato S, Borelli AC, Tomasini MC, Castelli MP, Pintori N, Cacciaglia R, et al. In vitro functional characterization of GET73 as possible negative allosteric modulator of metabotropic glutamate receptor 5. *Front Pharmacol*. 2018;9:327.
 249. Haass-Koffler CL, Goodyear K, Loche A, Long VM, Lobina C, Tran HH, et al. Administration of the metabotropic glutamate receptor subtype 5 allosteric modulator GET 73 with alcohol: a translational study in rats and humans. *J Psychopharmacol*. 2018;32(2):163–73.
 250. Haass-Koffler CL, Goodyear K, Long VM, Tran HH, Loche A, Cacciaglia R, et al. A phase I randomized clinical trial testing the safety, tolerability and preliminary pharmacokinetics of the mGluR5 negative allosteric modulator GET 73 following

- single and repeated doses in healthy volunteers. *Eur J Pharm Sci*. 2017;15(109):78–85.
251. Haass-Koffler CL, Perciballi R, Magill M, et al. An inpatient human laboratory study assessing the safety and tolerability, pharmacokinetics, and biobehavioral effect of GET 73 when co-administered with alcohol in individuals with alcohol use disorder. *Psychopharmacology (Berl)*. 2022;239:35–46.
 252. Maccioni P, Colombo G. Potential of gabab receptor positive allosteric modulators in the treatment of alcohol use disorder. *CNS Drugs*. 2019;33(2):107–23.
 253. Urwyler S. Allosteric modulation of family C G-protein-coupled receptors: from molecular insights to therapeutic perspectives. *Pharmacol Rev*. 2011;63(1):59–126.
 254. Gjoni T, Desrayaud S, Imobersteg S, Urwyler S. The positive allosteric modulator GS39783 enhances GABA(B) receptor-mediated inhibition of cyclic AMP formation in rat striatum in vivo. *J Neurochem*. 2006;96(5):1416–22.
 255. Agabio R, Colombo G. GABAB receptor ligands for the treatment of alcohol use disorder: preclinical and clinical evidence. *Front Neurosci*. 2014;8:140.
 256. Augier E, Dulman RS, Damadzic R, Pilling A, Hamilton JP, Heilig M. The GABAB positive allosteric modulator ADX71441 attenuates alcohol self-administration and relapse to alcohol seeking in rats. *Neuropsychopharmacology*. 2017;42(9):1789–99.
 257. Vengeliene V, Takahashi TT, Dravolina OA, Belozertseva I, Zvartau E, Besspalov AY, et al. Efficacy and side effects of baclofen and the novel GABAB receptor positive allosteric modulator CMPPE in animal models for alcohol and cocaine addiction. *Psychopharmacology*. 2018;235(7):1955–65.
 258. Maccioni P, Colombo G, Lorrai I, Zaru A, Carai MAM, Gessa GL, et al. Suppressing effect of COR659 on alcohol, sucrose, and chocolate self-administration in rats: involvement of the GABAB and cannabinoid CB1 receptors. *Psychopharmacology*. 2017;234(17):2525–43.
 259. Murai N, Kondo Y, Akuzawa S, Mihara T, Shiraishi N, Nakimoto S, et al. A novel GABAB receptor positive allosteric modulator, ASP8062, exerts analgesic effects in a rat model of fibromyalgia. *Eur J Pharmacol*. 2019;865:172750.
 260. Walzer M, Marek GJ, Wu R, Nagata M, Han D. Single- and multiple-dose safety, tolerability, and pharmacokinetic profiles of ASP8062: results from 2 phase 1 studies. *Clin Pharmacol Drug Dev*. 2020;9(3):297–306.
 261. Kojima M, Hosoda H, Date Y, Nakazato M, Matsuo H, Kangawa K. Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature*. 1999;402(6762):656–60.
 262. Müller TD, Nogueiras R, Andermann ML, Andrews ZB, Anker SD, Argente J, et al. Ghrelin. *Mol Metab*. 2015;4(6):437–60.
 263. Leggio L. Role of the ghrelin system in alcoholism: acting on the growth hormone secretagogue receptor to treat alcohol-related diseases. *Drug News Perspect*. 2010;23(3):157–66.
 264. Leggio L, Zywiak WH, Fricchione SR, Edwards SM, de la Monte SM, Swift RM, et al. Intravenous ghrelin administration increases alcohol craving in alcohol-dependent heavy drinkers: a preliminary investigation. *Biol Psychiatry*. 2014;76(9):734–41.
 265. Farokhnia M, Grodin EN, Lee MR, Oot EN, Blackburn AN, Stangl BL, et al. Exogenous ghrelin administration increases alcohol self-administration and modulates brain functional activity in heavy-drinking alcohol-dependent individuals. *Mol Psychiatry*. 2018;23(10):2029–38.
 266. Jerlhag E, Egecioglu E, Landgren S, Salomé N, Heilig M, Moechars D, et al. Requirement of central ghrelin signaling for alcohol reward. *Proc Natl Acad Sci USA*. 2009;106(27):11318–23.
 267. Kaur S, Ryabinin AE. Ghrelin receptor antagonism decreases alcohol consumption and activation of periculomotor urocortin-containing neurons. *Alcohol Clin Exp Res*. 2010;34(9):1525–34.
 268. Gomez JL, Cunningham CL, Finn DA, Young EA, Helpenstell LK, Schutte LM, et al. Differential effects of ghrelin antagonists on alcohol drinking and reinforcement in mouse and rat models of alcohol dependence. *Neuropharmacology*. 2015;97:182–93.
 269. Landgren S, Simms JA, Hyytiä P, Engel JA, Bartlett SE, Jerlhag E. Ghrelin receptor (GHS-R1A) antagonism suppresses both operant alcohol self-administration and high alcohol consumption in rats. *Addict Biol*. 2012;17(1):86–94.
 270. Stevenson JR, Francomacaro LM, Bohidar AE, Young KA, Pesarchick BF, Buirkle JM, et al. Ghrelin receptor (GHS-R1A) antagonism alters preference for ethanol and sucrose in a concentration-dependent manner in prairie voles. *Physiol Behav*. 2016;1(155):231–6.
 271. Kong J, Chuddy J, Stock IA, Loria PM, Straub SV, Vage C, et al. Pharmacological characterization of the first in class clinical candidate PF-05190457: a selective ghrelin receptor competitive antagonist with inverse agonism that increases vagal afferent firing and glucose-dependent insulin secretion ex vivo. *Br J Pharmacol*. 2016;173(9):1452–64.
 272. Lee MR, Tapocik JD, Ghareeb M, Schwandt ML, Dias AA, Le AN, et al. The novel ghrelin receptor inverse agonist PF-5190457 administered with alcohol: preclinical safety experiments and a phase 1b human laboratory study. *Mol Psychiatry*. 2018. <https://doi.org/10.1038/s41380-018-0064-y>.
 273. Denney WS, Sonnenberg GE, Carvajal-Gonzalez S, Tuthill T, Jackson VM. Pharmacokinetics and pharmacodynamics of PF-05190457: the first oral ghrelin receptor inverse agonist to be profiled in healthy subjects. *Br J Clin Pharmacol*. 2017;83(2):326–38.
 274. Laprairie RB, Bagher AM, Kelly MEM, Denovan-Wright EM. Cannabidiol is a negative allosteric modulator of the cannabinoid CB1 receptor. *Br J Pharmacol*. 2015;172(20):4790–805.
 275. Martínez-Pinilla E, Varani K, Reyes-Resina I, Angelats E, F F, Ferreira-Vera C, et al. Binding and signaling studies disclose a potential allosteric site for cannabidiol in cannabinoid CB2 receptors. *Frontiers in pharmacology [Internet]*. 2017 [cited 2021 Aug 30];8. <https://pubmed.ncbi.nlm.nih.gov/29109685/>.
 276. Pertwee RG. The diverse CB1 and CB2 receptor pharmacology of three plant cannabinoids: delta9-tetrahydrocannabinol, cannabidiol and delta9-tetrahydrocannabivarin. *Br J Pharmacol*. 2008;153(2):199–215.
 277. Russo EB, Burnett A, Hall B, Parker KK. Agonistic properties of cannabidiol at 5-HT1a receptors. *Neurochem Res*. 2005;30(8):1037–43.
 278. Kathmann M, Flau K, Redmer A, Tränkle C, Schlicker E. Cannabidiol is an allosteric modulator at mu- and delta-opioid receptors. *Naunyn Schmiedeberg Arch Pharmacol*. 2006;372(5):354–61.
 279. De Ternay J, Naassila M, Nourredine M, Louvet A, Bailly F, Sescousse G, et al. Therapeutic prospects of cannabidiol for alcohol use disorder and alcohol-related damages on the liver and the brain. *Front Pharmacol*. 2019;10:627.
 280. Turna J, Syan SK, Frey BN, Rush B, Costello MJ, Weiss M, et al. Cannabidiol as a novel candidate alcohol use disorder pharmacotherapy: a systematic review. *Alcohol Clin Exp Res*. 2019;43(4):550–63.
 281. Hurd YL, Spriggs S, Alishayev J, Winkel G, Gurgov K, Kudrich C, et al. Cannabidiol for the reduction of cue-induced craving and anxiety in drug-abstinent individuals with heroin use disorder: a double-blind randomized placebo-controlled trial. *Am J Psychiatry*. 2019;176(11):911–22.
 282. Freeman TP, Hindocha C, Baio G, Shaban NDC, Thomas EM, Astbury D, et al. Cannabidiol for the treatment of cannabis use disorder: a phase 2a, double-blind, placebo-controlled, randomised, adaptive Bayesian trial. *Lancet Psychiatry*. 2020;7(10):865–74.

283. Litten RZ, Falk DE, Ryan ML, Fertig JB. Discovery, development, and adoption of medications to treat alcohol use disorder: goals for the phases of medications development. *Alcohol Clin Exp Res*. 2016;40(7):1368–79.
284. Venegas A, Donato S, Meredith LR, Ray LA. Understanding low treatment seeking rates for alcohol use disorder: A narrative review of the literature and opportunities for improvement. *Am J Drug Alcohol Abuse*. 2021;31:1–16.
285. Volkow ND, Gordon JA, Koob GF. Choosing appropriate language to reduce the stigma around mental illness and substance use disorders. *Neuropsychopharmacology*. 2021;46(13):2230–2.
286. Falk DE, O'Malley SS, Witkiewitz K, Anton RF, Litten RZ, Slater M, et al. Evaluation of drinking risk levels as outcomes in alcohol pharmacotherapy trials: a secondary analysis of 3 randomized clinical trials. *JAMA Psychiat*. 2019;76(4):374–81.
287. Witkiewitz K, Kranzler HR, Hallgren KA, O'Malley SS, Falk DE, Litten RZ, et al. Drinking risk level reductions associated with improvements in physical health and quality of life among individuals with alcohol use disorder. *Alcohol Clin Exp Res*. 2018;42(12):2453–65.
288. Hasin DS, Wall M, Witkiewitz K, Kranzler HR, Falk D, Litten R, et al. Change in non-abstinent WHO drinking risk levels and alcohol dependence: a 3-year follow-up study in the US general population. *Lancet Psychiatry*. 2017;4(6):469–76.
289. Mann K, Vollstädt-Klein S, Reinhard I, Leménager T, Fauth-Bühler M, Hermann D, et al. Predicting naltrexone response in alcohol-dependent patients: the contribution of functional magnetic resonance imaging. *Alcohol Clin Exp Res*. 2014;38(11):2754–62.
290. Witkiewitz K, Roos CR, Mann K, Kranzler HR. Advancing precision medicine for alcohol use disorder: replication and extension of reward drinking as a predictor of naltrexone response. *Alcohol Clin Exp Res*. 2019;43(11):2395–405.
291. Litten RZ, Falk DE, Ryan ML, Fertig J, Leggio L. Advances in pharmacotherapy development: human clinical studies. *Handb Exp Pharmacol*. 2018;248:579–613.
292. Litten RZ, Ryan ML, Falk DE, Reilly M, Fertig JB, Koob GF. Heterogeneity of alcohol use disorder: understanding mechanisms to advance personalized treatment. *Alcohol Clin Exp Res*. 2015;39(4):579–84.
293. Agabio R, Pisanu C, Gessa GL, Franconi F. Sex differences in alcohol use disorder. *Curr Med Chem*. 2017;24(24):2661–70.
294. Heilig M, Leggio L. What the alcohol doctor ordered from the neuroscientist: therapeutic biomarkers for personalized treatments. *Prog Brain Res*. 2016;224:401–18.
295. Castillo-Carniglia A, Keyes KM, Hasin DS, Cerdá M. Psychiatric comorbidities in alcohol use disorder. *Lancet Psychiatry*. 2019;6(12):1068–80.
296. Leggio L, Lee MR. Treatment of alcohol use disorder in patients with alcoholic liver disease. *Am J Med*. 2017;130(2):124–34.
297. Guillot A, Ren T, Jourdan T, Pawlosky RJ, Han E, Kim S-J, et al. Targeting liver aldehyde dehydrogenase-2 prevents heavy but not moderate alcohol drinking. *Proc Natl Acad Sci USA*. 2019;116(51):25974–81.
298. Tampier L, Quintanilla ME, Israel Y. Tolerance to disulfiram induced by chronic alcohol intake in the rat. *Alcohol Clin Exp Res*. 2008;32(6):937–41.
299. Spanagel R, Hölter SM, Allingham K, Landgraf R, Zieglgänsberger W. Acamprosate and alcohol: I. Effects on alcohol intake following alcohol deprivation in the rat. *Eur J Pharmacol*. 1996;305(1–3):39–44.
300. Spanagel R, Putzke J, Steffler A, Schöbitz B, Zieglgänsberger W. Acamprosate and alcohol: II. Effects on alcohol withdrawal in the rat. *Eur J Pharmacol*. 1996;305(1–3):45–50.
301. Chester JA, Grahame NJ, Li TK, Lumeng L, Froehlich JC. Effects of acamprosate on sensitization to the locomotor-stimulant effects of alcohol in mice selectively bred for high and low alcohol preference. *Behav Pharmacol*. 2001;12(6–7):535–43.
302. Kurokawa K, Mizuno K, Shibasaki M, Higashioka M, Oka M, Hirouchi M, et al. Acamprosate suppresses ethanol-induced place preference in mice with ethanol physical dependence. *J Pharmacol Sci*. 2013;122(4):289–98.
303. Jacobsen JHW, Buisman-Pijlman FTA, Mustafa S, Rice KC, Hutchinson MR. The efficacy of (+)-Naltrexone on alcohol preference and seeking behaviour is dependent on light-cycle. *Brain Behav Immun*. 2018;1(67):181–93.
304. Jacobsen JHW, Buisman-Pijlman FT, Mustafa S, Rice KC, Hutchinson MR. Antagonising TLR4-TRIF signalling before or after a low-dose alcohol binge during adolescence prevents alcohol drinking but not seeking behaviour in adulthood. *Neuropharmacology*. 2018;1(128):460–73.
305. Wu Y, Lousberg EL, Moldenhauer LM, Hayball JD, Collier JK, Rice KC, et al. Inhibiting the TLR4-MyD88 signalling cascade by genetic or pharmacological strategies reduces acute alcohol-induced sedation and motor impairment in mice. *Br J Pharmacol*. 2012;165(5):1319–29.
306. Montesinos J, Gil A, Guerri C. nalmefene prevents alcohol-induced neuroinflammation and alcohol drinking preference in adolescent female mice: role of TLR4. *Alcohol Clin Exp Res*. 2017;41(7):1257–70.
307. Calleja-Conde J, Echeverry-Alzate V, Giné E, Bühler K-M, Nadal R, Maldonado R, et al. Nalmefene is effective at reducing alcohol seeking, treating alcohol-cocaine interactions and reducing alcohol-induced histone deacetylases gene expression in blood. *Br J Pharmacol*. 2016;173(16):2490–505.
308. Foo JC, Vengeliene V, Noori HR, Yamaguchi I, Morita K, Nakamura T, et al. drinking levels and profiles of alcohol addicted rats predict response to nalmefene. *Front Pharmacol*. 2019;10:471.
309. Colombo G, Gessa GL. Suppressing effect of baclofen on multiple alcohol-related behaviors in laboratory animals. *Front Psychiatry*. 2018;9:475.
310. Colombo G, Vacca G, Serra S, Brunetti G, Carai MAM, Gessa GL. Baclofen suppresses motivation to consume alcohol in rats. *Psychopharmacology*. 2003;167(3):221–4.
311. Maccioni P, Lorrai I, Contini A, Leite-Morris K, Colombo G. Microinjection of baclofen and CGP7930 into the ventral tegmental area suppresses alcohol self-administration in alcohol-preferring rats. *Neuropharmacology*. 2018;136(Pt A):146–58.
312. Colombo G, Addolorato G, Agabio R, Carai MAM, Pibiri F, Serra S, et al. Role of GABA(B) receptor in alcohol dependence: reducing effect of baclofen on alcohol intake and alcohol motivational properties in rats and amelioration of alcohol withdrawal syndrome and alcohol craving in human alcoholics. *Neurotox Res*. 2004;6(5):403–14.
313. Maccioni P, Pes D, Fantini N, Carai MAM, Gessa GL, Colombo G. gamma-Hydroxybutyric acid (GHB) suppresses alcohol's motivational properties in alcohol-preferring rats. *Alcohol*. 2008;42(2):107–13.