

# UC San Diego

## UC San Diego Previously Published Works

### Title

Phosphocholine as a pattern recognition ligand for CD36 1

### Permalink

<https://escholarship.org/uc/item/27s1k89b>

### Journal

Journal of Lipid Research, 46(5)

### ISSN

0022-2275

### Authors

Boullier, Agnès  
Friedman, Peter  
Harkewicz, Richard  
[et al.](#)

### Publication Date

2005-05-01

### DOI

10.1194/jlr.m400496-jlr200

Peer reviewed

# Phosphocholine as a pattern recognition ligand for CD36<sup>1</sup>,<sup>§</sup>

Agnès Boullier,<sup>2,\*</sup> Peter Friedman,<sup>2,†</sup> Richard Harkewicz,<sup>†</sup> Karsten Hartvigsen,<sup>\*</sup> Simone R. Green,<sup>\*</sup> Felicidad Almazan,<sup>\*</sup> Edward A. Dennis,<sup>†</sup> Daniel Steinberg,<sup>\*</sup> Joseph L. Witztum,<sup>\*</sup> and Oswald Quehenberger<sup>3,\*</sup>

Department of Medicine\* and Departments of Chemistry, Biochemistry, and Pharmacology,<sup>†</sup> University of California San Diego, La Jolla, CA 92093-0682

**Abstract** We have previously shown that CD36 recognizes oxidation products of phospholipids on oxidized LDL (OxLDL) such as 1-palmitoyl-2-(5'-oxovaleroyl)-sn-glycero-3-phosphocholine (POVPC). The current study was designed to examine whether the phosphocholine (PC) headgroup in POVPC constitutes an obligatory binding target for CD36. To examine the contribution of PC in the binding of POVPC to CD36, we used well-defined synthetic oxidized phospholipids (OxPLs) cross-linked to BSA or to a hexapeptide. The OxPL adducts were then tested for their ability to bind to CD36-transfected cells and for their ability to inhibit OxLDL binding to CD36. Both POVPC-BSA and POVPC-peptide adducts were high-affinity ligands for CD36 and potent inhibitors of OxLDL binding. Enzymatic removal of the entire PC moiety of the POVPC-peptide, or of the choline headgroup alone, as well as substitution of the choline headgroup by ethanolamine abrogated the inhibitory activity of POVPC. Interestingly, PC by itself or cross-linked to BSA did not show any intrinsic competition activity. **In conclusion, our data demonstrate that the PC headgroup of OxPL alone is sufficient for binding to CD36, but only if presented in the correct conformation as in OxPL of OxLDL or as in POVPC-peptide adducts.**—Boullier, A., P. Friedman, R. Harkewicz, K. Hartvigsen, S. R. Green, F. Almazan, E. A. Dennis, D. Steinberg, J. L. Witztum, and O. Quehenberger. **Phosphocholine as a pattern recognition ligand for CD36.** *J. Lipid Res.* 2005. 46: 969–976.

**Supplementary key words** scavenger receptor • oxidized phospholipids • atherosclerosis

Mounting evidence implicates oxidized low density lipoprotein (OxLDL) in the pathogenesis of atherosclerosis. The oxidative modifications are presumed to occur after the entry of plasma LDL into the intima and are catalyzed by the cellular constituents of the arterial wall. Although clearance of the highly cytotoxic OxLDL may protect the surrounding tissue from damage, the unregulated uptake of OxLDL by macrophages within the arterial wall leads to the formation of lipid-laden foam cells and the development of the fatty streak, the hallmark of early atherosclerosis.

The recognition and uptake of OxLDL by macrophages is mediated by specific cell surface scavenger receptors, including CD36, a class B scavenger receptor (1). CD36 is a heavily glycosylated protein with broad ligand specificity, a characteristic of pattern recognition receptors. It binds long-chain fatty acids (2), anionic phospholipids (3),  $\beta$ -amyloid (4), advanced glycation end products (5), and OxLDL (1). It also functions in the recognition and removal of *Plasmodium falciparum*-infected erythrocytes (6) and apoptotic cells (7). It serves as the primary receptor for platelet adhesion to collagen (8) and may mediate the antiangiogenic activity of thrombospondin (9).

Recent data derived from CD36-deficient mice support an important role of CD36 in foam cell formation and atherogenesis. A significant decrease in binding and degradation of OxLDL was observed in macrophages from null mice compared with those from control mice (10). Similarly, macrophages from CD36-deficient patients were less capable of binding and degrading OxLDL and accumulated less cholesteryl ester than macrophages from control subjects (11). The relative contribution of scavenger receptors in the uptake of OxLDL was further addressed in mice lacking both scavenger receptor A and CD36 (12). Binding and uptake studies demonstrated that scavenger receptor A and CD36 were the principal macrophage receptors responsible for the binding of OxLDL and the accumulation of cholesteryl ester derived from modified lipoproteins. Consistent with a principal pathogenic role of scavenger receptors in atherogenesis, the targeted disruption of these receptors in mice resulted in a marked reduction in atherosclerosis (13).

Abbreviations: Ac-TGTKGY, Ac-threonine-glycine-threonine-lysine-glycine-tyrosine; apoE, apolipoprotein E; KLH, keyhole limpet hemocyanin; OxLDL, oxidized low density lipoprotein; OxPL, oxidized phospholipid; PAMP, pathogen-associated molecular pattern; PC, phosphocholine; PLC, phospholipase C; PLD, phospholipase D; POVG, 1-palmitoyl-2-(5'-oxovaleroyl)-glycerol; POVPA, 1-palmitoyl-2-(5'-oxovaleroyl)-sn-glycero-3-phosphatidic acid; POVPC, 1-palmitoyl-2-(5'-oxovaleroyl)-sn-glycero-3-phosphocholine; POVPE, 1-palmitoyl-2-(5'-oxovaleroyl)-sn-glycero-3-phosphoethanolamine.

<sup>1</sup> Trudy Forte served as Guest Editor for this manuscript.

<sup>2</sup> A Boullier and P. Friedman contributed equally to this work.

<sup>3</sup> To whom correspondence should be addressed.

e-mail: oquehenberger@ucsd.edu

<sup>§</sup> The online version of this article (available at <http://www.jlr.org>) contains an additional table and two figures.

Manuscript received 16 December 2004 and in revised form 27 January 2005.

Published, JLR Papers in Press, February 1, 2005.

DOI 10.1194/jlr.M400496-JLR200

Copyright © 2005 by the American Society for Biochemistry and Molecular Biology, Inc.

This article is available online at <http://www.jlr.org>

tion of the CD36 gene markedly reduced lesion formation (10). Collectively, these data demonstrate that CD36 is importantly involved in the uptake of OxLDL by macrophages and in atherosclerotic lesion development.

In previous studies, we (13) and subsequently others (14, 15) have shown that the binding of OxLDL to CD36 is mediated in part by oxidized phospholipids (OxPLs). However, the specific epitopes on such OxPLs that trigger the binding of these lipids have not been fully defined. Natural autoantibodies against OxLDL have been isolated from hypercholesterolemic apolipoprotein E (apoE)-deficient mice (16). One of them, designated EO6, has been shown to specifically recognize the phosphocholine (PC) headgroup of PC-containing OxPLs, such as that found in 1-palmitoyl-2-(5'-oxovaleroyl)-sn-glycero-3-phosphocholine (POVPC) (17). The observation that EO6 inhibited the uptake of OxLDL by macrophages suggested that the PC group of OxPLs might be a specific binding motif on OxLDL recognized by macrophage scavenger receptors (13). The present study was designed to examine the involvement of the PC headgroup in the binding of POVPC to CD36 using well-defined synthetic lipids.



## METHODS

### Materials

COS-7 cells were purchased from the American Type Culture Collection (Manassas, VA). Dulbecco's modified Eagle's medium with 4.5 g/l glucose was from Bio-Whittaker (Walkersville, MD); fetal bovine serum was from Gemini Bioproducts, Inc. (Calabasas, CA). Penicillin-streptomycin, L-glutamine, and trypsin-EDTA were from Irvine Scientific (Santa Ana, CA). FuGene6 was purchased from Roche Applied Science (Indianapolis, IN). Anti-human CD36 antibody (clone FA6-152) was purchased from Immunotech (Marseille, France). Na<sup>125</sup>I (2,000 Ci/mmol) was from ICN (Costa Mesa, CA), and Iodogen was from Pierce (Rockford, IL). PC-keyhole limpet hemocyanin (KLH) and PC-BSA were from Bioresearch Technologies, Inc. (Novato, CA). All of the phospholipids were purchased from Avanti Polar Lipids (Alabaster, AL). Schiff's Reagent (for detection of aldehydes by thin-layer chromatography), Sigma Spray Reagent Molybdenum Blue (for detection of phosphates by thin-layer chromatography), BSA, PBS, phospholipase C (PLC) from *Bacillus cereus*, phospholipase D (PLD) from *Streptomyces species* and *Streptomyces chromofuscus*, ethanolamine, methyl sulfide, sodium cyanoborohydride, and sodium borohydride were purchased from Sigma-Aldrich (St. Louis, MO). Solvents and preparative as well as analytical thin-layer chromatography plates were purchased from Omnisolv, EMD Chemicals (Gibbstown, NJ). The peptide Ac-threonine-glycine-threonine-lysine-glycine-tyrosine (Ac-TGTKGY) was purchased from Synpep (Dublin, CA). Bio-Gel P-2 gel fine (45–90 μm) was from Bio-Rad (Hercules, CA).

### Transfection of COS-7 cells with CD36

COS-7 cells were transfected with CD36 DNA as previously described (13). COS-7 cells were maintained in glucose-containing Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum, penicillin (100 U/ml), streptomycin (100 μg/ml), and L-glutamine (2 mM). For transfection, cells were grown on a six-well plate to ~70–80% confluence and transfected with a mixture of 2 μg of human CD36 DNA subcloned into pSG5 (Stratagene) and 6 μl of FuGene6 (Roche). After 24 h, the cells were harvested with EDTA and plated on a 24-well plate at 0.2 × 10<sup>6</sup> cells/well. Nontransfected COS-7 cells were used as a con-



**Fig. 1.** Mass spectral analysis of the singly charged 1-palmitoyl-2-(5'-oxovaleroyl)-sn-glycero-3-phosphocholine (POVPC)-peptide [M<sup>+</sup>]. A: LC-enhanced resolution mass spectrum of POVPC-peptide. Note the 1 amu mass difference between adjacent isotopic peaks, indicating the 1<sup>+</sup> charge state. B: LC-MS/MS collision-induced fragmentation of the POVPC-peptide parent [M<sup>+</sup>]. All of the fragments were assigned to the POVPC-peptide parent, and the proposed molecular structures of two fragments are shown. The molecular structures of all other fragments are shown in supplementary Fig. I. Rel. Int., relative intensity.





the addition of 0.2 N NaOH. Aliquots were taken to measure the protein content by the method of Lowry et al. (19) and the cell-associated radioactivity using a  $\gamma$  spectrometer. All assays were done in triplicate.

**Binding of  $^{125}\text{I}$ -POVPC adducts.** The direct binding of  $^{125}\text{I}$ -POVPC-BSA was estimated by incubating the transfected cells with various concentrations of  $^{125}\text{I}$ -POVPC-BSA. The binding of POVPC-peptide was examined by inhibition binding analysis. Briefly, the transfected cells were incubated for 2 h at 4°C with trace amounts of  $^{125}\text{I}$ -POVPC-peptide in the presence of increasing concentrations of unlabeled POVPC-peptide as well as other OxPL peptides. Binding isotherms were generated and binding data were determined using Prism4 software (GraphPad Software, San Diego, CA).


## RESULTS

To examine the potential contribution of the PC moiety in the binding of OxPLs to CD36, we first characterized the binding of the POVPC-BSA adduct to CD36. As shown in **Fig. 3A**, the CD36-transfected cells bound  $^{125}\text{I}$ -POVPC-BSA with high affinity and in a saturable manner, characteristic of specific binding. Scatchard analysis of the equilibrium binding data revealed a binding affinity of  $0.19 \pm 0.02 \mu\text{M}$  for POVPC-BSA, expressed in moles of BSA. Under the conditions of synthesis used,  $\sim 32$  mol of POVPC was covalently bound per mole of BSA (23). On that basis, the affinity for POVPC presented on BSA was  $6.5 \pm 0.5 \mu\text{M}$ . Consistent with our previous data (13), POVPC-BSA proved to be an effective competitor and dose-dependently inhibited the binding of OxLDL to the CD36-transfected cells (**Fig. 3B**). These data suggest that POVPC and similar structures may constitute ligands on OxLDL that are recognized by scavenger receptors, including CD36. It should be noted here that the *sn*-2 aldehyde of POVPC was used to covalently link it to the amino groups of lysines of BSA via a reduced Schiff base adduct. Thus, the POVPC-BSA adduct did not contain any reactive carbonyl groups in the *sn*-2 position that could be potential ligands for CD36 (14). This suggests the possibility that the PC group itself, as presented in POVPC, was the essential ligand.


Because several moles of POVPC are covalently bound per mole of BSA, the OxPL-BSA adduct represents a multivalent ligand for CD36 in which individual epitopes may contribute to the binding either by interacting with multiple binding sites on CD36 or by receptor cross-linking, or both. To study the binding of POVPC as a monovalent ligand for CD36, we substituted a short peptide of six amino acid residues (Ac-TGTKGY), randomly chosen from the sequence of apoB, for BSA. This peptide contained only one available lysine residue to form the Schiff base, and the N terminus was blocked by acetylation so that each mole of peptide contained only a single mole of lipid. The structure of the POVPC-peptide was confirmed by mass spectroscopic analysis (**Fig. 1**). Compared with the POVPC-BSA adduct, the POVPC-peptide adduct appeared to be an even more potent inhibitor of OxLDL binding to the CD36-transfected cells (**Fig. 4**). However, it is important to note that the concentration of inhibitors in both cases was expressed

in terms of the concentration of POVPC (nanomoles of POVPC per milliliter), presented either as a BSA conjugate or as a hexapeptide conjugate. These results imply that some fraction of the POVPC molecules on the multiligand POVPC-BSA adduct are unavailable for binding to the receptors because of conformational or steric constraints.

To test more rigorously the role of the PC group of POVPC-peptide as a binding epitope for CD36, we sequentially hydrolyzed the functional group at the *sn*-3 position with PLD or PLC (**Fig. 2**). The PLD selectively hydrolyzed the choline group to yield POVPA-peptide, whereas the PLC selectively hydrolyzed the entire PC group in the *sn*-3 position to produce POVG-peptide (**Fig. 2**). In contrast to the intact POVPC-peptide, neither of these enzymatic hydrolysis products was an effective inhibitor for OxLDL



**Fig. 3.** Binding of POVPC-BSA to CD36. **A:** Direct binding of  $^{125}\text{I}$ -POVPC-BSA to CD36. CD36-transfected cells were incubated with  $^{125}\text{I}$ -POVPC-BSA (closed squares) or  $^{125}\text{I}$ -BSA (closed triangles) at the indicated protein concentrations for 2 h at 4°C. The values represent means  $\pm$  SD ( $n = 3$ ) of two independent experiments. The binding affinity of POVPC-BSA was determined by Scatchard analysis of the binding data (dissociation constant =  $0.19 \pm 0.02 \mu\text{M}$ ). On average, 32 mol of POVPC was covalently attached per mole of BSA. On that basis, the affinity for POVPC presented on BSA was  $6.5 \pm 0.5 \mu\text{M}$ . **B:** Inhibition of  $^{125}\text{I}$ -oxidized low density lipoprotein (OxLDL) binding to CD36 by the POVPC-BSA adduct. CD36-transfected cells (closed bars) and control cells (open bars) were incubated with  $5 \mu\text{g/ml}$   $^{125}\text{I}$ -OxLDL in the presence of the indicated concentrations of the POVPC-BSA adduct. As a positive control, a 30-fold excess of unlabeled OxLDL was used as a competitor. After 2 h of incubation at 4°C,  $^{125}\text{I}$ -OxLDL binding was determined. The values represent means  $\pm$  SD of triplicate determinations derived from three representative experiments.



**Fig. 4.** Inhibition of <sup>125</sup>I-OxLDL binding to CD36-transfected cells by enzymatically modified products of the POVPC-peptide. CD36-transfected cells were incubated with 5 μg/ml <sup>125</sup>I-OxLDL in the presence of the indicated concentrations of the various competitors. After 2 h of incubation at 4°C, OxLDL binding was determined. Data are presented as percentage binding of <sup>125</sup>I-OxLDL, 100% being the value of the binding without any competitor. Each value represents the mean ± SD of triplicate determinations from two separate experiments. POVPE, 1-palmitoyl-2-(5'-oxovaleroyl)-sn-glycero-3-phosphoethanolamine.

binding (Fig. 4). Similarly, the POVPE-peptide, in which the choline group of POVPC was replaced by ethanolamine, had no inhibitory effect on OxLDL binding to CD36.

The important role of the PC moiety as a binding motif for CD36 was further established by direct binding analysis. The <sup>125</sup>I-POVPC-peptide bound to the transfected cells with an affinity of  $4.81 \pm 1.25 \mu\text{M}$ , which was similar to the binding affinity of the POVPC-BSA adduct (Fig. 5). Removal or substitution of the headgroup as in POVPA, POVG, and POVPE abrogated the binding of these lipid-peptide adducts to CD36, and no inhibition of <sup>125</sup>I-POVPC-peptide binding was observed (data not shown). These data suggest that the PC headgroup is essential for the binding of POVPC to CD36 and imply a critical involvement of the choline group for the recognition of OxPL by CD36.


To examine whether PC by itself is recognized by CD36 or whether a specific conformational presentation is required for recognition, we examined the ability of PC to compete for the binding of OxLDL to CD36 when presented either as a free salt or covalently bound to KLH (PC-KLH) or BSA (PC-BSA). As shown in Fig. 6, neither free PC in solution (Fig. 6A) nor cross-linked to KLH or BSA (Fig. 6B) inhibited OxLDL binding. The failure of PC by itself to compete suggests that a specific conformational presentation of the headgroup is required for recognition by CD36.

## DISCUSSION

The role of CD36 as the principal scavenger receptor responsible for the uptake of OxLDL by macrophages and

foam cell formation has been solidly established (1, 11, 12). The present study was undertaken to identify and characterize the oxidation epitopes that may mediate the recognition and removal of modified lipoproteins by CD36. Of particular interest are the oxidation products derived from phosphatidylcholine, including POVPC. In a previous study, we made the observation that the POVPC-BSA adduct competed for the binding of OxLDL to CD36-transfected cells (13). We concluded from this study that OxPLs, present either in the lipid phase or covalently attached to apoB, constitute major binding epitopes and mediate the recognition of OxLDL by CD36. We have now extended these initial observations and identified the PC headgroup of oxidized phosphatidylcholine as an obligatory binding target for CD36.

During the oxidative modification of LDL, a variety of oxidized molecules are generated, including OxPLs (24). There is now a large body of evidence that such OxPLs have proinflammatory and proatherogenic properties and that their accumulation in the vessel wall contributes significantly to the pathological consequences of developing atherosclerotic lesions (24–30). Several bioactive lipids, including POVPC, have been demonstrated in lesion areas where they may regulate inflammatory functions of endothelial cells, such as the expression of adhesion proteins and chemokines (24, 31–33). In addition to the effects on the cellular constituents of the vessel wall, these oxidized lipids have also been implicated in the adaptive and innate immune response. We have previously cloned a series of natural autoantibodies from hypercholesterolemic mice directed against neoself antigens formed dur-




**Fig. 5.** Competition of  $^{125}\text{I}$ -POVPC-peptide binding to CD36. CD36-transfected cells were incubated with trace amounts of  $^{125}\text{I}$ -POVPC-peptide for 2 h at  $4^\circ\text{C}$  in the presence of the unlabeled of POVPC-peptide (closed triangles) and the unconjugated peptide (open triangles) at the indicated concentrations. Binding analysis of  $^{125}\text{I}$ -POVPC-peptide revealed a binding affinity of  $4.81 \pm 1.25 \mu\text{M}$ . Results represent means  $\pm$  SD ( $n = 3$ ) of two separate experiments.

ing oxidation of LDL (16). One of these natural antibodies, EO6, binds to the PC headgroup of OxPLs on OxLDL but not the PC headgroup of unoxidized phosphatidylcholine. In turn, EO6 blocks the uptake of OxLDL by macrophages (17, 22, 34).

These OxPLs also serve as ligands for CD36 and may mediate, at least in part, the recognition and removal of OxLDL by macrophages. Our data clearly implicate the PC headgroup of modified phosphatidylcholine in mediating the interaction with CD36. As was the case with the natural antibody EO6, CD36 also binds the PC headgroup only when other parts of the phospholipid have been modified by oxidation. Podrez et al. (14) have suggested that modifications of the *sn*-2 fatty acid may also contribute to the binding to CD36 and may represent a separate class of ligands on some OxPLs for CD36. To rule out any binding contribution of such oxidized polyunsaturated fatty acids in our studies and to reconstitute a ligand as it may be present on apoB of OxLDL, we conjugated the aldehyde in the *sn*-2 position of POVPC to BSA or to a lysine-containing peptide with subsequent reduction of the Schiff base to a secondary amine. Therefore, the binding of OxPLs to CD36 could only be attributed to the PC residue of the OxPL that became exposed and accessible for CD36 binding as a consequence of conformational changes associated with oxidative events. It is important to note that our data directly address the issue of the role of the PC headgroup as a ligand. Our data do not exclude the possibility that oxidized moieties in the *sn*-2 side chain of POVPC could also coordinate to other receptor binding sites of CD36 and, thus, could also be involved in the binding of the isolated POVPC lipid.

In support of a critical role of PC in the binding of OxPLs and OxLDL to CD36, the hydrolysis products of the POVPC-peptide adduct after treatment with PLD or PLC to remove choline or PC, respectively, were no longer competitors for the OxLDL ligand. Similarly, the substitution of the PC headgroup by phosphoethanolamine abrogated the inhibitory effect of POVPC-peptide. Furthermore,

Supplemental Material can be found at:  
<http://www.jlr.org/content/suppl/2005/04/27/M400496-JLR200.DC1.html>  
<http://www.jlr.org/content/suppl/2005/04/11/10.1194.jlr.M400496-JLR200.DC1.html>



**Fig. 6.** Competition of  $^{125}\text{I}$ -OxLDL binding to CD36 by free phosphocholine (PC) and PC-protein adducts. Control cells (open bars) and CD36-transfected cells (closed bars) were incubated with  $5 \mu\text{g}/\text{ml}$   $^{125}\text{I}$ -OxLDL for 2 h at  $4^\circ\text{C}$  in the absence or presence of PC presented as a salt in solution (A) or as protein adducts with keyhole limpet hemocyanin (KLH) or BSA (B). KLH and BSA were used as negative controls at  $10 \mu\text{M}$ . The values represent means  $\pm$  SD of triplicate determinations.

the PC group by itself when covalently bound to protein (KLH or BSA) did not compete for the binding of OxLDL to CD36, suggesting that the lipid environment and structural elements of the oxidized phosphatidylcholine molecule were necessary to maintain the appropriate conformation and presentation of PC to be recognized as a ligand.

CD36 developed as part of the innate immune system, and the scavenger function of CD36 is evolutionarily conserved (35, 36). Aside from its function as a receptor for OxLDL, CD36 is thought to play an important role in the recognition and removal of cells undergoing programmed cell death and of pathogens during infection, in cooperation with the phosphatidylserine receptor or the  $\alpha\text{v}\beta 3$  integrin (7, 37). The ligands for CD36 on apoptotic cells and pathogens are not well characterized, but the initial recognition of microbes as they enter the body is based on the presentation of common structural patterns. PC is known to be recognized by the innate immune system, and this may provide host defenses against microbial infection, such as by *Streptococcus pneumoniae*, in which PC is a dominant epitope of the cell wall polysaccharide. Furthermore, PC of OxPLs are also present on the outer leaflet of apoptotic cells (38, 39), and the PC of certain bacteria, of apoptotic cells, and of OxLDL all share molecular mimicry (34, 40). Thus, the PC moieties presented in these



contexts represent pathogen-associated molecular patterns (PAMPs), which are recognition ligands for pattern recognition innate immune receptors such as CD36. Consistent with this, we have previously shown that binding of both OxLDL and apoptotic cells to CD36-transfected cells and macrophages is mediated through a common epitope with structural similarities to POVPC (13, 39). Based on the data derived from EO6, which strongly blocked that binding, we propose that the PC moiety represents such a common PAMP.

In summary, our data strongly implicate a particular conformational presentation of PC as a PAMP, which is recognized by several facets of innate immunity, including natural antibodies, C-reactive protein (41), and pattern recognition receptors such as CD36 and likely scavenger receptor class B type I (42). In this context, the exposure of PC as a result of the oxidation of phospholipids present on OxLDL mediates the recognition of OxLDL by innate immune responses. The application of these findings with regard to the recognition of microbial pathogens and apoptotic cells by macrophage scavenger receptors is currently under study. **■**

This work was supported by National Institutes of Health Grants HL-56989 (Specialized Center of Research in Molecular Medicine and Atherosclerosis), HL-57505, HL-69464, GM-20501, and GM-069338 and by Tobacco-Related Disease Research Program Grant 10KT-0220. The authors thank Jennifer Pattison for her technical assistance in the isolation of the lipoproteins.

## REFERENCES

- Endemann, G., L. W. Stanton, K. S. Madden, C. M. Bryant, R. T. White, and A. A. Protter. 1993. CD36 is a receptor for oxidized low density lipoprotein. *J. Biol. Chem.* **268**: 11811–11816.
- Abumrad, N. A., M. Raafat El-Maghrabi, E. Amri, E. Lopez, and P. A. Grimaldi. 1993. Cloning of a rat adipocyte membrane protein implicated in binding or transport of long chain fatty acids that is induced during preadipocyte differentiation. Homology with human CD36. *J. Biol. Chem.* **268**: 17665–17668.
- Rigotti, A., S. L. Acton, and M. Krieger. 1995. The class B scavenger receptors SR-BI and CD36 are receptors for anionic phospholipids. *J. Biol. Chem.* **270**: 16221–16224.
- Moore, K. J., J. El Khoury, L. A. Medeiros, K. Terada, C. Geula, A. D. Luster, and M. W. Freeman. 2002. A CD36-initiated signaling cascade mediates inflammatory effects of beta-amyloid. *J. Biol. Chem.* **277**: 47373–47379.
- Ohgami, N., R. Nagai, M. Ikemoto, H. Arai, A. Kuniyasu, S. Horiuchi, and H. Nakayama. 2001. CD36, a member of the class B scavenger receptor family, as a receptor for advanced glycation end products. *J. Biol. Chem.* **276**: 3195–3202.
- Oquendo, P., E. Hundt, J. Lawler, and B. Seed. 1989. CD36 directly mediates cytoadherence of *Plasmodium falciparum* parasitized erythrocytes. *Cell*. **58**: 95–101.
- Fadok, V. A., M. L. Warner, D. L. Bratton, and P. M. Henson. 1998. CD36 is required for phagocytosis of apoptotic cells by human macrophages that use either a phosphatidylserine receptor or the vitronectin receptor (alpha v beta 3). *J. Immunol.* **161**: 6250–6257.
- Tandon, N. N., U. Kralisz, and G. A. Jamieson. 1989. Identification of glycoprotein IV (CD36) as a primary receptor for platelet-collagen adhesion. *J. Biol. Chem.* **264**: 7576–7583.
- Leung, L. L. K., W.-X. Li, J. L. McGregor, G. Albrecht, and R. J. Howard. 1992. CD36 peptides enhance or inhibit CD36-thrombospondin binding. *J. Biol. Chem.* **267**: 18244–18250.
- Febbraio, M., E. A. Podrez, J. D. Smith, D. P. Hajjar, S. L. Hazen, H. F.

- Hoff, K. Sharma, and R. L. Silverstein. 2000. Targeted disruption of the class B scavenger receptor CD36 protects against atherosclerotic lesion development in mice. *J. Clin. Invest.* **105**: 1049–1056.
- Nozaki, S., H. Kashiwagi, S. Yamashita, T. Nakagawa, B. Kostner, Y. Tomiyama, A. Nakata, M. Ishigami, J. Miyagawa, K. Kameda-Takemura, Y. Kurata, and Y. Matsuzawa. 1995. Reduced uptake of oxidized low density lipoprotein in monocyte-derived macrophages from CD36-deficient subjects. *J. Clin. Invest.* **96**: 1859–1865.
- Kunjathoor, V. V., M. Febbraio, E. A. Podrez, K. J. Moore, L. Anderson, S. Koehn, J. S. Rhee, R. L. Silverstein, H. F. Hoff, and M. W. Freeman. 2002. Scavenger receptors class A-I/II and CD36 are the principal receptors responsible for the uptake of modified low density lipoprotein leading to lipid loading in macrophages. *J. Biol. Chem.* **277**: 49982–49988.
- Boullier, A., K. L. Gillotte, S. Hörkö, S. R. Green, P. Friedman, E. A. Dennis, J. L. Witztum, D. Steinberg, and O. Quehenberger. 2000. The binding of oxidized low density lipoprotein to mouse CD36 is mediated in part by oxidized phospholipids that are associated with both the lipid and protein moieties of the lipoprotein. *J. Biol. Chem.* **275**: 9163–9169.
- Podrez, E. A., E. Poliakov, Z. Shen, R. Zhang, Y. Deng, M. Sun, P. J. Finton, L. Shan, B. Gugiu, P. L. Fox, H. F. Hoff, R. G. Salomon, and S. L. Hazen. 2002. Identification of a novel family of oxidized phospholipids that serve as ligands for the macrophage scavenger receptor CD36. *J. Biol. Chem.* **277**: 38503–38516.
- Podrez, E. A., E. Poliakov, Z. Shen, R. Zhang, Y. Deng, M. Sun, P. J. Finton, L. Shan, M. Febbraio, D. P. Hajjar, R. L. Silverstein, H. F. Hoff, R. G. Salomon, and S. L. Hazen. 2002. A novel family of atherogenic oxidized phospholipids promotes macrophage foam cell formation via the scavenger receptor CD36 and is enriched in atherosclerotic lesions. *J. Biol. Chem.* **277**: 38517–38523.
- Palinski, W., S. Hörkö, E. Miller, U. P. Steinbrecher, H. C. Powell, L. K. Curtiss, and J. L. Witztum. 1996. Cloning of monoclonal autoantibodies to epitopes of oxidized lipoproteins from apolipoprotein E-deficient mice. Demonstration of epitopes of oxidized low density lipoprotein in human plasma. *J. Clin. Invest.* **98**: 800–814.
- Hörkö, S., D. A. Bird, E. Miller, H. Itabe, N. Leitinger, G. Subbanagounder, J. A. Berliner, P. Friedman, E. A. Dennis, L. K. Curtiss, W. Palinski, and J. L. Witztum. 1999. Monoclonal autoantibodies specific for oxidized phospholipids or oxidized phospholipid-protein adducts inhibit macrophage uptake of oxidized low-density lipoproteins. *J. Clin. Invest.* **103**: 117–128.
- Havel, R. J., H. A. Eder, and J. H. Bragdon. 1955. The distribution and chemical composition of ultracentrifugally separated lipoproteins in human serum. *J. Clin. Invest.* **34**: 1345–1353.
- Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**: 265–275.
- Salacinski, P. R., C. McLean, J. E. Sykes, V. V. Clement-Jones, and P. J. Lowry. 1981. Iodination of proteins, glycoproteins, and peptides using a solid-phase oxidizing agent, 1,3,4,6-tetrachloro-3alpha,6alpha-diphenyl glycoluril (Iodogen). *Anal. Biochem.* **117**: 136–146.
- Yagi, K. 1976. A simple fluorometric assay for lipoperoxide in blood plasma. *Biochem. Med.* **15**: 212–216.
- Friedman, P., S. Hörkö, D. Steinberg, J. L. Witztum, and E. A. Dennis. 2002. Correlation of antiphospholipid antibody recognition with the structure of synthetic oxidized phospholipids. *J. Biol. Chem.* **277**: 7010–7020.
- Bird, D. A., K. L. Gillotte, S. Hörkö, P. Friedman, E. A. Dennis, J. L. Witztum, and D. Steinberg. 1999. Receptors for oxidized low-density lipoprotein on elicited mouse peritoneal macrophages can recognize both the modified lipid moieties and the modified protein moieties: implications with respect to macrophage recognition of apoptotic cells. *Proc. Natl. Acad. Sci. USA.* **96**: 6347–6352.
- Watson, A. D., N. Leitinger, M. Navab, K. F. Faull, S. Hörkö, J. L. Witztum, W. Palinski, D. Schwenke, R. G. Salomon, W. Sha, G. Subbanagounder, A. M. Fogelman, and J. A. Berliner. 1997. Structural identification by mass spectrometry of oxidized phospholipids in minimally oxidized low density lipoprotein that induce monocyte/endothelial interactions and evidence for their presence in vivo. *J. Biol. Chem.* **272**: 13597–13607.
- McIntyre, T. M., G. A. Zimmerman, and S. M. Prescott. 1999. Biologically active oxidized phospholipids. *J. Biol. Chem.* **274**: 25189–25192.
- Witztum, J. L., and J. A. Berliner. 1998. Oxidized phospholipids and isoprostanes in atherosclerosis. *Curr. Opin. Lipidol.* **9**: 441–448.
- Berliner, J. A., G. Subbanagounder, N. Leitinger, A. D. Watson,



- and D. Vora. 2001. Evidence for a role of phospholipid oxidation products in atherogenesis. *Trends Cardiovasc. Med.* **11**: 142–147.
28. Shih, D. M., Y. R. Xia, X. Wang, E. Miller, L. W. Castellani, G. Subbanagounder, H. Cheroutre, K. F. Faull, J. A. Berliner, J. L. Witztum, and A. J. Lusis. 2000. Combined serum paraoxonase knockout/apolipoprotein E knockout mice exhibit increased lipoprotein oxidation and atherosclerosis. *J. Biol. Chem.* **275**: 17527–17535.
29. Rozenberg, O., M. Rosenblat, R. Coleman, D. M. Shih, and M. Aviram. 2003. Paraoxonase (PON1) deficiency is associated with increased macrophage oxidative stress: studies in PON1-knockout mice. *Free Radic. Biol. Med.* **34**: 774–784.
30. Subbanagounder, G., N. Leitinger, and P. T. Shih. 1999. Evidence that phospholipid oxidation products and/or platelet-activating factor play an important role in early atherogenesis: in vitro and in vivo inhibition by WEB 2086. *Circ. Res.* **85**: 311–318.
31. Subbanagounder, G., Y. Deng, C. Borromeo, A. N. Dooley, J. A. Berliner, and R. G. Salomon. 2002. Hydroxy alkenal phospholipids regulate inflammatory functions of endothelial cells. *Vascul. Pharmacol.* **38**: 201–209.
32. Leitinger, N., T. R. Tyner, L. Oslund, C. Rizza, G. Subbanagounder, H. Lee, P. T. Shih, N. Mackman, G. Tigyi, M. C. Territo, J. A. Berliner, and D. K. Vora. 1999. Structurally similar oxidized phospholipids differentially regulate endothelial binding of monocytes and neutrophils. *Proc. Natl. Acad. Sci. USA.* **96**: 12010–12015.
33. Subbanagounder, G., N. Leitinger, D. Schwenke, J. Wong, H. Lee, C. Rizza, A. D. Watson, K. F. Faull, A. M. Fogelman, and J. A. Berliner. 2000. Determinants of bioactivity of oxidized phospholipids. Specific oxidized fatty acyl groups at the sn-2 position. *Arterioscler. Thromb. Vasc. Biol.* **20**: 2248–2254.
34. Binder, C. J., S. Hörkkö, A. Dewan, M. K. Chang, E. P. Kieu, C. S. Goodyear, P. X. Shaw, W. Palinski, J. L. Witztum, and G. J. Silverman. 2003. Pneumococcal vaccination decreases atherosclerotic lesion formation: molecular mimicry between *Streptococcus pneumoniae* and oxidized LDL. *Nat. Med.* **9**: 736–743.
35. Hart, K., and M. Wilcox. 1993. A drosophila gene encoding an epithelial membrane protein with homology to CD36/LIMPII. *J. Mol. Biol.* **234**: 249–253.
36. Franc, N. C., J. L. Dimarcq, M. Lagueux, J. Hoffman, and R. A. Ezekowitz. 1996. Croquemort, a novel *Drosophila* hemocyte/macrophage receptor that recognizes apoptotic cells. *Immunity.* **4**: 431–443.
37. Savill, J., N. Hogg, Y. Ren, and C. Haslett. 1992. Thrombospondin cooperates with CD36 and the vitronectin receptor in macrophage recognition of neutrophils undergoing apoptosis. *J. Clin. Invest.* **90**: 1513–1522.
38. Chang, M. K., C. J. Binder, Y. I. Miller, G. Subbanagounder, G. J. Silverman, J. A. Berliner, and J. L. Witztum. 2004. Apoptotic cells with oxidation-specific epitopes are immunogenic and proinflammatory. *J. Exp. Med.* **200**: 1359–1370.
39. Chang, M. K., C. Bergmark, A. Laurila, S. Hörkkö, K. H. Han, P. Friedman, E. A. Dennis, and J. L. Witztum. 1999. Monoclonal antibodies against oxidized low-density lipoprotein bind to apoptotic cells and inhibit their phagocytosis by elicited macrophages: evidence that oxidation-specific epitopes mediate macrophage recognition. *Proc. Natl. Acad. Sci. USA.* **96**: 6353–6358.
40. Binder, C. J., M. K. Chang, P. X. Shaw, Y. I. Miller, K. Hartvigsen, A. Dewan, and J. L. Witztum. 2002. Innate and acquired immunity in atherogenesis. *Nat. Med.* **8**: 1218–1226.
41. Chang, M. K., C. J. Binder, M. Torzewski, and J. L. Witztum. 2002. C-reactive protein binds to both oxidized LDL and apoptotic cells through recognition of a common ligand: phosphorylcholine of oxidized phospholipids. *Proc. Natl. Acad. Sci. USA.* **99**: 13043–13048.
42. Gillotte-Taylor, K., A. Boullier, J. L. Witztum, D. Steinberg, and O. Quehenberger. 2001. Scavenger receptor class B type I as a receptor for oxidized low density lipoprotein. *J. Lipid Res.* **42**: 1474–1482.