UC Berkeley
Berkeley Scientific Journal

Title
Bone is Listening: How Force Shapes the Skeleton

Permalink
https://escholarship.org/uc/item/282877q6

Journal
Berkeley Scientific Journal, 30(2)

ISSN
1097-0967

Author
Ha, Brandon


Publication Date
2026-11-05

DOI
10.5070/BS3.70132

Copyright Information
This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at https://creativecommons.org/licenses/by-nc-nd/4.0/

eScholarship.org Powered by the California Digital Library
University of California

Bone is Listening: How Force Shapes the Skeleton



BY: BRANDON HA

INTRODUCTION

In every step we take, waves of force are sent throughout the body. When a runner's foot strikes the concrete or a person simply walks across a room, mechanical forces travel upward through the skeletal system from the foot to the spine. These forces are distributed across bones, joints, and connective tissue.

Despite this constant mechanical activity, bones are often thought of as rigid, static structures. In reality, bone is a living tissue that continuously adapts to its environment. Unlike materials such as steel or concrete, which maintain fixed properties, bone undergoes constant remodeling through a balance of breakdown and regeneration. Certain cells remove old or damaged tissue while others build new bone, allowing the skeleton to adjust to the mechanical demands placed upon it. Thus, bone adaptation over time creates a structure that reflects an individual's unique patterns of movement.[1]

THE BONE AS A DYNAMIC TISSUE




Figure 1. Mechanical loading across the knee. Forces generated during movement are transmitted through the knee joint, placing stress on bone and surrounding structures. This mechanical loading is essential for maintaining bone strength as it stimulates adaptive remodeling in response to repeated use.

The relationship between mechanical force and bone structure has long been described by Wolff's Law, which states that bone adapts to the stresses placed upon

Thus bone adaptation over time creates a structure that reflects an individual's unique patterns of movement.

it. When mechanical loading increases through activities like walking, running, or weight-bearing motions, bone tissue responds by adapting and reinforcing itself. On the other hand, when loading decreases through prolonged immobility or exposure to microgravity (such as in outer space), bone density declines.[2]

This remodeling process is mainly governed by two primary types of cells: osteoclasts and osteoblasts. Osteoclasts are responsible for breaking down old or damaged bone, while osteoblasts form new bone tissue. These processes occur continuously and are tightly regulated and constantly updated. The balance between them is the main determining factor of whether bone becomes stronger or weaker in the body over time. Mechanical forces play a key role in maintaining this balance by initiating signaling pathways that regulate bone formation or resorption.[3]




Figure 3. Mechanotransduction in bone cells. Mechanical forces applied to bone are converted into biochemical signals through mechanotransduction pathways within cells. These signals regulate gene expression and cellular activity, allowing bone to adapt to changing mechanical demands.

OSTEOCYTES AS THE BONE'S INTERNAL SENSORS

At the center of this system are osteocytes, cells embedded within the mineralized bone matrix. Unlike osteoblasts and osteoclasts, which are located on bone surfaces, osteocytes are deep inside the tissue and form an interconnected network through microscopic channels known as canaliculi.[4]




Figure 2. Osteocyte and canaliculi network. Osteocytes are embedded within bone and connected through a network of tiny channels called canaliculi, as shown in this image. These channels allow cells to communicate and sense mechanical stress to help regulate bone remodeling.

Osteocytes function as the mechanosensors that are essential for changes in bone structure. When bone experiences mechanical stress, it undergoes small deformations that generate fluid movement within the surrounding matrix. Osteocytes detect these changes and respond by initiating signaling pathways that regulate bone remodeling.[4,5]

Through this process, osteocytes coordinate the activity of osteoblasts and osteoclasts. When mechanical loading is sufficient, signaling promotes bone formation. When loading is reduced, signaling shifts toward bone resorption. In this way, osteocytes act as a central regulatory system, continuously monitoring mechanical conditions and directing structural adaptation.

MECHANOTRANSDUCTION: CONVERTING FORCE INTO BIOLOGICAL SIGNALS

The ability of bone to respond to mechanical stress depends on a process known as mechanotransduction.

Mechanotransduction refers to the conversion of mechanical forces into biochemical signals within cells.

When force is applied to bone, it produces small deformations in the tissue. These microdeformations cause interstitial fluid within the bone matrix to move, generating shear stress on osteocytes. This mechanical stimulus is detected through cellular structures such as integrins and the cytoskeleton, which transmit the signal into the cell interior.[3,5]

Once activated, osteocytes initiate intracellular signaling pathways that regulate gene expression and protein synthesis. These pathways lead to release of signaling molecules that influence osteoblast and osteoclast activity. As a result, mechanical forces are translated into coordinated biological responses that guide bone growth, repair, and maintenance.[3]

This process allows mechanical forces to function as a form of biological information. Rather than simply exerting pressure on bone, forces provide signals that shape the structure and organization of skeletal tissue. Mechanotransduction is essential for maintaining bone integrity and adapting to changing mechanical demands.




Figure 4. Cellular regulation of bone remodeling. Bone remodeling is controlled by interactions between osteocytes, osteoblasts, and osteoclasts, which coordinate bone formation and resorption. Signaling molecules such as sclerostin help regulate this balance in response to mechanical conditions.

APPLICATION IN MOVEMENT AND BONE HEALTH

The effects of these cellular processes become most visible when comparing different loading conditions and their impact on skeletal health. Bone does not respond identically to every kind of movement; rather, its response depends on magnitude, frequency, and pattern of the forces it experiences. Weight-bearing and high-impact activities can support skeletal strength, but repetitive loading without enough recovery can also create problems.

In athletic settings, repeated strain can strengthen bone when recovery is sufficient. However, when loading accumulates faster than repair, microdamage can build up and increase the risk of stress fractures. These forces activate osteocytes, which promote osteoblast activity and increase bone density over time.[6] At the opposite extreme, non-weight-bearing activities provide less mechanical stimulation to the skeleton. While beneficial for cardiovascular health, they are less effective at promoting bone growth. Unloading the skeleton shows how dependent the skeleton is on continued mechanical input. In microgravity, astronauts can lose bone density because the usual weight-bearing forces acting on the skeleton are greatly reduced. Thus, interestingly, the same forces that can stimulate remodeling can also contribute to injury when the balance between loading and recovery is disrupted.




Figure 5. Bone fracture healing and remodeling. Bone healing occurs through a series of stages in which damaged tissue is gradually replaced and remodeled into new bone. Controlled mechanical loading supports this process, while imbalance in stress can delay recovery or lead to further injury.

INJURY AND REHABILITATION

Mechanotransduction also plays a key role in injury recovery. Bone healing requires carefully controlled mechanical loading. Insufficient loading can delay healing, while excessive force can disrupt the repair process.

Rehabilitation strategies often involve gradual reintroduction of mechanical stress to stimulate bone formation and restore structural integrity. Controlled loading has been shown to enhance fracture healing by promoting cellular activity and tissue regeneration.[7]

At the same time, excessive or repetitive loading without adequate recovery can lead to stress fractures. These injuries occur when the rate of bone damage exceeds the rate of repair, highlighting the importance of balance in the bone remodeling process.[8]

CONCLUSION

Bone is not a static framework but a dynamic system that continuously adapts to mechanical forces. Through the coordinated actions of osteocytes, osteoblasts, and osteoclasts, the skeleton is able to detect and respond to physical stress. Mechanotransduction serves as the link between mechanical forces and biological responses, allowing the body to translate movement into structural adaptation.

This process highlights the importance of physical activity in maintaining bone health. Everyday movements provide the mechanical stimuli necessary to sustain bone strength, while the absence of these forces can lead to significant structural declines. Ultimately, the structure of the human skeleton reflects not only genetic factors but also the cumulative effects of mechanical experience over time.

ACKNOWLEDGEMENTS

I would like to thank Morgan Pfaff for her thoughtful feedback and her willingness to help. Her insights greatly strengthened the clarity and scientific rigor of this work, as well as my knowledge and interest in the field.

REFERENCES

1. Turner, C. H. (1998). Three rules for bone adaptation to mechanical stimuli. Bone, 23(5), 399–407.
2. NASA. (2023). Bone Loss in Spaceflight.

3. Yavropoulou, M. P., & Yovos, J. G. (2016). The molecular basis of bone mechanotransduction. Journal of Musculoskeletal and Neuronal Interactions, 16(3), 221–236.
4. Bonewald, L. F. (2011). The amazing osteocyte. Journal of Bone and Mineral Research, 26(2), 229–238. https://doi.org/10.1002/jbmr.320
5. Robling, A. G., & Turner, C. H. (2009). Mechanical signaling for bone modeling and remodeling. Critical Reviews in Eukaryotic Gene Expression, 19(4), 319–338. https://doi.org/10.1615/CritRevEukarGeneExpr.v19.i4.50
6. Chang, X., Xu, S., & Zhang, H. (2022). Regulation of bone health through physical exercise: Mechanisms and types. Frontiers in Endocrinology, 13, 1029475. https://doi.org/10.3389/fendo.2022.1029475
7. Ma, Q., Miri, Z., Haugen, H. J., Moghanian, A., & Loca, D. (2023). Significance of mechanical loading in bone fracture healing, bone regeneration, and vascularization. Journal of Tissue Engineering, 14, 20417314231172573. https://doi.org/10.1177/20417314231172573
8. Hughes, J. M., Popp, K. L., Yanovich, R., Bouxsein, M. L., & Matheny, R. W., Jr. (2017). The role of adaptive bone formation in the etiology of stress fracture. Experimental Biology and Medicine, 242(9), 897–906. https://doi.org/10.1177/1535370216661646

IMAGE REFERENCES

Banner Image: Human Skeletal System SciePro GmbH, CC BY-SA 4.0 https://creativecommons.org/licenses/by-sa/4.0, via Wikimedia Commons
1. Joint and Bone pain InjuryMap, CC BY-SA 4.0 https://creativecommons.org/licenses/by-sa/4.0, via Wikimedia Commons
2. Osteocyte Matrix Eclairereno, CC BY 4.0 https://creativecommons.org/licenses/by/4.0, via Wikimedia Commons
3. Mechanotransduction Wu, D., Li, L., Wen, Z. et al., CC BY 4.0 https://creativecommons.org/licenses/by/4.0, via Wikimedia Commons
4. Mechanotransduction cascade in cells. Wang, N., Lu, Y., Rothrauff, B. B., Zheng, A., Lamb, A., Yan, Y., Lipa, K. E., Lei, G., & Lin, H. (2023). Mechanotransduction pathways in articular chondrocytes and the emerging role of estrogen receptor-α [Figure]. Bone Research, 11, 13. https://doi.org/10.1038/s41413-023-00248-x
5. Mechanotransduction pathways in articular chondrocytes and the emerging role of estrogen receptor-α [Figure]. Bone Research, 11, 13. https://doi.org/10.1038/s41413-023-00248-x