Lawrence Berkeley National Laboratory

Recent Work

Title

PHOTOSYNTHESIS OF ISOLATED CHLOROPLASTS. V. EFFECT ON FIXATION RATE AND METABOLITE TRANSPORT FROM THE CHLOROPLASTS CAUSED BY ADDED FRUCTOSE-1, 6-DIPHOSPHATASE.

Permalink

<https://escholarship.org/uc/item/28c5n37k>

Authors

Bassham, J.A.
El-Badry, A.M.
Kirk, M.R.
et al.

Publication Date

1970-07-01

e.2

RECEIVED
LAWRENCE
RADIATION LABORATORY

OCT 29 1970

LIBRARY AND
DOCUMENTS SECTION

PHOTOSYNTHESIS OF ISOLATED CHLOROPLASTS.
EFFECT ON FIXATION RATE AND METABOLITE
TRANSPORT FROM THE CHLOROPLASTS CAUSED
BY ADDED FRUCTOSE-1, 6-DIPHOSPHATASE

J. A. Bassham, A. M. El-Badry, M. R. Kirk,
H. C. J. Ottenheim, and H. Springer-Lederer

July 2, 1970

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

48
LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

e.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

PHOTOSYNTHESIS OF ISOLATED CHLOROPLASTS. V. EFFECT ON FIXATION
RATE AND METABOLITE TRANSPORT FROM THE CHLOROPLASTS CAUSED BY
ADDED FRUCTOSE-1,6-DIPHOSPHATASE

J. A. Bassham, A. M. El-Badry, M. R. Kirk, H.C.J. Ottenheim*
and H. Springer-Lederer**

Summary

Addition of fructose-1,6-diphosphatase (FDPase) plus Mg^{++} ions in increasing concentrations causes stimulation and then inhibition of photosynthesis in isolated spinach chloroplasts. The effects are similar to those previously found with leaf extract additions, including modifications of the effects caused by added inorganic pyrophosphate (PP_i). Under the conditions of pH, Mg^{++} ion concentration, and PP_i concentration in which these effects are seen, the added enzyme has only negligible activity as FDPase. Thus the enzyme appears to be exerting its effect through some function other than as an FDPase. It is tentatively suggested that this other function is in the transport of metabolites, especially fructose-1,6-diphosphate, out through the chloroplast outer membrane. This hypothesis is supported by the following observations:

1. Addition of FDPase plus Mg^{++} to chloroplasts photosynthesizing with $^{14}CO_2$ causes an increase in the ^{total}/labeled fructose-1,6-diphosphate /and dihydroxyacetone phosphate (which appear mostly in the suspending medium), and a decrease in the total labeled fructose-

* Present address: Department of Chemistry, Cornell University, Ithaca, New York.

** Present address: Institut für Physikalische Chemie, Universität Wien, Wien, Austria.

6-phosphate and of all compounds following fructose-1,6-diphosphate in the carbon reduction cycle, including the carboxylation substrate, ribulose-1,5-diphosphate. Furthermore, the amounts of all metabolites retained in the chloroplasts decrease.

2. Addition of Mg^{++} ions alone, in concentrations slightly higher than PP_i concentration, causes some loss of all compounds from the chloroplasts to the medium.

3. Addition of 5 mM PP_i decreases the loss of hexose and heptose monophosphates and of ribulose-1,5-diphosphate, but does not decrease the loss of FDP and of dihydroxyacetone phosphate from the chloroplasts to the medium.

Introduction

The rate of photosynthetic fixation of carbon dioxide is strongly affected by additions of small amounts of clarified spinach juice¹. The degree of inhibition or stimulation is a function of the ratio of amount of added spinach juice to the concentration of added inorganic pyrophosphate (PP_i). Fractionation of the juice by gel filtration gave two components which act synergistically to produce the stimulation or inhibition. The component of high molecular weight was identified as fructose-1,6-diphosphatase (D-fructose-1,6-diphosphate 1-phosphohydrolase, E.C.3.1.3.11)², hereafter referred to as FDPase. The component of low molecular weight, called "Factor B" in the previous reports^{1,2}, has now been found to

be an impure mixture of inorganic salts, of which the active constituent is Mg^{++} . The synergistic stimulation and inhibition, and their dependence on PP_1 , has been duplicated by additions to photosynthesizing chloroplasts of selected ratios of FDPase and Mg^{++} .

Addition of fructose-1,6-diphosphate ($Fru-1,6-P_2$) to relatively inactive isolated spinach chloroplasts stimulates their fixation rate³, while its addition to more active chloroplasts maintains a higher rate after 15 minutes of photosynthesis, when the rate begins to decline⁴. Other tested metabolites were generally inhibitory or without effect⁴. Thus the level of $Fru-1,6-P_2$ in chloroplasts might play some role in the regulation of the carboxylation of ribulose-1,5-diphosphate to give 3-phosphoglyceric acid.

The carboxylation reaction and the conversion of $Fru-1,6-P_2$ to $Fru-6-P$, as well as the conversion of sedoheptulose-1,6-diphosphate, appear to be metabolically-regulated steps in the photosynthetic carbon reduction cycle⁵⁻⁹. Both the carboxylase and the diphosphatases, when isolated and purified, exhibit a dependence of activity on pH and Mg^{++} concentration^{10,11} which could account at least in part for their regulation in vivo¹². Among the labeled metabolites formed in isolated spinach chloroplasts photosynthesizing with ¹⁴C-labelled bicarbonate, there are pronounced differences in the rates of movement out of the chloroplasts into the suspending medium¹³.

Those compounds which just precede the reactions mediated by the diphosphatases (Fru-1,6-P₂, dihydroxyacetone phosphate and glyceraldehyde phosphate) are found to a much greater extent outside the chloroplasts after a few minutes photosynthesis, whereas the hexose and heptose monophosphates, including Fru-6-P, and ribulose-1,5-diphosphate (Ribul-1,5-P₂), are very well retained. If Fru-1,6-P₂ moves out of the chloroplasts by means of facilitated transport, a part of the apparent regulation of FDPase in vivo could be in fact a regulation of this transport, provided FDPase in the cytoplasm is less active than that inside the chloroplasts.

We have now investigated the effects of FDPase, Mg⁺⁺ and PP_i, added together and separately, on the levels of labelled metabolites formed in photosynthesizing chloroplasts, and the effects of these additions on the movement of metabolites out of the chloroplasts. Most of the effects reported were under conditions of pH, Mg⁺⁺ concentration and PP_i level under which the added FDPase could have little or no activity for the conversion of Fru-1,6-P₂ to Fru-6-P. The effects in the medium of the added FDPase seem therefore to be indicative of another activity of this protein.

Experimental

Chloroplasts were prepared from fresh spinach as described previously¹⁴. Kinetic studies of the formation of labelled compounds and their movement from the chloroplasts to the medium during photosynthesis by the isolated chloroplasts with ¹⁴C-labelled bicarbonate were as described earlier¹³.

In some experiments, an attempt was made to remove enzymes external to the chloroplasts by "washing." The chloroplast pellet from the initial centrifugation¹⁴ was resuspended in 10 ml Solution B or Solution C and again centrifuged at 2000 x g for about 50 sec, after which the chloroplasts were stored for less than 10 min at 0°C until the beginning of the illumination at 20°.

In those experiments in which the distribution of metabolites between chloroplasts and medium was to be determined, a 100 µl sample of the suspension of photosynthesizing chloroplasts was removed at the indicated times, and this sample was subjected to 10 sec centrifugation with a Spinco microfuge. The pellet and the supernatant solution were separated and each was killed in 80% methanol. The elapsed time from sampling to killing was about 25 sec. Some samples were killed immediately, without separation of chloroplast from medium, to permit subsequent evaluation of metabolite conversion during centrifugation.

The supernatant solution could not be completely separated from the pellet, and a contamination of the pellet by 10% of

the supernatant solution is estimated. In those cases where a very large portion of a given compound is found in the supernatant solution, a correction has been made in the results. This correction, equal to 0.10 of the ^{14}C found in the compound in the supernatant solution, is subtracted from the value found for the pellet and added to the value found for the supernatant solution.

For determination of the rates of photosynthetic incorporation of $^{14}\text{CO}_2$, aliquot samples of the killed Δ -methanol-water mixtures were dried on filter paper, free CO_2 was driven off by addition of acetic acid, and the ^{14}C content was measured with a Geiger-Müller tube and scaler. Chlorophyll content was determined in a sample of the chloroplast suspension, and rate of photosynthesis was calculated from known specific radioactivity of bicarbonate used, ^{14}C content and chlorophyll content¹⁴. Products of photosynthesis were analyzed by two-dimensional paper chromatography and autoradiography, as previously described^{5,6}.

A solution of purified fructose diphosphatase was prepared from isolated spinach chloroplasts as reported earlier². The eluate from the gel filtration through Sephadex G-200 was dialyzed against "Solution C"¹⁴ minus PP_i and ascorbate, and the dialyzed enzyme solution was diluted 25 times in Solution C minus ascorbate and PP_i . Amounts of this diluted solution, ranging up to 30 μg of protein, were added to the chloroplasts in the illumination flasks, and the suspensions were made up to 0.5 ml for rate experiments or 1.0 ml for kine-

tic experiments, as indicated in the Results. Other additions of Mg^{++} , PP_1 , etc. are given under Results.

Results

The rate of photosynthesis by isolated spinach chloroplasts is stimulated 25% by the addition of small amounts of magnesium ion and purified FDPase and is strongly inhibited by the addition of greater amounts of Mg^{++} and FDPase (Fig. 1). The shape of this stimulation-inhibition curve and its dependence on the presence of 5 mM PP_1 are exactly analogous to the effects earlier observed upon the addition of various amounts of spinach juice to the isolated photosynthesizing chloroplasts¹.

The inhibition of photosynthetic rate by Mg^{++} and FDPase is reversed by higher levels of PP_1 (Fig. 2). As was the case with spinach juice¹, or with FDPase plus "Factor B"², the concentration of added PP_1 required to reverse inhibition is proportional to the amount of inhibiting substances added. Thus, it is clear that the effects formerly observed upon adding small amounts of spinach juice to the isolated chloroplasts, including dependence on PP_1 concentration, can be reproduced by the addition of purified FDPase and Mg^{++} .

Analysis of purified "Factor B"², by atomic absorption spectroscopy, showed that it contained approximately 10% Mg^{++} . In the presence of the usual suspending medium ($-Mg^{++}$) which contained 5 mM PP_1 , addition of 21 μg of purified FDPase caused 25% stimulation in the rate of photosynthesis (Table I), whereas addition of

enough Mg^{++} to make its concentration 2 mM, caused a 10% stimulation in the absence of added FDPase but an 85% inhibition in the presence of the FDPase. Since the total volume was 0.5 ml, the added Mg^{++} was 1 mM above the usual reaction mixture concentration and corresponded to 0.5 μmole , or 12 μg , of Mg^{++} . This may be compared with the synergistic effect found previously² upon the addition of 24 μg of fructose diphosphatase and 150 μg of "Factor B", corresponding to 15 μg of Mg^{++} .

At 1 mM PP_1 (Table I) the inhibition by fructose diphosphatase plus Mg^{++} is even more striking. It should be particularly noted that whereas the stimulation with fructose diphosphatase alone was 90%, and the stimulation by 1.5 mM Mg^{++} was 12%, the addition of both factors caused an inhibition of 96%.

When FDPase and Mg^{++} were added to photosynthesizing chloroplasts after 6 min of photosynthesis with ^{14}C -labelled bicarbonate, rather surprising effects on the levels of labelled metabolites occurred (Figs. 3 and 4). As compared with the levels of metabolites in the control chloroplasts, the level of Fru-1,6- P_2 in chloroplasts with added FDPase increased, while the level of Fru-6-P decreased. This behaviour has proved to be reproducible, as have other differences observed (Table II). These include an increased level of dihydroxyacetone phosphate, which is rapidly equilibrated with fructose-1,6-diphosphate in all of our experiments, due to high aldolase activity both in the chloroplasts and in the medium. The kinetic curves for

dihydroxyacetone phosphate are, in all cases, similar to the ones for fructose-1,6-diphosphate. Compounds with levels which decreased upon FDPase and Mg^{++} addition included glucose-6-phosphate, sedoheptulose-7-phosphate, and sedoheptulose-1,6-diphosphate. The level of sedoheptulose-1,7-diphosphate declined very rapidly upon FDPase and Mg^{++} addition (Fig. 5). The carboxylation substrate, ribulose-1,5-diphosphate (Ribul-1,5- P_2) also decreased considerably, and this could be partly responsible for the decreased rate of CO_2 fixation. However, it should be noted that the levels of pentose monophosphate, which is the sum of ribulose-5-phosphate, xylulose-5-phosphate and ribose-5-phosphate, were not changed by the addition of Mg^{++} and FDPase.

The effects of addition of FDPase and Mg^{++} on the distribution of metabolites between the chloroplasts and the suspending medium are very large (Table III). In the chloroplasts with these additions the level of fructose-1,6-diphosphate in the medium increased 11-fold, and the level of dihydroxyacetone phosphate increased 4-fold. At the same time, the level of fructose-6-phosphate in the pellet declined 75%, its level in the medium increased 2-fold, and its total level declined about 40%. Thus, the addition of FDPase and Mg^{++} ions has increased greatly the appearance of metabolites in the suspending medium. However, the added FDPase has not increased/ ^{conversion of} fructose-1,6-diphosphate to fructose-6-phosphate in the total, although some increased conversion in the medium may have occurred.

When enough FDPase and Mg^{++} are present from the start of photosynthesis to cause inhibition, the relative effects on the distribution of metabolites are similar to the effects when Mg^{++} and FDPase are added after several minutes of photosynthesis (Table IV). The ^{14}C levels in all compounds are, of course, decreased, but relatively greater decreases are seen in hexose and heptose monophosphate compared with dihydroxyacetone phosphate which is proportional to Fru-1,6-P₂.

A comparison of $^{14}CO_2$ fixation rates in the presence of added FDPase and bovine serum albumin (Table V), each added with Mg^{++} , shows that inhibition is not a general effect of protein addition.

The effects of addition of FDPase alone, without enough Mg^{++} ion to cause a synergistic effect in the presence of 5 mM PP_1 , were minor (Table VI). There was a slight stimulation of $^{14}CO_2$ fixation rate, but levels of individual compounds in chloroplasts and medium were virtually unchanged.

The effects of addition of 5 mM Mg^{++} alone, after 7 min photosynthesis in the normal medium (Solution C), were both qualitatively very similar to the effects of adding both FDPase and Mg^{++} in one experiment (Table VII). However, the photosynthetic rate in this experiment with Mg^{++} addition only was only 30 μ moles CO_2 fixed (mg chlorophyll hr)⁻¹, lower than usual, suggesting a greater than usual breakage of chloroplasts during the course of the experiment and, consequently,

the presence of considerable FDPase in the medium, coming from broken chloroplasts. This sort of difficulty can never be entirely overcome under the conditions which we use, and it is probable that there is always present in the medium some FDPase, as well as all of the other enzymes from the chloroplasts.

In another experiment (Table IV) Mg^{++} concentration was such that no inhibition was seen without added FDPase. In that case, no increased transport of metabolites was seen with Mg^{++} without added FDPase.

Although most of these enzymes retain some activity in the medium, we have found that there is no appreciable activity of either FDPase or ribulose diphosphate carboxylase (E.C.4.1.1.39) in the medium. In other words, although the FDPase protein is present in the medium, as a consequence of chloroplast breakage, its activity as FDPase is negligible. When chloroplasts were placed in the light in Solution C for 5 min and centrifuged from the medium, assay of the medium for FDPase activity showed an activity of less than 0.1 nmoles Fru-1,6- P_2 per min which may be compared with the CO_2 fixation rate of the chloroplasts of 20 nmoles per min, based on a rate of 60 μ moles (hr·mg chlorophyll) $^{-1}$ and 50 μ g chlorophyll in the chloroplasts in the flask. Also, it may be shown that the 10 μ g of FDPase added gave an activity of only 0.2 nmoles Fru-1,6- P_2 converted per min in Solution C, but would have an activity of 100 nmoles/min at pH 8.7 and 5 mM Mg^{++} .

When photosynthesis and metabolite transport of chloroplasts with 5 mM PP_1 are compared with that of chloroplasts without PP_1 (Solution C minus PP_1), the principal effects are seen to be the improved retention (with PP_1) in the chloroplasts of those metabolites which, under the usual conditions, tend to be retained, namely, hexose and heptose monophosphates and ribulose-1,5-diphosphate (Table VIII).

Additions of low concentrations of Fru-1,6- P_2 chloroplast suspensions (Table IX) before the start of photosynthesis result in some stimulation. Larger concentrations of added Fru-1,6- P_2 produce inhibition of CO_2 uptake. These effects also depend somewhat on the concentration of added PP_1 , the stimulation being somewhat more pronounced with 1 mM PP_1 than with 5 mM PP_1 whereas the inhibition is greater in the presence of 5 mM PP_1 .

Addition of 1 mM Fru-1,6- P_2 to chloroplasts after 5 min of photosynthesis with $^{14}CO_2$ (Table X) causes inhibition (1.9 μ moles $^{14}CO_2$ fixed per mg chlorophyll during the next 6 min compared with 6.7 μ moles fixed in the control and 7.7 μ moles fixed in the presence of added Fru-6-P). The effects of added Fru-6-P are rather minor, suggesting little penetration of this compound into the chloroplasts. The effects that are seen with added Fru-1,6- P_2 (note the "flushing of the ^{14}C from the Fru-6-P and other subsequent pools) suggest that the Fru-1,6- P_2 does enter the chloroplasts readily.

Discussion

The principal regulatory effects on the rate of photosynthesis by isolated chloroplasts previously ascribed to a substance of high molecular weight plus a substance of low molecular weight in the spinach juice¹ can be completely duplicated by the addition of appropriate amounts of fructose diphosphatase and of Mg^{++} . Fru-1,6- P_2 comes out rapidly from the isolated photosynthesizing spinach chloroplasts, while Fru-6-P is well retained¹³. It might have been postulated that added FDPase converts Fru-1,6- P_2 to Fru-6-P which cannot re-enter the chloroplast. This would remove carbon from the photosynthetic carbon reduction cycle, leading to a shortage of carboxylation substrate, ribulose-1,5-diphosphate. The fact that PP_i is an inhibitor of fructose diphosphatase¹⁶ while Mg^{++} is an activator of this enzyme, would be consistent with this assumption.

However, the data presented force the conclusion that the primary effect of added FDPase on photosynthesis by isolated chloroplasts is not produced by means of its function in converting Fru-1,6- P_2 to Fru-6-P.¹⁾ FDPase is nearly inactive under the conditions that exist in the suspending medium of the chloroplasts. 2) The level of labelled Fru-1,6- P_2 increases and the level of labelled Fru-6-P decreases on addition of the fructose diphosphatase. 3) Addition of FDPase leads to lower levels of hexose and heptose monophosphates and of Ribul-1,5- P_2 within the pellet and to an increased level of

Fru-1,6-P₂ in the supernatant solution. Similar effects are seen with added Mg⁺⁺ and the effects of Mg⁺⁺ and FDPase are synergistic.

The ratio of the volume of the supernatant solution to the chloroplast volume is of the order of 50:1. Thus, even for those metabolites which rapidly appear in the suspending medium, active transport is not required. Nevertheless, given the very considerable difference in rates of movement of metabolites from chloroplast to medium (e.g., Fru-1,6-P₂ vs. Fru-6-P), it seems clear that some kind of facilitated transport is required. We conclude that the facilitated transport of Fru-1,6-P₂ from the chloroplast to the suspending medium is enhanced by the addition of Fru-1,6-P₂ to the medium, and by the addition of Mg⁺⁺ to the medium. We tentatively conclude that the protein which can act as a fructose diphosphatase under suitable conditions is capable of being absorbed somewhere in the outer chloroplast double membrane and there functions as a transport enzyme. Such activity apparently would be stimulated by Mg⁺⁺ and inhibited by pyrophosphate, just as is its activity as a diphosphatase. Studies of metabolite transport and other physical and chemical properties of the chloroplast membrane probably will be required before such a postulate can be proved or disproved.

The effect of added pyrophosphate is to increase the levels of hexose and heptose monophosphates and of Ribul-1,5-P₂

within the chloroplasts and to diminish their appearance in the supernatant solution. However, the levels of Fru-1,6-P₂ and of dihydroxyacetone phosphate appearing and remaining in the supernatant solution are greater in the presence of pyrophosphate than in its absence. This may be due to the higher rate of photosynthesis and larger amount of reduced carbon when pyrophosphate is present.

It now appears that only a few steps in the photosynthetic carbon reduction cycle are necessarily restricted to the inside of the chloroplasts in the suspension of isolated chloroplasts which unavoidably contains broken chloroplasts and soluble enzymes in the medium. The steps which apparently occur exclusively inside the chloroplasts are, as might be expected, the rate-limiting steps or regulated steps, that is, the diphosphatase and the carboxylation reactions. Reactions mediated by aldolase, transketolase and isomerases appear to occur to some extent outside the chloroplasts. Probably the reduction of phosphoglyceric acid to triose phosphate and the phosphorylation of ribulose-5-phosphate to ribulose diphosphate also occur almost entirely in the chloroplasts.

It was found in earlier studies¹³ and again in the present investigation (see Table VIII) that levels of Fru-1,6-P₂ and dihydroxyacetone in the supernatant solution rise very high by about 6 to 10 min, and then decline between 10 and 20 min of photosynthesis in isolated chloroplasts. Apparently, the longer the concentration of these metabolites stays above a low level, the longer the rate of CO₂ uptake is maintained at

a high level. It was found earlier that addition of Fru-1,6-P₂ to the chloroplasts tends to maintain a high rate at a time when the rate would decline. The finding that small amounts of Fru-1,6-P₂ stimulates fixation of CO₂ while larger amounts inhibits provides further evidence for the regulatory effects of Fru-1,6-P₂.

The simplest explanation of the regulatory effects on CO₂ fixation rate of FDPase, Mg⁺⁺ and PP_i is based on the respective effects of these factors on the movement of metabolites out from the chloroplasts into the suspending medium. The retention of hexose and heptose monophosphates, and especially of ribulose-1,5-diphosphate in the chloroplasts, probably helps to maintain CO₂ fixation. In addition, it appears that there may be a direct regulatory effect of Fru-1,6-P₂ concentration on the rate of the carboxylation reaction in whole chloroplasts. The mechanism of such a regulation might operate either on the carboxylation enzyme or on the CO₂ transport into the chloroplasts.

It seems possible that FDPase in the outer chloroplast membrane might play an important role in regulation of photosynthetic metabolism in vivo. The postulated role would be in the movement of carbon compounds from the carbon reduction cycle in the chloroplast to other types of metabolism in the cytoplasm. The control of this "transport" activity could be largely mediated by Mg⁺⁺ ion and PP_i concentration inside and outside the chloroplast outer membrane.

Acknowledgements

This work was supported, in part, by the U.S. Atomic Energy Commission.

H. S.-L. was a public Health Service International Post-doctoral Research Fellow (number FO 5 TW 1094-01).

We are indebted to Dr. Tamotsu Kanazawa for valuable discussion and helpful suggestions during the course of these studies.

Laboratory of Chemical Biodynamics
Lawrence Radiation Laboratory
University of California
Berkeley, California 94720
U.S.A.

References

1. R. E. Moore, H. Springer-Lederer, H.C.J. Ottenheim and J. A. Bassham, Biochim. Biophys. Acta, 180 (1969) 368-376.
2. H. Springer-Lederer, A. M. El-Badry, H.C.J. Ottenheim and J. A. Bassham, Biochim. Biophys. Acta, 189 (1969) 464-467.
3. M. Gibbs and E. S. Bamberger, Plant Physiol. 37 (1962) 1Xii.
4. R. G. Jensen and J. A. Bassham, Biochim. Biophys. Acta, 153 (1968) 219-226.
5. T. A. Pedersen, M. Kirk and J. A. Bassham, Physiol. Plant. 19 (1966) 219-231.
6. T. A. Pedersen, M. Kirk and J. A. Bassham, Biochim. Biophys. Acta, 112 (1966) 189-203.
7. T. Kanazawa, K. Kanazawa, M. R. Kirk and J. A. Bassham, Plant & Cell Physiol. 11 (1970) 149-160.
8. R. G. Jensen and J. A. Bassham, Biochim. Biophys. Acta, 153 (1968) 227-234.
9. J. A. Bassham and G. H. Krause, Biochim. Biophys. Acta, 189 (1969) 207-221.
10. J. Preiss, M. L. Biggs and E. Greenberg, J. Biol. Chem. 242 (1967) 2292-2299.
11. J. A. Bassham, Pamela Sharp and Ian Morris, Biochim. Biophys. Acta, 153 (1968) 898-900.
12. J. A. Bassham and M. Kirk, in "Comparative Biochemistry and Biophysics of Photosynthesis", ed. K. Shibata, A. Takamiya, A. T. Jagendorf and R. C. Fuller, University of Tokyo Press, 1968, pp. 365-378.

13. J. A. Bassham, Martha Kirk and R. G. Jensen, Biochim. Biophys. Acta, 153 (1968) 211-218.
14. R. G. Jensen and J. A. Bassham, Proc. Natl. Acad. Sci. U.S. 56 (1966) 1095-1101.
15. I. Morris, Biochim. Biophys. Acta, 162 (1968) 462-464.

TABLE I

Effects of Mg^{++} and of FDPase on Rates of $^{14}CO_2$
Uptake by Isolated Spinach Chloroplasts

PPi (mM)	Enzyme (μ g)	Mg^{++} (mM)	Rate $\frac{\mu\text{moles } ^{14}C}{\text{mg. Chl. hr.}}$	% Stimulation (+) or Inhibition (-)
5	0	0	101	0
5	21	0	126	+25
5	0	1.0	115	+15
5	21	1.0	138	+38
5	0	2.0	110	+10
5	21	2.0	17	-83
5	0	5.0	48	-52
5	21	5.0	2.3	-98
1	0	0	83	--
1	8.4	0	158	+90
1	0	1.0	115	+38
1	8.4	1.0	108	+30
1	0	1.5	93	+12
1	8.4	1.5	3.5	-96

TABLE II

Effect of FDPase and Mg^{++} on ^{14}C Incorporation into Metabolites

Compound	$\mu\text{moles } ^{14}C/\text{mg. chlorophyll}$		
	After 5 min. PS	After 10 min. PS	
		W/O Mg^{++} & FDPase (control)	4 min. after addn. of Mg^{++} & FDPase
Total ^{14}C uptake	3.65	9.00	6.68
3-P-glycerate	1.38	3.18	2.39
dihydroxyacetone	0.28	0.42	1.15
Fru-1,6- P_2	0.07	0.265	0.37
Fru-6-P	0.166	0.25	0.134
Glc-6-P	0.115	0.375	0.150
Sed-7-P	0.20	0.325	0.060
Sed-1,7- P_2	0.073	0.08	0.009
pentose-P	0.070	0.22	0.22
Ribul-1,5-P	0.019	0.028	0.006
Origin	0.40	1.26	0.80
Misc. spots	0.363	0.528	0.39
Total on chromatograms	3.14	6.93	5.68
% accounted for	85.9	77.0	85.0
PS rate	43.8 $\mu\text{m}/\text{mg}/\text{hr}$	54.0 $\mu\text{m}/\text{mg}/\text{hr}$	40.1 $\mu\text{m}/\text{mg}/\text{hr}$

TABLE III
Effect of Added Mg^{++} and FDPase on Distribution of Metabolites between Chloroplasts and
Suspending Medium

Compound	$\mu\text{moles } ^{14}\text{C/mg chlorophyll}$									
	after 5 min. PS av. of A & B					after 12 min. PS & 6.5 min. after Mg^{++} and FDPase in B				
	Pellet	Supernatant	Total	Pellet		Supernatant		Total		
				A	B	A	B	A	B	Total
Total ^{14}C incorp.			6.50					13.72	15.61	
3-P-glycerate	0.55	2.32	3.53	0.75	0.40	3.08	3.75	7.92	5.70	
dihydroxyacetone-P	0.04	0.73	0.95	0.04	0.03	0.60	2.45	0.74	3.01	
Fru-1,6-P ₂	0.02	0.19	0.27	0.01	0.01	0.11	1.32	0.19	1.49	
Fru-6-P	0.094	0.024	0.186	0.202	0.054	0.054	0.110	0.303	0.184	
Glc-6-P	0.240	0.040	0.275	0.529	0.095	0.080	0.136	0.588	0.280	
Sed-7-P	0.36	0.03	0.034	0.715	0.185	0.090	0.075	0.675	0.190	
Sed-1,7-P ₂	0.015	0.012	0.135	0.014	0.004	0.005	0.029	0.115	0.066	
pentose-P	0.03	0.084	0.13	0.043	0.025	0.227	0.146	0.320	0.180	
Ribul-1,5-P ₂	0.016	0.005	0.046	0.038	0.013	0.008	0.109	0.078	0.166	
origin			0.80					1.73	1.55	
disaccharide	0.01	0.018	0.018	0.020	0.012	0.130	0.082	0.112	0.08	
glyceraldehyde-3-P	0.004	0.062	0.075	0.006	--	0.048	0.279	0.050	0.431	
miscellaneous								0.095	0.931	
Total from chromatograms			6.76					12.92	14.26	
% accounted for			104.0					94.1	91.3	
PS rate ($\mu\text{m } ^{14}\text{C/mg chl/hr}$)			78.0							

Flask A: control { 1 mM Mg^{++}
5 mM PP1

Flask B: { Same as Flask A for 5.5 min. of PS time
{ Mg^{++} to make 6 mM and 10.2 μg . FDPase added after 5.5 min. "washed chloroplasts"

TABLE IV

Effect of Mg^{++} and FDPase Added from the Beginning of Photosynthesis on the Level of Metabolites Seen after 5 min with $^{14}CO_2$
 1 mM Mg^{++} in control; 5 mM PP_i

Compound	% of total fixed					
	control	2 mM Mg^{++}	3 mM Mg^{++}	2 mM Mg^{++} 21 μg FDPase	3 mM Mg^{++} 21 μg FDPase	21 μg FDPase
3-P-glycerate	15	17	16	15	9.6	26
dihydroxyacetone-P	28	27	29	35	39	26
Frc-6-P	2.3	2.0	1.9	1.6	1.4	1.6
Sed + Glc-P	5.3	4.8	4.3	3.8	3.8	4.0
Pentose-P	1.2	1.2	1.3	1.2	1.3	1.3
$\mu moles \quad ^{14}C/mg \text{ chlorophyll}$						
Total ^{14}C fixed	5.43	5.68	5.39	3.83	0.770	4.32

TABLE V

Comparison of Effects of Addition of FDPase and of Bovine Serum Albumin on Photosynthesis Rate in Isolated Chloroplasts

5 min. photosynthesis

1 mM PP_i

1.5 mM Mg^{++}

μg protein/ flask	Relative rate of photosynthesis	
	FDPase	Bovine Serum Albumin
Control: "0"	1.0	1.0
8.4 μg	0.17	1.26

PS rate of control: 44.3 μm $^{14}C/mg.chl/hr$

TABLE VI

Effect of Added FDPase on Distribution of Metabolites between Chloroplasts and Suspending Medium

1 mM Mg⁺⁺
5 mM Ppi

"washed chloroplasts"

Compound	A: control		B: 30 μ g FDPase added after 5.5 min. PS		μ moles 14 C/mg chlorophyll					
	after 5 min. PS av. of A & B		after 12 min. PS-FDPase in B for 6.5 min.							
	Pellet	Supernatant	Total		Pellet		Supernatant		Total	
					A	B	A	B	A	B
Total 14 C incorp.			3.27						6.82	7.55
3-P-glycerate	0.35	1.09	1.52		0.460	0.460	2.28	2.64	2.49	3.16
dihydroxyacetone-P	0.060	0.390	0.60		0.115	0.030	0.615	0.560	0.953	0.816
Fru-1,6-P ₂	0.007	0.053	0.105		0.016	0.009	0.121	0.076	0.196	0.178
Fru-6-P	0.095	0.014	0.175		0.157	0.109	0.033	0.028	0.277	0.187
Glc-6-P	0.225	0.033	0.210		0.380	0.210	0.055	0.063	0.425	0.042
Sed-7-P	0.270	0.010	0.183		0.500	0.370	0.082	0.065	0.460	0.363
Sed-1,7-P ₂	0.017	0.003	0.085		0.020	0.007	0.005	0.003	0.116	0.022
pentose-P	0.025	0.085	0.125		0.058	0.041	0.245	0.320	0.380	0.392
Ribul-1,5-P ₂	0.016	0.002	0.027		0.017	0.007	0.003	0.003	0.038	0.050
misc. spots, inc. origin			0.344						1.235	1.339
Total from chromatograms			3.37						6.57	6.93
% accounted for			103.0						96.3	91.8
PS rate at 5 min (μ m 14 C/mg.chl/hr)			39.2							

TABLE VII

Effect of Addition of Mg^{++} (5 mM) on ^{14}C Incorporation into Metabolites in Chloroplasts and
 Suspending Medium "washed chloroplasts"

Compound	5 mM PPi $\mu\text{moles } ^{14}C/\text{mg chlorophyll}$									
	After 6 min. PS		no Mg^{++}		after 15 min. PS		8 min. after addn. of 5 mM		Mg^{++} fo flask B	
			av. of							
	Pellet	Supernatant	Total	2 flasks	Pellet	Supernatant	Total	A	B	Total
Total ^{14}C incorp.										
3-P-glycerate	0.127	1.22	3.32		0.25	0.10		3.75	3.00	8.6
dihydroxyacetone-P	0.033	0.48	0.58		0.03	0.01		0.27	0.59	5.0
Fru-1,6-P ₂	0.01	0.18	0.18		0.012	0.005		0.09	0.70	0.41
Fru-6-P	0.055	0.009	0.078		0.076	0.022		0.028	0.12	0.15
Glc-6-P	0.132	0.008	0.111		0.154	0.05		0.05	0.076	0.11
Sed-7-P	0.158	0.006	0.095		0.228	0.05		0.043	0.065	0.227
Sed-1,7-P ₂	0.015	0.008	0.087		0.021	0.003		0.005	0.019	0.244
pentose-P	0.021	0.048	0.073		0.025	0.008		0.13	0.165	0.100
Ribul-1,5-P ₂	0.007	0.001	0.017		0.02	0.006		0.003	0.002	0.17
origin			0.46							0.04
miscellaneous			0.83							1.41
Total from chromatograms			3.38							0.556
% accounted for			102							8.42
PS rate ($\mu\text{m } ^{14}C/\text{mg chl/hr}$)			33.2							97.9
										95.4

TABLE VIII
Effect of PPI on ^{14}C Incorporation into Metabolites in Chloroplasts and Suspending Medium

Compound	$\mu\text{moles } ^{14}\text{C/mg chlorophyll}$											
	after 6 min. PS						after 15 min. PS					
	Pellet			Supernatant			Pellet			Supernatant		
	+PPI	-PPI		+PPI	-PPI	Total	+PPI	-PPI		+PPI	-PPI	Total
Total ^{14}C incorp.	0.45	0.27		1.19	1.15	6.1	3.92					
3-P-glycerate	0.013	0.013		0.50	0.35	1.41	1.22		0.84	0.33	1.63	9.1
dihydroxyacetone-P	0.010	0.008		0.215	0.086	0.275	0.19		0.01	0.001	0.150	1.77
Fru-1,6-P ₂	0.102	0.043		0.018	0.035	0.165	0.162		0.003	0.001	0.020	0.10
Fru-6-P	0.282	0.10		0.030	0.020	0.290	0.133		0.145	0.020	0.028	0.05
Glc-6-P	0.390	0.145		0.022	0.010	0.261	0.125		0.40	0.050	0.067	0.198
Sed-7-P	0.006	0.001		0.007	0.012	0.033	0.024		0.34	0.090	0.085	0.494
Sed-1,7-P ₂	0.017	0.007		0.080	0.050	0.129	0.064		0.003	--	0.002	0.44
pentose-P	0.016	0.010		0.002	0.002	0.029	0.018		0.030	0.008	0.123	0.015
Ribul-1,5-P ₂	--	--		0.002	0.004	0.002	0.005		0.037	0.010	0.004	0.181
P-enolpyruvate									--	0.00140	0.0024	0.054
origin												0.003
disaccharide	0.014	0.010		0.028	0.023	0.036	0.028		0.037	0.022	0.180	1.08
miscellaneous												0.95
total on chromatograms												0.201
% accounted for												0.922
PS rate ($\mu\text{m } ^{14}\text{C/mg chl/hr}$)												1.012
												5.598
												4.213
												61.5
												69.1

TABLE IX

Effect of Concentration of Added Fru-1,6-P₂ on ¹⁴CO₂ Uptake During
5 min Photosynthesis by Isolated Chloroplasts

Conc. Fru-1,6-P ₂ added, mM	Fixation of ¹⁴ C	
	μmoles (mg.chlorophyll) ⁻¹	
	1 mM PP _i	5 mM PP _i
0	3.6	4.5
0.1	5.5	5.3
0.2	6.4	5.6
0.5	5.3	3.8
1.0	3.7	1.9

Rates (no Fru-1,6-P₂ added):

43.4 μm/mg/hr with 1.0 mM PP_i;

53.8 μm/mg/hr with 5.0 mM PP_i

Effect of Added Fru-1,6-P₂ and of Fru-6-P on Photosynthesis Rate
and Incorporation of ¹⁴C into Metabolites

Compound	μmoles of ¹⁴ C/mg chlorophyll			
	after 5 min PS no addns	control	6 min after addn 1 mM	6 min after addn 1 mM Fru-1,6-P ₂
Total ¹⁴ C incorp.	8.1	14.8	15.8	10.0
3-P-glycerate	2.65	4.75	4.60	2.70
dihydroxyacetone-P	0.93	1.19	1.27	1.38
Fru-1,6-P ₂	0.23	0.74	0.50	1.15
Fru-6-P	0.205	0.26	0.37	0.05
Glc-6-P	0.225	0.435	0.425	0.10
Sed-7-P	0.235	0.25	0.35	0.045
Sed-1,7-P ₂	0.115	0.145	0.460	0.075
pentose-P	0.120	0.180	0.66	0.11
Ribul-1,5-P ₂	0.035	0.064	0.062	0.01
origin	0.70	1.92	2.00	0.85
miscellaneous	0.425	0.896	1.028	0.81
Total on chromatogram	5.87	10.83	11.73	7.28
& accounted for	72.5	73.2	76.6	72.8
PS rate (μm ¹⁴ C/ mg.chl/hr) ⁻¹	97.2			

FIGURE CAPTIONS

Figure 1. Stimulation and inhibition of $^{14}\text{CO}_2$ fixation in isolated spinach chloroplasts by added fructose-1,6-diphosphatase and Mg^{++} . ○ 5 mM PP_i ; ● no PP_i . Relative rate of 1.0 obtained with 5 mM PP_i and 1.0 mM Mg^{++} . (Solution C) was 74.4 $\mu\text{moles } ^{14}\text{C}$ (mg chlorophyll hr) $^{-1}$. Mg^{++} and FDPase were added to the reaction flasks (total volumes 0.5 ml) at the start of the preillumination period to give the concentrations indicated. After 5 min preillumination, $\text{H}^{14}\text{CO}_3^-$ was added and the chloroplasts were allowed to photosynthesize for 5 min.

Figure 2. Dependence of inhibition of CO_2 fixation on concentration of added PP_i .

- ▲ 1 mM Mg^{++} , no FDPase
- 3 mM Mg^{++} , 4.2 μg FDPase
- 6 mM Mg^{++} , 10.5 μg FDPase

Control rate of 1.0, with 1 mM Mg^{++} , 5 mM PP_i (Solution C) was 120 $\mu\text{moles } ^{14}\text{CO}_2$ fixed (mg chlorophyll hr) $^{-1}$. Other conditions as in Fig. 1.

Figure 3. Effect of addition of FDPase and Mg^{++} on level of labelled Fru-1,6- P_2 in photosynthesizing spinach chloroplasts.

- Control (chloroplasts in Solution C)
- 17 μg FDPase and Mg^{++} to give 4 mM added after 6 min photosynthesis

Figure 4. Effect of addition of FDPase and Mg^{++} on level of labelled Fru-6-P in photosynthesizing spinach chloroplasts

- Control (chloroplasts in Solution C)
- 17 μ g FDPase and Mg^{++} to give 4 mM Mg^{++}

Figure 5. Effect of addition of FDPase and Mg^{++} on level of labelled Sedul-1,7- P_2 in photosynthesizing spinach chloroplasts

- Control (chloroplasts in Solution C)
- 17 μ g FDPase and Mg^{++} to give 4 mM Mg^{++}

XBL 706-5259

Fig. 1

XBL 705-5201

Fig. 2

XBL 704-5181

Fig. 3

XBL 704-5179

Fig. 4

XBL 704-5182

Fig. 5

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or*
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.*

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720