

UCLA

Proceedings of the UCLA Department of Medicine

Title

A Case of Septic Arthritis with Cryptococcus Neoformans in a Patient with Lupus and TB lymphadenitis

Permalink

<https://escholarship.org/uc/item/29b4h4gv>

Journal

Proceedings of the UCLA Department of Medicine, 14(1)

Authors

Lau, Christine

Taylor, David

Publication Date

2011-02-03

CLINICAL VIGNETTE

A Case of Septic Arthritis with *Cryptococcus Neoformans* in a Patient with Lupus and TB lymphadenitis

Christine Lau, M.D., and David Taylor, M.D.

Introduction

Septic arthritis is a medical emergency with an increasing frequency and a high associated mortality¹. Although the classic microbiologic presentation still represents the majority of cases², the increasing population of immunocompromised patients is changing the presentation³. In particular, these patients are at increased risk for fungal causes, which may present with a more indolent course and atypical picture⁴. Here we present a case of cryptococcal arthritis in an asplenic patient with systemic lupus erythematosus.

Case Description

A 44-year-old African-American woman with a medical history significant for systemic lupus erythematosus, idiopathic thrombocytopenia purpura requiring splenectomy, recurrent tuberculous lymphadenitis on therapy, and obsessive-compulsive disorder presented to the emergency department with one month of right upper extremity pain. The pain was worse with movement and was progressive in severity. She had been evaluated by her primary medical doctor with X-rays and an ultrasound of right upper extremity which were unrevealing. Physical exam revealed limited range of motion of right shoulder and elbow secondary to pain; this matched tenderness on palpation over the posterior mid-humerus. She was febrile to 101.3F and laboratory studies were remarkable for mild leukocytosis, anemia, and an elevated erythrocyte sedimentation rate (ESR) of 50 millimeters per hour (mm/hr). Repeat films of the right shoulder showed inferior subluxation of the humeral head with joint effusion. Magnetic resonance imaging (MRI) of shoulder showed a large joint effusion with marked edematous changes of the rotator cuff muscles, consistent with septic arthritis. Arthrocentesis revealed RBC 3900 and WBC 195,000 with a differential

of neutrophils 95%, lymphocytes 2%, and monocytes 3%. Gram stain was negative for bacteria. Orthopedic surgery debrided the shoulder, and cultures from both the aspiration and debridement grew yeast, subsequently identified as *Cryptococcus neoformans*. She was started on high-dose oral fluconazole. Lumbar puncture was performed and showed no evidence of cerebrospinal fluid (CSF) involvement.

Case Discussion

Recent studies show septic arthritis to be increasing in incidence with a frequency as high as 10 per 100,000 patients per year¹. This is thought to be primarily related to increased prevalence of risk factors such as immunocompromised state, diabetes, and prosthetic joints¹. Among culture positive cases, staphylococcus aureus or streptococci continue to represent the majority of cases; these pathogens have been the identified etiology in up to 91% in case series². Regardless of the organism, joint involvement results from direct inoculation from an exogenous source, hematogenous dissemination, or direct extension from an adjacent focus³. In this patient, her history of obsessive-compulsive disorder with dermatillomania with multiple associated skin excoriations led us to suspect that *S. aureus* and Streptococci were the most likely causative organisms of the septic arthritis. She was started on vancomycin and ceftriaxone empirically until culture results returned. Ultimately, when the cultures grew *Cryptococcus neoformans*, she was treated with a course of fluconazole with subsequent improvement.

Fungal arthritis is a rare condition that usually presents in an indolent fashion and is found typically in immunocompromised patients⁴. Diagnosis may be delayed as inflammation is

often less severe and because of the slow progression of disease⁴. Recognition of underlying medical problems that predispose to this condition is key in establishing an appropriate level of suspicion to make the diagnosis. This patient presented with a relative immunocompromised state with her history of SLE. In addition, her history of splenectomy predisposed her to opportunistic infection with encapsulated organisms, including *Streptococcus*, *Neisseria*, *Haemophilus*, and *Klebsiella*. *Cryptococcus neoformans*, an encapsulated yeast is also an opportunistic pathogen that can target splenectomized patients⁵. Splenectomy has been identified as a risk factor for *Cryptococcal* infection in 3% of cases⁶. Primary infection usually results in pulmonary disease⁴. In immunocompromised hosts and immunocompetent patients with CNS involvement, recommended induction therapy is with amphotericin-B and 5-fluorocytosine and fluconazole during consolidation⁷. In immunocompetent hosts, isolated pulmonary *Cryptococcus* may be carefully observed⁷. Symptomatic patients with non-CNS-isolated *Cryptococcus* can be treated with fluconazole for 3-6 months⁷. More severe disease requires amphotericin⁷. There is no established standard of care for the treatment of *cryptococcal* septic arthritis.

This case is more unusual because the patient also had concomitant TB lymphadenitis that was actively being treated. Literature search produced another case of *Cryptococcus* arthritis and TB lymphadenitis in the same patient⁸. *Cryptococcus* inhibits production of TNF alpha, IL-12, IL-18, and interferon gamma but stimulates production of IL-10. TNF alpha is known to play a role in the inhibition of pulmonary TB. This patient may have had inhibition of TNF alpha from the *Cryptococcal* infection, which reactivated the TB lymphadenitis. This patient is currently undergoing immunologic work-up to further evaluate the underlying etiology of her multiple infections.

Although rare, the diagnosis of *cryptococcal* septic arthritis should not have been surprising

based on key elements from her history: signs and symptoms suggestive of an infected joint; an immunocompromised state; a particular predisposition to encapsulated organisms; and the possible biological link between active *cryptococcal* infection and active tuberculosis. This case is an excellent example of the importance of tying apparent disparate elements from the history and physical exam into a unifying diagnosis.

REFERENCES

1. **Mathews CJ, Weston VC, Jones A, Field M, Coakley G.** Bacterial septic arthritis in adults. *Lancet*. 2010 Mar 6;375(9717):846-55. Review. PubMed PMID: 20206778.
2. **Gupta MN, Sturrock RD, Field M.** A prospective 2-year study of 75 patients with adult-onset septic arthritis. *Rheumatology (Oxford)*. 2001 Jan;40(1):24-30. PubMed PMID: 11157138.
3. **Pioro MH, Mandell BF.** Septic arthritis. *Rheum Dis Clin North Am*. 1997 May;23(2):239-58. Review. PubMed PMID: 9156391.
4. **Kohli R, Hadley S.** Fungal arthritis and osteomyelitis. *Infect Dis Clin North Am*. 2005 Dec;19(4):831-51. Review. PubMed PMID: 16297735.
5. **Qazzafi Z, Thiruchunapalli D, Birkenhead D, Bell D, Sandoe JA.** Invasive *Cryptococcus neoformans* infection in an asplenic patient. *J Infect*. 2007 Dec;55(6):566-8. Epub 2007 Oct 1. PubMed PMID: 17905439.
6. **Pappas PG, Perfect JR, Cloud GA, Larsen RA, Pankey GA, Lancaster DJ, Henderson H, Kauffman CA, Haas DW, Saccente M, Hamill RJ, Holloway MS, Warren RM, Dismukes WE.** Cryptococcosis in human immunodeficiency virus-negative patients in the era of effective azole therapy. *Clin Infect Dis*. 2001 Sep 1;33(5):690-9. Epub 2001 Jul 26. PubMed PMID: 11477526.
7. **Saag MS, Graybill RJ, Larsen RA, Pappas PG, Perfect JR, Powderly WG, Sobel JD, Dismukes WE.** Practice guidelines for the management of cryptococcal disease. Infectious Diseases Society of America. *Clin Infect Dis*. 2000 Apr;30(4):710-8. Epub 2000 Apr 20. PubMed PMID: 10770733.
8. **Al-Tawfiq JA, Ghandour J.** *Cryptococcus neoformans* abscess and osteomyelitis in an immunocompetent patient with tuberculous lymphadenitis. *Infection*. 2007 Oct;35(5):377-82. Epub 2007 Sep 20. PubMed PMID: 17885733.

Submitted on February 3, 2011