UC Merced

UC Merced Previously Published Works

Title

An isotopic study of dietary diversity in formative period Ancachi/Quillagua, Atacama Desert, northern Chile.

Permalink

<https://escholarship.org/uc/item/29d0q074>

Journal

American journal of physical anthropology, 170(4)

ISSN

0002-9483

Authors

Pinder, Danielle M
Gallardo, Francisco
Cabello, Gloria
et al.

Publication Date

2019-12-01

DOI

10.1002/ajpa.23922

Peer reviewed

**AN ISOTOPIC STUDY OF DIETARY DIVERSITY IN
FORMATIVE PERIOD ANCACHI/QUILLAGUA, ATACAMA
DESERT, NORTHERN CHILE**

Journal:	<i>American Journal of Physical Anthropology</i>
Manuscript ID	Draft
Wiley - Manuscript type:	Brief Communication
Date Submitted by the Author:	n/a
Complete List of Authors:	Pinder, Danielle; University of Miami, Department of Anthropology Gallardo, Fransisco ; Centro Interdisciplinario de Estudios Interculturales e Indígenas, Pontificia Universidad Católica de Chile Cabello, Gloria; Centro Interdisciplinario de Estudios Interculturales e Indígenas, Pontificia Universidad Católica de Chile Torres-Rouff, Christina; University of California, Merced, Anthropology; Universidad Católica del Norte, Instituto de Investigaciones Arqueológicas y Museo Pestle, William; University of Miami, Department of Anthropology
Key Words:	Stable Isotope Analysis, Formative Period, Atacama Desert, Southern Andes, Paleodiet
Subfield: Please select 2 subfields. Select the main subject first.:	Bioarchaeology [including forensics], Theory

SCHOLARONE™
Manuscripts

AN ISOTOPIC STUDY OF DIETARY DIVERSITY IN FORMATIVE PERIOD
ANCACHI/QUILLAGUA, ATACAMA DESERT, NORTHERN CHILE

Danielle M. Pinder¹, Francisco Gallardo², Gloria Cabello², Christina Torres-Rouff³, William J.
Pestle¹

¹ Department of Anthropology, University of Miami

² Centro Interdisciplinario de Estudios Interculturales e Indígenas, Pontificia Universidad
Católica de Chile

³ Department of Anthropology and Heritage Studies, University of California, Merced

Abstract

Objectives

To characterize the paleodiet of individuals from Formative Period (1500 B.C. - A.D. 400) Atacama Desert sites of Ancachi and Quillagua as a means of understanding the dietary and cultural impacts of regional systems of exchange.

Materials and Methods

Thirty-one bone samples recovered from the cemetery of Ancachi (02QU175) and in/around the nearby town of Quillagua, were subject of carbon and nitrogen stable isotope analysis and multi-source mixture modeling (FRUITS) of paleodiet. These individuals were compared with nearly 200 contemporary individuals from throughout the region to identify differences in dietary behaviors.

Results

80.6% (25/31) of the samples yielded sufficient well-preserved collagen and were included in the multi-source mixture model. The FRUITS model, which compared individuals with a robust database of available foods from the region, identified a wide diversity of diets in the Ancachi/Quillagua area (including both coastal and interior individuals), and, most notably, twenty-one individuals who consumed an average of $11.9 \pm 1.8\%$ terrestrial animals, $21.2 \pm 2.4\%$ legumes, and $20.0 \pm 4.1\%$ marine fauna, a balanced pattern of protein consumption distinct from both the coastal and inland individuals in our larger regional sample.

Conclusions

The combination of stable isotope analysis and multi-source mixture modeling permitted the characterization of dietary behavior of twenty-five individuals from nodal sites in the Atacama Desert, thus enhancing our understanding of the economic and social relationships that bound together Formative Period sites, populations, and individuals in this hyperarid region.

Introduction

The Atacama Desert is located in northern Chile and southern Peru, between ca. 18° and 30° South. The Atacama is the world’s driest desert, with less than 1 mm/yr falling in the study area (Houston 2006), and the region’s aridity seems to have been a persistent feature for millennia (Moreno 2009). A recent paleoclimatic study suggests that aridity equal to, if not exceeding, those seen in the present prevailed throughout the Holocene, with the exception of a somewhat wetter period between 1000-2000 years ago (Maldonado, et al. 2016). In order to survive these arid conditions, ancient peoples strategically chose settlement locations on the Pacific coast, at oases, in deep *quebradas* on the western slope of the Andes, and along the Loa River, the region’s only persistent river course (Castro, et al. 2016). To supply material needs and wants, Atacameños also developed systems of long-distance exchange, through which both essential and luxury goods moved (Pimentel 2013). Quillagua and Ancachi, the two localities at the heart of this research (Figure 1), are located at a nexus of these trade routes between coastal and inland/highland populations, and a variety of sources support the notion that the Ancachi/Quillagua area has functioned as a frontier zone between groups living to their east, north, and west for much of its inhabited history, from the Formative Period until the 18th century A.D. (Agüero et al. 1999, 2001, 2006; Paz Soldán 1878).

<Figure 1>

While borders or frontiers such as that seen at Ancachi/Quillagua are often conceived of as limiting the interaction of peoples from surrounding areas, a wealth of social science literature sees them instead as zones of cultural transition between the societies that lie on either side (Barth 1998, 2000; Newman, 2006; van Dommelen, 1997, 1998; White, 1991). Viewed as such, these frontier zones are judged to facilitate, rather than restrict, cross-cultural pollination. Indeed, people residing in these zones can become engines of cultural innovation and change. As discussed below, the “contact hypothesis”, a sociological precept that describes how perceptions and behavior change when groups of people with diverse backgrounds come into close contact, provides a powerful lens for understanding the conditions under which cultural change might occur. In this regard, the inhabitants of Ancachi and Quillagua serve as ideal indicators from which to develop a better understanding of the movement and exchange of resources within the Atacama Desert during the Formative, and the lived consequences of this exchange for their behaviors and lifeways.

The way in which individuals obtain the necessary nutrients to survive is one of the most fascinating aspects of human behavior (Schwarcz and Shoeninger 1991), and one of the most culturally enmeshed. Indeed, few aspects of human behavior are, simultaneously, as culturally bound and situationally responsive as diet, with the consequence that reconstruction of ancient dietary practices can offer insights into myriad aspects of ancient life (see, for example, the papers in Twiss 2007 [ed.]). The most well-established technique for reconstructing individual-level ancient human diet is stable isotope analysis, which provides high-fidelity data on long-

term (decadal) individual consumption patterns (Lee-Thorp 2008). Stable isotope analysis has been part of the archaeologist’s toolkit since the last quarter of the 20th century, and it has been proven to provide researchers with an accurate method for estimating the dietary composition of past people, assuming a series of pre-conditions are met.

When combined with multi-source mixture modeling (Fernandes et al. 2014), one can use stable isotope analysis to develop quantitative and probabilistic estimates of the lived behavior (diet) of past societies. In the present case, these techniques allow us to characterize individual-level diet at Ancachi/Quillagua and identify those persons who appear to have been behaving (consuming) in ways not seen elsewhere among their contemporaries or in the broader region. Through this process of estimating ancient diet, we contend that we can, in effect, measure the effects of interaction patterns in these individuals, and thus gauge the presence and effects of ancient frontiers in the archaeological record.

Materials and Methods

Cortical bone samples (~1 g) were obtained from thirty-one individuals from Formative Period burials in the tumulus cemetery of Ancachi and from pit/shaft tombs near the modern town of Quillagua (Agüero et al. 1999, Agüero and Uribe 2015, Gallardo et al. 1993, Latcham 1933). For these purposes, we consider the two sites, which sit approximately 10 km apart, jointly. The present work adds nineteen individuals to a previously published sample of twelve individuals from Ancachi/Quillagua (Pestle et al. 2019). Ultimately, we consider both the internal variability and structure of the paleodiet of these individuals and then position them against a larger regional sample of nearly 200 previously analyzed individuals from the Formative Period (Pestle et al. 2015a, 2015b; Pestle 2017).

The extraction of collagen and hydroxyapatite from human bone samples was performed at the Archaeological Stable Isotope Laboratory at the University of Miami. Each sample was individually ground by hand using a ceramic mortar and pestle. Samples were then separated into size fractions using geological screens. The collagen extraction protocol used was established by Longin (1971) and modified by Pestle (2010). For each bone sample, 0.5 grams of the 0.5-1.0 mm fraction was weighed and placed in 50 ml centrifuge tubes. The samples were demineralized in 30 mL of 0.2 M HCL on a spinning rotator for 24 hours. Samples were then rinsed to neutral through a process of centrifugation, decanting, and the addition of 30 mL of distilled water. Humic removal was accomplished by adding 30 mL of 0.0625 M NaOH to each sample for 20 hours. After time elapsed, the samples were again rinsed to neutral. The remaining collagen was then gelatinized in 10^{-3} M HCL at 90°C and filtered using single-use Millipore Steriflip® vacuum filters, condensed, frozen, and freeze dried. Start and end weights were recorded and used to calculate collagen yield (wt%) for each sample.

Hydroxyapatite extraction followed a protocol established in Lee-Thorp (1989) and Kruger (1991) and modified by Pestle (2010). Approximately 0.1 gram of the 0.125–0.25 mm. fraction was placed in a 50 mL centrifuge tube. After weighing, each sample underwent a 24 h oxidation of organics using 30 mL of 50% bleach. The bleach treatment was then repeated for an additional period for a total of 48 h of treatment. Samples then were rinsed to neutral. The final step in the protocol involved the samples undergoing an acid treatment for the removal of labile carbonates. This was accomplished by the addition of 30 mL of 0.1 M acetic acid to each centrifuge tube for a total of four hours with a 5 min vacuum treatment at the two-hour mark. After the acid treatment, each sample was rinsed again to neutral before being placed in a 50°C

oven overnight. Start and end weights were recorded for all hydroxyapatite samples and used to calculate the weight percent hydroxyapatite yield.

Collagen and hydroxyapatite isotopic analysis was performed in the Marine Geology and Geophysics Stable Isotope Laboratory and the Rosenstiel School of Marine and Atmospheric Science at the University of Miami. Collagen samples were packed into tin capsules and analyzed using PDZ Europa ANCA-GSL elemental analyzer interfaced to a PDZ Europa 20-20 isotope ratio mass spectrometer (IRMS). This analytical process yields information on elemental carbon and nitrogen composition as well as the stable isotopes of carbon and nitrogen ($\delta^{13}\text{C}_{\text{co}}$ and $\delta^{15}\text{N}_{\text{co}}$). Hydroxyapatite samples were analyzed using Kiel-IV Carbonate Device coupled to a Thermo Finnigan DeltaPlus IRMA, providing the $\delta^{13}\text{C}_{\text{ap}}$ values. Collagen results were calibrated using acetanilide and glycine. An in-house carbonate standard calibrated to NBS-19 was used for hydroxyapatite. Standards were analyzed in every sample set at the beginning and end of the run, as well as in-between the analyzed samples to ensure accuracy and instrumental stability.

With isotopic data in hand, the FRUITS (Food Reconstruction Using Isotopic Transferred Signals) model of Fernandes and colleagues (2014) was used to quantify individual dietary composition. This multi-source mixture modeling technique is one of several developed with the hope of better bounding estimates of food source contribution. Indeed, recent southern Andean attempts (Andrade, et al. 2015; Pestle, et al. 2019; Pestle, et al. 2016; Pestle, et al. 2017) at modeling have tended to use this, or similar, Bayesian approaches, which accommodate underdetermined systems (those with more than $n+1$ sources), and also allow for the incorporation of priors (Fernandes, et al. 2014; Moore and Semmens 2008; Parnell, et al. 2010). These approaches “offer a powerful means to interpret data because they can incorporate prior

information, integrate across sources of uncertainty and explicitly compare the strength of support for competing models or parameter values,” (Moore and Semmens 2008:471).

In order to generate consumer (human) data for the model, we first determined the consumer-foodstuff offset (and error) for $\delta^{13}\text{C}_{\text{co}}$ using the method of Pestle and colleagues (2015). The offset in $\delta^{13}\text{C}_{\text{ap}}$ was stipulated as $10.1 \pm 0.4\text{‰}$ (Fernandes et al. 2012). Finally, for $\delta^{15}\text{N}_{\text{co}}$, we employed a trophic fractionation value of $3.6 \pm 1.2\text{‰}$, as recommended by several experimental studies of omnivorous animals (Ambrose 2000; DeNiro et al. 1981; Hare et al. 1991; Howland et al. 2003; Sponheimer et al. 2003; Warinner and Tuross 2009).

Foodweb isotope values comprised the edible portions of eighty-nine archaeological and modern Atacameño plants and animals. The decision to restrict the foodweb sample to only those generated in the course of our work in the region was due to the isotopic dissimilarity between those samples and other previously published values. Any modern data included in this reference sample had $\delta^{13}\text{C}$ values corrected by $+1.5\text{‰}$ to account for recent fossil fuel burning (Keeling et al. 1979). Macronutrient composition of each food group was determined by reference to the USDA National Nutrient Database for Standard Reference (Agriculture 2013). Elemental composition (particularly %C) of each foodstuff/macronutrient group was based on formulae provided in Morrison et al. (2000). Digestibility was determined following Hopkins (1981). All nitrogen in bone collagen was stipulated as coming from dietary protein, the carbon in hydroxyapatite was stipulated as reflecting all dietary carbon, and the carbon composition of bone collagen was set as reflecting a 3:1 ratio of dietary protein to energy (Fernandes et al. 2012). Carbon isotope offsets between measured bulk food isotope values and the isotopic values of a foodstuff's fats (bulk-6‰) and carbohydrates (bulk+0.5‰) were based on data from Tieszen (1981). The carbon isotope signature of a measured bulk foodstuff's protein was

determined using a mass-balance equation, such that a proportional/weighted average of the $\delta^{13}\text{C}$ of protein and energy (fats and carbohydrates) would equal the measured $\delta^{13}\text{C}$ bulk value (corrected for the concentration of carbon in each macronutrient and foodstuff-appropriate macronutrient concentration).

Final food group isotope, macronutrient, and elemental concentration values used in the FRUITS simulations are presented in Table 1. We divided the available foodstuffs into five groups (C_3 plants, C_4 plants, legumes, terrestrial animals, and marine animals). Consumption of protein was limited to less than 45% of protein as energy (using the FRUITS a priori data option), reflecting the upper limit of possible human protein intake (World Health Organization 2007). All FRUITS simulations were performed using 10,000 iterations, as recommended by its developers.

Results

Sample preservation quality was determined using both chemical (collagen yield) and elemental (carbon and nitrogen yield, atomic C/N ratio) data. Only well-preserved (collagen yield >0.5 wt%, carbon yield >4.5 wt%, nitrogen yield >0.9 wt%, atomic C/N ratio between 2.9–3.6) samples were included in FRUITS calculations. Based on the arid environmental conditions of the region, samples that met these requirements also were assumed to have acceptable hydroxyapatite preservation (the lack of free water making the prospects of dissolution and recrystallization unlikely).

<Table 1>

As seen in Table 2, 80.6% (25/31) of the samples yielded sufficient well-preserved collagen to be considered as reflecting biogenic isotope signatures. The average collagen yield for those twenty-five samples was 13.7 ± 5.3 wt%, carbon yield averaged 40.8 ± 1.9 wt%, the average nitrogen yield was 14.2 ± 1.0 wt%, making the average atomic C/N ratio 3.4 ± 0.1 . Elemental values were not recorded for one sample (I-102), but due to its high collagen yield and unremarkable (non-outlying) isotope values, we nonetheless included it in later analysis. Similarly, sample I-101 had a slightly elevated atomic C:N ratio (3.7), but we retained the sample because its carbon and nitrogen yields were within acceptable ranges and it was not an isotopic outlier.

<Table 2>

Turning to isotopic results (Table 2), $\delta^{13}\text{C}_{\text{co}}$ for the twenty-five well preserved samples averaged $-15.5 \pm 0.9\text{‰}$ (range -17.1 – -13.4‰) and $\delta^{13}\text{C}_{\text{ap}}$ averaged $-11.4 \pm 0.8\text{‰}$ (range -13.0 – -9.8‰), which, when combined, yielded an average $\Delta^{13}\text{C}_{\text{ap-co}}$ of $4.2 \pm 0.6\text{‰}$ (range 3.2 – 5.2‰). $\delta^{15}\text{N}_{\text{co}}$ averaged $17.3 \pm 3.0\text{‰}$, and possessed an immense range of 10.9 – 25.8‰ . To begin with, then, there is notable isotopic variation within the sample, particularly in $\delta^{15}\text{N}_{\text{co}}$, suggesting similar diversity in patterns/sources of protein consumption.

Based on the results of FRUITS modeling (Table 3), C_3 plants were the largest average dietary contributor, providing an average of $37.2 \pm 4.4\%$ of calories, with a range of 28.5 – 46.2% . In comparison, C_4/CAM plants made up an average of only $8.7 \pm 1.4\%$ of diet, ranging between 6.2 – 13.0% . Turning to dietary protein, terrestrial animals contributed between 5.2 – 19.8% (average $12.0 \pm 2.7\%$), marine animals $19.5 \pm 7.0\%$ (range 7.7 – 41.3%), and legumes $22.5 \pm 6.0\%$

(range 10.8–39.2%). It is the variability of these protein sources, each of which show at least threefold variability, and in particular that of marine faunal sources (which shows a greater-than fivefold difference between minimum and maximum modeled contribution), that is most noteworthy.

<Table 3>

Discussion

Comparing these results to Formative Period individuals from coastal and interior sites in the Atacama (Figure 2), (at least) two distinct dietary regimes are evident among the individuals from Ancachi/Quillagua. On the one hand, certain individuals (e.g. J-86 from Quillagua [41.3±11.7% marine protein] and L-139 from Ancachi [10.7±7.7% marine]) possess modeled diets consistent with either a coastal (high marine protein intake) or interior (heavy terrestrial protein reliance) origin. This provides direct testament to the presence in Ancachi/Quillagua of people of presumably non-local origin, or at least people who ate in ways consistent with other far-removed locales. It is our contention that these individuals came to be buried in Ancachi/Quillagua as a consequence of their direct personal movement/involvement in systems of regional exchange. Like other Formative Period individuals who we have recovered from alongside trade routes through the Atacama (Knudson et al. 2012, Pimenetel et al. 2017, Torres-Rouff et al. 2012), these individuals were agents (travelers, traders) embedded in these region-wide systems of exchange.

<Figure 2>

1
2
3 A larger number of the Ancachi individuals (84%, 21/25), however, would appear to have
4 consumed a mixed diet (particularly in terms of protein composition/balance) unlike that seen in
5 almost any other site in the region (Villa Chuqicamata in the modern city of Calama being the
6 only other exception). These individuals consumed an average of $11.9 \pm 1.8\%$ terrestrial animals,
7 $21.2 \pm 2.4\%$ legumes, and $20.0 \pm 4.1\%$ marine fauna, a balanced pattern of protein consumption
8 that is notably distinct from both the coastal and inland individuals in our larger regional sample.
9 Since Ancachi/Quillagua were located at a centralized location between coastal and interior
10 populations, in a border/frontier space, we contend that the unique dietary pattern seen in this
11 population evince the types of interactions and cultural innovation that only (or most often) occur
12 in such border spaces. Individuals of diverse origin and food culture were coming together at
13 Ancachi/Quillagua, interacting via meaningful economic and social exchanges, and (one of) the
14 products of this interaction was new dietary practices, new cultural forms.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 In Sociology, the Contact Hypothesis has been employed for over sixty years as means of
32 explaining how attitudes and behaviors can change as a consequence of long-term meaningful
33 (equal-status, non-transactional) interaction between groups of distinct interest/origin. While this
34 notion was originally developed in the context of racial prejudice reduction in the United States
35 of the mid-20th century (Allport 1954), decades of further study has validated its prediction that
36 under circumstances of prolonged equal-status co-existence and interaction, common experience
37 will shape and sway the opinions and worldview of even the most entrenched actors (Kende et
38 al. 2018; Mirwaldt 2010, Pettigrew and Troop 2006, Pettigrew et al. 2011). When these
39 interactions extend beyond the transactional, to the kinds of more profound egalitarian
40 relationships that emerge when diverse individuals interact and coexist for long periods of times
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 while engaged in mutually-beneficial activities, individuals begin to exhibit real social and
4
5 cultural exchange, and new and hybrid behaviors emerge.
6
7

8 That a form of eating unlike anything else seen in the Formative Period Atacama would
9
10 emerge in a space like Ancachi and Quillagua would suggest that beyond simply functioning as
11
12 economic nodes, these sites acted as locations of social exchange and interculturalization. People
13
14 were not only passing through these spaces in pursuit of material needs, but the positioning of
15
16 these sites as a nexus or node in the Formative Period’s regional exchange network would appear
17
18 to have facilitated the transculturation of individuals involved, and the emergence of new ways
19
20 of eating, if not new ways of living. These processes would likely have been similar/the same
21
22 that produced, for instance, new regional stylistic conventions and symbolic vocabularies during
23
24 the Formative (Castro et al. 2016).
25
26
27
28
29
30
31
32

33 *Conclusions*
34
35

36 Dietary patterns are fundamental to an individual’s identity, and their reconstruction can
37
38 serve as a powerful tool for understanding past cultural and ethnic differences and identity
39
40 formation. In this present work, stable isotope analysis and multi-source mixture modeling
41
42 permitted the characterization of dietary behavior of twenty-five individuals buried in a region
43
44 thought to be central to a vast regional exchange system. Our results suggest that the diets of
45
46 these Ancachi/Quillagua individuals were strongly influenced by the kind of exchange systems
47
48 that surrounded them in life. One possible explanation for the novel dietary patterns we observed
49
50 is that these systems of economic exchange had fostered meaningful social relationships among
51
52 different cultural groups. Through these interactions, some of the individuals studied here
53
54
55
56
57
58
59
60

1
2
3 adopted new cultural lifestyles and behaviors, consuming resources from both coastal and
4
5 interior cultural patterns, in an entirely new way of living otherwise not seen in the surrounding
6
7 region. Further analysis of additional human remains recovered from Ancachi/Quillagua
8
9 cemetery should be performed to validate and develop this notion, but based on the data
10
11 presented here, it is clear that something novel, and indeed phenomenal, was taking place in this
12
13 portion of Atacama Desert region more than 2,000 years ago.
14
15
16
17
18
19

20 **Acknowledgements**

21
22
23 Financial support for this research came from CONICYT, FONDAP 15110006 and FONDECYT
24
25 160045. The authors wish to thank Dr. Peter Swart and the staff of the Marine Geology and
26
27 Geophysics Stable Isotope Laboratory at the University of Miami's Rosenstiel School of Marine
28
29 and Atmospheric Science.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References Cited

Andrade, P., Fernandes, R., Codjambassis, K., Urrea, J., Olguín, L., Rebolledo, S., Lira, F., Aravena, C., Berríos, M. (2015). Subsistence continuity linked to consumption of marine protein in the Formative Period in the interfluvic coast of northern Chile: re-assessing contacts with agropastoral groups from highlands, *Radiocarbon* 57, 679-688.

Agüero, C., Ayala, P., Uribe, M., Carrasco, C., Dases, B. (2006). El periodo formativo desde Quillagua, in: Lechman, H. (Ed.), *Esferas de interacción prehistóricas y fronteras nacionales modernas, IEP, Lima, Peru*, pp. 73-118.

Agüero, C., Uribe, M. (2015). Tombs and tumuli on the coast and Pampa of Tarapaca: explaining the Formative Period in northern Chile (South-Central Andes). In: Eeckhout, P., Owens, L.S. (Eds.), *Funerary Practices and Models in the Ancient Andes. The Return of the Living Dead*. Cambridge University Press, Cambridge, UK, pp. 152–183.

Agüero, C., Uribe, M., Ayala, P., Cases, B., Carrasco, C. (2001). Ceremonialismo del periodo formativo en Quillagua, Norte Grande De Chile, *Boletín de la Sociedad Chilena de Arqueología* 32, 24-34.

Agüero, C., Uribe, M., Ayala, P., Dases, B. (1999). Una aproximación arqueológica a la etnicidad: el rol de los textiles en la construcción de la identidad cultural en los cementerios de Quillagua (Norte de Chile). *Gaceta Arqueológica Andina* 25, 167–198.

Agriculture, U.S.D.o. (2013). National Nutrient Database for Standard Reference, Release 27.

Allport, G.W. (1954). *The Nature of Prejudice*. Addison-Wesley, Cambridge MA.

Ambrose, S.H. (2000). Controlled diet and climate experiments on nitrogen isotope ratios of rats, in: Ambrose, S.H., Katzenberg, M.A. (Eds.), *Biogeochemical approaches to paleodietary analysis*, Kluwer Academic/Plenum Publishers, New York, pp. 243–259.

Barth, F. (1998). *Ethnic Groups and Boundaries: The Social Organization of Culture Difference*. Waveland press, Prospect Heights, Illinois.

Barth, F. (2000). Boundaries and connections. In Cohen, A.P. (Ed.) *Signifying Identities*. Routledge, London, pp. 17-36.

Castro, V., Berenguer, J., Gallardo, F., Llagostera, A., Salazar, D. (2016). Vertiente Occidental Circumpuñena. Desde Las Sociedades Posarcaicas Hasta Las Preincas (Ca. 1.500 años A.C. A 1.470 años D.C.), in: Falabella, F., Uribe, M., Sanhueza, L., Aldunate, C., Hidalgo, J. (Eds.), *Prehistoria En Chile: Desde Sus Primeros Habitantes Hasta Los Incas*, Editorial Universitaria, Santiago de Chile, pp. 239–283.

DeNiro, M.J., Epstein, S. (1981). Influence of diet on the distribution of nitrogen isotopes in animals, *Geochimica et Chosmochimica Acta*, 45, 341-351.

Fernandes, R., Nadeau, M-J., & Grootes, P.M. (2012). Macronutrient-based model for dietary carbon routing in bone collagen and bioapatite. *Archaeological and Anthropological Sciences*, 4, 291-301.

Fernandes, R., Millard, A.R., Barabec, M., Nadeau, M-J., & Grootes, R. (2014). Food reconstruction using isotopic transferred signals (FRUITS): A Bayesian model for diet reconstruction, *PLOS One*, 9(2), e87436.

Gallardo, F., Cornejo, L., Sánchez, R., Casses, B., Román, A., Deza, A. (1993). Una aproximación a la cronología y el asentamiento en el oasis de Quillagua (río Loa, II región). *Actas del XII Congreso Nacional de Arqueología*. vol. 4. Boletín del Museo Regional de la Araucanía, Temuco, pp. 41–60.

Hare, P.E., Fogel, M.L., Stafford, T.W., Mitchell, A.D., Hoering, T.C. (1991). The isotopic composition of carbon and nitrogen in individual amino acids isolated from modern and fossil proteins, *Journal of Archaeological Science*, 18, 277–292.

Hopkins, D.T. (1981). Effects of variations in protein digestibility, in: Bodwell, C.E., Adkins, J.S., Hopkins, D.T. (Eds.), *Protein quality in humans: assessment and in vitro estimation*, AVI Publishing, Westport, CT, pp. 178-181.

Houston, J. (2006). Variability of precipitation in the Atacama Desert: It causes an hydrological impact. *International Journal of Climatology*, 26, 2181-2198.

Howland, M.R., L.T. Corr, S. M. M. Young, V. Jones, S. Jim, N. J. van der Merwe, A.D. Mitchell, and R.P. Evershed. (2003). Expression of the dietary isotope signal in the compound-specific C^{13} values of pig bone lipids and amino acids. *International Journal of Osteoarcheology*, 13:54-65.

Kende, J., Phalet, K., Van den Noortgate, W., Kara, A., Fischer, R. (2018). Equality revisited: A cultural meta-analysis of intergroup contact and prejudice, social psychological and personality. *Science* 9, 887-895.

Keeling, C.D., W.G. Mook, and P.P Tans. (1979). Recent trends in the C^{13}/C^{12} ratio of atmospheric carbon dioxide. *Nature*, 277 (5692): 121-123.

Knudson, K.J., Pestle, W.J., Torres-Rouff, C., Pimentel, G. (2012). Assessing the life history of an Andean traveller through biogeochemistry: stable and radiogenic isotope analyses of archaeological human remains from Northern Chile, *International Journal of Osteoarchaeology* 22, 435-451.

1
2
3 Krueger, H.W. (1991). Exchange of carbon with biological apatite. *Journal of Archaeological*
4
5 *Science*, 18, 355-362.
6
7

8
9
10 Latcham, R. (1933). Notas preliminares de un viaje arqueológico a Quillagua. vol. XXXVII.
11
12 *Revista Chilena de Historia Natural*, pp. 130–138.
13
14

15
16
17 Lee-Thorp, J.A. (1989). Stable carbon isotopes in deep time: the diets of fossil fauna and
18
19 hominids. Unpublished Ph.D. Dissertation, Department of Archaeology, University of Cape
20
21 Town.
22
23

24
25
26 Lee-Thorp, J.A. (2008). On Isotopes and Old Bones. *Archaeometry*, 50(6), 925-950.
27
28

29
30
31 Longin, R. (1971). New method of collagen extraction for radiocarbon dating. *Nature*, 230, 241-
32
33 242.
34
35

36
37
38 Maldonado, A., de Porra, M.E., Zamora, A., Rivadeneira, M., Abarzúa, A.M. (2016). El
39
40 Escenario Geográfico Y Paleoambiental De Chile, in: Falabella, F., Uribe, M., Sanhueza, L.,
41
42 Aldunate, C., Hidalgo, J. (Eds.), Prehistoria En Chile: Desde Sus Primeros Habitantes Hasta Los
43
44 Incas, Editorial Universitaria, Santiago de Chile, pp. 23-70.
45
46

47
48
49 Mirwaldt, K. (2010). Contact, conflict and geography: what factors shape cross-border citizen
50
51 relations? *Political Geography*, 29 (8), 434–443.
52
53

1
2
3 Moore, J.W., Semmens, B.X., 2008. Incorporating Uncertainty and Prior Information into Stable
4 Isotope Mixing Models, *Ecology Letters* 11, 470-480.
5
6
7
8
9

10 Moreno, A., Santoro, C.M., Latorre, C., 2009. Climate Change and Human Occupation in the
11 Northernmost Chilean Altiplano over the Last Ca. 11,500 cal. a Bp, *Journal of Quaternary*
12 *Science* 24, 373-382.
13
14
15
16
17
18

19 Morrison, D.J., Dodson, B., Slater, C., Preston, T. (2000). ^{13}C Natural abundance in the British
20 diet: implications for ^{13}C breathe tests, *Rapid Communications in Mass Spectrometry*, 14, 1312-
21 1324.
22
23
24
25
26
27

28 Newman, D. (2006). The lines that continue to separate us: borders in our 'borderless' world.
29 *Progress in Human Geography*, 30 (2), 143–161.
30
31
32
33
34

35 Parnell, A.C., Inger, R., Bearhop, S., Jackson, A.L. (2010). Source Partitioning Using Stable
36 Isotopes: Coping with Too Much Variation, *PloS one* 5, e9672.
37
38
39
40
41

42 Paz Soldán, M. (1878). Verdaderos límites entre el Perú y Bolivia. Imprenta Liberal, Lima, Peru.
43
44
45
46

47 Pestle, W.J. (2010). Diet and society in prehistoric Puerto Rico, an isotopic approach.
48 Unpublished PhD. Dissertation, Department of Anthropology, University of Illinois at Chicago.
49
50
51
52
53
54
55
56
57
58
59
60

Pestle, W.J., Hubbe, M., Smith, E.K., Stevenson, J.M. (2015). A linear model for predicting $\Delta^{13}\text{C}_{\text{protein}}$, *American Journal of Physical Anthropology*, 157, 694-703.

Pestle, W.J., Torres-Rouff, C., Hubbe, M. (2016). Modeling diet in times of change: the case of Quitor, San Pedro De Atacama, Chile, *Journal of Archaeological Science: Reports* 7, 82-93.

Pestle, W.J., Torres-Rouff, C., Hubbe, M., Smith, E.K. (2017). Eating in or dining out: modeling diverse dietary strategies in middle period San Pedro De Atacama, Chile, *Archaeological and Anthropological Sciences* 9, 1363-1377.

Pestle, W.J. (2017). Living, eating, and dying in the Formative Period Atacama, in: Gallardo, F. (Ed.), *Monumentos funerarios de la costa del desierto De Atacama: Contribuciones sobre el intercambio de bienes e información entre cazadores-recolectores marinos (Norte De Chile)*, Centro Interdisciplinario de Estudios Interculturales e Indígenas, Sociedad Chilena de Arqueología, Santiago de Chile, pp. 209-222.

Pestle, W.J., Torres-Rouff, C., Gallardo, F., Ballester, B., Clarot, A. (2015). Mobility and exchange among marine hunter-gatherer and agropastoralist communities in the Formative Period Atacama Desert, *Current Anthropology* 56, 121-133.

Pestle, W.J., Torres-Rouff, C., Hubbe, M., Santana, F., Pimentel, G., Gallardo, F., Knudson, K.J., (2015). Explorando la diversidad dietética en la prehistoria del desierto de Atacama: Un acercamiento a los patrones regionales, *Chungará (Arica)* 47, 201-209.

Pestle, W.J., Torres-Rouff, C., Gallardo Ibanez, F., Andrea Cabello, G., and E.K. Smith. (2019). The interior frontier: exchange and interculturalism in the formative period (1000 B.C.-A.D. 400) of Quillagua, Antofagasta Region, northern Chile. *Quaternary International*.
<https://doi.org/10.1016/j.quaint.2019.03.014>

Pettigrew, T.F., Tropp, L.R. (2006). A meta-analytic test of intergroup contact theory. *Journal of Personality and Social Psychology*. 90 (5), 751–783.

Pettigrew, T.F., Tropp, L.R., Wagner, U., Christ, O. (2011). Recent advances in intergroup contact theory. *International Journal of Intercultural Relations*, 35, 271–280.

Pimentel, G. (2013). Redes Viales Prehispánicas En El Desierto De Atacama: Movilidad, Viajeros E Intercambio. Unpublished Ph.D. Dissertation. Instituto de Investigaciones arqueológicas y Museo Gustavo Le Paige S.J. , Universidad Católica del Norte y Universidad de Tarapacá, San Pedro de Atacama, Chile.

Pimentel, G., Ugarte, F., Blanco, J.F., Torres-Rouff, C., Pestle, W.J. (2017). Calate: de lugar desnudo a laboratorio arqueológico de la movilidad y el tráfico intercultural prehispánico en el desierto de Atacama (Ca. 7000 Ap-550 Ap), Estudios atacameños, 21-56.

Schwarcz, H. P., Schoeninger, M.J. (1991). Stable isotope analysis in human nutritional ecology. *Yearbook of Physical Anthropology*, 34:283-321.

Sponheimer, M., Robinson, T., Ayliffe, L., Roeder, B., Hammer, J., Passey, B., West, A., Cerling, T., Dearing, D., Ehleringer, J. (2003). Nitrogen isotopes in mammalian herbivores: Hair $\delta^{15}\text{N}$ values from a controlled feeding study, *International Journal of Osteoarchaeology*, 13, 80-87.

Tieszen, L.L. (1981). Natural variation in the carbon isotope values of plants: implications for archaeology, ecology, and paleoecology, *Journal of Archaeological Science* 18, 227-248.

Torres-Rouff, C., Pimentel, G., Ugarte, M., (2012). ¿Quiénes Viajaban? Investigando la muerte de viajeros Prehispánicos en el desierto de Atacama (Ca. 800 Ac–1536 Dc), *Estudios atacameños* 43, 167-186.

Twiss, K.C. (2007). The archaeology of food and identity, Center for Archaeological Investigations, Southern Illinois University Carbondale, Carbondale.

van Dommelen, P. (1997). Colonial constructs: colonialism and archaeology in the Mediterranean. *World Archaeology*, 28 (3), 305–323.

1
2
3 van Dommelen, P. (1998). On colonial grounds: a comparative study of colonialism and rural
4 settlements in first millennium BC west central Sardinia. Doctoral Thesis. Faculty of
5
6 Archaeology, Leiden University, Leiden.
7
8
9

10
11
12 Warinner, C., Tuross, N. (2009). Alkaline cooking and stable isotope tissue-diet spacing in
13 swine: archaeological implications, *Journal of Archaeological Science*, 36, 1690-1697.
14
15
16
17

18
19 White, R. (1991). The middle ground: Indians, empires and republics in the Great Lakes region.
20 Cambridge University Press, Cambridge, pp. 1650–1815.
21
22
23
24

25
26 World Health Organization. 2007. Protein and Amino Acid Requirements in Human Nutrition
27 vol. 935. WHO Press, Geneva.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Figure and Table Captions
4

5
6
7 Figure 1: Map of study region, with sites mentioned in text noted.
8

9
10
11 Figure 2: FRUITS modeled consumption of marine animals and C3 plants for Ancachi/Quillagua
12 individuals and comparative interior and coastal populations.
13

14
15
16 Table 1: Macronutrient, isotopic, and elemental data for food groups used in FRUITS multi-
17 source mixture model.
18

19
20
21 Table 2: Chemical, elemental, and isotopic data for all Ancachi/Quillagua individuals.
22

23
24
25 Table 3: Results of FRUITS multi-source mixture modeling for Ancachi/Quillagua individuals.
26

Figure 1: Map of study region, with sites mentioned in text noted.

Figure 2: FRUITS modeled consumption of marine animals and C3 plants for Ancachi/Quillagua individuals and comparative interior and coastal populations.

241x174mm (300 x 300 DPI)

		Macronutrient concentration (%)				%C				Tissue $\delta^{13}\text{C}$ (‰)					Tissue $\delta^{15}\text{N}$ (‰)	
Food grouping	Group <i>n</i>	Protein	Fat	Carbohydrates	Energy	Protein	Fat	Carbohydrates	Energy	Bulk	Protein	Fat	Carbohydrates	Energy	Bulk	Protein
Terrestrial animals	24	83±12	16±12	1±3	17±12	43±12.7	12±12.1	0±4.2	13±12.7	-16.5±3.8	-14.8±3.8	-22.5±3.8	-16±3.8	-22.3±3.8	9.8±2.4	9.8±2.4
Marine animals	31	74±15	19±16	7±9	26±15	39±15.3	15±16.6	3±9.6	18±15.3	-14.6±2.9	-12.4±2.9	-20.6±2.9	-14.1±2.9	-19.5±2.9	20.9±3.4	20.9±3.4
C ₃ plants	17	10±5	5±4	84±7	89±5	5±6.9*	4±6.7	37±8.5	41±6.9	-23.7±2.0	-22.3±2.0	-29.7±2.0	-23.2±2.0	-23.8±2.0	8.6±5.4	8.6±5.4
C ₄ /CAM plants	13	10±5	5±4	84±7	89±5	5±6.9*	4±6.7	37±8.5	41±6.9	-11.2±1.6	-9.8±1.6	-17.2±1.6	-10.7±1.6	-11.3±1.6	12±5.6	12±5.6
Legumes	4	24±2	2±1	71±3	72±2	13±5.6*	1±5.3	31±5.8	33±5.6	-23.5±1.7	-24.2±1.7	-29.5±1.7	-23.0±1.7	-23.2±1.7	0.7±3.0	0.7±3.0

* assumes 87.4% digestibility of plant protein as compared to animal protein

Sample	Site	Burial	Collagen yield	%C	%N	Atomic C:N	d ¹³ C _{co} (‰)	d ¹⁵ N _{co} (‰)	d ¹³ C _{ap} (‰)	Δ ¹³ C _{ap-co} (‰)
I-99*	Ancachi	UR-3	18.4%	39.6	13.6	3.4	-14.8	20.4	-11.6	3.2
I-100*	Ancachi	UR-3	23.9%	43.5	14.9	3.4	-15.1	17.9	-11.4	3.7
I-101*	Ancachi	UR-1	14.6%	38.6	12.0	3.7	-15.4	18.5	-10.5	4.9
I-102*	Ancachi	UR-6	24.5%	-	-	-	-15.3	16.9	-11.5	3.8
I-103*	Ancachi	UR-2	14.6%	39.6	13.4	3.4	-15.4	16.7	-11.8	3.6
I-105*	Ancachi	UR-4	20.2%	39.8	14.0	3.3	-14.6	16.9	-11.2	3.4
J-84*	Quillagua	Museo, caja 3-3	16.0%	44.4	15.6	3.3	-15.1	17.7	-11.6	3.5
J-86*	Quillagua	Torre 203, Qui.1, museo, exhibido	7.0%	41.9	13.6	3.6	-13.4	25.8	-10	3.4
J-92*	Quillagua	Torre 203, Qui 02	16.6%	44.9	16.3	3.2	-16.5	10.9	-11.5	5.0
J-93*	Quillagua	Qui. Res. 2013-1	3.2%	38.9	13.4	3.4	-17.1	11.2	-12.1	5.0
L-133	Ancachi	12140, 343, 399	7.8%	37.9	13.2	3.3	-16.4	16.8	-11.9	4.5
L-134	Ancachi	12146, 344, 400	5.7%	38.1	13.6	3.3	-15.1	17.5	-10.6	4.5
L-135	Ancachi	12065, 340, 401	13.4%	41.0	14.6	3.3	-14.7	17.7	-10.7	4.0
L-136	Ancachi	12141, 342, 403	11.1%	39.9	14.6	3.2	-17.1	16.0	-13.0	4.1
L-137	Ancachi	12137, 345, 402	16.8%	41.1	15.0	3.2	-14.8	20.2	-10.9	3.9
L-138	Ancachi	12139, 347, 398	15.1%	41.6	14.7	3.3	-16.0	15.3	-11.4	4.6
L-139	Ancachi	12148, 1899, 403	12.6%	40.7	14.8	3.2	-14.6	12.3	-10.1	4.5
L-140	Ancachi	12152, 348, 397	12.9%	41.0	14.7	3.2	-15.8	16.7	-12.1	3.8
L-141	Ancachi	ANC16	16.7%	40.8	13.6	3.5	-16.0	19.0	-11.7	4.3
L-144	Ancachi	ANC-30-I1	15.1%	42.2	15.0	3.3	-15.5	16.4	-11.7	3.9
L-148	Ancachi	ANC12	7.4%	39.9	14.1	3.3	-16.0	16.5	-11.2	4.8
L-149	Ancachi	ANC7	8.1%	42.2	14.1	3.5	-16.9	16.7	-12.5	4.4
L-150	Ancachi	ANC6-I1	12.9%	41.4	14.3	3.4	-16.4	19.4	-12.9	3.5
L-151	Ancachi	ANC13	10.8%	37.3	12.7	3.4	-15.4	19.1	-11.2	4.1
L-152	Ancachi	ANC-2-I1	16.4%	42.2	15.1	3.3	-15.0	19.6	-9.8	5.2

* individuals were previously published in Pestle et.al 2019

Sample	Site	Burial	Terrestrial animals	sd	C ₃ plants	sd	C ₄ /CAM plants	sd	Legumes	sd	Marine animals	sd
I-99*	Ancachi	UR-3	8.4%	7.6%	39.6%	18.2%	7.7%	6.4%	16.7%	13.7%	27.6%	11.3%
I-100*	Ancachi	UR-3	12.3%	10.3%	38.0%	18.5%	8.7%	7.1%	20.1%	15.2%	20.9%	10.4%
I-101*	Ancachi	UR-1	11.0%	9.5%	36.2%	18.4%	9.6%	7.6%	20.0%	15.0%	23.2%	11.1%
I-102*	Ancachi	UR-6	13.0%	10.9%	36.8%	19.0%	8.8%	7.2%	22.9%	16.1%	18.5%	9.5%
I-103*	Ancachi	UR-2	13.3%	11.0%	37.9%	19.2%	8.3%	6.8%	23.1%	16.7%	17.3%	9.3%
I-105*	Ancachi	UR-4	14.0%	11.5%	34.4%	18.3%	9.8%	7.7%	22.5%	15.3%	19.3%	10.0%
J-84*	Quillagua	Museo, caja 3-3	12.7%	10.4%	38.5%	18.6%	8.4%	6.9%	20.4%	15.9%	20.0%	10.0%
J-86*	Quillagua	Torre 203, Qui.1, museo, exhibido	5.2%	4.6%	32.8%	14.7%	10.0%	8.0%	10.8%	9.4%	41.3%	11.7%
J-92*	Quillagua	Torre 203, Qui 02	13.4%	10.5%	30.8%	19.1%	9.6%	7.4%	38.5%	18.1%	7.7%	6.3%
J-93*	Quillagua	Qui. Res. 2013-1	13.9%	11.1%	30.9%	19.8%	8.3%	6.6%	39.2%	19.1%	7.7%	5.9%
L-133	Ancachi	12140, 343, 399	12.6%	10.7%	40.3%	19.1%	7.5%	6.3%	23.3%	17.2%	16.4%	9.1%
L-134	Ancachi	12146, 344, 400	13.4%	11.1%	33.7%	18.4%	9.8%	8.0%	22.0%	15.6%	21.0%	10.6%
L-135	Ancachi	12065, 340, 401	12.3%	9.9%	36.5%	18.4%	10.3%	8.1%	20.2%	15.1%	20.8%	10.1%
L-136	Ancachi	12141, 342, 403	11.7%	10.5%	45.3%	21.3%	6.2%	5.2%	23.1%	17.5%	13.7%	8.5%
L-137	Ancachi	12137, 345, 402	9.1%	8.6%	36.9%	17.5%	8.7%	7.1%	17.8%	13.8%	27.5%	11.0%
L-138	Ancachi	12139, 347, 398	14.5%	11.4%	37.1%	20.0%	8.8%	6.9%	25.3%	17.4%	14.4%	8.8%
L-139	Ancachi	12148, 1899, 403	19.8%	12.8%	28.5%	17.5%	13.0%	8.9%	28.1%	15.7%	10.7%	7.7%
L-140	Ancachi	12152, 348, 397	12.8%	10.8%	39.9%	20.3%	7.5%	6.3%	22.7%	16.7%	17.0%	9.3%
L-141	Ancachi	ANC16	9.4%	8.5%	42.1%	18.6%	7.5%	6.4%	18.5%	14.6%	22.5%	10.7%
L-144	Ancachi	ANC-30-I1	13.4%	11.1%	37.9%	19.7%	8.2%	6.5%	24.0%	16.9%	16.5%	9.0%
L-148	Ancachi	ANC12	13.3%	10.8%	37.0%	19.1%	9.0%	7.3%	23.4%	16.4%	17.4%	9.5%
L-149	Ancachi	ANC7	12.1%	11.1%	43.3%	20.1%	7.0%	5.8%	22.1%	16.3%	15.6%	8.9%
L-150	Ancachi	ANC6-I1	8.8%	8.6%	46.2%	18.8%	6.4%	5.6%	17.5%	14.9%	21.1%	10.1%
L-151	Ancachi	ANC13	10.0%	9.0%	38.4%	18.6%	8.5%	7.0%	19.6%	14.6%	23.6%	10.5%
L-152	Ancachi	ANC-2-I1	10.6%	9.5%	32.3%	17.4%	10.3%	8.1%	19.9%	14.7%	26.9%	11.1%

* individuals were previously published in Pestle et.al 2019