

UC San Diego

UC San Diego Previously Published Works

Title

Better executive function is independently associated with full HIV suppression during combination therapy.

Permalink

<https://escholarship.org/uc/item/2bc336q3>

Journal

AIDS, 33(15)

ISSN

0269-9370

Authors

Anderson, Albert M
Pérez-Santiago, Josué
Zheng, Ziduo
et al.

Publication Date

2019-12-01

DOI

10.1097/qad.0000000000002348

Peer reviewed



Published in final edited form as:

AIDS. 2019 December 01; 33(15): 2309–2316. doi:10.1097/QAD.0000000000002348.

Better executive function is independently associated with full HIV suppression during combination therapy

Albert M. ANDERSON¹, Josué PÉREZ-SANTIAGO^{2,3}, Ziduo ZHENG⁴, Eugene HUANG⁴, Donald FRANKLIN², Jennifer IUDICELLO², David J. MOORE², Ronald J. ELLIS⁵, Robert K. HEATON², Scott L. LETENDRE^{2,6}

¹Department of Medicine, Emory University School of Medicine

²Department of Psychiatry, University of California at San Diego School of Medicine

³University of Puerto Rico Comprehensive Cancer Center

⁴Department of Biostatistics and Bioinformatics, Emory University Rollins School of Public Health

⁵Department of Neurology, University of California at San Diego School of Medicine

⁶Department of Medicine, University of California at San Diego School of Medicine

Abstract

Objective—HIV-associated neurocognitive impairment continues to be prevalent and clinically relevant. We examined the relationship between neurocognition and full plasma HIV RNA suppression among study participants over a 15 year period at a large research program.

Design/Methods—We analyzed the combined prospective studies of the HIV Neurobehavioral Research Program (HNRP) at the University of California at San Diego. Participants were eligible for analysis if on three drug cART with comprehensive neuropsychological testing results. Participants who reported recent non-adherence were excluded. The primary outcome was plasma HIV RNA of ≤ 50 copies/ml. Generalized estimating equation was used to assess for associations with full virologic suppression taking into account longitudinal visits.

Results—There were 1943 participants at baseline, of whom 69.4% had plasma HIV RNA ≤ 50 copies/ml. Participants with full suppression were slightly older, less likely to abuse cocaine, and had significantly better executive function. Multivariate analysis with incorporation of longitudinal visits (total= 5555) confirmed current cocaine abuse to be strongly associated with lack of virologic suppression (odds ratio= 0.45, 95% confidence interval= 0.31–0.63). In contrast, increasing age, increasing years of HIV infection, and increasing executive function (odds ratio= 1.18 for T score change of 10, 95% confidence interval= 1.07–1.30) were associated with full virologic suppression. Lack of virologic suppression at baseline was associated with a significant subsequent decline in executive function.

Conclusions—In a 15 year research cohort of almost 2000 HIV-infected individuals on cART, better executive function was associated with full virologic suppression, possibly as a result rather than a cause.

Keywords

HIV; AIDS; Sustained virologic response; Neurocognitive disorders

Introduction

Infection with the human immunodeficiency virus (HIV) has become a treatable condition in the combination antiretroviral therapy (cART) era, and virologic suppression is commonly achieved with current cART regimens.^[1] The World Health Organization (WHO) has proposed a goal that at least 90% of HIV-infected (HIV+) individuals on cART attain full virologic suppression.^[2] Some countries appear to have reached this goal. Sweden, for example, recently reported that 94.7% of HIV+ individuals on cART had achieved full virologic suppression.^[3] Many other countries, however, have not yet reached such a high level of virologic suppression. For example, the WHO recently reported that across Latin America and the Caribbean, the median rate of virologic suppression among HIV+ individuals on cART was 66%.^[2] Overall, the United States also has not yet achieved 90% virologic suppression. For example, a very large United States study published in 2017 of 339, 515 HIV+ individuals receiving care showed that 72.6% had durable virologic suppression.^[4] Lack of virologic suppression predisposes infected individuals to increased rates of opportunistic infections, end organ disease, and death.^[5] Additionally, HIV transmission is more likely when plasma HIV RNA levels are not suppressed.^[6] Therefore, the causes for and consequences of lack of full virologic suppression continue to merit study. Lack of virologic suppression during cART has been associated with multiple factors. These include younger age, differences in race, and substance abuse.^[7–10] To achieve the goals of optimal health and zero transmission, a fuller understanding of the factors that may affect virologic suppression during cART is needed.

HIV-associated neurocognitive disorder (HAND) specifically and neurocognitive impairment more broadly continue to be common in the cART era, particularly if asymptomatic impairment is included. The prevalence of HAND during the cART era varies widely depending on study, but has been found to be as high as 30–40%.^[11, 12] HAND in the cART era is associated with decreased quality of life and increased risk of mortality.^[13, 14] Additionally, there may be a link between worse neurocognitive performance and lack of virologic suppression. Previous studies have reported a significant association between diminished neuropsychological (NP) performance (including within individual domains such as executive function) and cART non-adherence.^[15–17] Additionally, individuals with decreased NP performance may believe that they are taking cART correctly, when in actuality they may not be. Alternatively, HIV+ individuals without full virologic suppression may have worse NP performance due to the detrimental effects of uncontrolled viral replication. To better understand the relationship between NP performance and virologic suppression, we undertook an analysis of a large research cohort in which all participants received comprehensive neuropsychological testing. We hypothesized that worse global NP performance would be associated with lack of full virologic suppression despite adherence by self-report. Our secondary hypothesis was that worse performance on individual NP domains such as executive function would be associated with lack of full virologic suppression despite adherence by self-report.

Methods

Data were pooled from HIV Neurobehavioral Research Program (HNRP) observational research studies at the University of California at San Diego from a fifteen year period in the cART era (years 2000–2015). The analysis included data from the Central Nervous System Antiretroviral Therapy Effects Research (CHARTER) study (which includes multiple geographic sites across the United States), the California NeuroAIDS Tissue Consortium study, and other federally and locally funded studies. For the purposes of this analysis, only participants who were prescribed three drug cART, reported recent adherence, and had full neuropsychological assessment were considered. The neuropsychological assessment by definition included at least two tests from each domain per HNRP protocol: 1) Executive Function (Wisconsin Card Sorting Test-64 item, Trail Making Test Part B, or Stroop Color Word interference); 2) Working Memory/Attention (WMS-III Spatial Span and Paced Auditory Serial Addition Task); 3) Speed of Information Processing (WAIS III Digit Symbol, WAIS III Symbol Search, Trail Making Test Part A, or Stroop Color Naming); 4) Learning (Hopkins Verbal Learning Test and Brief Visuospatial Memory Test); 5) Memory (Hopkins Verbal Learning Test-delayed recall and Brief Visuospatial Memory Test-delayed recall); 6) Verbal Fluency (Controlled Oral Word Association Test and Categories- Animals and Actions); and 7) Fine motor skill (Grooved Pegboard Test dominant and non-dominant hands). Raw scores on these tests were converted to mean T scores based on published demographic norms with mean= 50 and standard deviation= 10.^[18–22] These were then used to compute neurocognitive domain T-scores and a global mean T-score. Commercially available HIV RNA tests were used for all of the studies. Due to the different timeframes of the studies, these tests included lower limits of detection of 40, 48, and 50 copies/milliliter (ml). Therefore for the purposes of this analysis, the primary end point was defined as plasma HIV RNA \geq 50 copies/ml. Adherence was assessed using the AIDS Clinical Trials Group (ACTG) interview form. Only participants who reported recent cART adherence were eligible for inclusion in this analysis. Therefore, participants who answered “yes” to missing medication during any of the following intervals were excluded: “in the last four days”, “within the last week”, or “one to two weeks ago”. Instrumental activities of daily living (IADLs) were measured with a modified Lawton-Brody questionnaire.^[23] Depression symptoms were assessed with version two of the Beck Depression Inventory.^[24] Current substance abuse in the last 30 days was defined by diagnostic and statistical manual of mental disorders version four (DSM-IV) criteria.^[25] Longitudinal assessments were also included when available for participants with multiple study visits. Score adjustment for practice effects was also made for longitudinal visits by using median practice effect data from previous work.^[26]

Statistical Approach

Continuous variables were assessed for normality with the Shapiro-Wilk test. When conditions for normality were not met, the Wilcoxon rank sum test was used for differences in continuous variables. Otherwise, student’s t-test was used for differences in continuous variables. Chi-squared test was used to examine differences in categorical variables. Cohen’s d was reported for effect size. Risk factors for lack of full virologic suppression were then identified using logistic regression model via generalized estimating equation (GEE) to

accommodate multiple visits. Observations were weighted by inverse number of visits, and covariates included demographic and medical characteristics as well as neuropsychological measures. The following covariates were screened during the modeling process: age, sex, race (white versus non-white), years of HIV infection, current and nadir CD4+ as well as current CD4+/CD8+ ratio, months on current cART regimen, hepatitis C virus seropositivity, medical history (hypertension, stroke, and diabetes), current tobacco smoker or current abuse of alcohol, marijuana, cocaine, methamphetamine, or opiates, Beck Depression Inventory scale, and activities of daily living dependency. Individual neuropsychological domain T scores as well global T score were examined as continuous variables. Additionally, an exploratory multivariate analysis of domains found to be significant as continuous variables in the primary multivariate analysis was performed by transforming those domains into impaired/not impaired (T score ≤ 40 versus >40) categorical variables. Covariates that were significant at the 0.05 level in univariate analysis were considered in the multivariate analysis and the final model was obtained using backward stepwise selection. Covariate values were updated for each visit analyzed, as opposed to limiting the analysis to covariate baseline values. While the primary analytic outcome was plasma HIV RNA ≤ 50 copies/ml, a secondary GEE modeling analysis was performed using the outcome of plasma HIV RNA ≤ 200 copies/ml, which has been recently suggested to define virologic suppression in HIV primary care guidelines.^[27] For domains that were found to have a statistically significant association, we then examined change in performance over time based on virologic suppression and vice versa. This was performed with Chi squared test.

Results

The larger cohort from which the current analysis was derived encompassed 4605 individuals with a total of 20,511 study visits. Of the 4595 participants from this larger cohort with full demographic data, the mean age at the first visit was 40.8 years and 83.8% were male. In terms of race/ethnicity, 49.6% were white, 29.3% were black, and 17.3% were hispanic. The vast majority of participant exclusions from this larger cohort resulted from: 1. not taking combination antiretroviral therapy, 2. comprehensive neuropsychological testing results not available, 3. report of recent non-adherence or 4. missing data in laboratory parameters such as CD4+ count or HIV RNA level.

The final analysis cohort of 1943 participants were older, more likely to be female, and more likely to be black compared to participants who were excluded (all $p < 0.05$). This final analysis cohort had a total of 5555 study visits between the years 2000 and 2015. Mean number of months on the current cART regimen at the baseline visit was 20.6. Demographic and disease characteristics based on baseline plasma HIV RNA status ($n=1943$) are summarized in Table 1, while demographic and disease characteristics of the cohort with all visits combined ($n=5555$) are summarized in Table 2. 30.6% of participants had lack of full virologic suppression (plasma HIV RNA >50 copies/ml) at baseline. These participants were slightly younger, more likely to be of non-white race, and had lower current and nadir CD4+ T-cell counts (404 versus 546 cells/microliter and 150 versus 175 cells/microliter, respectively) compared to the group with full virologic suppression (plasma HIV RNA ≤ 50 copies/ml). Participants in the two groups had similar frequencies of hepatitis C seropositive status, diabetes, and stroke history, but participants without full suppression were on their

current cART regimen for less time and were less likely to be hypertensive. Participants without full suppression were less likely to be current tobacco smokers (11.3% versus 15.9%, $p=0.008$) but more likely to be current abusers of cocaine (12.0% versus 4.9%, $p<0.001$). There were no differences between the two groups in terms of current abuse of other drugs. Neuropsychological testing showed that executive function was significantly lower in the group without full suppression (mean T score 44.6 versus 46.4, $p=0.001$), but that verbal performance was significantly higher in this group (mean T score 49.2 versus 47.9, $p=0.018$). However, the effect sizes for these differences in cognitive performance were relatively small. Specifically, Cohen's d effect size was 0.13 for difference in executive function and 0.18 for difference in verbal. Central nervous system penetration effectiveness (CPE) score was not significantly different between the two groups (mean 7.5 in non-suppressed group versus mean 7.6 in group with HIV RNA <50 copies/ml, $p=0.8$). 83.6% of participants were on a regimen of protease inhibitor (PI) plus nucleoside reverse transcriptase inhibitor (NRTI) or a regimen of non-nucleoside reverse transcriptase inhibitor (NNRTI) plus NRTI at baseline. 1078 participants had multiple visits (mean number of visits= 4.4 and mean number of days between visits= 266). Table 2 shows that when all visits were taken into account ($n=5555$) as opposed to baseline only, most differences were similar between groups based on virologic suppression.

Significant variables that were associated with HIV RNA <50 copies/ml in univariate analyses by logistic regression along with GEE are shown in the top half of Table 3. The bottom half of Table 3 shows the multivariate GEE analysis. Specifically, this multivariate analysis indicated older age (odds ratio [OR] = 1.14, 95% confidence interval [95% CI] = 1.07–1.21 for five year increase), years since HIV diagnosis (OR=1.09, 95% CI=1.01–1.18 for five year increase), and executive function (OR=1.18, 95% CI 1.07–1.30 for T score increase of 10) were significantly associated with full virologic suppression. CD4+/CD8+ ratio was also significantly associated with full virologic suppression (OR=1.19, 95% CI 1.15–1.24 for every 0.1 increase). Conversely, current cocaine abuse was negatively associated with full virologic suppression (OR= 0.45, 95% CI 0.31–0.63). In the secondary analysis using <200 copies/ml as the cutoff for full virologic suppression, the multivariate GEE analysis revealed similar results. Specifically, current cocaine abuse was again negatively associated with full virologic suppression (OR= 0.42, 95% CI 0.29–0.61). Similarly, executive function (OR= 1.35, 95% CI= 1.19–1.53 for T score increase of 10), older age (OR = 1.19, 95% CI= 1.10–1.28 for five year increase), and CD4+/CD8+ ratio (OR = 1.27, 95% CI= 1.19–1.35 for 0.1 increase) were again positively associated with full virologic suppression. The exploratory multivariate analysis using executive function as a categorical predictor (impaired versus not impaired as defined by T score <40 versus T score >40) yielded a similar outcome. Specifically, executive function T score >40 was significantly associated with full virologic suppression (OR= 1.57, 95% CI= 1.25–1.97, $p<0.001$).

To further evaluate direction of association, we then examined change in executive function based on virologic suppression (HIV RNA <50 copies/ml) and vice versa. Data on 908 participants who had a second study visit at least 30 days after the baseline visit were available for analysis. 71.7% had stable executive function and 8.7% had a significant increase in executive function where the domain T score increased by at least ten, which is

equivalent to one standard deviation. A small proportion (19.6%) had a significant decrease in executive function of at least ten at the second visit. Of those 281 participants who were not suppressed at baseline, 71 (25.3%) had a decrease in executive function T score of at least ten at the second visit. Meanwhile, of those 627 who were suppressed at baseline and had a second visit, 107 (17.1%) had a decrease in executive function T score of at least ten, which was a significantly smaller proportion ($p=0.004$). Among participants with plasma HIV RNA ≤ 50 copies/ml at their first visit who then went on to have a second visit, 140 of 573 (24.4%) had executive impairment (defined by T score ≥ 40) at the first visit. Of this 140, 17 individuals (12.1%) had loss of virologic suppression at the second visit. Of the 433 without executive impairment, 55 (12.7%) had loss of virologic suppression at the second visit ($p=0.86$ for difference in proportions between the two groups).

Discussion

Given that there is no vaccine available for HIV, the most effective means to prevent the spread of the virus is virologic suppression with cART. Virologic suppression through cART is also the most effective means to optimize the health of people living with HIV (PLHIV).^[5] However, certain countries have not yet reached the WHO stated goal of at least 90% virologic suppression among HIV+ on cART.^[2] More research is needed to better understand why HIV-infected individuals do not achieve full virologic suppression as well as the clinical impact of incomplete virologic suppression. In this analysis of multiple prospective studies, we investigated the relationship between NP performance and full HIV suppression among a large cohort of PLHIV who reported recent adherence to their cART regimen. We found that better executive function was significantly associated with full virologic suppression in multivariate analyses. By linking NP performance with an important laboratory outcome, this extends the findings of other studies showing that better cART adherence is associated with better NP performance (particularly in the domain of executive function).^[16, 17] The association between full virologic suppression and better executive function was significant for both the 50 copies/ml and 200 copies/ml suppression cutoffs. We excluded participants who reported recent non-adherence in this analysis. Despite this, a protective association between full virologic suppression and better executive function was found.

Based on our longitudinal analysis, it is more likely that worse executive function is a result of lack of virologic suppression rather than a cause of it. Specifically, we found in a subgroup analysis that lack of virologic suppression at baseline was associated with a higher frequency of significant executive performance decrease over time. Individuals with uncontrolled HIV may be more likely to develop impairment, as evidenced by the fact that HIV-associated dementia was much more common before the era of cART.^[11] Therefore, the association between worse executive function and lack of virologic suppression could reflect the impact of uncontrolled viral replication on neurocognition. Conversely, there was no significant difference in loss of virologic suppression over time based on executive function. While this subgroup analysis was limited by small sample size, the finding argues against the possibility that executive dysfunction comes first. It should also be noted that the 19.6% of individuals with executive function decline at a second visit may be higher than some clinical settings, particularly in clinics with very high virologic suppression rates.

Like some other studies, we found that younger age, longer time since HIV diagnosis, and current cocaine abuse were significantly associated with lack of full virologic suppression.^[7–10] Other factors such as race, while significant in univariate analyses, were not significant in multivariate analyses. We also found that increasing CD4+/CD8+ ratio was associated with full virologic suppression. While we suspect this may be the result of full virologic suppression (as opposed to the cause of), this is not completely clear. Specifically, higher plasma HIV RNA levels and shorter duration of HIV suppression have both been associated with lower CD4/CD8 ratio in previous studies.^[28, 29]

We acknowledge the limitations of our study. The integrase inhibitor class has become the standard of care for initial cART. However, few study visits (10% of total) in the analysis involved integrase inhibitor- based cART, which is reflective of the timeframe analyzed. Recent studies of the integrase inhibitors have shown that therapy with this drug class may increase the likelihood of undetectable HIV RNA.^[30] With this in mind, if the study were conducted among a population of mostly integrase inhibitor treated individuals, it is possible that the overall rate of full virologic suppression would have been higher and that the association with executive function would not have been found. We also acknowledge that this study did not address functional symptoms. Studies in the cART era have shown that a large proportion of individuals with HAND have asymptomatic neurocognitive impairment.^[12] It is possible that the findings of the study would have been different had we limited the study to individuals with functional symptoms. We acknowledge that certain countries have been able to achieve almost 95% virologic suppression among PLHIV on cART.^[3] Our findings may not be generalizable to settings where essentially complete virologic suppression has been achieved. We also acknowledge that because over 80% of the cohort was male, the study findings may not be generalizable to HIV-infected women.

We did not find an association between global NP performance and virologic suppression (our primary hypothesis). We did find that better executive function was associated with full virologic suppression (a secondary hypothesis), but we did not find this association with other domains. While better working memory was associated with full virologic suppression in the univariate analysis, this association became non-significant in the multivariate analysis. It is somewhat surprising that none of the other domains were significant in the final analysis. However, there is evidence in the cART era that the executive function domain is more affected than other domains during HIV infection.^[31] Therefore, it may be more likely to find significant associations between this domain and other outcomes. Alternatively, it is possible that adherence is more closely linked to executive function than other domains. One study has shown that while impairment in both executive function and memory domains is associated with worse cART adherence, the effect size appears to be larger for the executive function domain.^[32] Assessment of adherence in this study was performed with the ACTG adherence questionnaire. While this questionnaire has been widely used,^[33] self-report may not be as accurate for measuring adherence as other measures such as electronic cap measurements, pharmacy records, and therapeutic drug monitoring.^[34] Going forward, further research on the clinical implications of both neurocognitive impairment and lack of virologic suppression are needed in studies of PLHIV.

REFERENCES

1. Palella FJ Jr., Delaney KM, Moorman AC, Loveless MO, Fuhrer J, Satten GA, et al. Declining morbidity and mortality among patients with advanced human immunodeficiency virus infection. HIV Outpatient Study Investigators. *The New England journal of medicine* 1998; 338(13):853–860. [PubMed: 9516219]
2. UNAIDS. 90–90–90 - An ambitious treatment target to help end the AIDS epidemic. <http://www.unaids.org/en/resources/documents/2017/90-90-90>. 2017.
3. Gisslen M, Svedhem V, Lindborg L, Flamholz L, Norrgren H, Wendahl S, et al. Sweden, the first country to achieve the Joint United Nations Programme on HIV/AIDS (UNAIDS)/World Health Organization (WHO) 90–90–90 continuum of HIV care targets. *HIV medicine* 2017; 18(4):305–307. [PubMed: 27535540]
4. Crepez N, Tang T, Marks G, Hall HI. Changes in viral suppression status among US HIV-infected patients receiving care. *Aids* 2017; 31(17):2421–2425. [PubMed: 29028661]
5. Strategies for Management of Antiretroviral Therapy Study G, El-Sadr WM, Lundgren J, Neaton JD, Gordin F, Abrams D, et al. CD4+ count-guided interruption of antiretroviral treatment. *The New England journal of medicine* 2006; 355(22):2283–2296. [PubMed: 17135583]
6. Cohen MS, Chen YQ, McCauley M, Gamble T, Hosseinipour MC, Kumarasamy N, et al. Antiretroviral Therapy for the Prevention of HIV-1 Transmission. *The New England journal of medicine* 2016; 375(9):830–839. [PubMed: 27424812]
7. Katz IT, Leister E, Kacanek D, Hughes MD, Bardeguet A, Livingston E, et al. Factors associated with lack of viral suppression at delivery among highly active antiretroviral therapy-naïve women with HIV: a cohort study. *Annals of internal medicine* 2015; 162(2):90–99. [PubMed: 25599347]
8. McFall AM, Dowdy DW, Zelaya CE, Murphy K, Wilson TE, Young MA, et al. Understanding the disparity: predictors of virologic failure in women using highly active antiretroviral therapy vary by race and/or ethnicity. *Journal of acquired immune deficiency syndromes* 2013; 64(3):289–298. [PubMed: 23797695]
9. Cescon AM, Cooper C, Chan K, Palmer AK, Klein MB, Machouf N, et al. Factors associated with virological suppression among HIV-positive individuals on highly active antiretroviral therapy in a multi-site Canadian cohort. *HIV medicine* 2011; 12(6):352–360. [PubMed: 21059167]
10. O'Connor J, Smith C, Lampe FC, Johnson MA, Chadwick DR, Nelson M, et al. Durability of viral suppression with first-line antiretroviral therapy in patients with HIV in the UK: an observational cohort study. *Lancet HIV* 2017; 4(7):e295–e302. [PubMed: 28479492]
11. Heaton RK, Clifford DB, Franklin DR Jr., Woods SP, Ake C, Vaida F, et al. HIV-associated neurocognitive disorders persist in the era of potent antiretroviral therapy: CHARTER Study. *Neurology* 2010; 75(23):2087–2096. [PubMed: 21135382]
12. Sacktor N, Skolasky RL, Seaberg E, Munro C, Becker JT, Martin E, et al. Prevalence of HIV-associated neurocognitive disorders in the Multicenter AIDS Cohort Study. *Neurology* 2016; 86(4):334–340. [PubMed: 26718568]
13. Tozzi V, Balestra P, Galgani S, Murri R, Bellagamba R, Narciso P, et al. Neurocognitive performance and quality of life in patients with HIV infection. *AIDS Res Hum Retroviruses* 2003; 19(8):643–652. [PubMed: 13678465]
14. Tozzi V, Balestra P, Serraino D, Bellagamba R, Corpolongo A, Piselli P, et al. Neurocognitive impairment and survival in a cohort of HIV-infected patients treated with HAART. *AIDS Res Hum Retroviruses* 2005; 21(8):706–713. [PubMed: 16131310]
15. Andrade AS, Deutsch R, S AC, Duarte NA, Marcotte TD, Umlauf A, et al. Relationships among neurocognitive status, medication adherence measured by pharmacy refill records, and virologic suppression in HIV-infected persons. *Journal of acquired immune deficiency syndromes* 2013; 62(3):282–292. [PubMed: 23202813]
16. Ettenhofer ML, Foley J, Castellon SA, Hinkin CH. Reciprocal prediction of medication adherence and neurocognition in HIV/AIDS. *Neurology* 2010; 74(15):1217–1222. [PubMed: 20220123]
17. Waldrop-Valverde D, Jones DL, Weiss S, Kumar M, Metsch L. The effects of low literacy and cognitive impairment on medication adherence in HIV-positive injecting drug users. *AIDS care* 2008; 20(10):1202–1210. [PubMed: 19012081]

18. Diehr MC, Cherner M, Wolfson TJ, Miller SW, Grant I, Heaton RK, et al. The 50 and 100-item short forms of the Paced Auditory Serial Addition Task (PASAT): demographically corrected norms and comparisons with the full PASAT in normal and clinical samples. *Journal of clinical and experimental neuropsychology* 2003; 25(4):571–585. [PubMed: 12911108]
19. Heaton R, Taylor M, Manly J. Demographic effects and use of demographically corrected norms with the WAIS-III and WMS-III Clinical Interpretation of the WAIS-III and WMS-III. San Diego, CA: Academic Press, 2002.
20. Heaton RK, Miller SW, Taylor MJ, Grant I. Revised Comprehensive norms for an expanded Halstead- Reitan Battery: Demographically adjusted neuropsychological norms for African American and Caucasian adults scoring program Lutz, FL: Psychological Assessment Resources; 2004.
21. Norman MA, Moore DJ, Taylor M, Franklin D Jr., Cysique L, Ake C, et al. Demographically corrected norms for African Americans and Caucasians on the Hopkins Verbal Learning Test-Revised, Brief Visuospatial Memory Test-Revised, Stroop Color and Word Test, and Wisconsin Card Sorting Test 64-Card Version. *Journal of clinical and experimental neuropsychology* 2011; 33(7):793–804. [PubMed: 21547817]
22. Woods SP, Scott JC, Sires DA, Grant I, Heaton RK, Troster AI, et al. Action (verb) fluency: test-retest reliability, normative standards, and construct validity. *J Int Neuropsychol Soc* 2005; 11(4): 408–415. [PubMed: 16209421]
23. Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist* 1969; 9(3):179–186. [PubMed: 5349366]
24. Beck AT, Steer RA, Brown GK. Beck depression inventory II Manual. San Antonio, TX.: Psychological Corp. (1996)
25. WorldHealthOrganization. Composite International Diagnostic Interview, Version 2.1. 1997.
26. Cysique LA, Franklin D Jr., Abramson I, Ellis RJ, Letendre S, Collier A, et al. Normative data and validation of a regression based summary score for assessing meaningful neuropsychological change. *Journal of clinical and experimental neuropsychology* 2011; 33(5):505–522. [PubMed: 21391011]
27. Aberg JA, Gallant JE, Ghanem KG, Emmanuel P, Zingman BS, Horberg MA, et al. Primary care guidelines for the management of persons infected with HIV: 2013 update by the HIV medicine association of the Infectious Diseases Society of America. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 2014; 58(1):e1–34.
28. Caby F, Guihot A, Lambert-Niclot S, Guiguet M, Boutolleau D, Agher R, et al. Determinants of a Low CD4/CD8 Ratio in HIV-1-Infected Individuals Despite Long-term Viral Suppression. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 2016; 62(10):1297–1303. [PubMed: 26908792]
29. Serrano-Villar S, Gutierrez C, Vallejo A, Hernandez-Novoa B, Diaz L, Abad Fernandez M, et al. The CD4/CD8 ratio in HIV-infected subjects is independently associated with T-cell activation despite long-term viral suppression. *The Journal of infection* 2013; 66(1):57–66. [PubMed: 23046968]
30. Molina JM, Clotet B, van Lunzen J, Lazzarin A, Cavassini M, Henry K, et al. Once-daily dolutegravir is superior to once-daily darunavir/ritonavir in treatment-naïve HIV-1-positive individuals: 96 week results from FLAMINGO. *J Int AIDS Soc* 2014; 17(4 Suppl 3):19490. [PubMed: 25393999]
31. Heaton RK, Franklin DR, Ellis RJ, McCutchan JA, Letendre SL, Leblanc S, et al. HIV-associated neurocognitive disorders before and during the era of combination antiretroviral therapy: differences in rates, nature, and predictors. *Journal of neurovirology* 2011; 17(1):3–16. [PubMed: 21174240]
32. Hinkin CH, Hardy DJ, Mason KI, Castellon SA, Durvasula RS, Lam MN, et al. Medication adherence in HIV-infected adults: effect of patient age, cognitive status, and substance abuse. *Aids* 2004; 18 Suppl1:S19–25.
33. Schackman BR, Ribaldo HJ, Krambrink A, Hughes V, Kuritzkes DR, Gulick RM. Racial differences in virologic failure associated with adherence and quality of life on efavirenz-containing regimens for initial HIV therapy: results of ACTG A5095. *Journal of acquired immune deficiency syndromes* 2007; 46(5):547–554. [PubMed: 18193496]

34. Lam WY, Fresco P. Medication Adherence Measures: An Overview. *Biomed Res Int* 2015; 2015:217047.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1:
Baseline characteristics of participants by plasma HIV RNA suppression.

Variable Reported as mean (standard deviation) or number (%)	<i>Baseline Plasma HIV RNA status</i>		P value
	HIV RNA >50 (n=595)	HIV RNA ≤50 (n=1348)	
Male Gender	495 (83.3%)	1090 (81.0%)	0.22
Race= white	268 (45.1%)	682 (50.7%)	0.02
Years of age	43.6 (8.8)	45.9 (9.7)	<0.001
Years since HIV diagnosis	10.7 (7.1)	11.8 (7.2)	0.002
Current CD4+ (cells/μl)	404 (259)	546 (290)	<0.001
Nadir CD4+ (cells/μl)	150 (143)	175 (161)	0.005
Current CD4+/CD8+ ratio	0.5 (0.3)	0.7 (0.4)	<0.001
Hepatitis C seropositive	142 (24.3%)	340 (25.7%)	0.51
Stroke History	13 (2.2%)	49 (3.7%)	0.09
Hypertension	104 (17.8%)	341 (25.8%)	<0.001
Diabetes	54 (9.2%)	141 (10.7%)	0.34
Global Mean NP score	45.7 (7.0)	45.9 (7.4)	0.53
Verbal Mean T	49.2 (10.5)	47.9 (10.0)	0.018
Executive Function Mean T	44.6 (10.0)	46.4 (10.2)	0.001
Information process Mean T	48.5 (9.4)	47.9 (9.2)	0.25
Learning Mean T	43.1 (8.6)	43.3 (9.3)	0.73
Recall Mean T	44.5 (8.7)	44.3 (9.3)	0.72
Working memory Mean T	45.4 (8.7)	46.1 (9.3)	0.16
Motor Mean T	45.6 (10.2)	45.5 (11.1)	0.89
Beck Depression Inventory	12.4 (10.3)	12.3 (10.4)	0.64
Current smoker	66 (11.3%)	210 (15.9%)	0.008
Current Alcohol abuse	1 (0.2%)	8 (0.6%)	0.23
Current Marijuana abuse	134 (24.3%)	307 (23.3%)	0.65
Current Cocaine abuse	66 (12.0%)	64 (4.9%)	<0.001
Current Methamphetamine abuse	17 (3.1%)	30 (2.3%)	0.31
Current Opiate abuse	35 (6.3%)	73 (5.5%)	0.50
IADL dependent	241 (44.6%)	574 (44.8%)	0.96
Months current regimen	14.2 (22.4)	23.3 (24.5)	<0.001

CPE= central nervous system penetration effectiveness score; IADL= Instrumental Activities of Daily Living.

P values <0.05 are in bold.

Table 2:
Characteristics of participants by plasma HIV RNA suppression: all visits combined

Variable Reported as mean (standard deviation) or number (%)	<i>Plasma HIV RNA status at all visits (n= 5555)</i>		P value
	HIV RNA >50 (n=1348)	HIV RNA ≤ 50 (n=4207)	
Male Gender	1139 (84.6%)	3457 (82.2%)	0.77
Race= white	596 (44.3%)	2241 (53.3%)	<0.001
Years of age	45.4 (9.0)	48.2 (9.9)	<0.001
Years since HIV diagnosis	12.3 (7.0)	13.4 (7.3)	<0.001
Current CD4+ (cells/μl)	437 (273)	581 (297)	<0.001
Nadir CD4+ (cells/μl)	136 (133)	161 (151)	<0.001
Current CD4+/CD8+ ratio	0.5 (0.4)	0.7 (0.4)	<0.001
Hepatitis C seropositive	399 (30.7%)	1114 (28.1%)	0.99
Stroke History	40 (3.1%)	174 (4.4%)	0.1
Hypertension	324 (30.0%)	1347 (34%)	<0.001
Diabetes	174 (13.4%)	616 (15.5%)	0.19
Global Mean NP score	46.3 (7.4)	47.0 (7.6)	0.07
Verbal Mean T	49.0 (10.4)	48.6 (10.0)	0.08
Executive Function Mean T	45.5 (10.6)	48.0 (10.8)	<0.001
Information process Mean T	48.8 (10.0)	49.1 (9.8)	0.94
Learning Mean T	44.1 (9.2)	44.3 (9.7)	0.30
Recall Mean T	45.3 (9.4)	44.9 (9.7)	0.67
Working memory Mean T	46.4 (9.4)	47.5 (9.5)	0.01
Motor Mean T	45.0 (10.5)	45.5 (11.0)	0.65
Beck Depression Inventory	11.1 (10.5)	10.7 (10.0)	0.06
Current smoker	134 (10.3%)	639 (16.1%)	0.002
Current Alcohol abuse	4 (0.3%)	28 (0.7%)	0.02
Current Marijuana abuse	324 (26.3%)	936 (22.9%)	0.59
Current Cocaine abuse	171 (13.9%)	182 (4.4%)	<0.001
Current Methamphetamine abuse	66 (5.4%)	134 (3.3%)	0.06
Current Opiate abuse	106 (8.6%)	239 (5.8%)	0.09
IADL dependent	497 (41.9%)	1577 (41.1%)	0.97
Months current regimen	18.0 (23.2)	28.9 (27.3)	<0.001

CPE= central nervous system penetration effectiveness score; IADL= Instrumental Activities of Daily Living.

P values <0.05 are in bold.

Table 3

(top half) Significant variables associated with HIV RNA ≥ 50 copies/milliliter by weighted univariate analysis using generalized estimating equation (GEE). OR= Odds ratio and CI= confidence interval (bottom half): Multivariate Adjusted (Adj) Odds ratio (OR) for plasma HIV RNA ≥ 50 copies/milliliter using GEE.

Significant univariate GEE results			
Variable	Units	OR (95% CI)	P value
Race= white		1.50 (1.25– 1.80)	<0.001
Years of age	5	1.15 (1.10– 1.21)	<0.001
Years since HIV diagnosis	5	1.12 (1.06– 1.19)	<0.001
CD4+ nadir	50	1.07 (1.03– 1.10)	<0.001
CD4+/CD8+ Ratio	0.1	1.22 (1.17– 1.26)	<0.001
Mean Executive function	10	1.24 (1.13– 1.36)	<0.001
Mean Working Memory	10	1.14 (1.03– 1.26)	0.012
Current cocaine abuse		0.33 (0.24– 0.45)	<0.001
Significant multivariate GEE results			
Variable	Units	Adj OR (95% CI)	P value
Years of age	5	1.14 (1.07– 1.21)	<0.001
Years since HIV diagnosis	5	1.09 (1.01– 1.18)	0.037
CD4/CD8 Ratio	0.1	1.19 (1.15– 1.24)	<0.001
Mean Executive function	10	1.18 (1.07– 1.30)	0.001
Current Cocaine abuse		0.45 (0.31– 0.63)	<0.001