

UC Irvine

UC Irvine Previously Published Works

Title

A gene regulatory program controlling early *Xenopus* mesendoderm formation: Network conservation and motifs

Permalink

<https://escholarship.org/uc/item/2bf5b9nw>

Journal

Seminars in Cell and Developmental Biology, 66

ISSN

1084-9521

Authors

Charney, Rebekah M

Paraiso, Kitt D

Blitz, Ira L

et al.

Publication Date

2017-06-01

DOI

10.1016/j.semcdb.2017.03.003

Peer reviewed



Published in final edited form as:

Semin Cell Dev Biol. 2017 June ; 66: 12–24. doi:10.1016/j.semcdb.2017.03.003.

A gene regulatory program controlling early *Xenopus* mesendoderm formation: network conservation and motifs

Rebekah M. Charney¹, Kitt D. Paraiso¹, Ira L. Blitz, and Ken W. Y. Cho

Department of Developmental and Cell Biology, Ayala School of Biological Sciences, University of California, Irvine, CA 92697, USA

Abstract

Germ layer formation is among the earliest differentiation events in metazoan embryos. In triploblasts, three germ layers are formed, among which the endoderm gives rise to the epithelial lining of the gut tube and associated organs including the liver, pancreas and lungs. In frogs (*Xenopus*), where early germ layer formation has been studied intensively, the process of endoderm specification involves the interplay of dozens of transcription factors. Here, we review the interactions between these factors, summarized in a transcriptional gene regulatory network (GRN). We highlight regulatory connections conserved between *Xenopus*, zebrafish, mouse, and human endodermal lineages. Especially prominent is the conserved role and regulatory targets of the Nodal signaling pathway and the T-box transcription factors, Vegt and Eomes. Additionally, we highlight some network topologies and motifs, and speculate on their possible roles in development.

Keywords

Xenopus; endoderm; gene regulatory network; network motifs; transcription factors; evolutionary conservation

1. Vertebrate mesendoderm formation

During early embryogenesis, cell fate specification proceeds through the combinatorial interactions of several signaling pathways and numerous transcription factors (TFs), which function within a broader chromatin landscape. The integration of these factors ('inputs') leads to a specific transcriptome profile ('outputs') that determines the identity of a particular cell. Critical genomic regions for this integration are cis-regulatory modules (CRMs) – combinations of regulatory elements, such as enhancers, where TFs bind to specific sequence motifs and recruit the necessary co-factors [1]. CRMs are critical for the proper implementation of gene regulatory programs in development, because they modulate

Correspondence to: Ken W. Y. Cho.

¹Authors contributed equally to this work

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

the rate of gene transcription and control when a gene is turned “on” or “off” in both time and space. These complex programs can be organized into gene regulatory networks (GRNs) and visualized through logic maps [2,3]. Elucidating the GRNs will enhance our mechanistic understanding of developmental processes, and will enable comparisons across different organ systems, and across different species. These insights will also enhance our understanding of the causes of developmental defects.

Germ layer specification is one of the earliest developmental events in metazoan organisms, preceding the establishment of the organ and tissue primordia that form the complex adult organism. Cells of the three primary germ layers – the ectoderm, mesoderm, and endoderm – become further specified along distinct lineages. Ectodermal cells form the epidermis and nervous system; mesodermal cells become blood, muscle, kidneys, notochord, and connective tissue; and endodermal cells become the gastrointestinal and respiratory tracts. In amphibian development, the germ layers form along the animal to vegetal axis, which is established during oogenesis. The ectoderm develops from the animal region, while the endoderm arises from the opposite, vegetal pole. The mesoderm forms from the equatorial cells. During early stages, there is overlap between endodermal and mesodermal domains – and therefore it is common to refer to these jointly as the ‘mesendoderm.’ Two major advantages of the *Xenopus* system are the ease of obtaining thousands of synchronously developing embryos from a single clutch of eggs, and the ease of performing both gain- and loss-of-function studies to ascertain gene function. These advantages, combined with the relatively close evolutionary distance between *Xenopus* and other vertebrates, makes *Xenopus* a powerful model for elucidating the mechanisms underlying cell fate specification.

A comprehension of the complex GRN architecture that contributes to the specification of the germ layers *in vivo* is a critical unanswered question in developmental and evolutionary biology. Since germ layer formation, like most biological processes, is controlled by a hierarchy of regulatory steps, examining the earliest inputs in the regulation of germ layer development is important. This is underscored by the realization that germ layer specification is not programmed by molecules acting in a linear fashion, but instead is controlled by a set of transcription factors (TFs) acting in a complex network. In addition, the study of GRN in amphibian germ layer specification will enable powerful comparisons across different developmental systems, and across evolutionary taxa, to identify core GRN structures that are conserved, as well as subnetworks that were modified during evolution. Beyond animal development, numerous human congenital diseases result from abnormal formation of the mesoderm and endoderm [4,5]. Furthermore, recent advances in regenerative medicine and stem cell biology bring the promise of a new era of personalized medicine, aided by highly efficient *in vitro* differentiation techniques [5]. These goals can be significantly aided by a strong understanding of complex *in vivo* cellular differentiation programs, namely “GRN science.”

A little over a decade ago, efforts were made to compile the available molecular data into GRNs describing mesendoderm [6] and Spemann organizer [7] development. Since then, the widespread use of high-throughput technologies (e.g. microarrays, sequencing) provides us with the capacity to significantly broaden the number of network connections and, therefore,

our understanding of the structure of the mesendoderm GRN. For this review, we have utilized recent findings to update the *Xenopus* mesendoderm GRN through the beginning of gastrulation, linking together critical signaling pathways with transcriptional targets. We discuss the network's structure and its motifs, and review areas of conservation across vertebrates.

2. Generation of the mesendoderm gene regulatory network

2.1 Mesendoderm factors

We have assembled a *Xenopus* mesendoderm GRN that reflects data obtained from both *Xenopus laevis*, and the closely related diploid species *Xenopus tropicalis*. Each species greatly contributes to the GRN assembly, as *X. laevis* has been traditionally used in the study of mesendoderm formation, while *X. tropicalis* has been adapted more recently for genomic approaches. The ‘inputs’ in this GRN represent the TFs and signaling molecules (transduced via intracellular TFs) important for mesendoderm formation and early endoderm patterning, many of which have been elucidated [8,9]. Recently, genome-wide approaches have identified additional localized maternal and zygotic transcripts encoding transcription factors [10-16]. Based on a comprehensive catalogue of the *X. tropicalis* TFs [14], 130 TFs are found to be enriched vegetally (in comparison to the animal pole), and we have focused on the ~50 TFs expressed at relatively high abundance in the vegetal tissue (Transcripts Per Million values ≥ 50). This corresponds approximately to the expression level of *siamois1* – a critical and localized mesendoderm TF.

2.2 Criteria for identifying transcriptional targets

In order to build the mesendoderm GRN, we have taken the following rigorous approach to determine direct connections between the above TF ‘inputs’ and their downstream target genes [17]. This approach is similar to that previously taken by both Koide et al. [7] and Loose and Patient [6]. First, we enforced that there should be a strong correlation between perturbation of a regulatory TF and the expression changes of the suspected target genes. Regulation can be measured following gain- and/or loss-of-function experiments (e.g., injection of RNA encoding a TF, or a translation blocking antisense morpholino oligonucleotide, respectively) by analyzing changes in target RNA expression (e.g., RNA-seq, RT-PCR, northern blotting, *in situ* hybridization).

Second, we required that the ‘inputs’ and ‘outputs’ be expressed in a spatiotemporally consistent manner with regulation and the proposed direction (activation or repression). For a proposed activating (‘positive’) connection, the upstream TF and proposed target gene must be expressed in the same or overlapping regions, and during overlapping developmental time. Conversely, a proposed inhibitory (‘negative’) connection should find the target excluded from the spatiotemporal domain of the repressor. However, we note the possibility of finding some exceptions to this rule based on the known biological properties of TFs, and these knowledge-based connections were also included in the network. One example includes a negative autoregulatory feedback loop by *gsc* [18], where a gene modulates its own expression.

Third, we required demonstration of a direct physical interaction between the TF and the regulatory region of the proposed target. As perturbation experiments alone are insufficient to distinguish between direct and indirect effects, we find this criterion essential. This criterion was satisfied experimentally through chromatin immunoprecipitation (ChIP), gel electromobility shift assay (EMSA), DNase footprinting, or reporter gene assays (containing appropriate mutations). Only connections that satisfied all three criteria were defined as direct. However, we note that DNA binding is only suggestive of functional regulation, and that the ‘gold standard’ evidence is to mutate the binding site and examine the effect on gene expression. While the rise in the use of ChIP-seq – ChIP coupled with high-throughput sequencing (HTS) – has produced large datasets of physical connections, the vast majority of these sites have not been subjected to laborious mutagenesis assays. In building the network, we therefore distinguished between functionally validated CRMs and physical interactions lacking this support. Finally, we have also looked to experiments in which protein synthesis is inhibited (e.g. cycloheximide) as a way to determine ‘directness.’ Regulatory connections proposed based on perturbation analyses that were validated in the presence of cycloheximide (CHX), although lacking in physical binding evidence, were identified as ‘putative direct’ targets.

2.3 Building network connections

To assemble the updated network, we have analyzed hundreds of manuscripts published over the approximately 25 year history of the investigation into the molecular basis of *Xenopus* mesendoderm formation. Building upon the networks presented by Koide et al. [7] and Loose and Patient [6] over a decade ago, we have made extensive use of recently published HTS data using *X. tropicalis* in multiple aspects. First, RNA-seq transcriptome profiling studies have revealed, in great detail, the timing of gene activation [19-22]. These data have allowed us to incorporate, to a greater extent, temporal information into a graphical organization of our network. Additionally, HTS data revealed vegetally enriched transcripts [14-16], which can be difficult to visualize by whole mount *in situ* hybridization.

Second, gain- and loss-of-function experiments, coupled with HTS or microarrays, have provided a wealth of regulatory connections. Additionally, ChIP-seq allows for the considerable improvement in the identification of direct target genes *in vivo*. Genome-wide binding of β -catenin [23,24], T-box TFs [25], Smad2/3 [26,27], Foxh1 [26], and several organizer-specific TFs [28] have all been investigated in *X. tropicalis*. Due to differences in data formatting across publications, we have mapped all available ChIP-seq datasets performed from blastula to early gastrula (Supplemental Table S1) to the version 9 *Xenopus tropicalis* genome (www.xenbase.org) using Bowtie2 [29], and identified peaks using the software MACS2 [30]. We then used Bedtools [31] to associate ChIP-seq peaks with TFs in our network, where peaks are filtered using a q-value of 0.01 and a peak distance of 10 kilobases (kb) from the gene body.

Using the criteria described above, we present a model of the GRN contributing to *Xenopus* mesendoderm development from fertilization through early gastrula (Nieuwkoop-Faber stage 10.5) (Figure 1). In total, this network includes 23 TFs and 12 growth factors. As we have focused on direct transcriptional responses, we have not included well-characterized

secreted signaling antagonists such as *chrd*, *nog* or *dkk1* in this network. However, we have chosen to include connections into the multi-signaling antagonist *cer1*, as the regulation of this gene has been extensively characterized [32,33]. We present the supporting evidence for these connections in Table 1, and in Supplemental Table S2 which includes additional putative connections between mesendodermal genes that did not satisfy our criteria. Thus, the directness of these connections is uncertain, and this represents an area of future investigation.

2.4 Organization of the network diagram

Using Biotapestry [34], we have built the GRN as a single bird's-eye view from the full genome [2] (Figure 1). In this visualization, all connections are displayed at once, regardless of time and space. We have made an effort to arrange the genes vertically based on approximate activation time [22], and horizontally from right to left across the dorsal to ventral axis [14]. At the top of the network, maternally inherited TFs (e.g., *Vegt*, *Foxh1*) and signaling ligands (e.g., *Gdf1*, *Wnt11b*) are shown. Maternally inherited TFs are displayed as diamonds, and ligands are shown as circles. All targets of signaling pathways are connected through chevrons indicating cell-surface receptors to their intracellular signal transducers. Zygotically activated growth factor ligands (i.e. *Nodals*, *Bmp4*) are connected back through the same signal transducer, so that all connections from a given pathway feed through a single TF node (e.g. *Smad2/3*).

We have compared the connections in this network with two previous GRNs [6,7] and find that a major improvement in the current network is the identification of more bone-fide direct transcriptional connections between TFs (Supplemental Table S3). Our network contains a total of 96 direct network connections - 82 positive and 14 negative. Direct connections are displayed in the GRN as a solid line connecting the upstream transcription factor to its downstream target. We also identify 36 putative direct interactions, which are displayed as dashed lines. Comparatively, Koide et al. [7] and Loose and Patient [6] previously reported a total of 41 and 60 direct network connections, respectively. Based on differing criteria from the current analysis, some of the direct connections identified by Loose and Patient [6] were here considered putative. Below, we review key features of the mesendoderm network.

3. Maternal factors and the initiation of the network

3.1 Vegetally-localized maternal transcription factors: *Vegt* and *Sox7*

Maternal factors play a critical role in the activation of the mesendoderm GRN. The best characterized TF important for the initiation of the *Xenopus* zygotic mesendoderm gene program is the T-box factor *Vegt*, which is asymmetrically localized vegetally [35-37]. The maternal knockdown of *vegt* in *Xenopus laevis* results in the loss of both mesoderm and endoderm [38,39]. Several studies have revealed *Vegt* to be a master regulator of the endoderm lineage through transcriptional regulation along two parallel routes: the zygotic activation of the Nodal genes, and activation of endodermal TFs [38-41]. Importantly, the *Vegt* LOF phenotype can be rescued by the injection of RNA encoding various Nodal ligands, indicating that a critical function of *Vegt* is the zygotic activation of the Nodal

signaling pathway [41]. Consistent with this, *nodal1*, *nodal5* and *gdf3* are direct targets of Vegt, and *nodal* is putatively direct based on protein synthesis inhibition. The expression of *nodal6* and *nodal2* are also regulated by Vegt, but it remains unclear whether these activations are direct.

In addition to regulating the pSmad2/3 signaling pathway, Vegt is important for the direct activation of core mesendodermal genes. TFs *sox17a*, *sox7*, and *gsc* are bona-fide Vegt direct targets; and *sox17b*, *mixer*, *mix1*, *hhex*, and *ventx1* are additional *putative* direct targets. The *cer1* gene, encoding a secreted Bmp/Wnt/Nodal antagonist [42], is also a direct target. At present, we have very little data to fully understand how these targets interact to establish the mesendoderm GRN.

In zebrafish and human, it appears that the T-box transcription factor Eomes plays a role similar to frog maternal Vegt in the activation of the endodermal gene regulatory program. Interestingly, functional studies of *Xenopus* Eomes add further support to the notion that this TF can perform an overlapping role with Vegt in specifying early mesendoderm. Eomes gain-of-function in naive animal caps results in expression of the mesendodermal genes *mix1*, *xbra*, *wnt8*, *sox17a*, *foxa4*, and *gsc* [43,44]. Furthermore, zygotic Vegt and Eomes cooperate in mesoderm formation in the late-gastrula embryo [25]. In zebrafish, perturbation analysis reveals that Eomesa is required for the activation of *sox17*, as well as the two nodal ligands *squint* and *cyclops* [45]. ChIP analyses confirm that Eomesa physically binds to the *squint* locus [46,47]. Eomesa also binds regulatory regions near *sox17*, and additional binding was observed for *mix11*, *foxa*, *foxa3*, *vent*, and *gsc* [46].

Eomes ChIP-seq and shRNA knockdown, in combination with microarray analysis, during the *in vitro* differentiation of human embryonic stem (ES) cells to definitive endoderm (DE) reveals that EOMES regulates a similar set of target genes as *Xenopus* Vegt [48]. The genes *MIXL1*, *GDF3*, *CER1*, *SOX17*, *FOXA1*, *FOXA2*, *FOXA3*, AND *VENTX* (with the exception of *FOXA3*, which is not found in *Xenopus*) are likely direct EOMES targets based on the application of our criteria. Finally, the NODAL gene – a *Xenopus* Vegt target – appears to have a regulatory region associated with EOMES binding, but is unaffected by the knockdown [48]. Overall, the Vegt T-box transcription factor has a highly evolutionarily conserved relationship with Eomes orthologs in some other species, sitting at the top of the gene regulatory hierarchy to function as a master regulator of endoderm formation.

In *Xenopus*, Sox7, a maternal SoxF type TF, has been implicated in the activation of mesendoderm targets. Overexpression of Sox7 mRNA into naive animal caps reveals that *nodal*, *nodal5*, and *nodal6* are putative direct target genes [49]. Since Vegt also putatively activates zygotic *sox7* expression [49], this indicates the likelihood of the *nodal* genes to be co-regulated by maternal Vegt and both maternal and zygotic Sox7. This network structure illustrates the importance of understanding gene regulation as a network, instead of activation mechanisms at the single gene/activator level. We also note that recent transcriptome profiling of blastomeres from 8-cell stage embryos identified 65 genes reproducibly enriched in the vegetal pole [12]. Among these, *otx1*, *pbx1*, *sox7* and *vegt* are highly enriched in the *Xenopus* vegetal tissue, most of them with poorly characterized roles in endoderm formation.

3.2 Nodal signaling and maternal Foxh1

It is clear that, in addition to directly activating endodermal TFs, a major function of Vegt is to activate expression of the *nodal* genes. The loss of Nodal signaling in *Xenopus* results in the disruption of endoderm and mesoderm formation, a severe delay or complete disruption of gastrulation movements, resulting in an embryo with a shortened anteroposterior (A-P) axis [26,50-54]. These findings are consistent with an analysis of Nodal-deficient zebrafish embryos (*sqt;cyc* double mutants) where mesodermal and endodermal markers are not expressed and gastrulation is abnormal [55]. Mice have a single *Nodal* gene, and *Nodal-null* embryos fail to form a primitive streak and do not undergo proper gastrulation [56]. Taken together, Nodal signaling is necessary for mesendoderm development in vertebrates.

How this pathway functions in the early formation of the mesendoderm has been the focus of intense study. The maternal TGF β ligand Gdf1 (previously known as Vg1) is also localized to the vegetal cells [57], and likely plays a role in anterior mesendoderm formation [58]; however, the endogenous role of Gdf1 has remained understudied, in part due to the inefficient conversion of the ligand precursor into its mature form [59]. Importantly, expression of *nodal5* and *nodal6* prior to the mid-blastula transition indicate that these ligands contribute to the earliest activation of the Nodal signaling pathway [60,61], and the early onset of *gdf3* suggests that it also contributes to the early phosphorylation of Smad2/3. The Nodal signaling cascade activates transcription in the blastula through Smad2/3 (in complex with the maternal partner to all R-Smad signaling, Smad4). Overall, the vegetal localization of Nodals and Gdf1 is consistent with the model in which high levels of Nodal promote endoderm and low levels promote mesoderm, which has been observed in *Xenopus* explant experiments [40].

As it is difficult to distinguish between the output of different Nodal and TGF β ligands, we have therefore fed all 7 ligands through a single Smad2/3 node. The mesendoderm GRN presented here contains 24 direct targets activated by Nodal signaling (Figure 1). These targets include core endoderm TFs such as *gata4*, *gata6*, *mix1*, *mixer*, and *foxa4*, dorsal endoderm genes such as *hhex* and *cer1*, the organizer gene *gsc* and the pan-mesodermal gene *t* (also known as *brachyury*), among others. These targets validate the notion that Nodal signaling contributes broadly in gene activation in the mesoderm and endoderm germ layers. Additionally, positive autoregulation of *nodal1* and *nodal2* promotes further enhanced expression of the signal [62].

The activated Smad2/3-Smad4 complex regulates target genes in concert with co-TFs, and to date, identified Smad2/3 co-factors include Foxh1, Eomes, Foxh1.2, Gtf2i, Gtf2ird1, Mixer, Tcf3 (also known as E2a) and Tp53 [48,63-71]. Of these, the transcriptional regulation via Smad2/3 interactions with maternal Foxh1 has been extensively investigated in *Xenopus* [26,66,72], zebrafish [70,73,74], mouse [75,76], and differentiated human ES cells [71]. Our *Xenopus* GRN contains 9 direct targets activated via Smad/Foxh1, including the growth factors, *nodal1*, *nodal2*, and *wnt8a*; the BMP/Wnt/Nodal antagonist *cer1*; and the TFs *gsc*, *otx2*, *mix1*, *hhex*, and *pitx2*. Among these, to date, *gsc* and *pitx2* regulation via Foxh1/Smad2 is conserved across mouse, fish and frog [62,77]. Finally, in human definitive endoderm differentiation *CER1*, *PITX2*, *GSC*, and *MIX1* are also induced by FOXH1/SMAD2 [78].

It has been well-characterized in mouse, fish and frog that the loss of Foxh1 does not fully recapitulate the loss of Nodal signaling – indicating the necessity for additional Smad2/3 binding partners. To this extent, in zebrafish, Eomes has been implicated as the Smad2/3 co-factor responsible for the remaining Nodal-mediated regulation that occurred in the Foxh1-null [70]. Our network suggests *gata4*, *gata6*, *eomes*, and *foxa2* are also regulated by Nodal-signaling via a Foxh1-independent mechanism. It will be necessary to investigate whether Eomes and Mixer regulate these targets in a Foxh1-independent fashion.

3.3 Wnt/ β -catenin signaling

In addition to germ layer specification along the animal-vegetal axis, Nodal signaling is critical in patterning the embryo along the dorsal-ventral axis. While *vegt* mRNA appears to be distributed across the vegetal tissue, Nodal signaling is higher in the dorsal mesendoderm of the *Xenopus* blastula [52,79]. This pattern is attributed to high levels of Wnt/ β -catenin signaling on the dorsal side of the embryo. While a detailed discussion of dorsal-ventral patterning of mesendoderm is beyond the scope of this review, it is useful to discuss in brief the role of maternal Wnt/ β -catenin signaling. Maternal *wnt11b*, is localized to the vegetal pole in the egg, relocated to the dorsal vegetal cells following cortical rotation, and activates Wnt signaling to specific dorsal fate [80-82]. Dorsal nuclear β -catenin directly regulates *sia1* and *sia2*, two homeobox genes that control dorso-anterior specification [83,84] and many other genes, via the canonical Wnt cascade feeding through β -catenin-Lef/Tcf complexes. Maternal Wnt/ β -catenin signaling also activates the expression of all Nodal genes in the dorsal mesendoderm, in particular the early activation of *nodal5* and *nodal6* [61,85]. Both Nodal and Wnt signaling are critical for the formation of the Nieuwkoop center and Spemann organizer [84,86,87], and the network reveals substantial overlap in the regulation of dorsal mesendoderm target genes such as *hhx*, *lhx1*, *otx2*, *cer1* and *gsc*. Consistent with this crosstalk model, there is a substantial co-occurrence between Foxh1 (a major Smad2/3 co-factor) and β -catenin ChIP-seq peaks [24,26]. It should be noted that recent β -catenin ChIP-seq performed by Nakamura et al. [24] reveals β -catenin binding associated with target genes previously thought to be indirectly regulated by Wnt/ β -catenin via Sia, such as *hhx* and *gsc* [84,166]. While the biological activities of the putative enhancers remain to be validated, this finding suggests that dorsal mesendoderm targets are regulated through complex network motifs (see section 5). Taken together, the mesendoderm GRN is highly controlled by maternal Vegt, and the signaling inputs from Nodal and Wnt signaling pathways, the activation of which coincide with the onset of zygotic gene transcription.

4. Core zygotic mesendoderm transcription factors

A number of zygotic TFs have been identified as critical for the formation of the mesendoderm. Here, we discuss the roles of the Mix family, Gata4/5/6, Foxa and Sox17.

4.1 The Mix family

The critical role of the Mix family TFs in mesendoderm development has been investigated across numerous model systems [88]. In the two *Xenopus* species, the single mammalian gene encoding Mix-like 1 (Mixl1) is represented by *mix1*, *mixer*, and species-specific expansions and losses of genes referred to as *bix* [89].

In our network, we have examined connections into and from *mix1* and *mixer* based on integrated data from *X. laevis* and *X. tropicalis*. Of these, *mix1* is the earliest to be activated at the mid-blastula transition via direct regulation by Smad/Foxh1, and putative regulation by Vegt. To date, evidence also supports the direct activation of *gsc* and *cer1*, and the repression of *t* by Mix1. This network supports the notion that Mix1 is critical for the activation of dorsoanterior mesendoderm, and the exclusion of *t* from the dorsal organizer [90-92]. This places Mix1 at the top of a negative feed-forward loop, together with Gsc [93], which also represses *t* expression (see section 5.2). This loop may be conserved in mammals, as differentiating Mix1-null mouse ES cells reveal a down-regulation of *gsc* and an up-regulation of *t* [94]. It is conceivable that Mixer – which also directly activates *gsc* – functions in a similar capacity to repress ventrolateral mesoderm. Supporting this, Mixer deficient *Xenopus* embryos showed up-regulation of mesoderm genes including *eomes*, *fgf3*, *fgf8*, *not*, and *gata2* [95]. Similarly, *t* expression was mildly reduced in Mixer morphants. Finally, while the relationship between *X. laevis* and *X. tropicalis* Bix TFs is unclear, current evidence suggests that *bix1*, in *X. laevis*, is directly activated by T [96], and in *X. tropicalis*, is activated by Nodal/Smad2 [26]. In *X. laevis*, *bix2* (previously known as *milk*) is putatively activated by Nodal/Smad2 [97], and in turn directly activates *gsc* [65,97]. Further elucidating the direct targets of Mix TFs in amphibians, fish, and mammals will be crucial to teasing out these subnetworks.

4.2 Gata family

The Gata transcription factors are highly conserved regulators of endoderm formation across metazoan model systems. In invertebrates, Gata transcription factors play crucial roles in the formation of *Drosophila* midgut [98,99], in the E lineage during *C. elegans* germ layer patterning [100-102], and the sea urchin mesendoderm [2,103]. In vertebrates, the Gata4/5/6 subfamily of Gata factors play a role in the formation of mouse extraembryonic endoderm [104,105], and in the formation of *Xenopus* [106,107] and zebrafish [108,109] endoderm.

Despite their importance, little is known about their molecular targets. In *Xenopus*, the putative direct targets of Gata4/5/6 are endodermal genes *hnf1b* and *sox17a*. Evidence in mouse ES cells also suggests that both *hnf1b* and *sox17a* are Gata4/6 targets [104,105], and zebrafish *gata5* mutants show a reduction in *sox17* expression [110]. While the directness of these interactions is unclear in the mouse and zebrafish, conservation of gene activation suggests similar network topologies are likely operating between frog, zebrafish and mouse.

A common feature of the Gata factors is the extensive mutual regulation among these three genes. In *Xenopus*, *gata5* gain-of-function upregulates *gata4* expression, while *gata4* gain-of-function upregulates *gata6* expression [107]. In zebrafish, loss of *gata5* downregulates the expression of *gata6* and vice versa [109]. Overexpression of Gata4, Gata5, or Gata6 in mouse ES cells results in the upregulation of all three factors [104,105,111], and *Gata4*-null mice show reduced *Gata6* expression [112].

Presently, it is unclear whether these connections are direct; however, functional analyses of a *Gata4* cis-regulatory module in mouse supports direct co-regulation. Gata4 was shown to bind to an enhancer controlling foregut and midgut expression, suggesting an autoregulatory loop [113]. A second enhancer that controls *Gata4* expression in the septum transversum and

the mesenchyme surrounding the liver are bound by all three Gata factors [114]. These data suggest a direct positive relationship between the three factors.

4.3 Sox17

Sox17 is a highly conserved endodermal transcription factor across vertebrates. Sox17 plays a role in *Xenopus* [115] and zebrafish [116] endoderm formation, and in both mouse extraembryonic and definitive endoderm formation [117], and in the definitive endoderm in human ES cell assays [118]. In *Xenopus*, little is known about the direct targets of Sox17, but putative direct targets include *foxa1*, *foxa2*, and *hnf1b* [119-121]. Sox17 also directly regulates the expression of *foxa2* orthologs in mouse and human extraembryonic and definitive endoderm [118,122]. In addition, Sox17 targets the extraembryonic endoderm marker *Hnf1b* in mouse ES cells [122]. However, it's not known if either of these targets are direct. Finally, functional evidence shows that Sox17 genes are subjected to a positive feedback loop [119,121].

4.4 Foxa family

Foxa TFs are critical for endoderm development across diverse organisms [123,124]. Of the three Foxa TFs in mouse (*Foxa1*, *Foxa2*, and *Foxa3*), Foxa2 is required for early development. Foxa2-null mice display defects in the node (the equivalent of the Spemann organizer) and later gut tube [125]. Both Foxa1 and 2 bind to liver-specific enhancers in mouse pluripotent gut endoderm, well before these genes become transcriptionally active [126], and genetic analyses indicated that these TFs function together in hepatic development [127]. Since they have the capacity to bind to and open compact chromatin [128], Foxa TFs have been deemed 'pioneer factors' for gut development [129].

The *Xenopus tropicalis* genome encodes three Foxa TFs (*foxa1*, *foxa2*, and *foxa4*), which are zygotically transcribed. Gain-of-function analyses indicate that Foxa2 inhibits mesoderm and anterior endoderm formation in the gastrula embryo [130]. Loss-of-function analyses in sea urchin also support the conservation of this mechanism [131]. However, as the overexpression of VP16-Foxa2 fusion protein phenocopied the overexpression of Foxa2, Foxa2 presumably functions as an activator [130], and the authors postulate that Foxa2 activates a key repressor of mesodermal cell fate. At present, the direct targets of Foxa in the early embryo are unclear. Finally, in *Xenopus*, *foxa4* is the earliest and most abundantly expressed *foxa* gene during early mesendoderm specification. However, Foxa4's role in early mesendoderm development is not known, although by early neurula stages it promotes notochord formation and inhibits prechordal and paraxial mesoderm [132]. Our network analysis reveals that *foxa4* is activated by Smad2/3, via a Foxh1-independent mechanism [26]. Since *foxa4* expression is repressed directly by the Smad1 target *ventx1* [133], these connections support exclusion of *foxa4* expression from the ventrolateral mesoderm.

5. Network motifs in the *Xenopus* mesendodermal GRN

Network motifs are a subgroup of patterns found in GRN architectures. Here we analyzed the network motifs – representing autoregulatory, feedback, and feedforward loops [134] – found in the network presented here, as well as those previously reported [6,7]. Due to the

increase in the number of direct connections presented in this review over previous networks, we identified significantly more motifs (Figure 2 and Supplemental Table S3), which we will discuss below.

5.1 Autoregulatory and feedback loops

Autoregulatory loops involve the self-regulation of a transcription factor or a signaling pathway (Figure 2A and Supplemental Table S4). Based on our criteria, current evidence supports 4 direct autoregulatory loops. These include the positive autoregulation of Nodal signaling in the endoderm and dorsal mesoderm [26,62,135]; the positive autoregulation of *ventx2* on the ventral side of the embryo [136]; the negative autoregulation of *gsc* [18,28,137]; and the exclusion of *wnt8a* expression in the dorsal mesendoderm through the action of Tcf1/Ctnnb1 [24,138].

We next computationally interrogated our network for feedback loops – a motif that involves mutual regulation between two genes X and Y (Figure 2B and Supplemental Table S4). In a negative feedback loop, gene X positively regulates gene Y, while gene Y negatively regulates gene X; a double negative feedback loop is defined by mutual inhibition between the two genes. A positive feedback loop involves a mutual activation between genes X and Y. We identified five feedback loops in our network – one negative feedback, one double negative feedback and three positive feedback loops.

Based on gain- and loss-of-function analyses, a double negative feedback loop between the dorsal organizer gene *gsc* and the ventral gene *ventx2* has been proposed [139]. While reporter assays and EMSA experiments had confirmed *gsc* as a direct target of Ventx2 [140], it was only recently via ChIP-seq that Gsc binding to *ventx2* cis-regulatory regions has been confirmed [28]. This type of feedback loop enables the formation of sharp expression boundaries between cell lineages, as computationally demonstrated in modeling a double negative feedback between *gsc* and *t* ([141,142].

The network reveals a negative feedback loop between Gsc and Wnt/ β -catenin (Ctnnb1), whereby *gsc* – activated by maternal Wnt11 b/ β -catenin [82,85,138,143] – represses the expression of zygotic *wnt8a* from the dorsal organizer [28,137]. This type of feedback loop can be useful in cases where the initial activator becomes unnecessary in the control of later gene expression.

Positive feedback loops enable continuous expression of two genes that are important for the same lineage. On the ventral side of the embryo, Smad1 mediating BMP signaling binds the regulatory region of *ventx2* and activates its expression, which in turn appears to regulate the expression of *bmp4* and increases the production of Bmp4 ligand [144]. Finally, in the mesendoderm, the Nodal and Wnt/ β -catenin signaling pathways positively feedback into each other [24,26,85,119,145-147]. In this motif, maternal Wnt11b/ β -catenin activates the expression of all *nodal* genes in the dorsal mesendoderm. In turn, Smad2/3 (mediated by Foxh1), activates zygotic *wnt8a*, which is excluded from the dorsal organizer via the *gsc* gene as discussed above.

5.2 Feedforward loops

One important feature found in the network presented is the prominent presence of feedforward loops in three-gene network motifs. Of the eight different types of feedforward loops [148] (Figure 2C, Supplemental Table S4), the coherent feedforward type I is the most abundant. The relative abundance of coherent feedforward type I loops compared to other types of feedforward loops is a feature found in a variety of transcriptional GRNs, including *E. coli* [148], *P. aeruginosa* [149], *S. cerevisiae* [148], and *D. melanogaster* embryogenesis [150,151]. In our network, approximately three-quarters of identified feedforward loops were coherent type I (Supplemental Table S4). In this type of loop, a positive regulator (gene X) and its target (gene Y) both positively regulate the expression of a common downstream gene (gene Z). In the *Xenopus* mesendoderm GRN, this loop appears frequently where gene X is maternal, gene Y is a primary activated zygotic gene and gene Z is a secondary activated zygotic gene (Figure 2D). In the majority of cases, the initial activator appears to be either β -catenin, Foxh1, Smad2/3, or Vegt (gene X). These maternal factors activate the expression of early and mid-blastula zygotic genes such as *wnt8a*, *sia1*, *sia2*, *mix1*, *gsc*, and the *nodal* genes (gene Y), which, in turn, activate the expression a larger number of later expressed mesendodermal genes (gene Z). Some examples of gene Z include *cer1*, *eomes*, *ventx1*, *hhex*, and *pitx2*. The benefits of this type of loop depend on whether the co-regulation of Z by both X and Y is an ‘AND-gate’ (where both factors are required to activate factor Z) or an ‘OR-gate’ (where either factor can activate gene Z) [134]. An ‘AND-gate’ can be beneficial in the tight control of factor Z expression, as factor Z is only activated once factor Y is expressed. On the other hand, an ‘OR-gate’ enables the sustained expression of factor Z, despite the loss of the initial factor X [134]. Such would be a critical motif functioning in early *Xenopus* embryogenesis, where maternal factors that act as the initial activators are degraded during blastula stages. Then, their direct, primary, activated zygotic targets can function to maintain the expression of later, secondary, activated genes. It's not clear as to whether the coherent type I feedforward loops we identified are controlled by AND-gates or OR-gates. It will be important to address this type of question as we further refine our understanding of gene regulation in the early *Xenopus* mesendoderm.

The other types of feedforward loops involve a negative regulation between genes X, Y and Z [148] (Figure 2C, Supplemental Table S4). In the *Xenopus* system, these types of loops appear enriched among genes required for dorsal-ventral patterning of the mesendoderm. For example, Smad2/3 signaling promotes both *t* and *gsc* expression [26,43,87,93,152]. In turn, *gsc* represses *t* from the dorsal organizer. In a similar loop, *otx2* also functions in the restriction of *t* expression [90,153]. These loops, along with negative and double negative feedback loops, appear to be particularly useful in regionalizing the dorsal and ventral sides of the mesendoderm. It is likely that more of these types of loops will be identified in the future, as we better define the expression domains of more transcription factors during *Xenopus* gastrulation [14-16]. Also, as this review focused primary on mesendodermal genes, the molecular mechanism for the regionalization of the embryo between mesendoderm and prospective ectoderm is not reflected on this network. However, as some animal pole factors repressing Nodal signaling have been identified (e.g., maternal Sox3) [154], similar network motifs might be used in animal-vegetal patterning.

6. Prospects

We have generated a comprehensive gene regulatory network governing *Xenopus* mesendoderm development. New findings, and the increased accessibility of HTS technologies, have contributed greatly to the number of direct regulatory interactions between critical factors, and have revealed many more possible players whose functions remain unclear. We predict that the network will exponentially increase as more datasets are generated. In addition, advances in the use of CRISPR/Cas9 in *Xenopus* provides the opportunity to modify the endogenous interactions between TFs and their target CRMs. This will enable in-depth investigations into the role of CRMs in gene regulation, ultimately aiding in addressing one of the most critical questions in biology: how mutations in regulatory regions influence overall gene expression levels.

The mesendoderm network is initiated in the blastula embryo at the onset of zygotic gene activation, and the network presented here – which extends through the beginning of gastrulation – covers a timespan of approximately 3 hours. During this time of rapid developmental transitions, as maternal factors – important for the initiation of the network – are degraded, and zygotic transcription ramps up, we expect the network to be highly dynamic. While single-stage analyses may be sufficient in identifying direct target genes, the investigation of TF targets over time provides valuable kinetic information to uncover the complexities of the dynamic regulatory network.

In addition to transcription factors and signaling molecules, screens for non-coding RNAs, including microRNAs [155-159], and long non-coding RNAs [160,161], have identified many more potential regulators of gene expression. One example is through the negative regulation by mir-427 of the Nodal ligands *nodal5* and *nodal6*, and the nodal antagonist *lefty* [162]. Loss of function of mir-427 leads to mesodermal patterning defects. Interestingly, the interaction between Nodal signaling, Lefty and mir-427 generates an incoherent type II feedforward loop. As we learn more about the roles of these non-coding RNAs, we will have to integrate their regulatory roles into the GRN diagram.

As the GRN increases in connectivity and becomes more complete, this will enable future researchers to investigate the GRN from a systems level perspective. Identification of various network motifs can provide some new and interesting hypotheses based on the theoretical properties of these motifs [134], which can be experimentally tested *in vivo*. Additionally, quantitative modeling of these network can delve deeper into the nature of the regulatory relationships between transcription factors, as has been done in smaller networks [141,142,163]. Some of the challenging questions in the future are to parse which mesendodermal factors play major roles in regulating network function and maintaining network output robustness; as well as the role of redundancy in network regulation, and feedback/feedforward loop regulation. The *Xenopus* mesendoderm GRN, with its rich history and amenability to modern genomic tools, presents itself as one of the best systems to study these types of network questions *in vivo*. We hope that this network will provide a useful framework in moving towards a greater understanding of the complex GRN controlling early mesendoderm development, and as well as the formation of later endodermal derivatives.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Xenbase (<http://www.xenbase.org/>; RRID:SCR_003280) for *Xenopus* resources, the members of the Cho laboratory for useful discussion, and the UC Irvine Office of Information Technology for ongoing support of the High Performance Computing Cluster. We apologize to researchers whose work we were unable to include due to space limitations. This work was supported by NIH HD073179 to K.W.Y.C.

References

1. Levine M 1. Transcriptional enhancers in animal development and evolution. *Curr Biol*. 2010; 20:R754–63. DOI: 10.1016/j.cub.2010.06.070 [PubMed: 20833320]
2. Davidson EH 1, Rast JP, Oliveri P, Ransick A, Calestani C, Yuh CH, et al. A genomic regulatory network for development. *Science*. 2002; 295:1669–1678. DOI: 10.1126/science.1069883 [PubMed: 11872831]
3. Levine M 1, Davidson EH. Gene regulatory networks for development. *Proc Natl Acad Sci USA*. 2005; 102:4936–4942. DOI: 10.1073/pnas.0408031102 [PubMed: 15788537]
4. Sakabe M 1, Matsui H, Sakata H, Ando K, Yamagishi T, Nakajima Y. Understanding heart development and congenital heart defects through developmental biology: a segmental approach. *Congenit Anom (Kyoto)*. 2005; 45:107–118. DOI: 10.1111/j.1741-4520.2005.00079.x [PubMed: 16359490]
5. Spence JR 1, Wells JM. Translational embryology: using embryonic principles to generate pancreatic endocrine cells from embryonic stem cells. *Dev Dyn*. 2007; 236:3218–3227. DOI: 10.1002/dvdy.21366 [PubMed: 17973329]
6. Loose M 1, Patient R. A genetic regulatory network for *Xenopus* mesendoderm formation. *Developmental Biology*. 2004; 271:467–478. DOI: 10.1016/j.ydbio.2004.04.014 [PubMed: 15223347]
7. Koide T 1, Hayata T, Cho KWY. *Xenopus* as a model system to study transcriptional regulatory networks. *Proc Natl Acad Sci USA*. 2005; 102:4943–4948. DOI: 10.1073/pnas.0408125102 [PubMed: 15795378]
8. Heasman J 1. Patterning the early *Xenopus* embryo. *Development*. 2006; 133:1205–1217. DOI: 10.1242/dev.02304 [PubMed: 16527985]
9. Zorn AM 1, Wells JM. Vertebrate endoderm development and organ formation. *Annu Rev Cell Dev Biol*. 2009; 25:221–251. DOI: 10.1146/annurev.cellbio.042308.113344 [PubMed: 19575677]
10. Flachsova M 1, Sindelka R, Kubista M. Single blastomere expression profiling of *Xenopus laevis* embryos of 8 to 32-cells reveals developmental asymmetry. *Sci Rep*. 2013; 3doi: 10.1038/srep02278
11. Grant PA 1, Yan B, Johnson MA, Johnson DLE, Moody SA. Novel animal pole-enriched maternal mRNAs are preferentially expressed in neural ectoderm. *Dev Dyn*. 2014; 243:478–496. DOI: 10.1002/dvdy.24082 [PubMed: 24155242]
12. De Domenico E 1, Owens NDL, Grant IM, Gomes-Faria R, Gilchrist MJ. Molecular asymmetry in the 8-cell stage *Xenopus tropicalis* embryo described by single blastomere transcript sequencing. *Developmental Biology*. 2015; 408:252–268. DOI: 10.1016/j.ydbio.2015.06.010 [PubMed: 26100918]
13. Gaur S 1, Mandelbaum M, Herold M, Majumdar HD, Neilson KM, Maynard TM, et al. Neural transcription factors bias cleavage stage blastomeres to give rise to neural ectoderm. *Genesis*. 2016; 54:334–349. DOI: 10.1002/dvg.22943 [PubMed: 27092474]
14. Blitz IL 1, Paraiso KD, Patrushev I, Chiu WTY, Cho KWY, Gilchrist MJ. A catalog of *Xenopus tropicalis* transcription factors and their regional expression in the early gastrula stage embryo. *Developmental Biology*. (in press). doi: 10.1016/j.ydbio.2016.07.002.

15. Ding Y 1, Colozza G, Zhang K, Moriyama Y, Ploper D, Sosa EA, et al. Genome-wide analysis of dorsal and ventral transcriptomes of the *Xenopus laevis* gastrula. *Developmental Biology*. in press.
16. Popov IK 1, Kwon T, Crossman DK, Crowley MR, Wallingford JB, Chang C. Identification of new regulators of embryonic patterning and morphogenesis in *Xenopus* gastrulae by RNA sequencing. *Developmental Biology*. in press.
17. Li E 1, Davidson EH. Building developmental gene regulatory networks. *Birth Defects Res C Embryo Today*. 2009; 87:123–130. DOI: 10.1002/bdrc.20152 [PubMed: 19530131]
18. Danilov V 1, Blum M, Schweickert A, Campione M, Steinbeisser H. Negative autoregulation of the organizer-specific homeobox gene goosecoid. *Journal of Biological Chemistry*. 1998; 273:627–635. [PubMed: 9417125]
19. Paranjpe SS 1, Jacobi UG, van Heeringen SJ, Veenstra GJC. A genome-wide survey of maternal and embryonic transcripts during *Xenopus tropicalis* development. *BMC Genomics*. 2013; 14:762. doi:10.1186/1471-2164-14-762. [PubMed: 24195446]
20. Tan MH 1, Au KF, Yablonovitch AL, Wills AE, Chuang J, Baker JC, et al. RNA sequencing reveals a diverse and dynamic repertoire of the *Xenopus tropicalis* transcriptome over development. *Genome Res*. 2013; 23:201–216. DOI: 10.1101/gr.141424.112 [PubMed: 22960373]
21. Collart C 1, Owens NDL, Bhaw-Rosun L, Cooper B, De Domenico E, Patrushev I, et al. High-resolution analysis of gene activity during the *Xenopus* mid-blastula transition. *Development*. 2014; 141:1927–1939. DOI: 10.1242/dev.102012 [PubMed: 24757007]
22. Owens NDL 1, Blitz IL, Lane MA, Patrushev I, Overton JD, Gilchrist MJ, et al. Measuring Absolute RNA Copy Numbers at High Temporal Resolution Reveals Transcriptome Kinetics in Development. *Cell Reports*. 2016; 14:632–647. DOI: 10.1016/j.celrep.2015.12.050 [PubMed: 26774488]
23. Kjolby RAS 1, Harland RM. Genome-wide identification of Wnt/ β -catenin transcriptional targets during *Xenopus* gastrulation. *Developmental Biology*. in press.
24. Nakamura Y 1, de Paiva Alves E, Veenstra GJC, Hoppler S. Tissue- and stage-specific Wnt target gene expression is controlled subsequent to β -catenin recruitment to cis-regulatory modules. *Development*. 2016; 143:1914–1925. DOI: 10.1242/dev.131664 [PubMed: 27068107]
25. Gentsch GE 1, Owens NDL, Martin SR, Piccinelli P, Faial T, Trotter MWB, et al. In Vivo T-Box Transcription Factor Profiling Reveals Joint Regulation of Embryonic Neuromesodermal Bipotency. *Cell Rep*. 2013; 4:1185–1196. DOI: 10.1016/j.celrep.2013.08.012 [PubMed: 24055059]
26. Chiu WT 1, Le R Charney, Blitz IL, Fish MB, Li Y, Biesinger J, et al. Genome-wide view of TGF / Foxh1 regulation of the early mesendoderm program. *Development*. 2014; 141:4537–4547. 1. DOI: 10.1242/dev.107227 [PubMed: 25359723]
27. Gupta R 1, Wills A, Ucar D, Baker J. Developmental enhancers are marked independently of zygotic Nodal signals in *Xenopus*. *Developmental Biology*. 2014; 395:38–49. DOI: 10.1016/j.ydbio.2014.08.034 [PubMed: 25205067]
28. Yasuoka Y 1, Suzuki Y, Takahashi S, Someya H, Sudou N, Haramoto Y, et al. Occupancy of tissue-specific cis-regulatory modules by Otx2 and TLE/Groucho for embryonic head specification. *Nature Communications*. 2014; 5:4322. doi: 10.1038/ncomms5322
29. Langmead B 1, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Meth*. 2012; 9:357–359. DOI: 10.1038/nmeth.1923
30. Zhang Y 1, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-based analysis of ChIP-Seq (MACS). *Genome Biol*. 2008; 9:R137. doi: 10.1186/gb-2008-9-9-r137 [PubMed: 18798982]
31. Quinlan AR 1, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics*. 2010; 26:841–842. DOI: 10.1093/bioinformatics/btq033 [PubMed: 20110278]
32. Yamamoto S 1, Hikasa H, Ono H, Taira M. Molecular link in the sequential induction of the Spemann organizer: direct activation of the cerberus gene by Xlim-1, Xotx2, Mix.1, and Siamois, immediately downstream from Nodal and Wnt signaling. *Developmental Biology*. 2003; 257:190–204. DOI: 10.1016/S0012-1606(03)00034-4 [PubMed: 12710967]

33. Sudou N 1, Yamamoto S, Ogino H, Taira M. Dynamic in vivo binding of transcription factors to cis-regulatory modules of *cer* and *gsc* in the stepwise formation of the Spemann-Mangold organizer. *Development*. 2012; 139:1651–1661. DOI: 10.1242/dev.068395 [PubMed: 22492356]
34. Longabaugh WJR 1, Davidson EH, Bolouri H. Computational representation of developmental genetic regulatory networks. *Developmental Biology*. 2005; 283:1–16. DOI: 10.1016/j.ydbio.2005.04.023 [PubMed: 15907831]
35. Lustig KD 1, Kroll KL, Sun EE, Kirschner MW. Expression cloning of a *Xenopus* T-related gene (*Xombi*) involved in mesodermal patterning and blastopore lip formation. *Development*. 1996; 122:4001–4012. [PubMed: 9012520]
36. Stennard F 1, Carnac G, Gurdon JB. The *Xenopus* T-box gene, *Antipodean*, encodes a vegetally localised maternal mRNA and can trigger mesoderm formation. *Development*. 1996; 122:4179–4188. [PubMed: 9012537]
37. Zhang J 1, King ML. *Xenopus* VegT RNA is localized to the vegetal cortex during oogenesis and encodes a novel T-box transcription factor involved in mesodermal patterning. *Development*. 1996; 122:4119–4129. [PubMed: 9012531]
38. Zhang J 1, Houston DW, King ML, Payne C, Wylie C, Heasman J. The role of maternal VegT in establishing the primary germ layers in *Xenopus* embryos. *Cell*. 1998; 94:515–524. [PubMed: 9727494]
39. Kofron M 1, Demel T, Xanthos J, Lohr J, Sun B, Sive H, et al. Mesoderm induction in *Xenopus* is a zygotic event regulated by maternal VegT via TGFβ growth factors. *Development*. 1999; 126:5759–5770. [PubMed: 10572051]
40. Clements D 1, Friday RV, Woodland HR. Mode of action of VegT in mesoderm and endoderm formation. *Development*. 1999; 126:4903–4911. [PubMed: 10518506]
41. Xanthos JB 1, Kofron M, Wylie C, Heasman J. Maternal VegT is the initiator of a molecular network specifying endoderm in *Xenopus laevis*. *Development*. 2001; 128:167–180. [PubMed: 11124113]
42. Piccolo S 1, Agius E, Leyns L, Bhattacharyya S, Grunz H, Bouwmeester T, et al. The head inducer *Cerberus* is a multifunctional antagonist of Nodal, BMP and Wnt signals. *Nature*. 1999; 397:707–710. DOI: 10.1038/17820 [PubMed: 10067895]
43. Ryan K 1, Garrett N, Mitchell A, Gurdon JB. *Eomesodermin*, a Key Early Gene in *Xenopus* Mesoderm Differentiation. *Cell*. 1996; 87:989–1000. DOI: 10.1016/S0092-8674(00)81794-8 [PubMed: 8978604]
44. Conlon FL, Fairclough L, Price BM, Casey ES, Smith JC. Determinants of T box protein specificity. *Development*. 2001; 128:3749–3758. [PubMed: 11585801]
45. Bjornson CRR 1, Griffin KJP, Farr GH, Terashima A, Himeda C, Kikuchi Y, et al. *Eomesodermin* is a localized maternal determinant required for endoderm induction in zebrafish. *Developmental Cell*. 2005; 9:523–533. DOI: 10.1016/j.devcel.2005.08.010 [PubMed: 16198294]
46. Nelson AC 1, Cutty SJ, Niini M, Stemple DL, Flicek P, Houart C, et al. Global identification of Smad2 and *Eomesodermin* targets in zebrafish identifies a conserved transcriptional network in mesendoderm and a novel role for *Eomesodermin* in repression of ectodermal gene expression. *BMC Biol*. 2014; 12:81.doi: 10.1186/s12915-014-0081-5 [PubMed: 25277163]
47. Xu P 1, Zhu G, Wang Y, Sun J, Liu X, Chen YG, et al. Maternal *Eomesodermin* regulates zygotic nodal gene expression for mesendoderm induction in zebrafish embryos. *J Mol Cell Biol*. 2014; 6:272–285. DOI: 10.1093/jmcb/mju028 [PubMed: 24924767]
48. Teo AKK 1, Arnold SJ, Trotter MWB, Brown S, Ang LT, Chng Z, et al. Pluripotency factors regulate definitive endoderm specification through *eomesodermin*. *Genes & Development*. 2011; 25:238–250. DOI: 10.1101/gad.607311 [PubMed: 21245162]
49. Zhang C 1, Basta T, Klymkowsky MW. *SOX7* and *SOX18* are essential for cardiogenesis in *Xenopus*. *Dev Dyn*. 2005; 234:878–891. DOI: 10.1002/dvdy.20565 [PubMed: 16193513]
50. Osada SI 1, Wright CV. *Xenopus* nodal-related signaling is essential for mesendodermal patterning during early embryogenesis. *Development*. 1999; 126:3229–3240. DOI: 10.1073/pnas.88.15.6511 [PubMed: 10375512]

51. Sun BI 1, Bush SM, Collins-Racie LA, LaVallie ER, DiBlasio-Smith EA, Wolfman NM, et al. derrière: a TGF-beta family member required for posterior development in *Xenopus*. *Development*. 1999; 126:1467–1482. [PubMed: 10068640]
52. Agius E 1, Oelgeschläger M, Wessely O, Kemp C, De Robertis EM. Endodermal Nodal-related signals and mesoderm induction in *Xenopus*. *Development*. 2000; 127:1173–1183. [PubMed: 10683171]
53. Onuma Y 1, Takahashi S, Yokota C, Asashima M. Multiple nodal-related genes act coordinately in *Xenopus* embryogenesis. *Developmental Biology*. 2002; 241:94–105. DOI: 10.1006/dbio.2001.0493 [PubMed: 11784097]
54. Ho DM 1, Chan J, Bayliss P, Whitman M. Inhibitor-resistant type I receptors reveal specific requirements for TGF-beta signaling in vivo. *Developmental Biology*. 2006; 295:730–742. DOI: 10.1016/j.ydbio.2006.03.050 [PubMed: 16684517]
55. Feldman B 1, Gates MA, Egan ES, Dougan ST, Rennebeck G, Sirotkin HI, et al. Zebrafish organizer development and germ-layer formation require nodal-related signals. *Nature*. 1998; 395:181–185. DOI: 10.1038/26013 [PubMed: 9744277]
56. Conlon FL 1, Lyons KM, Takaesu N, Barth KS, Kispert A, Herrmann B, et al. A primary requirement for nodal in the formation and maintenance of the primitive streak in the mouse. *Development*. 1994; 120:1919–1928. DOI: 10.1016/0168-9525(94)90026-4 [PubMed: 7924997]
57. Weeks DL 1, Melton DA. A maternal mRNA localized to the vegetal hemisphere in *Xenopus* eggs codes for a growth factor related to TGF-beta. *Cell*. 1987; 51:861–867. [PubMed: 3479264]
58. Birsoy B 1, Kofron M, Schaible K, Wylie C, Heasman J. Vg 1 is an essential signaling molecule in *Xenopus* development. *Development*. 2006; 133:15–20. DOI: 10.1242/dev.02144 [PubMed: 16308332]
59. Kiecker C, Bates T, Bell E. Molecular specification of germ layers in vertebrate embryos. *Cell Mol Life Sci*. 2015; 73:923–947. DOI: 10.1007/s00018-015-2092-y [PubMed: 26667903]
60. Skirkanich J 1, Luxardi G, Yang J, Kodjabachian L, Klein PS. An essential role for transcription before the MBT in *Xenopus laevis*. *Developmental Biology*. 2011; 357:478–491. DOI: 10.1016/j.ydbio.2011.06.010 [PubMed: 21741375]
61. Yang J 1, Tan C, Darken RS, Wilson PA, Klein PS. β -Catenin/Tcf-regulated transcription prior to the midblastula transition. *Development*. 2002; doi: 10.1242/dev.00150
62. Osada SI 1, Saijoh Y, Frisch A, Yeo CY, Adachi H, Watanabe M, et al. Activin/nodal responsiveness and asymmetric expression of a *Xenopus* nodal-related gene converge on a FAST-regulated module in intron 1. *Development*. 2000; 127:2503–2514. [PubMed: 10804190]
63. Chen X 1, Rubock MJ, Whitman M. A transcriptional partner for MAD proteins in TGF- β signalling. *Nature*. 1996; 383:691–696. doi:10.1038/383691 a. [PubMed: 8878477]
64. Chen X 1, Weisberg E, Fridmacher V, Watanabe M, Naco G, Whitman M. Smad4 and FAST-1 in the assembly of activin-responsive factor. *Nature*. 1997; 389:85–89. DOI: 10.1038/38008 [PubMed: 9288972]
65. Germain S, Howell M, Esslemont GM, Hill CS. Homeodomain and winged-helix transcription factors recruit activated Smads to distinct promoter elements via a common Smad interaction motif. *Genes & Development*. 2000; 14:435–451. DOI: 10.1101/gad.14.4.435 [PubMed: 10691736]
66. Howell M 1, Inman GJ, Hill CS. A novel *Xenopus* Smad-interacting forkhead transcription factor (XFast-3) cooperates with XFast-1 in regulating gastrulation movements. *Development*. 2002; 129:2823–2834. [PubMed: 12050132]
67. Ring C 1, Ogata S, Meek L, Song J, Ohta T, Miyazono K, et al. The role of a Williams-Beuren syndrome-associated helix-loop-helix domain-containing transcription factor in activin/nodal signaling. *Genes & Development*. 2002; 16:820–835. DOI: 10.1101/gad.963802 [PubMed: 11937490]
68. Cordenonsi M 1, Dupont S, Maretto S, Insinga A, Imbriano C, Piccolo S. Links between Tumor Suppressors p53 Is Required for TGF- β Gene Responses by Cooperating with Smads. *Cell*. 2003; 113:301–314. DOI: 10.1016/S0092-8674(03)00308-8 [PubMed: 12732139]
69. Ku M 1, Sokol SY, Wu J, Tussie-Luna MI, Roy AL, Hata A. Positive and Negative Regulation of the Transforming Growth Factor β /Activin Target Gene goosecoid by the TFII-I Family of

- Transcription Factors. *Mol Cell Biol.* 2005; 25:7144–7157. DOI: 10.1128/MCB.25.16.7144-7157.2005 [PubMed: 16055724]
70. Slagle CE 1, Aoki T, Burdine RD. Nodal-dependent mesendoderm specification requires the combinatorial activities of FoxH1 and Eomesodermin. *PLoS Genet.* 2011; 7:e1002072.doi: 10.1371/journal.pgen.1002072 [PubMed: 21637786]
 71. Yoon SJ 1, Wills AE, Chuong E, Gupta R, Baker JC. HEB and E2A function as SMAD/FOXH1 cofactors. *Genes & Development.* 2011; 25:1654–1661. DOI: 10.1101/gad.16800511 [PubMed: 21828274]
 72. Kofron M 1, Puck H, Standley H, Wylie C, Old R, Whitman M, et al. New roles for FoxH1 in patterning the early embryo. *Development.* 2004; 131:5065–5078. DOI: 10.1242/dev.01396 [PubMed: 15459100]
 73. Pogoda HM 1, Solnica-Krezel L, Driever W, Meyer D. The zebrafish forkhead transcription factor FoxH1/Fast1 is a modulator of Nodal signaling required for organizer formation. *Current Biology.* 2000; 10:1041–1049. DOI: 10.1016/S0960-9822(00)00669-2 [PubMed: 10996071]
 74. Sirotkin HI 1, Gates MA, Kelly PD, Schier AF, Talbot WS. Fast1 is required for the development of dorsal axial structures in zebrafish. *Curr Biol.* 2000; 10:1051–1054. DOI: 10.1016/S0960-9822(00)00679-5 [PubMed: 10996072]
 75. Hoodless PA 1, Pye M, Chazaud C, Labbé E, Attisano L, Rossant J, et al. FoxH1 (Fast) functions to specify the anterior primitive streak in the mouse. *Genes & Development.* 2001; 15:1257–1271. DOI: 10.1101/gad.881501 [PubMed: 11358869]
 76. Yamamoto M 1, Meno C, Sakai Y, Shiratori H. The transcription factor FoxH1 (FAST) mediates Nodal signaling during anterior-posterior patterning and node formation in the mouse. *Genes & Development.* 2001; 15:1242–1256. DOI: 10.1101/gad.883901 [PubMed: 11358868]
 77. Norris DP 1, Brennan J, Bikoff EK, Robertson EJ. The Foxh1-dependent autoregulatory enhancer controls the level of Nodal signals in the mouse embryo. *Development.* 2002; 129:3455–3468. [PubMed: 12091315]
 78. Kim SW 1, Yoon SJ, Chuong E, Oyolu C, Wills AE, Gupta R, et al. Chromatin and transcriptional signatures for Nodal signaling during endoderm formation in hESCs. *Developmental Biology.* 2011; 357:492–504. DOI: 10.1016/j.ydbio.2011.06.009 [PubMed: 21741376]
 79. Faure S 1, Lee MA, Keller T, ten Dijke P, Whitman M. Endogenous patterns of TGFbeta superfamily signaling during early Xenopus development. *Development.* 2000; 127:2917–2931. [PubMed: 10851136]
 80. Ku M 1, Melton DA. Xwnt-11: a maternally expressed Xenopus wnt gene. *Development.* 1993; 119:1161–1173. [PubMed: 8306880]
 81. Larabell CA 1, Torres M, Rowning BA, Yost C, Miller JR, Wu M, et al. Establishment of the dorso-ventral axis in Xenopus embryos is presaged by early asymmetries in beta-catenin that are modulated by the Wnt signaling pathway. *J Cell Biol.* 1997; 136:1123–1136. [PubMed: 9060476]
 82. Tao Q 1, Yokota C, Puck H, Kofron M, Birsoy B, Yan D, et al. Maternal wnt11 activates the canonical wnt signaling pathway required for axis formation in Xenopus embryos. *Cell.* 2005; 120:857–871. DOI: 10.1016/j.cell.2005.01.013 [PubMed: 15797385]
 83. Brannon M 1, Gomperts M, Sumoy L, Moon RT, Kimelman D. A beta-catenin/XTcf-3 complex binds to the siamois promoter to regulate dorsal axis specification in Xenopus. *Genes & Development.* 1997; 11:2359–2370. [PubMed: 9308964]
 84. Laurent MN 1, Blitz IL, Hashimoto C, Rothbächer U, Cho KW. The Xenopus homeobox gene twin mediates Wnt induction of goosecoid in establishment of Spemann's organizer. *Development.* 1997; 124:4905–4916. [PubMed: 9428427]
 85. Xanthos JB 1, Kofron M, Tao Q, Schaible K, Wylie C, Heasman J. The roles of three signaling pathways in the formation and function of the Spemann Organizer. *Development.* 2002; 129:4027–4043. [PubMed: 12163406]
 86. Kimelman D 1, Christian JL, Moon RT. Synergistic principles of development: overlapping patterning systems in Xenopus mesoderm induction. *Development.* 1992; 116:1–9. [PubMed: 1483380]

87. Watabe T 1, Kim S, Candia A, Rothbacher U, Hashimoto C, Inoue K, et al. Molecular mechanisms of Spemann's organizer formation: conserved growth factor synergy between *Xenopus* and mouse. *Genes & Development*. 1995; 9:3038–3050. [PubMed: 8543150]
88. Pereira LA 1, Wong MS, Lim SM, Stanley EG, Elefanty AG. The Mix family of homeobox genes —Key regulators of mesendoderm formation during vertebrate development. *Developmental Biology*. 2012; 367:163–177. DOI: 10.1016/j.ydbio.2012.04.033 [PubMed: 22580160]
89. Watanabe M 1, Yasuoka Y, Mawaribuchi S, Kureitani A, Ito M, Kondo M, et al. Conservatism and variability of gene expression profiles among homeologous transcription factors in *Xenopus laevis*. *Developmental Biology*. in press.
90. Latinkic BV 1, Umbhauer M, Neal KA, Lerchner W, Smith JC, Cunliffe V. The *Xenopus* Brachyury promoter is activated by FGF and low concentrations of activin and suppressed by high concentrations of activin and by paired-type homeodomain proteins. *Genes & Development*. 1997; 11:3265–3276. [PubMed: 9389657]
91. Lemaire P 1, Darras S, Caillol D, Kodjabachian L. A role for the vegetally expressed *Xenopus* gene Mix.1 in endoderm formation and in the restriction of mesoderm to the marginal zone. *Development*. 1998; 125:2371–2380. [PubMed: 9609820]
92. Colas A 1, Cartry J, Buisson I, Umbhauer M, Smith JC, Riou JF. Mix.1/2-dependent control of FGF availability during gastrulation is essential for pronephros development in *Xenopus*. *Developmental Biology*. 2008; 320:351–365. DOI: 10.1016/j.ydbio.2008.05.547 [PubMed: 18614163]
93. Artinger M 1, Blitz I, Inoue K, Tran U, Cho KW. Interaction of goosecoid and brachyury in *Xenopus* mesoderm patterning. *Mech Dev*. 1997; 65:187–196. [PubMed: 9256355]
94. Pereira LA 1, Wong MS, Mossman AK, Sourris K, Janes ME, Knezevic K, et al. *Pdgfra* and *Flk1* are direct target genes of *Mixl1* in differentiating embryonic stem cells. *Stem Cell Res*. 2012; 8:165–179. DOI: 10.1016/j.scr.2011.09.007 [PubMed: 22265737]
95. Kofron M 1, Wylie C, Heasman J. The role of Mixer in patterning the early *Xenopus* embryo. *Development*. 2004; 131:2431–2441. DOI: 10.1242/dev.01132 [PubMed: 15128672]
96. Tada M 1, Casey ES, Fairclough L, Smith JC. *Bix1*, a direct target of *Xenopus* T-box genes, causes formation of ventral mesoderm and endoderm. *Development*. 1998; 125:3997–4006. [PubMed: 9735361]
97. Ecochard V 1, Cayrol C, Rey S, Foulquier F, Caillol D, Lemaire P, et al. A novel *Xenopus* mix-like gene milk involved in the control of the endomesodermal fates. *Development*. 1998; 125:2577–2585. [PubMed: 9636073]
98. Okumura T 1, Matsumoto A, Tanimura T, Murakami R. An endoderm-specific GATA factor gene, *dGATAe*, is required for the terminal differentiation of the *Drosophila* endoderm. *Developmental Biology*. 2005; 278:576–586. DOI: 10.1016/j.ydbio.2004.11.021 [PubMed: 15680371]
99. Okumura T 1, Tajiri R, Kojima T, Saigo K, Murakami R. GATAe-dependent and -independent expressions of genes in the differentiated endodermal midgut of *Drosophila*. *Gene Expression Patterns*. 2007; 7:178–186. DOI: 10.1016/j.modgep.2006.07.001 [PubMed: 16914392]
100. Sommermann EM 1, Strohmaier KR, Maduro MF, Rothman JH. Endoderm development in *Caenorhabditis elegans*: the synergistic action of ELT-2 and -7 mediates the specification→differentiation transition. *Developmental Biology*. 2010; 347:154–166. DOI: 10.1016/j.ydbio.2010.08.020 [PubMed: 20807527]
101. Boeck ME 1, Boyle T, Bao Z, Murray J, Mericle B, Waterston R. Specific roles for the GATA transcription factors end-1 and end-3 during *C. elegans* E-lineage development. *Developmental Biology*. 2011; 358:345–355. DOI: 10.1016/j.ydbio.2011.08.002 [PubMed: 21854766]
102. Maduro MF 1. Gut development in *C. elegans*. *Seminars in Cell and Developmental Biology*. in press.
103. Peter IS 1, Davidson EH. The endoderm gene regulatory network in sea urchin embryos up to mid-blastula stage. *Developmental Biology*. 2010; 340:188–199. DOI: 10.1016/j.ydbio.2009.10.037 [PubMed: 19895806]
104. Fujikura J 1, Yamato E, Yonemura S, Hosoda K, Masui S, Nakao K, et al. Differentiation of embryonic stem cells is induced by GATA factors. *Genes & Development*. 2002; 16:784–789. doi:10.1101 /gad.968802. [PubMed: 11937486]

105. Shimosato D 1, Shiki M, Niwa H. Extra-embryonic endoderm cells derived from ES cells induced by GATA factors acquire the character of XEN cells. *BMC Dev Biol.* 2007; 7:80.doi: 10.1186/1471-213X-7-80 [PubMed: 17605826]
106. Weber H 1, Symes CE, Walmsley ME, Rodaway AR, Patient RK. A role for GATA5 in *Xenopus* endoderm specification. *Development.* 2000; 127:4345–4360. [PubMed: 11003835]
107. Afouda BA 1, Ciau-Uitz A, Patient R. GATA4, 5 and 6 mediate TGFbeta maintenance of endodermal gene expression in *Xenopus* embryos. *Development.* 2005; 132:763–774. DOI: 10.1242/dev.01647 [PubMed: 15659482]
108. Lou X 1, Deshwar AR, Crump JG, Scott IC. Smarcd3b and Gata5 promote a cardiac progenitor fate in the zebrafish embryo. *Development.* 2011; 138:3113–3123. DOI: 10.1242/dev.064279 [PubMed: 21715426]
109. Tseng WF 1, Jang TH, Huang CB, Yuh CH. An evolutionary conserved kernel of gata5, gata6, otx2 and prdm1a operates in the formation of endoderm in zebrafish. *Developmental Biology.* 2011; 357:541–557. DOI: 10.1016/j.ydbio.2011.06.040 [PubMed: 21756893]
110. Reiter JF 1, Kikuchi Y, Stainier DY. Multiple roles for Gata5 in zebrafish endoderm formation. *Development.* 2001; 128:125–135. [PubMed: 11092818]
111. Turbendian HK 1, Gordillo M, Tsai SY, Lu J, Kang G, Liu TC, et al. GATA factors efficiently direct cardiac fate from embryonic stem cells. *Development.* 2013; 140:1639–1644. DOI: 10.1242/dev.093260 [PubMed: 23487308]
112. Watt AJ 1, Zhao R, Li J, Duncan SA. Development of the mammalian liver and ventral pancreas is dependent on GATA4. *BMC Dev Biol.* 2007; 7:37.doi: 10.1186/1471-213X-7-37 [PubMed: 17451603]
113. Rojas A 1, Schachterle W, Xu SM, Black BL. An endoderm-specific transcriptional enhancer from the mouse Gata4 gene requires GATA and homeodomain protein-binding sites for function in vivo. *Dev Dyn.* 2009; 238:2588–2598. DOI: 10.1002/dvdy.22091 [PubMed: 19777593]
114. Rojas A 1, De Val S, Heidt AB, Xu SM, Bristow J, Black BL. Gata4 expression in lateral mesoderm is downstream of BMP4 and is activated directly by Forkhead and GATA transcription factors through a distal enhancer element. *Development.* 2005; 132:3405–3417. DOI: 10.1242/dev.01913 [PubMed: 15987774]
115. Hudson C 1, Clements D, Friday RV, Stott D, Woodland HR. Xsox17 α and - β Mediate Endoderm Formation in *Xenopus*. *Cell.* 1997; 91:397–405. DOI: 10.1016/S0092-8674(00)80423-7 [PubMed: 9363948]
116. Alexander J 1, Stainier D. A molecular pathway leading to endoderm formation in zebrafish. *Current Biology.* 1999; doi: 10.1016/S0960-9822(00)80016-0
117. Kanai-Azuma M 1, Kanai Y, Gad JM, Tajima Y, Taya C, Kurohmaru M, et al. Depletion of definitive gut endoderm in Sox17-null mutant mice. *Development.* 2002; 129:2367–2379. [PubMed: 11973269]
118. Séguin CA 1, Draper JS, Nagy A, Rossant J. Establishment of endoderm progenitors by SOX transcription factor expression in human embryonic stem cells. *Cell Stem Cell.* 2008; 3:182–195. DOI: 10.1016/j.stem.2008.06.018 [PubMed: 18682240]
119. Sinner D 1, Rankin S, Lee M, Zorn AM. Sox17 and beta-catenin cooperate to regulate the transcription of endodermal genes. *Development.* 2004; 131:3069–3080. DOI: 10.1242/dev.01176 [PubMed: 15163629]
120. Dickinson K 1, Leonard J, Baker JC. Genomic profiling of Mixer and Sox17 β targets during *Xenopus* endoderm development. *Dev Dyn.* 2006; 235:368–381. DOI: 10.1002/dvdy.20636 [PubMed: 16278889]
121. Howard L 1, Rex M, Clements D, Woodland HR. Regulation of the *Xenopus* Xsox17 α (1) promoter by co-operating VegT and Sox17 sites. *Developmental Biology.* 2007; 310:402–415. DOI: 10.1016/j.ydbio.2007.07.028 [PubMed: 17719026]
122. Shimoda M 1, Kanai-Azuma M, Hara K, Miyazaki S, Kanai Y, Monden M, et al. Sox17 plays a substantial role in late-stage differentiation of the extraembryonic endoderm in vitro. *Journal of Cell Science.* 2007; 120:3859–3869. DOI: 10.1242/jcs.007856 [PubMed: 17940068]

123. Friedman JR 1, Kaestner KH. The Foxa family of transcription factors in development and metabolism. *Cell Mol Life Sci.* 2006; 63:2317–2328. DOI: 10.1007/s00018-006-6095-6 [PubMed: 16909212]
124. de-Leon SBT 1. The conserved role and divergent regulation of foxa, a pan-eumetazoan developmental regulatory gene. *Developmental Biology.* 2011; 357:21–26. DOI: 10.1016/j.ydbio.2010.11.027 [PubMed: 21130759]
125. Weinstein DC 1, Ruiz A, Altaba i, Chen WS, Hoodless P, Prezioso VR, Jessell TM, et al. The winged-helix transcription factor HNF-3 beta is required for notochord development in the mouse embryo. *Cell.* 1994; 78:575–588. [PubMed: 8069910]
126. Gualdi R 1, Bossard P, Zheng M, Hamada Y, Coleman JR, Zaret KS. Hepatic specification of the gut endoderm in vitro: cell signaling and transcriptional control. *Genes & Development.* 1996; 10:1670–1682. [PubMed: 8682297]
127. Lee CS 1, Friedman JR, Fulmer JT, Kaestner KH. The initiation of liver development is dependent on Foxa transcription factors. *Nature.* 2005; 435:944–947. DOI: 10.1038/nature03649 [PubMed: 15959514]
128. Cirillo LA 1, Lin FR, Cuesta I, Friedman D, Jarnik M, Zaret KS. Opening of compacted chromatin by early developmental transcription factors HNF3 (FoxA) and GATA-4. *Mol Cell.* 2002; 9:279–289. DOI: 10.1016/S1097-2765(02)00459-8 [PubMed: 11864602]
129. Iwafuchi-Doi M 1, Zaret KS. Cell fate control by pioneer transcription factors. *Development.* 2016; 143:1833–1837. DOI: 10.1242/dev.133900 [PubMed: 27246709]
130. Suri C 1, Harembaki T, Weinstein DC. Inhibition of mesodermal fate by *Xenopus* HNF3 β /FoxA2. *Developmental Biology.* 2004; 265:90–104. DOI: 10.1016/j.ydbio.2003.09.017 [PubMed: 14697355]
131. Oliveri P 1, Walton KD, Davidson EH, Mc Clay DR. Repression of mesodermal fate by foxa, a key endoderm regulator of the sea urchin embryo. *Development.* 2006; 133:4173–4181. DOI: 10.1242/dev.02577 [PubMed: 17038513]
132. Murgan S 1, Castro Colabianchi AM, Monti RJ, Boyadjian López LE, Aguirre CE, Stivala EG, et al. FoxA4 favours notochord formation by inhibiting contiguous mesodermal fates and restricts anterior neural development in *Xenopus* embryos. *PLoS ONE.* 2014; 9:e110559.doi: 10.1371/journal.pone.0110559 [PubMed: 25343614]
133. Friedle H 1, Rastegar S, Paul H, Kaufmann E, Knochel W. Xvent-1 mediates BMP-4-induced suppression of the dorsal-lip-specific early response gene XFD-1' in *Xenopus* embryos. *Embo J.* 1998; 17:2298–2307. DOI: 10.1093/emboj/17.8.2298 [PubMed: 9545242]
134. Alon U 1. Network motifs: theory and experimental approaches. *Nat Rev Genet.* 2007; 8:450–461. DOI: 10.1038/nrg2102 [PubMed: 17510665]
135. Luxardi G 1, Marchal L, Thomé V, Kodjabachian L. Distinct *Xenopus* Nodal ligands sequentially induce mesendoderm and control gastrulation movements in parallel to the Wnt/PCP pathway. *Development.* 2010; 137:417–426. DOI: 10.1242/dev.039735 [PubMed: 20056679]
136. Henningfeld KA 1, Friedle H, Rastegar S, Knöchel W. Autoregulation of Xvent-2B; direct interaction and functional cooperation of Xvent-2 and Smad1. *Journal of Biological Chemistry.* 2002; 277:2097–2103. DOI: 10.1074/jbc.M108524200 [PubMed: 11704665]
137. Sander V 1, Reversade B, De Robertis EM. The opposing homeobox genes Goosecoid and Vent1/2 self-regulate *Xenopus* patterning. *Embo J.* 2007; 26:2955–2965. DOI: 10.1038/sj.emboj.7601705 [PubMed: 17525737]
138. Schohl A 1, Fagotto F. A role for maternal β -catenin in early mesoderm induction in *Xenopus*. *Embo J.* 2003; 22:3303–3313. DOI: 10.1093/emboj/cdg328 [PubMed: 12839992]
139. Onichtchouk D 1, Gawantka V, Dosch R, Delius H, Hirschfeld K, Blumenstock C, et al. The Xvent-2 homeobox gene is part of the BMP-4 signalling pathway controlling [correction of controlling] dorsoventral patterning of *Xenopus* mesoderm. *Development.* 1996; 122:3045–3053. [PubMed: 8898218]
140. Trindade M 1, Tada M, Smith JC. DNA-binding specificity and embryological function of Xom (Xvent-2). *Developmental Biology.* 1999; 216:442–456. DOI: 10.1006/dbio.1999.9507 [PubMed: 10642784]

141. Middleton AM, King JR, Loose M. Bistability in a model of mesoderm and anterior mesendoderm specification in *Xenopus laevis*. *J Theor Biol.* 2009; 260:41–55. DOI: 10.1016/j.jtbi.2009.05.016 [PubMed: 19490918]
142. Saka Y, Smith JC. A mechanism for the sharp transition of morphogen gradient interpretation in *Xenopus*. *BMC Dev Biol.* 2007; 7:47. doi:10.1186/1471-213X-7-47. [PubMed: 17506890]
143. Zorn AM 1, Butler K, Gurdon JB. Anterior Endomesoderm Specification in *Xenopus* by Wnt/ β -catenin and TGF- β Signalling Pathways. *Developmental Biology.* 1999; 209:282–297. DOI: 10.1006/dbio.1999.9257 [PubMed: 10328921]
144. Schuler-Metz A 1, Knochel S, Kaufmann E, Knochel W. The homeodomain transcription factor Xvent-2 mediates autocatalytic regulation of BMP-4 expression in *Xenopus* embryos. *Journal of Biological Chemistry.* 2000; 275:34365–34374. DOI: 10.1074/jbc.M003915200 [PubMed: 10938274]
145. Rex M 1, Hilton E, Old R. Multiple interactions between maternally-activated signalling pathways control *Xenopus* nodal-related genes. *Int J Dev Biol.* 2002; 46:217–226. [PubMed: 11934150]
146. Cha SW 1, Tadjuidje E, Tao Q, Wylie C, Heasman J. Wnt5a and Wnt11 interact in a maternal Dkk1-regulated fashion to activate both canonical and non-canonical signaling in *Xenopus* axis formation. *Development.* 2008; 135:3719–3729. DOI: 10.1242/dev.029025 [PubMed: 18927149]
147. Blythe SA 1, Cha SW, Tadjuidje E, Heasman J, Klein PS. β -Catenin primes organizer gene expression by recruiting a histone H3 arginine 8 methyltransferase. *Prmt2*, *Developmental Cell.* 2010; 19:220–231. DOI: 10.1016/j.devcel.2010.07.007 [PubMed: 20708585]
148. Mangan S 1, Alon U. Structure and function of the feed-forward loop network motif. *Proc Natl Acad Sci USA.* 2003; 100:11980–11985. DOI: 10.1073/pnas.2133841100 [PubMed: 14530388]
149. Galán-Vázquez E 1, Luna B, Martínez-Antonio A. The Regulatory Network of *Pseudomonas aeruginosa*. *Microb Inform Exp.* 2011; 1:3. doi: 10.1186/2042-5783-1-3 [PubMed: 22587778]
150. Kim MS 1, Kim JR, Cho KH. Dynamic network rewiring determines temporal regulatory functions in *Drosophila melanogaster* development processes. *Bioessays.* 2010; 32:505–513. DOI: 10.1002/bies.200900169 [PubMed: 20486137]
151. Kim MS 1, Kim JR, Kim D, Lander AD, Cho KH. Spatiotemporal network motif reveals the biological traits of developmental gene regulatory networks in *Drosophila melanogaster*. *BMC Syst Biol.* 2012; 6:31. doi: 10.1186/1752-0509-6-31 [PubMed: 22548745]
152. Latinkic BV 1, Smith JC. Goosecoid and mix.1 repress Brachyury expression and are required for head formation in *Xenopus*. *Development.* 1999; 126:1769–1779. [PubMed: 10079237]
153. Carron C 1, Bourdelas A, Li HY, Boucaut JC, Shi DL. Antagonistic interaction between IGF and Wnt/JNK signaling in convergent extension in *Xenopus* embryo. *Mech Dev.* 2005; 122:1234–1247. DOI: 10.1016/j.mod.2005.06.007 [PubMed: 16169711]
154. Zhang C 1, Basta T, Jensen ED, Klymkowsky MW. The β -catenin/VegT-regulated early zygotic gene *Xnr5* is a direct target of SOX3 regulation. *Development.* 2003; 130:5609–5624. DOI: 10.1242/dev.00798 [PubMed: 14522872]
155. Faunes F 1, Almonacid LI, Melo F, Larraín J. Characterization of small RNAs in *X. tropicalis* gastrulae. *Genesis.* 2012; 50:572–583. DOI: 10.1002/dvg.22027 [PubMed: 22566284]
156. Harding JL 1, Horswell S, Heliot C, Armisen J, Zimmerman LB, Luscombe NM, et al. Small RNA profiling of *Xenopus* embryos reveals novel miRNAs and a new class of small RNAs derived from intronic transposable elements. *Genome Res.* 2014; 24:96–106. DOI: 10.1101/gr.144469.112 [PubMed: 24065776]
157. Ahmed A 1, Ward NJ, Moxon S, Lopez-Gomollon S, Viaut C, Tomlinson ML, et al. A Database of microRNA Expression Patterns in *Xenopus laevis*. *PLoS ONE.* 2015; 10:e0138313. doi: 10.1371/journal.pone.0138313 [PubMed: 26506012]
158. Liu C 1, Lou CH, Shah V, Ritter R, Talley J, Soibam B, et al. Identification of microRNAs and microRNA targets in *Xenopus* gastrulae: The role of miR-26 in the regulation of Smad1. *Developmental Biology.* 2016; 409:26–38. DOI: 10.1016/j.ydbio.2015.11.005 [PubMed: 26548531]

159. Shah VV 1, Soibam B, Ritter RA, Benham A, Oomen J, Sater AK. MicroRNAs and ectodermal specification I. Identification of miRs and miR-targeted mRNAs in early anterior neural and epidermal ectoderm. *Developmental Biology*. in press.
160. Necsulea A 1, Soumillon M, Warnefors M, Liechti A, Daish T, Zeller U, et al. The evolution of lncRNA repertoires and expression patterns in tetrapods. *Nature*. 2014; 505:635–640. DOI: 10.1038/nature12943 [PubMed: 24463510]
161. Forouzmand E 1, Owens ND, Blitz IL, Paraiso KD, Khokha MK, Gilchrist MJ, et al. Developmentally regulated long non-coding RNAs in *Xenopus tropicalis*. *Developmental Biology*. in press.
162. Rosa A 1, Spagnoli FM, Brivanlou AH. The miR-430/427/302 family controls mesendodermal fate specification via species-specific target selection. *Developmental Cell*. 2009; 16:517–527. DOI: 10.1016/j.devcel.2009.02.007 [PubMed: 19386261]
163. Paulsen M 1, Legewie S, Eils R. Negative feedback in the bone morphogenetic protein 4 (BMP4) synexpression group governs its dynamic signaling range and canalizes development. 2011; doi: 10.1074/jbc.M113.509331
164. Heasman J 1, Kofron M, Wylie C. β Catenin Signaling Activity Dissected in the Early *Xenopus* Embryo: A Novel Antisense Approach. *Developmental Biology*. 2000; 222:124–134. DOI: 10.1006/dbio.2000.9720 [PubMed: 10885751]
165. Hilton E 1, Rex M, Old R. VegT activation of the early zygotic gene *Xnr5* requires lifting of Tcf-mediated repression in the *Xenopus* blastula. *Mech Dev*. 2003; 120:1127–1138. DOI: 10.1016/j.mod.2003.08.004 [PubMed: 14568102]
166. Rankin SA 1, Kormish J, Kofron M, Jegga A, Zorn AM. A gene regulatory network controlling *hhx* transcription in the anterior endoderm of the organizer. *Developmental Biology*. 2011; 351:297–310. DOI: 10.1016/j.ydbio.2010.11.037 [PubMed: 21215263]
167. Houston DW 1, Kofron M, Resnik E, Langland R, Destree O, Wylie C, et al. Repression of organizer genes in dorsal and ventral *Xenopus* cells mediated by maternal XTcf3. *Development*. 2002; 129:4015–4025. [PubMed: 12163405]
168. White RJ 1, Sun BI, Sive HL, Smith JC. Direct and indirect regulation of *derrière*, a *Xenopus* mesoderm-inducing factor, by VegT. *Development*. 2002; 129:4867–4876. [PubMed: 12361977]
169. Clements D 1, Cameleyre I, Woodland HR. Redundant early and overlapping larval roles of *Xsox17* subgroup genes in *Xenopus* endoderm development. *Mech Dev*. 2003; 120:337–348. DOI: 10.1016/S0925-4773(02)00450-1 [PubMed: 12591603]
170. Taverner NV 1, Kofron M, Shin Y, Kabitschke C, Gilchrist MJ, Wylie C, et al. Microarray-based identification of VegT targets in *Xenopus*. *Mech Dev*. 2005; 122:333–354. DOI: 10.1016/j.mod.2004.10.010 [PubMed: 15763211]
171. Zhang C 1, Basta T, Fawcett SR, Klymkowsky MW. SOX7 is an immediate-early target of VegT and regulates Nodal-related gene expression in *Xenopus*. *Developmental Biology*. 2005; 278:526–541. DOI: 10.1016/j.ydbio.2004.11.008 [PubMed: 15680368]
172. Cao Y 1, Siegel D, Donow C, Knöchel S, Yuan L, Knöchel W. POU- V factors antagonize maternal VegT activity and β -Catenin signaling in *Xenopus* embryos. *Embo J*. 2007; 26:2942–2954. DOI: 10.1038/sj.emboj.7601736 [PubMed: 17541407]
173. Labbé E 1, Silvestri C, Hoodless PA, Wrana JL, Attisano L. Smad2 and Smad3 positively and negatively regulate TGF beta-dependent transcription through the forkhead DNA-binding protein FAST2. *Mol Cell*. 1998; 2:109–120. [PubMed: 9702197]
174. Yeo CY 1, Chen X, Whitman M. The role of FAST-1 and Smads in transcriptional regulation by activin during early *Xenopus* embryogenesis. *Journal of Biological Chemistry*. 1999; 274:26584–26590. [PubMed: 10473623]
175. Shiratori H 1, Sakuma R, Watanabe M, Hashiguchi H, Mochida K, Sakai Y, et al. Two-Step Regulation of Left–Right Asymmetric Expression of *Pitx2*. *Mol Cell*. 2001; 7:137–149. DOI: 10.1016/S1097-2765(01)00162-9 [PubMed: 11172719]
176. Sinner D 1, Kirilenko P, Rankin S, Wei E, Howard L, Kofron M, et al. Global analysis of the transcriptional network controlling *Xenopus* endoderm formation. *Development*. 2006; 133:1955–1966. DOI: 10.1242/dev.02358 [PubMed: 16651540]

177. Reid CD 1, Zhang Y, Sheets MD, Kessler DS. Transcriptional integration of Wnt and Nodal pathways in establishment of the Spemann organizer. *Developmental Biology*. 2012; 368:231–241. DOI: 10.1016/j.ydbio.2012.05.018 [PubMed: 22627292]
178. Gawantka V 1, Delius H, Hirschfeld K, Blumenstock C, Niehrs C. Antagonizing the Spemann organizer: role of the homeobox gene Xvent-1. *Embo J*. 1995; 14:6268–6279. [PubMed: 8557046]
179. Lagna G 1, Hata A, Hemmati-Brivanlou A, Massagué J. Partnership between DPC4 and SMAD proteins in TGF- β signalling pathways. *Nature*. 1996; 383:832–836. DOI: 10.1038/383832a0 [PubMed: 8893010]
180. Candia AF 1, Watabe T, Hawley SH, Onichtchouk D, Zhang Y, Derynck R, et al. Cellular interpretation of multiple TGF-beta signals: intracellular antagonism between activin/BVg1 and BMP-2/4 signaling mediated by Smads. *Development*. 1997; 124:4467–4480. DOI: 10.1101/gad.10.15.1880 [PubMed: 9409665]
181. Ladher R 1, Mohun TJ, Smith JC, Snape AM. Xom: a Xenopus homeobox gene that mediates the early effects of BMP-4. *Development*. 1996; 122:2385–2394. [PubMed: 8756284]
182. Zhang C 1, Evans T. BMP-like signals are required after the midblastula transition for blood cell development. *Dev Genet*. 1996; 18:267–278. DOI: 10.1002/(SICI)1520-6408(1996)18:3<267::AID-DVG7>3.0.CO;2-7 [PubMed: 8631160]
183. Henningfeld KA 1, Rastegar S, Adler G, Knochel W. Smad1 and Smad4 are components of the bone morphogenetic protein-4 (BMP-4)-induced transcription complex of the Xvent-2B promoter. *Journal of Biological Chemistry*. 2000; 275:21827–21835. DOI: 10.1074/jbc.M000978200 [PubMed: 10791953]
184. Lee HS 1, Park MJ, Lee SY, Hwang YS, Lee H, Roh DH, et al. Transcriptional regulation of Xbr-1a/Xvent-2 homeobox gene: analysis of its promoter region. *Biochem Biophys Res Commun*. 2002; 298:815–823. DOI: 10.1016/S0006-291X(02)02570-6 [PubMed: 12419328]
185. Miyanaga Y 1, Torregroza I, Evans T. A maternal Smad protein regulates early embryonic apoptosis in *Xenopus laevis*. *Mol Cell Biol*. 2002; 22:1317–1328. DOI: 10.1128/mcb.22.5.1317-1328.2002 [PubMed: 11839799]
186. Peiffer DA 1, von Bubnoff A, Shin Y, Kitayama A, Mochii M, Ueno N, et al. A Xenopus DNA microarray approach to identify novel direct BMP target genes involved in early embryonic development. *Dev Dyn*. 2005; 232:445–456. DOI: 10.1002/dvdy.20230 [PubMed: 15614776]
187. Torregroza I 1, Evans T. Tid1 is a Smad-binding protein that can modulate Smad7 activity in developing embryos. *Biochem J*. 2006; 393:311–320. DOI: 10.1042/BJ20050785 [PubMed: 16156721]
188. Wills A 1, Dickinson K, Khokha M, Baker JC. Bmp signaling is necessary and sufficient for ventrolateral endoderm specification in *Xenopus*. *Dev Dyn*. 2008; 237:2177–2186. DOI: 10.1002/dvdy.21631 [PubMed: 18651654]
189. Kessler DS 1. Siamois is required for formation of Spemann's organizer. *Proc Natl Acad Sci USA*. 1997; 94:13017–13022. DOI: 10.1073/pnas.94.24.13017 [PubMed: 9371792]
190. Yamamoto S 1, Hikasa H, Ono H, Taira M. Molecular link in the sequential induction of the Spemann organizer: direct activation of the cerberus gene by Xlim-1, Xotx2, Mix.1, and Siamois, immediately downstream from Nodal and Wnt signaling. *Developmental Biology*. 2003; 257:190–204. DOI: 10.1016/S0012-1606(03)00034-4 [PubMed: 12710967]
191. Bae S 1, Reid CD, Kessler DS. Siamois and Twin are redundant and essential in formation of the Spemann organizer. *Developmental Biology*. 2011; 352:367–381. DOI: 10.1016/j.ydbio.2011.01.034 [PubMed: 21295564]
192. Klein SL 1, Moody SA. Early neural ectodermal genes are activated by Siamois and Twin during blastula stages. *Genesis*. 2015; 53:308–320. DOI: 10.1002/dvg.22854 [PubMed: 25892704]
193. Mead PE 1, Brivanlou IH, Kelley CM, Zon LI. BMP-4-responsive regulation of dorsal–ventral patterning by the homeobox protein Mix.1. *Nature*. 1996; 382:357–360. DOI: 10.1038/382357a0 [PubMed: 8684465]
194. Hayata T 1, Tanegashima K, Takahashi S. Overexpression of the secreted factor Mig30 expressed in the Spemann organizer impairs morphogenetic movements during *Xenopus* gastrulation. *Mech Dev*. 2002; 112:37–51. DOI: 10.1016/S0925-4773(01)00638-4 [PubMed: 11850177]

195. Randall RA 1, Germain S, Inman GJ. Different Smad2 partners bind a common hydrophobic pocket in Smad2 via a defined proline-rich motif. *The EMBO* 2002
196. Ya J, Kessler DS 1. Goosecoid promotes head organizer activity by direct repression of Xwnt8 in Spemann's organizer. 2001:1–13.
197. Mochizuki T 1, Karavanov AA, Curtiss PE, Ault KT, Sugimoto N, Watabe T, et al. Xlim-1 and LIM Domain Binding Protein 1 Cooperate with Various Transcription Factors in the Regulation of the goosecoid Promoter. *Developmental Biology*. 2000; 224:470–485. DOI: 10.1006/dbio.2000.9778 [PubMed: 10926781]
198. Brickman JM 1, Jones CM, Clements M, Smith JC, Beddington RS. Hex is a transcriptional repressor that contributes to anterior identity and suppresses Spemann organiser function. *Development*. 2000; 127:2303–2315. [PubMed: 10804173]
199. Smithers LE 1, Jones CM. Xhex-expressing endodermal tissues are essential for anterior patterning in *Xenopus*. *Mech Dev*. 2002; 119:191–200. DOI: 10.1016/S0925-4773(02)00361-1 [PubMed: 12464432]
200. Zamparini AL 1. Hex acts with -catenin to regulate anteroposterior patterning via a Groucho-related co-repressor and Nodal. *Development*. 2006; 133:3709–3722. DOI: 10.1242/dev.02516 [PubMed: 16936074]
201. Afouda BA 1, Martin J, Liu F, Ciau-Uitz A, Patient R, Hoppler S. GATA transcription factors integrate Wnt signalling during heartdevelopment. *Development*. 2008; 135:3185–3190. DOI: 10.1242/dev.026443 [PubMed: 18715946]
202. Friedle H 1, Knöchel W. Cooperative interaction of Xvent-2 and GATA-2 in the activation of the ventral homeobox gene Xvent-1B. *Journal of Biological Chemistry*. 2002; 277:23872–23881. DOI: 10.1074/jbc.M201831200 [PubMed: 11964398]
203. Drews C 1, Senkel S, Ryffel GU. The nephrogenic potential of the transcription factors *osr1*, *osr2*, *hnf1b*, *lhx1* and *pax8* assessed in *Xenopus* animal caps. *BMC Dev Biol*. 2011; 11:5.doi: 10.1186/1471-213X-11-5 [PubMed: 21281489]
204. Onichtchouk D 1, Glinka A, Niehrs C. Requirement for Xvent-1 and Xvent-2 gene function in dorsoventral patterning of *Xenopus* mesoderm. *Development*. 1998; 125:1447–1456. [PubMed: 9502725]
205. Schuler-Metz A 1, Knochel S, Kaufmann E, Knochel W. The homeodomain transcription factor Xvent-2 mediates autocatalytic regulation of BMP-4 expression in *Xenopus* embryos. *Journal of Biological Chemistry*. 2000; 275:34365–34374. DOI: 10.1074/jbc.M003915200 [PubMed: 10938274]

Highlights

- Comprehensive *Xenopus* mesendoderm gene regulatory network (GRN)
- Conserved regulatory connections in frog, fish, mouse, and human endoderm are revealed
- Coherent feedforward loops are the most common network motif in the GRN




Figure 1. Xenopus mesendoderm gene regulatory network from fertilization through early-gastrula

The network is comprised of 23 transcription factors and 12 growth factors. Maternal proteins are represented as diamonds, and signaling ligands as circles. Connections are drawn from the transcriptional regulator to the cis-regulatory region of the target gene. Direct connections are indicated as solid lines, and putative connections as dashed lines. Activating connections are represented by arrowheads, and repressive connections as a flat line. Connections from secreted ligands pass through a chevron, and are mediated by their respective intracellular transcription factors (e.g. Smad2/3, β -catenin). Approximate spatiotemporal information is provided from top to bottom (egg through early-gastrula) and from right to left (dorsal to ventral), with some exceptions (e.g. *xbra/t*). The activation time of zygotic *sox7* (boxed) is unknown. All direct connections are annotated for TF binding (blue diamond), reporter assay (pink diamond), and TF binding plus functional validation (maroon diamond). For additional connection details, including experimental evidence and references, see Table 1 and Table S2. Zygotic genome activation, ZGA.




Figure 2. Network motifs found in the *Xenopus* mesendoderm GRN

(A) Autoregulatory loop, for example by nodal signaling (B) Positive feedback loop, for example between ventral genes Nodal and Wnt signaling (C) Coherent and incoherent feedforward loops and their regulatory structure (D) Type I feedforward loop, which appears to be the most common type of feedforward loop, frequently appears in the structure such that X is a maternal factor, Y is an early zygotic gene, and Z is either an early or late zygotic gene.

Table 1
Summary of direct and putative connections between network transcription factors

See also Table S2 for additional connection details. Putative direct targets are denoted as [P].

ctnnb1/β-catenin (Wnt)	cer1, eomes, foxa1, foxa2, gsc, hhhex, hnf1b, lhx1, nodal1, nodal2, nodal, nodal5, nodal6, otx2, sebox, sia1, sia2, sox17a, t, vegt, ventx1, wnt8a	[23,24,82-85,119,138,145,147,164-166]
vegt	cer1, gdf3, gsc, mix1 [P], mixer [P], nodal1, nodal [P], nodal5, sia1, snai1 [P], sox17a, sox17b1 [P], sox7 [P], ventx1 [P]	[33,35,36,38,41,85,121,165,167-172]
foxx1	cer1, gata2, gdf3, gsc, hhhex, lhx1, mix1, nodal1, nodal2, otx1, otx2, pitx2, sebox, wnt8a	[26,62,63,66,72,166,173-175]
sox7	nodal [P], nodal5 [P], nodal6 [P]	[49,171]
smad2/3 (Nodal)	bix1, bix2, cer1, eomes, foxa4, gata4, gata6, gsc, hhhex, hnf1b, lhx1, mix1, mixer, nodal1, nodal2, osr2, otx2, pitx2, sia1, snai1, sox17b, t, vegt, wnt8a, zic2	[26,27,62,63,72,87,90,166,176,177]
smad1 (BMP)	gata2, ventx1, ventx2	[61,178-188]
sia1	cer1, gsc, hhhex, zic2 [P]	[33,84,91,92,152,189,190]
sia2	gsc, hhhex, zic2 [P]	[84,166,191,192]
mix1	cer1, gsc, t	[33,90-92,152,190,193]
mixer	gsc	[65,194,195]
bix2 (laevis only)	gsc	[65,97]
gsc	gsc, otx2 [P], pitx2 [P], t, ventx1, ventx2, wnt8a	[18,28,90,137,196]
otx2	cer1, gsc, hhhex, t	[32,90,153,166,197]
lhx1	cer1, gsc, hhhex	[166,190,197]
hhhex	gsc, nodal1 [P], nodal2 [P]	[198-200]
gata4	hnf1b [P]	[106,107,201]
gata5	hnf1b [P], sox17a [P]	[106,107]
gata6	hnf1b [P], sox17a [P]	[107,201]
gata2	ventx1	[202]
sox17b	foxa1 [P], foxa2 [P], hnf1b [P], sox17a	[119-121,176]
hnf1b	lhx1	[203]
t	bix1	[96]
ventx1	foxa4	[133]
ventx2	bmp4, gsc, hhhex, ventx1, ventx2	[136,140,166,202,204,205]