

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

THE CRYSTAL STRUCTURE OF XENON (II) FLUORIDE FLUOROSULFATE, FXeO<sub>2</sub>S<sub>2</sub>F

### Permalink

<https://escholarship.org/uc/item/2dm4p553>

### Authors

Bartlett, Neil.  
Wechsberg, M.  
Jones, G.R.  
et al.

### Publication Date

1972

Submitted to Inorganic Chemistry

LBL-170  
Preprint c.1

THE CRYSTAL STRUCTURE OF XENON (II) FLUORIDE  
FLUROSULFATE,  $\text{FXeOSO}_2\text{F}$

Neil Bartlett, M. Wechsberg, G. R. Jones  
and R. D. Burbank

January 1972

AEC Contract No. W-7405-eng-48



**For Reference**

Not to be taken from this room

LBL-170

c.1

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

A Contribution from The Lawrence Berkeley Laboratory and the Department of Chemistry, University of California, Berkeley, California 94720; The Chemistry Department, Princeton University, Princeton, New Jersey 08540; and Bell Telephone Laboratories, Murray Hill, New Jersey 07974.

THE CRYSTAL STRUCTURE OF XENON(II) FLUORIDE

FLUOROSULFATE,  $\text{FXeOSO}_2\text{F}$

by

Neil Bartlett \*

Inorganic Materials Research Division, Lawrence Berkeley Laboratory;  
and Department of Chemistry, University of California  
Berkeley, California 94720

and

M. Wechsberg

Chemistry Department, Princeton University  
Princeton, New Jersey 08540

and

G. R. Jones <sup>1</sup> and R. D. Burbank

Bell Telephone Laboratories, Incorporated  
Murray Hill, New Jersey 07971

\* To whom correspondence should be addressed.

ABSTRACT

The crystal structure of xenon(II) fluoride fluorosulfate has been determined from three-dimensional x-ray data. The compound crystallizes in the orthorhombic system, with eight molecules in a unit cell of dimensions  $a = 9.88(1)$ ,  $b = 10.00(1)$  and  $c = 10.13(1)$  Å. The space group is  $Pbca$  and refinement has proceeded satisfactorily in this space group, with a final conventional R factor of 0.045 for 838 nonzero reflections. The structure analysis has established the existence of discrete  $\text{FXeOSO}_2\text{F}$  molecules. The xenon atom is approximately linearly coordinated by an oxygen atom of the fluorosulfate group and a fluorine atom. The angle F-Xe-O is  $177.4(3)^\circ$ , and the interatomic distances are Xe-F =  $1.940(8)$  and Xe-O =  $2.155(8)$  Å. The fluorosulfate group is similar to that observed in the alkali salts, with the difference that, in this structure, the group is distorted as a consequence of one oxygen atom being linked to the xenon atom. This oxygen atom is longer-bonded to the sulfur atom and subtends lower angles to its neighboring atoms of the  $-\text{OSO}_2\text{F}$  group, than the other oxygen atoms.

Introduction

Xenon difluoride can act as a fluoride ion donor, forming salts with strong fluoride ion acceptors, such as arsenic pentafluoride and metal pentafluorides.<sup>2,3,4,5</sup>

It also forms 1:1 molecular addition compounds with xenon tetrafluoride<sup>6</sup>, iodine pentafluoride<sup>7</sup>, and xenon oxide

tetrafluoride.<sup>8</sup> A third type of complex is obtained by the interaction of the difluoride with fluorosulfonic and perchloric acids.<sup>9</sup> The last type of  $\text{XeF}_2$  derivative was the subject of an earlier communication<sup>10</sup> in which we briefly reported the preparation and some properties of  $\text{FXeOSO}_2\text{F}$ ,  $\text{Xe}(\text{OSO}_2\text{F})_2$ ,  $\text{FXeOClO}_3$  and  $\text{Xe}(\text{OClO}_3)_2$ . In this paper we give our detailed x-ray single crystal structural analysis for  $\text{FXeOSO}_2\text{F}$ . The related compounds are discussed in the accompanying paper.<sup>11</sup>

### Experimental

Xenon(II) fluoride fluorosulfate was prepared by treating  $\text{XeF}_2$  with the correct molar quantity of fluorosulfonic acid at  $-75^\circ$ <sup>11</sup>, the hydrogen fluoride, formed in the reaction, being removed under vacuum at temperatures below  $-30^\circ$ . Material, powdered at  $\sim -10^\circ$ , was sealed in thin-walled quartz capillaries. Crystals were grown by sublimation at room temperature.

### Crystal Data

Crystals of  $\text{FXeOSO}_2\text{F}$ ,  $M = 249.4$ , are orthorhombic  
 $a = 9.88 \pm 0.01$ ,  $b = 10.00 \pm 0.01$ ,  $c = 10.13 \pm 0.01$  Å,  
 $V = 1001$  Å<sup>3</sup>,  $z = 8$ ,  $D_c = 3.30 \pm 0.02$  g cm<sup>-3</sup>,  $F(000) = 896$ .

Single crystal precession photographs of the  $h0l$ ,  $h1l$ ,  $0kl$ ,  $hkk$ , and  $hhl$  levels showed absences for  $0kl$ , with  $k$  odd,  $h0l$  with  $l$  odd,  $hk0$  with  $h$  odd,  $Ok0$  with  $k$  odd, and  $00l$  with  $l$  odd. This indicated that the most probable

space group was  $\text{Pbca} = \text{D}_{2h}^{15}$ . Complete indexing of the powder data, on the basis of the single crystal parameters proved that the crystals were representative of the bulk material.

### X-Ray Measurements

Data were recorded, at a temperature of  $0 \pm 2^\circ$ , using a manually operated single crystal orienter, with a low-temperature attachment,<sup>12</sup> on a G.E. diffractometer using Zr filtered MoK $\alpha$  radiation,  $\lambda = 0.7107 \text{ \AA}$ . The crystal was without well defined faces, edges or corners and was roughly spherical, the diameter at the outset being  $\sim 0.13 \text{ mm}$ . It was mounted with the (001) direction parallel to the  $\phi$  axis of the orienter. High order h00, 0k0 and 00l reflections were used to determine the unit cell constants. Intensities were measured, both for reflections and background, by the stationary crystal-stationary counter technique with 10 second counts. The background measurements were taken at  $\pm 1.0^\circ 2\theta$  for reflections up to  $20^\circ 2\theta$  and  $\pm 2^\circ 2\theta$  beyond this point. Counting rates were kept below 10,000 counts/second by the use of zirconium filters of known attenuation. Measurements were made on 1453 unique reflections occurring in the range  $2\theta \leq 60^\circ$ , of which 849 were considered to be observable above the background. The criterion for presence of a reflection was  $I > 3\sigma(I)$ , where  $\sigma(I)$  was determined from counting statistics, i.e.,  $\sigma(I) = \sqrt{C_P + C_B}$  where  $C_P$  is the peak count and  $C_B$  is the background count. Several standard reflections were monitored during the experiment at frequent intervals. There was an over-all intensity decrease of about 20% during the collection period. The raw data were divided into 8 batches and each batch was corrected by a different scale factor. The 8 experimental scale factors were taken directly proportional to the monitored intensities measured at 8 different time intervals. The data were then refined as a single problem, yielding a single R value. However, 8 scale factor parameters were included in the refinement to serve as a measure of the appropriateness of

the original choice of relative scale factors. At the end of the refinement the scale parameters had the following values:

.7248972  
.7683301  
.7608877  
.7318397  
.7841719  
.7549182  
.7309175  
.7207349

where the chronology of the experimental data is from top to bottom. The parameters have a mean of .7470872, maximum deviation from the mean of .0370847, and standard deviation of .0216772. There is no systematic trend in the deviations which is as it should be if we have treated the problem properly. The standard deviation is only 2.9% of the mean. We consider this to be a very satisfactory resolution of a difficult experimental problem. The small size and near-spherical nature of the crystal permitted a spherical sample absorption correction to be applied, for which  $\mu R$  was taken to be  $\leq 0.5$ .

Structure Analysis<sup>13</sup>

The position of the xenon atom was derived from the three-dimensional Patterson function and six cycles of full-matrix least-squares refinement of the scale, positional and thermal parameters, for this atom in an eight-fold position gave a value of 0.29 for the conventional  $R$  factor for all reflections. A three-dimensional electron density summation showed the position of the sulfur atom and subsequent refinement for Xe and S yielded  $R = 0.21$ . The light atom positions were obtained from a three-dimensional difference synthesis. Refinement by least-squares methods was continued, with scattering factors for neutral Xe, S, F, O obtained from the International Tables.<sup>14</sup> A correction for the real part of the anomalous dispersion effect<sup>14</sup> was made for xenon. The longer bonded terminal ligand of the  $SO_3F$  group was assumed to be the fluorine atom. In the final stages of refinement, anisotropic temperature factors were introduced for all atoms and the unobserved reflections were given zero weight in the analysis. The criteria for the latter were  $I_{unobs} = 1.5\sigma(I)$ . There were 604 unobserved reflections in a total of 1453. Nine weak, high order, reflections which appeared to be greatly in error, and two strong, low order, reflections, which were probably subject to extinction effects, were discarded. The final parameter shifts were all less than 0.1 $\sigma$  and the final agreement for 838 observed reflections was  $R = 0.0448$  and  $R' = 0.0441$  where  $R' = \sqrt{\sum w(F_o - F_c)^2} / \sqrt{\sum w F_o^2}$ . Unit weights were used throughout, except when the unobserved reflections were discarded (given zero weight). The standard deviation of an observation of unit weight with this weighting scheme was 2.36. The positional

and thermal parameters are listed in Table I. The  $F_o$  and  $F_c$  data are given in the microfilm version of this paper.

### Discussion

The structural analysis shows that crystals of  $\text{FXeOSO}_2\text{F}$  each consist of an ordered assembly of the monomer units illustrated in Figure 1. None of the (shorter) intermolecular contacts listed in Table II are short enough to demand special comment. All distances are compatible with the close packing of somewhat dipolar molecules. The arrangement of the molecules in the lattice is illustrated in Figure 2.

The molecule of  $\text{FXeOSO}_2\text{F}$  consists of a xenon atom approximately linearly coordinated to a fluorine atom on one side and an approximately tetrahedral fluorosulfate group on the other. The fluorosulfate group is coordinated to the Xe atom by way of an oxygen atom. The bond distances and angles are given in Table III.

The near-linear arrangement of  $\text{F}(1)\text{-Xe-O}(1)$  is typical of the coordination geometry previously observed in  $\text{Xe}(\text{II})$  compounds. Relevant structural features of xenon difluoride and some of its derivatives are given in Table IV. Although the Xe-F bond in  $\text{FXeOSO}_2\text{F}$  is shorter than in  $\text{XeF}_2$  it is larger than the terminal bonds in  $\text{Xe}_2\text{F}_3^+$ . The Xe-F bond is evidently much more  $\text{XeF}_2$ -like than in any of the other derivatives listed.

It is generally agreed that the Xe-F bond in  $\text{XeF}_2$  is less than an electron-pair bond. The simple molecular orbital bonding model, given first by Pimentel<sup>15</sup> and Rundle,<sup>16</sup> depicts the three atoms as bound by one electron pair. In a formally different model, Bilham and Linnett<sup>17</sup> have represented the binding of each fluorine atom to the xenon atom by a single electron bond. The valence-bond treatment advocated by Coulson<sup>18</sup> presents a similar picture. Coulson argues that the major canonical forms in the resonance hybrid for  $\text{XeF}_2$  are  $(\text{F-Xe})^+\text{F}^-$  and  $\text{F}^-(\text{Xe-F})^+$  (both ion species are classical octet species). Again, one bonding electron pair serves for all three atoms. As with the other models mentioned, the valence-bond representation suggests high bond polarity; indeed  ${}^{-\frac{1}{2}}\text{F-Xe}^+{}^{-\frac{1}{2}}\text{F}$  appears to be at least an approximate representation. The valence-bond approach is probably the most suitable one for the discussion of the  $\text{FXeOSO}_2\text{F}$  structure.

It is reasonable to assume that the major canonical forms in the  $\text{FXeOSO}_2\text{F}$  resonance hybrid are  $(\text{F-Xe})^+(\text{SO}_3\text{F})^-$  and  $\text{F}^-(\text{XeOSO}_2\text{F})^+$ . The bond length of 1.94 Å, for Xe-F, suggests that the  $(\text{Xe-F})^+$  weight is greater than in  $\text{XeF}_2$  and therefore indicates that  $(\text{F-Xe})^+(\text{SO}_3\text{F})^-$  is more dominant than  $\text{F}^-(\text{Xe-OSO}_2\text{F})^+$  in the resonance hybrid. A more quantitative measure of this dominance is of interest and can be derived from consideration of the Xe-F and Xe-O bond lengths.

-7-

The shortest observed<sup>14</sup> Xe(II)-F bond is the terminal bond in the compound  $\text{FXeFSb}_2\text{F}_{10}$ . This bond, which has a length of 1.84 Å is shorter than the bond in the I-F molecule, which is reported<sup>19</sup> to be 1.906 Å. Furthermore, the stretching frequency  $\nu(\text{Xe-F}) = 621 \text{ cm}^{-1}$  in  $\text{FXeFSb}_2\text{F}_{10}$ , is greater than given<sup>19</sup> for  $\nu(\text{I-F}) = 610 \text{ cm}^{-1}$ . This suggests that the Xe-F species in  $\text{FXeFSb}_2\text{F}_{10}$  is, at least approximately, the cation  $(\text{Xe-F})^+$ . (The cation is isoelectronic with I-F.) Both of these species may be represented in conventional bonding models, as electron-pair-bonded, octet, species. If we assign the electron-pair bond as possessing bond order unity, then the bond order in  $(\text{Xe-F})^+$  is 1 and in  $\text{XeF}_2$  it is 0.5.<sup>3</sup>

Although the relationship between bond order and bond length is not easily resolved from purely theoretical considerations, Pauling has given<sup>20</sup> an empirical relationship for fractional bonds:

$$D(n) = D(1) - 0.60 \log n$$

where  $D(n)$  is the bond length for bond of order  $n$  and  $D(1)$  is the bond length for order unity. Solving for the latter, assuming (from the  $\text{XeF}_2$  data)  $D(0.5) = 2.01 \text{ Å}$ , the Xe-F bond length  $D(1) = 1.83 \text{ Å}$ , which is in excellent agreement with our hypothesis. Continuing on this basis, the bond order in the terminal Xe-F bond in  $\text{FXeFSb}_2\text{F}_{10}$  is

0.96, whereas the terminal bonds in  $\text{Xe}_2\text{F}_3^+$  have a bond order of 0.76. The XeF bond order in  $\text{FXeOSO}_2\text{F}$  is only 0.63. If this result and our assumptions concerning the major canonical forms are valid, the canonical form  $(\text{F-Xe})^+(\text{OSO}_2\text{F})^-$  has a 63:37 dominance over the  $^-\text{F}(\text{XeOSO}_2\text{F})^+$  form.

The Xe-O bond length is larger than any Xe-O bond previously described, but a treatment analogous to that given in the Xe-F case cannot be made, since this is the first case of a Xe(II)-O bond length. It should be noted, however, that the bond lengths in both  $\text{XeO}_3$  and  $\text{XeO}_4$ , which are 1.76 and 1.74 Å, respectively,<sup>21,22</sup> are much shorter than the bond observed here. The Xe-O bonds in  $\text{XeO}_3$  and  $\text{XeO}_4$  can either be described as double bonds (i.e.,  $\text{Xe}=\text{O}_3$  and  $\text{Xe}=\text{O}_4$ ) or as semi-ionic bonded species (i.e.,  $(\text{Xe}^+:\rightarrow\text{O}^-)_3$  and  $(\text{Xe}^+:\rightarrow\text{O}^-)_4$ ). Either representation indicates that it would not be realistic to take 1.74 Å as the bond length for bond order unity. It is however possible to make a rough estimate of the bond order if we assume<sup>23</sup> that the bond length in the molecule  $\text{Xe}:\text{O}$  is akin to that in  $\text{I}:\text{F}$ , namely 1.91 Å. On this basis, the 2.16 Å Xe-O bond in  $\text{FXeOSO}_2\text{F}$  has a bond order of 0.38. This is in close agreement with the dominance of the  $(\text{F-Xe})^+(\text{OSO}_2\text{F})^-$  canonical form derived earlier.

The fluorosulfate geometry is compatible with the partial ionic bonding just discussed. It should first be

-9-

noted that the shape of the  $-\text{OSO}_2\text{F}$  group is fully consistent with the assignment of the fluorine atom to the position shown. The F(2)-S bond is not only the longest in the  $-\text{OSO}_2\text{F}$  group, but the bond angles, which this bond subtends to the other bonds in this group, are in the range  $100-106^\circ$ , and are, on the whole, less than the angles subtended by the other bonds to adjacent bonds. It is also impressive that the plane defined by the atoms F(2), S and O(1) is not significantly different from a mirror plane, as far as the  $-\text{OSO}_2\text{F}$  group is concerned. The F-Xe-O(1) group of atoms does not lie in the plane just defined, but there is no reason to anticipate restricted rotation about either the Xe-O(1) or O(1)-S bonds. Therefore the disposition adopted in this lattice is presumably one which results from the best packing and crystal energy.

The greater bond angles for O-S-O (e.g.  $120^\circ$ ) compared with O-S-F ( $106^\circ$ ) may be attributed to the greater repulsive effect of oxygen atoms. This may either be due to double bonding of oxygen to sulfur (i.e. to four-electron bonding) or due to high bond polarity (a consequence of a semi-ionic linkage  $\text{S}^+ \rightarrow \text{O}^-$ ). Many object to the major involvement of sulfur  $3d$  orbitals in bonding.<sup>24</sup> For them, the latter model for the S-O bond is appropriate. With this representation, each of the terminal S-O bonds is a semi-ionic linkage (involving one electron pair) and the S-F bond is a

(covalent) electron-pair bond. Clearly for the isolated  $\text{SO}_3\text{F}^-$  group we should anticipate three, equivalent, semi-ionic S-O bonds and this appears to be the case in  $\text{KSO}_3\text{F}$ <sup>25</sup> and  $\text{NH}_4\text{SO}_3\text{F}$ .<sup>26</sup> Such a situation is not observed in  $\text{FXeOSO}_2\text{F}$ . We see rather, that the third oxygen ligand of the sulfur atom (O(1)) subtends smaller angles to the other oxygen atoms (112, 111°) than they do to one another (120°). Furthermore, the fluorine atom (F(2)), subtends smaller angles to O(1) (101°) than to the other oxygen ligands (105.8, 105.3°). This means that the repulsive effect of O(1) is less than for O(2) or O(3), although evidently greater than for F(2). This is compatible with a decrease in the net negative charge borne by the ligands in the sequence  $\text{O}(2) = \text{O}(3) > \text{O}(1) > \text{F}(2)$ . The greater polarity of the S-O (term.) bonds compared with S-O (bridge) may also account for the former bonds being significantly shorter than the latter.

It is instructive to compare the  $-\text{OSO}_2\text{F}$  group observed here with the  $\text{SO}_3\text{F}^-$  ion observed in the potassium<sup>25</sup> and ammonium<sup>26</sup> salts. Although there is evidently disordering of the O and F placement of the sulfur ligands in the anion in the potassium salt, and partial disordering in the ammonium salt, ion dimensions were determined for each case, assuming  $\text{C}_{3v}$  symmetry<sup>25,26</sup> of the disordered ion.

The  $-\text{OSO}_2\text{F}$  and  $\text{SO}_3\text{F}^-$  species are compared in Table V.

Evidently the S-F and S-O (terminal) bonds in the xenon compound are essentially the same as in the simple salts. Indeed even the bond angles are remarkably akin, but for those subtended by the S-O(1) bond. Departure of the -OSO<sub>2</sub>F group geometry from the ionic (SO<sub>3</sub>F<sup>-</sup>) ideal, may be attributed solely to a change in the character of the oxygen atom, O(1), linked to xenon. The atom O(1) has evidently lost electron density to the Xe-F group. This fits rather conveniently into the description of the SO<sub>3</sub>F<sup>-</sup> species as a sulfur atom semi-ionic bonded (S<sup>+</sup>: → O<sup>-</sup>) to each oxygen atom and electron-pair bonded to the F atom (S : F). But in the xenon compound, atom O(1) possesses less electron density, as a consequence of the contributing canonical form (FO<sub>2</sub>SO-Xe)<sup>+</sup>F<sup>-</sup>, in which O(1) is bicovalent.

#### Acknowledgments

The authors are grateful to the National Science Foundation for grant GP-7153X, for support of that part of this work which was carried out at Princeton University.

References

1. Royal Radar Establishment, Malvern, England.
2. F. O. Sladky, P. A. Bulliner, N. Bartlett, B. G. DeBoer, and A. Zalkin, Chem. Comm., 1968, 1048.
3. F. O. Sladky, P. A. Bulliner, and N. Bartlett, J. Chem. Soc. (A), 1969, 2179.
4. V. M. McRae, R. D. Peacock, and D. R. Russell, Chem. Comm., 1969, 62.
5. J. G. Knowles and J. H. Holloway, J. Chem. Soc. (A), 1969, 756.
6. J. H. Burns, R. D. Ellison, and H. A. Levy, Acta Cryst., 18, 1965, 11.
- 7a. G. R. Jones, R. D. Burbank, and Neil Bartlett, Inorg. Chem., 9, 1970, 2264.
- 7b. N. Bartlett and F. O. Sladky, J. Chem. Soc. (A), 1969, 2188.
8. N. Bartlett and M. Wechsberg, Z. anorg. u. allgem. Chemie, 1971, in press.
9. N. Bartlett and F. O. Sladky, paper presented at 2nd European Fluorine Chemistry Symposium, Göttingen, August 28-31, 1968.
10. Neil Bartlett, M. Wechsberg, F. O. Sladky, P. A. Bulliner, G. R. Jones and R. D. Burbank, Chem. Comm., 1969, 703.
11. N. Bartlett, M. Wechsberg, and P. A. Bulliner, accompanying paper.

12. R. D. Burbank and S. S. DeBalla, to be published.
13. Programs employed during the analysis included FOUR (C. J. Fritchie, unpublished, modified by L. Guggenberger and P. B. Jemison), ORFLS (W. R. Busing, K. O. Martin, and H. A. Levy, ORNL Dept. TM-305, 1962, modified by B. B. Cetlin and W. C. Hamilton), ORFFE (W. R. Busing, K. O. Martin, and H. A. Levy, ORNL Dept. TM-306, 1964), and ORTEP (C. K. Johnson, ORNL Dept. 3794, 1965, modified by R. L. Kornegay). All computations were carried out on a G.E. 635 computer.
- 14a. International Tables for x-ray Crystallography, Vol. III Kynoch Press, Birmingham, p. 202.
- 14b. ibid, p. 216.
15. G. Pimentel, J. Chem. Phys., 19, 446 (1951).
16. R. E. Rundle, J. Am. Chem. Soc., 85, 112 (1963).
17. J. Bilham and J. W. Linnett, Nature, 201, 1323 (1964).
18. C. A. Coulson, J. Chem. Soc., 1442 (1964).
19. R. A. Durie, Proc. Roy. Soc., A, 207, 388 (1951).
20. L. Pauling, The Nature of the Chemical Bond, 3rd Edition, Cornell University Press, 1960, Ithaca, New York, p. 255.
21. D. H. Templeton, A. Zalkin, J. D. Forrester, and S. M. Williamson, J. Am. Chem. Soc., 85, 817 (1963).
22. G. Gunderson, K. Hedberg, and J. L. Huston, Acta Cryst., 25, 124 (1969).

23. The combined nuclear charges for XeO and IF are the same and both may be represented as electron pair bonded species.
24. R. E. Rundle, Record of Chemical Progress, 23, 195 (1962); E. H. Wiebenga, E. E. Havinga, and K. H. Boswijk, Advances in Inorganic and Radiochemistry, 2, 155 (1961); Various authors, in Noble Gas Compounds (H. H. Hyman, ed.), Chicago University Press, Chicago and London, 1963, pp. 315-387.
25. K. O'Sullivan, R. C. Thompson, and James Trotter, J. Chem. Soc., A, 2026 (1967).
26. K. O'Sullivan, R. C. Thompson, and James Trotter, J. Chem. Soc., A, 1814 (1970).

TABLE I

## Final Positional and Thermal Parameters

Atom	x	y	z	$\beta_{11}$ (a)	$\beta_{22}$	$\beta_{33}$	$\beta_{12}$	$\beta_{13}$	$\beta_{23}$
Xe	0.6757(1) <sup>b</sup>	0.4519(1)	0.6686(1)	0.0071(1)	0.0070(1)	0.0086(1)	-0.0008(1)	0.001(1)	-0.1007(1)
S	0.5340(3)	0.2269(3)	0.4849(3)	0.0075(3)	0.0062(3)	0.0080(3)	-0.0002(2)	0.001(3)	-0.1001(3)
F <sub>1</sub>	0.8095(8)	0.5394(9)	0.7791(9)	0.0112(10)	0.0139(1)	0.0169(12)	-0.0027(9)	-0.0023(9)	-0.0033(10)
F <sub>2</sub>	0.4555(9)	0.1398(9)	0.5842(9)	0.0156(12)	0.0123(10)	0.0142(11)	-0.0056(1)	0.0013(10)	0.0039(9)
O <sub>1</sub>	0.5205(9)	0.3618(8)	0.5484(9)	0.0077(9)	0.0059(8)	0.0139(12)	0.0012(7)	-0.0014(9)	-0.0030(8)
O <sub>2</sub>	0.6701(9)	0.1784(9)	0.4886(11)	0.0094(10)	0.0099(10)	0.0151(13)	0.0024(9)	-0.0028(11)	-0.0022(10)
O <sub>3</sub>	0.4598(9)	0.2206(9)	0.3656(8)	0.0102(10)	0.0125(12)	0.0078(9)	0.001(9)	-0.0029(9)	-0.0026(9)

<sup>a</sup>The form of the anisotropic thermal ellipsoid is  $\exp \left\{ -(\beta_{11}h^2 + \beta_{22}k^2 + \beta_{33}l^2 + \beta_{12}hk + 2\beta_{13}hl + 2\beta_{23}kl) \right\}$

<sup>b</sup>Number in parentheses is estimated standard deviation in the least significant digit .

TABLE II

Intermolecular Contacts Less Than 4.0 Å

Xe...F(1 <sup>VIII</sup> )**	3.760(8)*	Xe...O(2 <sup>III</sup> )	3.283(9)
Xe...F(1 <sup>IX</sup> )	3.949(9)	Xe...O(2 <sup>IV</sup> )	3.494(10)
Xe...F(2 <sup>II</sup> )	3.879(9)	Xe...O(3 <sup>II</sup> )	3.313(9)
Xe...F(2 <sup>V</sup> )	3.388(8)	Xe...O(3 <sup>IV</sup> )	3.392(9)
Xe...O(1 <sup>I</sup> )	3.473(9)		
S...O(2 <sup>X</sup> )	3.728(10)	S...O(3 <sup>IV</sup> )	3.960(10)
F(1)...F(2 <sup>III</sup> )	3.209(13)	F(1)...O(2 <sup>III</sup> )	3.262(14)
F(1)...F(2 <sup>V</sup> )	3.127(12)	F(1)...O(2 <sup>VII</sup> )	3.536(13)
F(1)...O(1 <sup>VI</sup> )	3.249(12)	F(1)...O(2 <sup>IV</sup> )	3.338(13)
F(1)...O(1 <sup>VII</sup> )	3.352(13)	F(1)...O(3 <sup>II</sup> )	3.334(12)
F(2)...F(2 <sup>XI</sup> )	3.392(19)	F(2)...O(3 <sup>IV</sup> )	3.173(13)
F(2)...O(2 <sup>X</sup> )	3.435(14)	F(2)...O(3 <sup>XI</sup> )	3.735(13)
F(2)...O(2 <sup>XI</sup> )	3.494(12)		
O(1)...O(3 <sup>IV</sup> )	3.370(13)	O(1)...O(2 <sup>X</sup> )	3.506(13)
O(1)...O(1 <sup>I</sup> )	2.961(16)		
O(2)...O(3 <sup>II</sup> )	3.376(13)		

\* Estimated standard deviations in parentheses.

\*\* The crystal-chemical unit is at xyz and the Roman numbers refer to equivalent positions: I(1-x, 1-y, 1-z), II( $\frac{1}{2}+x$ ,  $\frac{1}{2}-y$ , 1-z), III( $1\frac{1}{2}-x$ ,  $\frac{1}{2}+y$ , z), IV(x,  $\frac{1}{2}-y$ ,  $\frac{1}{2}+z$ ), V(1-x,  $\frac{1}{2}+y$ ,  $1\frac{1}{2}-z$ ), VI( $\frac{1}{2}+x$ , y,  $1\frac{1}{2}-z$ ), VII( $1\frac{1}{2}-x$ , 1-y,  $\frac{1}{2}+z$ ), VIII( $x-\frac{1}{2}$ , y,  $1\frac{1}{2}-z$ ), IX( $1\frac{1}{2}-x$ , 1-y,  $z-\frac{1}{2}$ ), X( $x-\frac{1}{2}$ ,  $\frac{1}{2}-y$ , 1-z), XI(1-x, y, 1-z).

TABLE III

Intramolecular Distances (Å) and Angles (°) for FXeOSO<sub>2</sub>F

Xe - F(1)	1.940(8)	Xe - O(1)	2.155(8)
S - O(1)	1.501(8)	S - O(2)	1.430(9)
S - O(3)	1.415(9)	S - F(2)	1.540(9)
F(2) - O(1)	2.340(11)	F(2) - O(2)	2.363(13)
F(2) - O(3)	2.359(12)	O(1) - O(2)	2.432(11)
O(1) - O(3)	2.405(12)	O(2) - O(3)	2.459(13)
F(1) - Xe - O(1)	177.4(0.3)	Xe - O(1) - S	123.7(0.5)
O(1) - S - O(2)	112.1(0.5)	O(1) - S - O(3)	111.1(0.6)
O(1) - S - F(2)	100.6(0.5)	F(2) - S - O(2)	105.3(0.6)
F(2) - S - O(3)	105.8(0.5)	O(2) - S - O(3)	119.6(0.6)



TABLE V

Comparison of the  $-\text{OSO}_2\text{F}$  Dimensions ( $\text{\AA}$  and degrees) with those of  $\text{SO}_3\text{F}^-$  in  $\text{KSO}_3\text{F}^{(a)}$  and  $\text{NH}_4\text{SO}_3\text{F}^{(b)}$

	<u><math>-\text{OSO}_2\text{F}</math></u>		<u><math>\text{SO}_3\text{F}^-</math></u>
F(2)—S	1.54(1)	F—S	$\left\{ \begin{array}{l} 1.58(2)^{(a)} \\ 1.55(1)^{(b)} \end{array} \right.$
O(2)—S	1.43(1)	O—S	$\left\{ \begin{array}{l} 1.43(1)^{(a)} \\ 1.45(1)^{(b)} \end{array} \right.$
O(3)—S	1.42(1)		
O(1)—S	1.50(1)		
F(2)—S—O(3)	105.8(5)	F—S—O	$\left\{ \begin{array}{l} 106.0(5)^{(a)} \\ 105.8(7)^{(b)} \end{array} \right.$
F(2)—S—O(2)	105.3(6)		
F(2)—S—O(1)	100.6(5)		
O(2)—S—O(3)	119.6(6)	O—S—O	$\left\{ \begin{array}{l} 112.9(7)^{(a)} \\ 113.0(5)^{(b)} \end{array} \right.$
O(2)—S—O(1)	112.1(5)		
O(3)—S—O(1)	111.1(6)		

(a) Ref. 25.

(b) Ref. 26.

NOTICE FOR PRINTER

This table is to be included in the microfilm version of this paper but excluded from the version for regular issue.

TABLE VI

Calculated and Observed Structure Factors for  $\text{FXeOSO}_2\text{F}$   
(Scale unity)




NOTICE FOR PRINTER: This table is to be included in the microfilm version of this paper but excluded from the version for regular use.

TABLE VII


The root-mean-square component of thermal displacement along principal axis R in Angstrom units.

Atom	R	RMS Displacement
Xe	1	0.1765 ± 0.0013
	2	0.1949 ± 0.0013
	3	0.2146 ± 0.0010
S	1	0.1761 ± 0.0039
	2	0.1927 ± 0.0040
	3	0.2037 ± 0.0038
F(1)	1	0.1999 ± 0.0118
	2	0.2747 ± 0.0116
	3	0.3137 ± 0.0135
F(2)	1	0.1766 ± 0.0123
	2	0.2847 ± 0.0113
	3	0.3183 ± 0.0117
O(1)	1	0.1543 ± 0.0123
	2	0.1924 ± 0.0121
	3	0.2812 ± 0.0118
O(2)	1	0.1893 ± 0.0130
	2	0.2228 ± 0.0132
	3	0.3000 ± 0.0129
O(3)	1	0.1629 ± 0.0132
	2	0.2363 ± 0.0120
	3	0.2678 ± 0.0122



XBL 717-6931

Figure 1. The FXeOSO<sub>2</sub>F molecule (distances in Å units and angles in degrees).



XBL 723-6063

Figure 2. A view of the molecular arrangement in FXeOSO<sub>2</sub>F.

LEGAL NOTICE

*This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.*

TECHNICAL INFORMATION DIVISION  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720