Lawrence Berkeley National Laboratory

Recent Work

Title

ALPHA-PARTICLE EMISSION FROM THE DEEP-INELASTIC REACTION: 1354 MeV ^{165}Ho + ^{181}Ta

Permalink

<https://escholarship.org/uc/item/2dn7f39j>

Author

Sobotka, L.G.

Publication Date

1981-10-01

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

NOV 4 1981

Submitted to Physical Review C

LIBRARY AND
DOCUMENTS SECTION

ALPHA-PARTICLE EMISSION FROM THE DEEP-INELASTIC REACTION:
1354 MeV $^{165}\text{Ho} + ^{181}\text{Ta}$

L.G. Sobotka, R.J. McDonald, G.J. Wozniak, D.J. Morrissey,
A.J. Pacheco, and L.G. Moretto

October 1981

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782

LBL-13319
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Alpha-particle Emission from the Deep-Inelastic Reaction:

$$1354 \text{ MeV } ^{165}\text{Ho} + ^{181}\text{Ta}$$

L.G. Sobotka, R.J. McDonald, G.J. Wozniak, D.J. Morrissey,
A.J. Pacheco, and L.G. Moretto

Nuclear Science Division
Lawrence Berkeley Laboratory
University of California
Berkeley, CA 94720

The in-plane correlation between projectile-like fragments and α -particles has been studied over a broad range of angles for the system 1354 MeV $^{165}\text{Ho} + ^{181}\text{Ta}$. These data unambiguously indicate that the dominant sources of α -particles are the fully accelerated fragments.

This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics and by Nuclear Sciences of the Basic Energy Sciences Program of the U.S. Department of Energy under Contract W-7405-ENG-48.

The study of light particles emitted in low energy (≤ 10 MeV/u) heavy-ion reactions in coincidence with deep-inelastic or relaxed exit channels has been conducted in two steps. First, one has tried to identify the emission source of the light particles and then, after verification of the emitting sources, the emitted particles have been used as a probe of the temperature and spin of the emitter. So far, these studies have not yielded a consensus on the dominant source of light particles. For example, the reactions $96 \text{ MeV } ^{16}\text{O} + ^{58}\text{Ni}$ ¹, $280 \text{ MeV } ^{40}\text{Ar} + ^{58}\text{Ni}$ ², $400 \text{ MeV } ^{40}\text{Ar} + ^{93}\text{Nb}$ ³, and $664 \text{ MeV } ^{84}\text{Kr} + \text{nat Ag}$ ^{4,5} have indicated that the dominant sources of α -particles are the fully accelerated fragments. On the other hand, studies of 222 , 274 , and $340 \text{ MeV } ^{40}\text{Ar} + ^{116}\text{Sn}$, ^{154}Sm , ^{164}Dy , ^{197}Au ^{6,7}, and $480 \text{ MeV } ^{56}\text{Fe} + \text{nat Ag}$ ⁸ have indicated that the bulk of the α -particle emission occurs prior to scission of the di-nuclear system. In addition to evaporative components, some studies have observed a fast forward component not easily explained in terms of evaporation.⁶ With this communication we present evidence that the bulk of α -particle emission, from the $1354 \text{ MeV } ^{165}\text{Ho} + ^{181}\text{Ta}$ system, occurs from the fully accelerated fragments.

A natural Ta target (1.4 mg/cm^2) was bombarded with $8.2 \text{ MeV/u } ^{165}\text{Ho}$ (~ 10 ena) from the Lawrence Berkeley Laboratory SuperHILAC. A solid state detector ($300 \text{ }\mu\text{m}$) positioned at the grazing angle (29°) was used to detect the projectile-like fragment and to define the reaction plane. On the other side of the beam, five solid state ΔE -E telescopes ($40 \text{ }\mu\text{m}$, 5 mm) were used to detect the α -particles. While both in- and out-of-plane angular distributions were obtained, only the in-plane data will be presented in this communication.

Laboratory α -particle energy spectra are shown in Fig. 1 for angles between 30° and 115° . These spectra were obtained by requiring a coincidence between an α -particle at the laboratory angles indicated in Fig. 1 and a heavy-ion detected at -29° with an energy in the deep-inelastic region. A notable feature of these spectra is the occurrence of two separate peaks at the most forward angle, suggesting the presence of two emission sources. The data at larger angles show essentially a single component, which shifts towards lower energy as the in-plane angle increases.

In order to determine the emission sources, we have plotted in Fig. 2a the experimentally extracted root-mean-square velocity (v^{rms}) of the α -particles. Also indicated on this figure are the velocity vectors for the detected projectile-like fragment (gated on the deep-inelastic events), the calculated velocity of the undetected fragment, and the velocity of the system center of mass. As Fig. 2a shows, the α -particle velocities are centered around the end of the velocity vector of the target-like fragment. This agrees with the assumption that the α -particles are emitted from the fully accelerated target-like fragment, rather than from a system moving with the center of mass velocity.

Further evidence for fragment emission can be obtained by determining the Q-value dependence of v_α^{rms} . In Fig. 2b we have plotted the average vector diagram for three different Q-value bins (all in the deep-inelastic region) and the corresponding v_α^{rms} . A systematic motion of the loci of v_α^{rms} values is seen that can only be explained by a source that has a Q-dependent velocity. This trend is explained by the change of the velocity of the target-like fragment with Q value, as shown in the figure.

Finally, utilizing two body kinematics the events were reconstructed and transformed event by event into the rest frame of the target recoil. In this frame the α -particle spectral shape is found to be independent of angle with the exception of the most forward angle. In addition, this spectral shape is reproduced quite well by an evaporation formalism that accounts for exit channel shape polarization and fluctuations.⁹ The importance of such deformation effects can be readily appreciated by comparing the effective Coulomb barrier (B_c) observed in the present data, calculated from $\langle \epsilon_\alpha \rangle = B_c + 2T$ with observed entrance Coulomb barriers in α -particle + nucleus reactions.¹⁰ The former effective barrier is approximately 25% lower than the entrance channel value.

As has been seen in previous studies of α -particles emitted in heavy ion reactions⁵, a high energy contaminant to the major evaporative component was observed at the most forward angle. Unfortunately, insufficient data were obtained at forward angles to determine the source of these α -particles.

In conclusion, by use of in-plane correlation techniques we have shown that for the system 1354 MeV $^{165}\text{Ho} + ^{181}\text{Ta}$, the dominant sources of α -particles are the fully accelerated fragments.

We would like to thank the authors of references 7 and 8 for providing us with preprints of their work. We would also like to acknowledge the staff of the SuperHILAC for delivering an excellent ^{165}Ho beam. This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics and by Nuclear Sciences of the Basic Energy Sciences Program of the U.S. Department of Energy under Contract W-7405-ENG-48.

References

1. H. Ho, R. Albrecht, H. Damjantschitsch, F.J. Demond, W. Kühn, J. Slemmer, J.P. Wurm, D. Disdier, V. Rauch, F. Scheibling and T. Dossing, Z. Phys. A300, 205 (1981).
2. R. Babinet, B. Cauvin, J. Girard, J.M. Alexander, T.H. Chiang, J. Galin, B. Gatty, D. Guerreau and X. Tarrago, Z. Phys. A295, 153 (1980).
3. W. Kühn, R. Albrecht, H. Damjantschitsch, H. Ho, R.M. Ronningen, J. Slemmer, J.P. Wurm, I. Rode and F. Scheibling, Z. Phys. A298, 95 (1980).
4. L.G. Sobotka, C.C. Hsu, G.J. Wozniak, G.U. Rattazzi, R.J. McDonald, A.J. Pacheco and L.G. Moretto, Phys. Rev. Lett. 46, 887 (1981).
5. L.G. Sobotka, C.C. Hsu, G.J. Wozniak, D.J. Morrissey and L.G. Moretto, Nucl. Phys. A (1981), in press.
6. D. Logan, H. Delagrange, M.F. Rivet, M. Rajogopalan, J.M. Alexander, M. Kaplan, M.S. Zisman and E. Duck, Phys. Rev. C22, 1080 (1980).
7. M.F. Rivet, D. Logan, J.M. Alexander, E. Duck, M.S. Zisman and M. Kaplan (preprint 1981).
8. D. Guerreau, D. Logan, M.S. Zisman, J.M. Alexander, E. Duck and M. Kaplan (preprint 1981).
9. L.G. Moretto, Nucl. Phys. A247, 211 (1975).
10. M.A. McMahan and J.M. Alexander, Phys. Rev. C21 (1980).

Figure Captions

Fig. 1. Coincident laboratory α -particle energy spectra. Laboratory angles are indicated.

Fig. 2. a) Velocity diagram for the $^{181}\text{Ta} + ^{165}\text{Ho}$ (1354 MeV) system.

The open circles are the rms velocities extracted from the coincident laboratory α -particle energy spectra. The solid circles indicate the rms velocities of the two separate peaks that appear in the most forward data. The full large rings indicate the loci of expected α -particle velocities⁷ from the three different rest frames. For the target-like fragment, the locus of velocities for a 10% reduction in the expected average emission energy is indicated by a partial ring. The detection threshold is shown as a dashed arc. The letters P and T stand for projectile-like and target-like, respectively.

b) The velocity diagrams for three different Q-bins (all in the deep-inelastic region). The rms α -particle velocities for each bin are indicated. The smallest energy loss bin is indicated by triangles and the largest energy loss data by squares. The three partial rings are drawn to guide the eye. They have the same radius and are centered on the three different recoil velocities.

XBL 818-2462

Fig. 1

$^{181}\text{Ta} + ^{165}\text{Ho} \text{ (1354 MeV)}$

(b)

XBL 818-2456A

Fig. 2

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720