

UC Riverside

UC Riverside Previously Published Works

Title

Shallow Genome Sequencing for Phylogenomics of Mycorrhizal Fungi from Endangered Orchids

Permalink

<https://escholarship.org/uc/item/2f34b5qf>

Authors

Unruh, Sarah A

Pires, J Chris

Zettler, Lawrence

et al.

Publication Date

2019-12-03

DOI

10.1101/862763

Peer reviewed

1 **Shallow Genome Sequencing for Phylogenomics of Mycorrhizal Fungi from**
2 **Endangered Orchids**

3
4 Sarah A. Unruh¹, J. Chris Pires¹, Lawrence Zettler², Luigi Erba², Igor Grigoriev³, Kerrie Barry³,
5 Christopher Daum³, Anna Lipzen³, and Jason E. Stajich⁴

6
7 1. Division of Biological Sciences, University of Missouri, Columbia, MO, United States

8
9 Sarah Unruh Roles: Conceptualization, Data Curation, Formal Analysis, Funding Acquisition,
10 Investigation, Project Administration, Resources, Visualization, Writing – Original Draft
11 Preparation, Writing – Review and Editing

12
13 J. Chris Pires Roles: Conceptualization, Funding Acquisition, Project Administration, Resources,
14 Supervision, Writing – Review and Editing

15
16 2. Department of Biology, Illinois College, Jacksonville, IL, United States

17
18 Lawrence W. Zettler Roles: Funding acquisition, Resources

19
20 Luigi Erba Roles: Investigation

21
22 3. United States of America Department of Energy Joint Genome Institute, Walnut Creek, CA,
23 United States

24
25 Igor Grigoriev Roles: Project Administration, Resources

26
27 Kerrie Barry Roles: Project Administration

28
29 Christopher Daum: Resources

30
31 Anna Lipzen: Investigation

32
33 4. Microbiology and Plant Pathology, University of California-Riverside, Riverside, CA, United
34 States

35
36 Jason E. Stajich Roles: Data Curation, Formal analysis, Investigation, Methodology, Resources,
37 Software, Visualization, Writing – Review and Editing

ABSTRACT

41
42
43 Most plant species form symbioses with mycorrhizal fungi and this relationship is especially
44 important for orchids. Fungi in the genera *Tulasnella*, *Ceratobasidium*, and *Serendipita* are
45 critically important for orchid germination, growth and development. The goals of this study are
46 to understand the phylogenetic relationships of mycorrhizal fungi and to improve the taxonomic
47 resources for these groups. We identified 32 fungal isolates with the internal transcribed spacer
48 region and used shallow genome sequencing to functionally annotate these isolates. We
49 constructed phylogenetic trees from 408 orthologous nuclear genes for 50 taxa representing 14
50 genera, 11 families, and five orders in Agaricomycotina. While confirming relationships among
51 the orders Cantharellales, Sebaciales, and Auriculariales, our results suggest novel relationships
52 between families in the Cantharellales. Consistent with previous studies, we found the genera
53 *Ceratobasidium* and *Thanatephorus* of Ceratobasidiaceae to not be monophyletic. Within the
54 monophyletic genus *Tulasnella*, we found strong phylogenetic signals that suggest a potentially
55 new species and a revision of current species boundaries (e.g. *Tulasnella calospora*); however it
56 is premature to make taxonomic revisions without further sampling and morphological
57 descriptions. There is low resolution of *Serendipita* isolates collected . More sampling is needed
58 from areas around the world before making evolutionary-informed changes in taxonomy. Our
59 study adds value to an important living collection of fungi isolated from endangered orchid
60 species, but also informs future investigations of the evolution of orchid mycorrhizal fungi.
61

INTRODUCTION

62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85

Fungi are more than mere decomposers, they form symbioses with every other group of organisms on Earth. Fungal interactions span the entire symbiotic spectrum, from parasitism to mutualism. Most intertwined with plants, may have even enabled development/existence of land plants (Lutzoni et al., 2018). As a result of this long-term association, fungi are essential symbionts to almost every plant species on Earth. The fungi live in plant roots are called mycorrhizal fungi and associate with more than 85% of plant species (Smith and Read, 2008). Mycorrhizal fungi are critical for plant health and function by helping obtain and retain water, mediating defense responses, participating in signaling between roots, and facilitating the exchange of nutrients like carbon, phosphorus, and nitrogen (Barto et al., 2012; Jung et al., 2012; Peterson and Massicotte, 2004; Wang et al., 2017; Yoder et al., 2010). The plant group that relies the most on their mycorrhizal fungi are orchids.

Orchids rely on their mycorrhizal symbionts to stimulate plant development during seed germination by providing carbon resources (Kuga et al., 2014). Orchid mycorrhizal fungi (ORM) form hyphal coils termed pelotons inside the cells of orchid embryos and in the adult roots, tubers, or rhizomes (McCormick et al., 2016; Rasmussen et al., 2015). These pelotons are the sites of nutrient exchange and the molecular nature of this marketplace remains poorly understood though exciting new research shedding light (Fochi et al., 2017a; Fochi et al., 2017b; Kuga et al., 2014). Most orchids associate with mycobionts belonging to the basidiomycete groups Sebacinales, Ceratobasidiaceae and Tulasnellaceae. In addition to orchid mycorrhizal fungi, these groups contain saprotrophs, plant pathogens, and ectomycorrhizal representing a wide array of metabolic capabilities (Kohler et al., 2015; Nagy et al., 2016). Furthermore, molecular studies have revealed simultaneous root colonization by multiple fungal partners in

86 both photosynthetic terrestrial and epiphytic orchids (Martos et al., 2012). Concluding sentence
87 that makes the argument that there are many dynamics we need to better understand so we need
88 to characterize the diversity of these fungi to untangle their interactions and mechanisms.

89 Although fungi play critical roles, they are rarely visible on the landscape. The number of
90 extant fungal species on Earth ranges from 2-5 million (Blackwell, 2011; Hawksworth and
91 Lücking, 2017) up to 166 million species (Larsen et al., 2017). Most species are microscopic and
92 over the last few decades species identification has relied on molecular methods. Historically,
93 these methods often have used a single molecular marker such as ITS (Nilsson et al., 2014).
94 However modern genome sequencing methods are important tools to discover and describe
95 taxonomic, phylogenetic and functional diversity. The use of different, new analytical tools has
96 also greatly benefited our knowledge of the below-ground ecology of orchids and orchid
97 mycorrhizal fungi. On the right track with multiple markers and Bayesian species delimitations
98 (Ruibal et al., 2014; Ruibal et al., 2013; Whitehead et al., 2017). New species of *Tulasnella* and
99 relatives are constantly being identified (Linde et al., 2017). Continue to combine sequencing
100 with taxonomic knowledge to provide a comprehensive description of the species that associate
101 with orchids.

102 The genera of orchid fungi we have sampled belong to two orders, Cantharellales and
103 Sebaciales, in the Agaricomycetes. Cantharellales is sister to the rest of class Agaricomycetes
104 and comprises seven families total (Ceratobasidiaceae, Tulasnellaceae, Botryobasidiaceae,
105 Cantharellaceae, Clavulinaceae, Hydnaceae, and Aphelariaceae), though Hibbett et al., (2014),
106 define Cantharellaceae and Clavulinaceae as synonymous with Hydnaceae and the status of
107 Aphelariaceae is unknown (Kirk et al., 2008; Leacock, 2018). Ceratobasidiaceae has two genera
108 (*Ceratobasidium* and *Rhizoctonia/Thanatephorus*) that have been demonstrated to be

109 polyphyletic (Veldre et al., 2013). In fact, the type specimen for *Ceratobasidium* has since been
110 reclassified as a member of the order Auriculariales based on the characters like the shape of the
111 basidia and the dolipore (specialized hyphal septa) ultrastructure , leading Oberwinkler et al.,
112 (2013a) to restrict *Ceratobasidium* and Ceratobasidiaceae to the type specimen and reclassifying
113 *Ceratobasidium* spp. as *Rhizoctonia* (Kirk et al., 2008). Tulasnellaceae contains 3 genera and c.
114 50 sp (Kirk et al., 2008). In addition to these described families, the genus *Sisotrema* is known to
115 be polyphyletic with members in Auriculariales as well as Cantharellales. Successively sister to
116 the rest of the Agaricomycetes is the order Sebaciniales which includes two families – the
117 Sebacinaceae and Serendipitaceae (Weiss et al., 2016). Though this order comprises a wide
118 swath of diversity, it remains difficult to adequately describe species due to a high volume of
119 environmental sequence data without information about morphological characters (Oberwinkler
120 et al., 2013b; Weiss et al., 2016).

121 In this study, our primary goal is to shallowly sequence a rich living collection of fungi
122 isolated from orchid roots and seedlings to provide a phylogenetic framework for future genome-
123 enabled evolutionary and functional studies. Our secondary goal, with the addition of key
124 outgroups, is to answer a series of nested phylogenetic questions about the relationships among
125 the orders, families and genera of Agaricomycetes, with a focus on Ceratobasidiaceae,
126 Tulasnellaceae, and Sebacinaceae. We screened taxa using ITS sequencing, and after
127 contaminants were removed we chose 32 taxa for shallow genome sequencing. A total of 50 taxa
128 were analyzed and we extracted 408 orthologous genes. Two highly-supported phylogenetic
129 trees were constructed with RAxML and ASTRAL-III that were overall highly congruent. We
130 discuss how our study provides new insight into the relationships of these orchid mycorrhizal
131 fungi, highlights areas for taxonomic attention and we suggest future research directions.

132
133
134

2. MATERIALS AND METHODS

135 2.1 Taxonomic sampling

136 32 environmental samples were isolated from endangered orchids. These samples span
137 three genera in three families in two orders. Outgroup genomes were chosen from the repository
138 in MycoCosm to capture the breadth of taxonomic diversity (Grigoriev et al., 2014). Two super
139 outgroups (*Kockovaella* sp and *Calocera* sp) were chosen from the successively sister classes
140 outside the ingroup class Agaricomycetes [Tremellomycetes, [Dacrymycetes,
141 [Agaricomycetes]]]. In the Cantharellales we sampled the three genomes in Ceratobasidiaceae
142 (*Rhizoctonia solani*, *Thanatephorus cucumeris*, and *Ceratobasidium* sp AG1), the two genomes
143 in Tulasnellaceae (*Tulasnella calospora* AL13/4D, and *Tulasnella calospora* UAMH9824), and
144 one genome each from 4 of the remaining 5 families *Botryobasidium botryosum*
145 (Botryobasidiaceae), *Clavulina* sp (Clavulinaceae), *Cantharellus anzutake* (Cantharellaceae), and
146 *Hydnum rufescens* (Hydnaceae) . We also included three genomes in Serendipitaceae
147 (Sebacinales) *Sebacina vermifera* (syn. *Serendipita vermifera*), *Piriformospora indica* (syn.
148 *Serendipita indica*), and *Serendipita* sp. 407. We sampled representatives from the order
149 Auriculariales to capture the entire diversity of these sequences (*Oliveonia pauxilla*, *Auricularia*
150 *subglabra*, *Aporpium caryae*, and *Exidia glandulosa*).

151 2.2 Fungal Isolates

152 The 32 fungal samples used in this study were isolated from roots or protocorms (the
153 seedling stage) of endangered orchid species in areas spanning from Hawaii to Florida, with a
154 focus on the Midwest and the Florida Panther National Wildlife Refuge (Table 1). For the full
155 description of the isolation techniques used, see Zettler and Corey (2018). Briefly, root tissue

156 was surface-sterilized then placed in a petri dish with sterile water and finely diced with a
157 scalpel. Fungal Isolation Media (Clements et al., 1986) was poured on the diced root tissue and
158 left at ambient temperature. After 24-48 hours, the plates were examined with a dissecting
159 microscope to identify fungal growth. Mycelia were excised and placed on Difco Potato
160 Dextrose Agar (PDA; Becton Dickinson and Co., Sparks, MD, Mfr # BD 213400). Those fungi
161 with morphological characteristics consistent with fungi in the form genus *Rhizoctonia* as
162 identified in Currah et al., (1997) were retained for identification with ITS sequencing (Figure
163 1).

164 Fungi were grown in flasks with 75ml of full strength Difco Potato Dextrose Broth
165 (Difco Becton Dickinson and Co., Sparks, MD, Mfr # BD 254920) on a shaker table until there
166 was enough tissue for extraction. Depending on the isolate this took 2-6 weeks. Often multiple
167 flasks of each isolate were grown at one time to speed up this process. For extraction, the entire
168 contents of each flask was poured into a 150mL Polystyrene Bottle Top Filter 0.45um (Corning
169 Incorporated, Corning, USA, Cat # 430627) and washed with DI water. These samples were
170 weighed to determine how many samples could be processed from each sample (minimum of 0.2
171 grams filtered weight/tube). Fungi were isolated with either the Bacterial/Fungal DNA extraction
172 kit (Zymo Research, Irvine, USA, Cat # D6005, Lot # ZRC201856) according to manufacturer
173 protocol or a CTAB, phenol chloroform isoamyl procedure (Supplemental Figure S1). When the
174 Zymo kit was used, fungi were added to lysis tubes and put on bead beater for two rounds of four
175 minutes. If the CTAB extraction was employed, fungal tissue was ground with liquid Nitrogen in
176 ceramic mortar and pestle. Extracted DNA was assayed on a NanoDrop 2000 (ThermoFisher
177 Scientific, USA, cat # ND-2000) and on a Qubit 2.0 Fluorometer (ThermoFisher Scientific,
178 USA, cat # Q32866) with the Qubit double-stranded DNA High Sensitivity Assay kit

179 (ThermoFisher Scientific, USA, cat # Q32851). We followed JGI instruction for sample
180 submission by submitting approximately 500 ng of each sample in a total volume of 25-35 uL in
181 one 96-well plate provided by JGI.

182 **2.3 ITS sequencing for Species Identification**

183 To determine species identity, we sequenced the internal transcribed spacer (ITS) region
184 of the rDNA. We used the same DNA extraction methods referenced above. We used the primer
185 pairs ITS1/ITS4-Tul or ITS1-OF/ITS4-OF for isolates presumed to be *Tulasnella* as the ITS
186 sequences in this genus are highly divergent and not captured well with other primers (Taylor
187 and McCormick, 2008). For the genera *Ceratobasidium* and *Serendipita*, the general primers
188 ITS1/ITS4 or ITS1-OF/ITS4-OF were used and if these did not successfully amplify the ITS
189 region of *Serendipita* isolates the primer pair ITS3Seb/NL4 (Bellemain et al., 2010; Ray et al.,
190 2015; White et al., 1990). The amplified DNA was cleaned with the DNA Clean and
191 Concentrator-25 kit (Zymo Research, Irvine, USA, cat # D4033). These PCR products were
192 assessed on a 1.5% agarose gel and Sanger sequencing was performed at the University of
193 Missouri DNA Core Facility. These sequences were evaluated for confidence in base calling and
194 edited by trimming low quality bases from the beginning and end of each sequence in Geneious
195 9.1.8 (<http://www.geneious.com/>). These trimmed sequences were queried against NCBI's
196 default nucleotide-nucleotide database as well as the UNITE database for species identification
197 (Nilsson et al., 2019). These sequences were generated for the purpose of accurate species ID
198 before sending DNA samples for shallow genome sequencing.

199 200 **2.4 Shallow Genome Sequencing and Quality Control**

201 Shallow genome sequencing of 32 samples, quality control, and filtering were performed
202 at the Joint Genome Institute (JGI) under a Community Sequencing Proposal (#2000). Samples

203 were run on an Illumina NovaSeq with 2x151 base pair (bp) reads. The quality control and
204 filtering at the JGI use BBmap to remove contamination and remove low quality reads (Bushnell
205 B., BBMap. <http://sourceforge.net/projects/bbmap/>). Three samples were sequenced at the
206 University of Missouri’s DNA Core Facility which were run on an Illumina NextSeq 500
207 machine on one lane with 45 other samples generating 2x150 bp reads.

208 **2.5 Shallow Genome Assembly and Annotation**

209 All cleaned and filtered sequences from the Joint Genome Institute and the University of
210 Missouri were assembled with the AAFTF pipeline for read assembly, remove vector
211 contamination and duplicate contigs, contig sequence polishing and sorting the contigs by length
212 (Stajich, JE., Automatic Assembly For the Fungi. <https://github.com/stajichlab/AAFTF>). The
213 pipeline performs assembly with Spades 3.10.0 using default parameters which consider 3 kmer
214 values and select the optimal assembly based on summary statistics (Nurk et al., 2013). As a
215 measure to assess genome completeness, all samples were run through BUSCO 3.0.2 using the
216 Basidiomycota database (Simao et al., 2015). For most samples, RNA sequence data was used to
217 facilitate annotation. When samples were too distantly related to map efficiently to the RNA
218 sequencing reads, these taxa were annotated without aligning to the RNA sequences (Table 5).
219 The RNA sequences used for reference were also generated from JGI CSP #2000 and will be
220 published as part of a separate study.

221 All samples were then prepared for gene prediction using Funannotate 1.6.0 (Palmer JP,
222 Stajich JE. 2018, <https://github.com/nextgenusfs/funannotate>), which performs all the steps
223 necessary for genome annotation from gene prediction training to final gene consensus model,
224 functional prediction, and dataset preparation for deposition into GenBank. The tool first runs
225 RepeatMasker 4.0.7 (<http://www.repeatmasker.org>). This “softmasks” the genome by converting

226 repetitive elements into lowercase letters in the assembly files. This step is necessary for the gene
227 prediction steps that follow. After masking, each assembly is run through a training step to
228 provide the initial models for the *ab initio* gene prediction programs AUGUSTUS 3.3.0 (Keller
229 et al., 2011; Stanke and Waack, 2003), SNAP (Korf, 2004), CodingQuarry (Testa et al.,
230 2015), and GeneMark-ES/ET 4.38.0 (Lomsadze et al., 2014). Protein sequences are also aligned
231 with diamond (Buchfink et al., 2015) and gene models polished with exonerate (Slater and
232 Birney, 2005). When RNASeq reads were available for a strain, these were applied as part of a
233 training step which first aligned short RNASeq reads, followed by assembly of these reads into
234 contig with Trinity. Finally these assembled transcripts were aligned to the genome to produce
235 gene models which were used for gene predictor training. Table 5 has the strains which were
236 able to use the RNASeq data as support for gene model training and prediction. These combined
237 evidence of these gene predictions, both *ab initio* and protein and transcript sequence based,
238 were combined with EvidenceModeler to use combined evidence to predict a final set of protein
239 coding genes. In addition tRNA gene predictions were performed with tRNAScan-SE (Lowe and
240 Eddy, 1997). The resulting predicted protein files were then used for the phylogenetic analyses.

241 **2.6 Phylogenomic analysis**

242 We used the pipeline PHYling 1.0 (<https://doi.org/10.5281/zenodo.1257001>) developed
243 by the Stajich lab, to extract orthologous genes from the predicted proteins of our taxa (Spatafora
244 et al., 2016). PHYling uses Hmmer3 (v3.2.1) to compare our predicted proteins to a list of
245 Profile-Hidden-Markov models of phylogenetically informative markers. The list we used is the
246 434 orthologous gene set (<https://doi.org/10.5281/zenodo.1251476>) constructed by the 1000
247 Fungal Genomes Project and identified as single-copy in orthologous gene clusters available
248 from the Joint Genome Institute's MycoCosm repository (Grigoriev et al., 2014). We used

249 hmmsearch to compare each sample's proteome to the 434 gene list. The protein sequence
250 homologs we identified were aligned to the marker-profile HMM with hmalign. These
251 alignments were concatenated to run a phylogenetic analysis with RAxML 8.2.12 (Stamatakis,
252 2006; Stamatakis et al., 2008). The model of evolution was determined automatically and
253 bootstrapped with 100 replicates. The gene trees generated from RAxML were used to construct
254 a consensus tree with ASTRAL-III 5.6.3 (Mirarab et al., 2014; Zhang et al., 2017).

255 **2.7 Data accessibility**

256 Isolates with UAMH numbers are stored in the UAMH Centre for Global Microfungal
257 Biodiversity repository. Raw DNA sequence data have been deposited in SRA and are associated
258 with BioProjects listed in Table 3. Scripts used for these analyses and all alignments, trees, and
259 intermediate files will be made available in a Dryad repository upon publication. BioProject IDs
260 and JGI Mycocosm repositories are summarized in Table 3.

261

262

3. RESULTS

263

3.1 ITS identifications

264

265

266

267

268

For the 35 isolates studied, ITS identifications, primers used and the length of each
sequence are summarized in Table 2. One sample sent to the Joint Genome Institute was not
sequenced due to poor DNA quality. Two isolates were identified as contaminants (isolates 420
and 422) and were excluded from further analysis (Table 2). Only four out of 35 isolates were
identified to species.

269

3.2 Shallow genome sequencing and annotation

270

271

272

Shallow genome sequencing of 32 fungal isolates resulted in a wide range in the number
of genes annotated in each individual genome. The isolate *Serendipita sp* 396 has the least
number of annotated genes at 8,285 and *Ceratobasidium sp* 428 has the most at 25,099. The

273 BUSCO completeness scores ranged from 54.2% to 96.6% of the 1335 orthologues in the
274 BUSCO dataset. For assembly statistics see Table 4 and for BUSCO completeness scores see
275 Table 5. Out of 435 orthologous genes, 429 had enough significant hits for further analysis. The
276 number of genes present for each taxa ranged from 299 in *Tulasnella sp* 408 to 425 in the
277 outgroups *Auricularia subglabra* and *Botryobasidium botryosum*. For full matrix occupancy see
278 Table 6. The outgroup *Kockovaella imperatae* contained 408 of the 429 genes so those 408
279 sequences were included in the phylogenetic analyses. The concatenated alignment has 128,774
280 distinct alignment patterns and is 14.31% gaps.

281 **3.3 Phylogenetic analysis**

282 The best concatenated tree likelihood is -3406977.36. The bootstrap (BS) support is
283 overall very high with the majority of branches at 100 (Figure 2). Eight branches have bootstrap
284 values below 100, and, of those, only three are below 75. The ASTRAL-III tree shows high
285 congruence with the concatenated tree and all but five branches are supported with 0.7 local
286 posterior probability or higher (Figure 3). The two phylogenies have the exact same topology on
287 the class, order, and family level and recapitulate with high support previously published
288 relationships between orders in the Agaricomycetes [Cantharellales, [Sebacinales,
289 [Auriculariales]]]. The phylogenies are highly congruent within Cantharellales, however, the
290 relationships between *Serendipita* isolates are quite different as discussed below.

291 Within the Cantharellales, we have strong support (94 BS, 0.99 posterior probability) for
292 Ceratobasidiaceae as sister to the rest of the order. Within Ceratobasidiaceae, the *Ceratobasidium*
293 isolates cluster together with very strong support with the exception of *Ceratobasidium sp* 423,
294 which is nested within *Rhizoctonia solani* and *Thanatephorus cucumeris*. The only difference
295 between the ML and ASTRAL-III in the family is the placement of *Ceratobasidium sp* 370. In

296 the ML tree, 370 is sister to a clade of [414, [394+UAMH11750]] and in the ASTRAL-III tree,
297 370 is sister with isolate 414 and equally related to 394+UAMH11750. There is no phylogenetic
298 signal based on orchid source, geographic location (Figure 3, Table 1). Both trees show
299 Tulasnellaceae as sister to the clade [Botryobasidium, [Clavulina, [Cantherellus + Hydnum]]]
300 with 100BS and 1.0 pp. The relationships in *Tulasnella* are highly supported with all but one
301 branch with 100 BS values and all but two branches with pps less than 1.0. Notably, the genome
302 sequence and the shallow genome sequence data for *Tulasnella calospora* UAMH 9824 are sister
303 to each other in the tree, though two other isolates are included in a clade with *Tulasnella*
304 *calospora* AL13.

305 The samples in the Sebacinales are not as well-resolved. The *Serendipita* isolates have the
306 least support overall due to the short branches of all isolates aside from *Serendipita* 399, which is
307 sister to the rest. All *Serendipita* spp in this study are most closely related to *Serendipita*
308 (=Piriformospora) *indica* with 100 BS/1.0. It is important to note our inclusion of the reference
309 genome *Serendipita* 407 (*Serendipita* sp._407_v1.0) and a shallow genome sequence of the same
310 isolate (*Serendipita*_sp_407.Orchid). In our dataset these two samples are not sister to each other.
311 In the quartet-based ASTRAL-III tree, *Serendipita* 400 and 411 are sister to each other with 0.77
312 posterior probability, whereas in the concatenated tree, the genome of isolate 407 was sister to
313 the rest of the *Serendipita* isolates aside from 399. The short branches in this group indicate a
314 small number of changes in the alignment in the ML tree and a high degree of discordance in the
315 ASTRAL-III tree. All of the *Serendipita* isolates are from epiphytic orchids in the Florida
316 Panther National Wildlife Refuge (Figure 3, Table 1). In the ASTRAL tree, *Serendipita* spp tend
317 to cluster with orchid source compared to the ML tree.

318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341

4. DISCUSSION

4.1 Overview

The primary goal of this study was to use shallow genome sequencing and phylogenetic methods to uncover the evolutionary relationships in a collection of fungal isolates that interact with endangered orchid species. The secondary goal was to leverage current genomic resources to investigate relationships among the orders, families and genera of Agaricomycetes, with a focus on Ceratobasidiaceae, Tulasnellaceae, and Sebacinaceae. Understanding of species in the fungal genera that facilitate orchid germination is extremely poor, as the number of formally described species is much lower than the diversity of fungi revealed from metagenomic or environmental sequencing. The results of this study add to our understanding of the genetic diversity of these fungal taxa and provide an example of how sequence data can be incorporated with taxonomic expertise to better describe fungal species.

The fungi that help germinate orchids were first categorized under one “form genus” called *Rhizoctonia* (Currah et al., 1997). This classification is not phylogenetically informative and today we know many orchid symbionts come from two orders (Cantharellales and Sebaciales) in the class Agaricomycetes (Hibbett, 2006). However, the taxonomy remains to be fully resolved. One reason classification can be difficult in these taxa is that these isolates do not sporulate or make sexual structures in laboratory conditions. Another is that traditionally, fungi were classified under two different names – the sexual stage (teleomorph) or vegetative state (anamorph). This policy ended during the 2011 International Botanical Congress when the Nomenclature Section voted to eliminate this dual nomenclature system (Hibbett and Taylor, 2013). Many of the names published in literature are no longer considered the correct taxonomy though in many cases these changes are not strongly reinforced. This study examines the

342 phylogenetic relationships of a collection of isolates so that the genetic distance of these strains
343 is known and to provide a framework for future evolutionary questions. Data from these
344 phylogenies can also provide evidence for new species or to revise current species concepts.
345 Understanding of taxonomy and species relationships is critical for testing evolutionary
346 hypotheses. Increased sampling within taxonomic groups and from sites around the globe is
347 necessary for future studies.

348

349 **4.2 Relationships among Orders and Families**

350 We used shallow genome sequencing for phylogenomics to describe the evolutionary
351 relationships among a collection of orchid mycorrhizal fungi. We also included numerous
352 outgroups to span the amount of biodiversity represented by these fungi. The large number of
353 coding genes allowed us to provide strong evidence for relationships between orders and a novel
354 result within the families of Cantharellales. Our results show strong support for the relationships
355 [Cantharellales, [Sebacinales, [Auriculariales]]]. This is consistent with previously reported
356 studies (Nagy et al., 2016). Within Cantharellales, the taxonomy is less certain and is still
357 undergoing changes. For example, Dictionary of the Fungi lists seven families while Hibbett et
358 al., (2014) claim four by defining Clavulinaceae and Cantharellaceae as synonymous with
359 Hydnaceae. This decision seems to be based on the authors' interpretations as the data in the
360 papers they cite don't support this conclusion (Leacock 2018). González et al., (2016), found
361 some support for the relationships [Tulasnellaceae, [Ceratobasidiaceae +Botryobasidiaceae,
362 [Hydnaceae]]] based on the markers ITS-LSU, rpb2, tef1, and atp6. They did state that multiple
363 coding genes would be necessary to see if their result was robust (González et al., 2016). Our
364 results show strong support (99 BS and .94 posterior probability) for Ceratobasidiaceae as the

365 sister family to [Tulasnellaceae, [Botryobasidiaceae + rest of Cantharellales]]. We did only
366 include one sample from the four groups besides Ceratobasidiaceae and Tulasnellaceae so more
367 sampling is needed in this group of fungi to produce a robust and consistent phylogenetic
368 inference.

369 **4.3 Relationships in Ceratobasidiaceae**

370 The *Ceratobasidium* samples are closely related with the exception of isolate
371 *Ceratobasidium* sp 423 that is nested within *Rhizoctonia solani* and *Thanatephorus cucumeris*
372 (Figures 2, 3). These results are consistent with Veldre et al., (2013), who found that the genera
373 *Ceratobasidium* and *Thanatephorus* are polyphyletic. Given the type specimen for
374 *Ceratobasidium* has since been placed in the Auriculariales, Oberwinkler et al., (2013a)
375 recommended *Ceratobasidium* should be renamed *Rhizoctonia*. Given these taxonomic
376 conundrums, attention is needed to make a robust classification system. Something we found
377 affirming was the close relationship of isolates *Ceratobasidium* 11750 and *Ceratobasidium* 394.
378 Based on a nearly identical ITS sequence alignment, these isolates were assumed to be very
379 closely related. This result is noteworthy because they have differential abilities to germinate
380 seeds from the endangered Ghost orchid, *Dendrophylax lindenii*. 394 can germinate seeds but
381 379 does not. More sampling is needed to compare how the isolates included in our study are
382 related to other *Ceratobasidium* spp. that are in defined Anastomosis Groups.

383

384 **4.4 Relationships in Tulasnellaceae**

385 Our *Tulasnella* isolates show a well-supported monophyletic clade in both phylogenetic
386 trees (Figures 2 and 3). Without further targeted sampling, it is premature to delimit species
387 boundaries; however, one species that could use revision is *Tulasnella calospora*. In both the

388 concatenated and coalescent phylogenies, the two *T. calospora* genomes are not sister to each
389 other but include the isolates 408 and 417, which were not identified as *T. calospora* based on
390 the ITS sequence. This result could be a function of the relatively low number of orthologous
391 genes that we recovered from 408 and 417, 291 and 330 out of 434, respectively (Tables 5 and
392 6). However, others have voiced concern over the species concept (Melissa McCormick, pers.
393 comm.).

394 Three isolates in this analysis are from the Hawaiian island of Molokai (330, 331, and
395 332; Table 1). These isolates cluster very closely in both phylogenies and are sister to three
396 isolates of *Tulasnella inquilina*. These isolates turn pink when exposed to light and have highly
397 divergent ITS sequences from the other *Tulasnella* isolates in this analysis. The strong support
398 for the monophyly of these Hawaiian samples, and their placement in the tree, suggest a
399 potentially new species. With increased sampling, more robust methods to delineate species
400 boundaries such as those used in (Whitehead et al., 2017) and we will have the power to better
401 describe the diversity of orchid mycorrhizal fungi.

402 **4.5 Relationships in Sebacinales**

403 All of the *Serendipita* isolates in this analysis are from the Florida National Wildlife
404 Panther Refuge (NWPR) in Florida and they are associated with three different epiphytic orchid
405 species (Table 1). In both phylogenetic analyses, *Serendipita* 399 is sister to the rest of our
406 samples. Growing on PDA, 399 looks morphologically distinct from the other *Serendipita sp* due
407 to a darker orange pigment and a crustose layer on the surface of the agar. This isolate also
408 grows much more slowly than other *Serendipita* taxa, it would take longer than four weeks for
409 the fungus to grow to the edge of a standard petri dish. For the remaining samples, it could be,
410 that there is one main species or population of *Serendipita* that grows in orchid roots in the

411 NWPR as their relationships are poorly resolved in the RAxML phylogeny and highly
412 incongruent between the two phylogenies. However, in the ASTRAL analysis, the *Serendipita*
413 isolates cluster somewhat closely by the orchid species from which they were isolated though
414 this is not a strong signal (Figure 3). A more thorough and targeted analysis is required to
415 determine the number of distinct populations of these fungi in the Florida National Panther
416 Wildlife Refuge similar to that conducted by Ruibal et al., (2017) to describe the population
417 structure of *Tulasnella prima* in Australia. It would be interesting to survey the fungi growing in
418 the roots of all plant species in the NWPR to determine the genetic diversity of *Serendipita*
419 across the landscape. Such an experiment would show whether orchids are using a narrow
420 distribution of fungi or if the plants are less discerning but the genetic diversity of the fungi is
421 simply very low.

422 Another result from our analysis shows that these fungal strains are most closely related
423 to *Piriformospora indica*, a known ectomycorrhizal fungus species (Varma et al., 2001). Many
424 fungi in the order Sebaciales are ecologically characterized as ectomycorrhizal fungi and
425 interact with a wide diversity of plant species (Kohler et al., 2015). Indeed, researchers are
426 isolating fungi in the Sebaciales from plants like switchgrass (*Panicum virgatum*) to determine
427 the benefit of these fungi for applications in agriculture (Craven and Ray, 2019). Orchids might
428 contribute to this effort, as it took more than one year for the Craven lab to isolate one strain of
429 *Sebacina vermifera ssp. bescii* from switchgrass; similar fungi are much more easy to isolate
430 from orchid roots (Prasun Ray, pers. comm.). Orchids could be environmental filters for fungi
431 that could be beneficial in many plant-fungal interactions.

432

433 **4.6 Future directions**

434 The next steps stemming from this study are to combine the phylogenetic relationships
435 with taxonomic expertise to name new species or to revisit problematic species concepts like
436 *Tulasnella calospora*. Additionally, it would be beneficial to sequence the genome of the type
437 specimens for many of these genera and species. Being able to compare the genetic sequences of
438 the type specimens would be extremely beneficial for fungal species that do not present sexual
439 characteristics in the lab. A set of fifteen isolates from the collection have been sequenced on the
440 PacBio platform and will be assembled into reference genomes as part of another aim of the
441 Community Sequencing Proposal (Table 3).

442

443

ACKNOWLEDGEMENTS

444

445 I thank the Zettler lab students who worked in the field and the lab to isolate these beautiful
446 fungi. We are grateful to the teams of Dr. Greg Bonito, Dr. Daniel Lindner, and Dr. Francis
447 Martin including the ‘Mycorrhizal Genomics Initiative’ consortium and the 1KFG project for
448 access to unpublished genome data. The genome sequence data were produced by the US
449 Department of Energy Joint Genome Institute in collaboration with the user community. The
450 work conducted by the U.S. Department of Energy Joint Genome Institute, a DOE Office of
451 Science User Facility, is supported by the Office of Science of the U.S. Department of Energy
452 under Contract No. DE-AC02-05CH11231. Computations were performed using the computer
453 clusters and data storage resources of the University of California-Riverside HPCC, which were
454 funded by grants from NSF (MRI-1429826) and NIH (1S10OD016290-01A1). Funding for this
455 work is from Joint Genome Institute Community Sequencing Proposal (JGI CSP 2000), National

456 Science Foundation, IOS 1339156, and the Department of Energy (DOE), Defense Threat
457 Reduction Agency (DTRA).
458
459

LITERATURE CITED

- 460
461
462 Barto, E.K., Weidenhamer, J.D., Cipollini, D., Rillig, M.C., 2012. Fungal superhighways: do
463 common mycorrhizal networks enhance below ground communication? *Trends in Plant*
464 *Science* 17, 633-637.
- 465 Bellemain, E., Carlsen, T., Brochmann, C., Coissac, E., Taberlet, P., Kauserud, H., 2010. ITS as
466 an environmental DNA barcode for fungi: an in silico approach reveals potential PCR
467 biases. *BMC microbiology* 10, 189.
- 468 Blackwell, M., 2011. The Fungi: 1, 2, 3 ... 5.1 million species? *American Journal of Botany* 98,
469 426-438.
- 470 Buchfink, B., Xie, C., Huson, D.H., 2015. Fast and sensitive protein alignment using
471 DIAMOND. *Nature Methods* 12, 59-60.
- 472 Bushnell, B., 2014. BBmap.
- 473 Clements, M., Muir, H., Cribb, P., 1986. A preliminary report on the symbiotic germination of
474 European terrestrial orchids. *Kew Bulletin*, 437-445.
- 475 Craven, K.D., Ray, P., 2019. More than Serendipity: The Potential to Manage Soil Carbon and
476 Emissions While Promoting Low-Input Agriculture with Serendipitoid Mycorrhizae.
477 *Phytobiomes Journal* 3, 161-164.
- 478 Currah, R., Zelmer, C., Hambleton, S., Richardson, K., 1997. Fungi from orchid mycorrhizas.
479 *Orchid Biology*. Springer, pp. 117-170.
- 480 Fochi, V., Chitarra, W., Kohler, A., Voyron, S., Singan, V.R., Lindquist, E.A., Barry, K.W.,
481 Girlanda, M., Grigoriev, I.V., Martin, F., Balestrini, R., Perotto, S., 2017a. Fungal and
482 plant gene expression in the *Tulasnella calospora*-*Serapias vomeracea* symbiosis provides
483 clues about nitrogen pathways in orchid mycorrhizas. *New Phytol* 213, 365-379.
- 484 Fochi, V., Falla, N., Girlanda, M., Perotto, S., Balestrini, R., 2017b. Cell-specific expression of
485 plant nutrient transporter genes in orchid mycorrhizae. *Plant Sci* 263, 39-45.
- 486 González, D., Rodríguez-Carres, M., Boekhout, T., Stalpers, J., Kuramae, E.E., Nakatani, A.K.,
487 Vilgalys, R., Cubeta, M.A., 2016. Phylogenetic relationships of Rhizoctonia fungi within
488 the Cantharellales. *Fungal Biol* 120, 603-619.
- 489 Grigoriev, I.V., Nikitin, R., Haridas, S., Kuo, A., Ohm, R., Otilar, R., Riley, R., Salamov, A.,
490 Zhao, X., Korzeniewski, F., Smirnova, T., Nordberg, H., Dubchak, I., Shabalov, I., 2014.
491 MycoCosm portal: gearing up for 1000 fungal genomes. *Nucleic Acids Research* 42,
492 D699-D704.
- 493 Hawksworth, D.L., Lücking, R., 2017. Fungal Diversity Revisited: 2.2 to 3.8 Million Species.
494 *The Fungal Kingdom*. American Society of Microbiology.
- 495 Hibbett, D.S., 2006. A phylogenetic overview of the Agaricomycotina. *Mycologia* 98, 917-925.
- 496 Hibbett, D.S., Bauer, R., Binder, M., Giachini, A.J., Hosaka, K., Justo, A., Larsson, E., Larsson,
497 K.H., Lawrey, J.D., Miettinen, O., Nagy, L.G., Nilsson, R.H., Weiss, M., Thorn, R.G.,
498 2014. 14 Agaricomycetes. In: McLaughlin, D.J., Spatafora, J.W. (Eds.), *Systematics and*
499 *Evolution: Part A*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 373-429.
- 500 Hibbett, D.S., Taylor, J.W., 2013. Fungal systematics: is a new age of enlightenment at hand?
501 *Nat Rev Microbiol* 11, 129-133.
- 502 Jung, S.C., Martinez-Medina, A., Lopez-Raez, J.A., Pozo, M.J., 2012. Mycorrhiza-Induced
503 Resistance and Priming of Plant Defenses. *Journal of Chemical Ecology* 38, 651-664.
- 504 Keller, O., Kollmar, M., Stanke, M., Waack, S., 2011. A novel hybrid gene prediction method
505 employing protein multiple sequence alignments. *Bioinformatics* 27, 757-763.

506 Kirk, P.M., Ainsworth, G.C., Bisby, G.R., 2008. Ainsworth & Bisby's dictionary of the fungi.
507 CABI.

508 Kohler, A., Kuo, A., Nagy, L.G., Morin, E., Barry, K.W., Buscot, F., Canback, B., Choi, C.,
509 Cichocki, N., Clum, A., Colpaert, J., Copeland, A., Costa, M.D., Dore, J., Floudas, D.,
510 Gay, G., Girlanda, M., Henrissat, B., Herrmann, S., Hess, J., Hogberg, N., Johansson, T.,
511 Khouja, H.R., LaButti, K., Lahrmann, U., Levasseur, A., Lindquist, E.A., Lipzen, A.,
512 Marmeisse, R., Martino, E., Murat, C., Ngan, C.Y., Nehls, U., Plett, J.M., Pringle, A.,
513 Ohm, R.A., Perotto, S., Peter, M., Riley, R., Rineau, F., Ruytinx, J., Salamov, A., Shah,
514 F., Sun, H., Tarkka, M., Tritt, A., Veneault-Fourrey, C., Zuccaro, A., Mycorrhizal
515 Genomics Initiative, C., Tunlid, A., Grigoriev, I.V., Hibbett, D.S., Martin, F., 2015.
516 Convergent losses of decay mechanisms and rapid turnover of symbiosis genes in
517 mycorrhizal mutualists. *Nat Genet* 47, 410-415.

518 Korf, I., 2004. Gene finding in novel genomes. *BMC Bioinformatics* 5, 59.

519 Kuga, Y., Sakamoto, N., Yurimoto, H., 2014. Stable isotope cellular imaging reveals that both
520 live and degenerating fungal pelotons transfer carbon and nitrogen to orchid protocorms.
521 *New Phytol* 202, 594-605.

522 Larsen, B.B., Miller, E.C., Rhodes, M.K., Wiens, J.J., 2017. Inordinate Fondness Multiplied and
523 Redistributed: the Number of Species on Earth and the New Pie of Life. *The Quarterly*
524 *Review of Biology* 92, 229-265.

525 Leacock, P.R., 2018. Cantharellales - MycoGuide.

526 Linde, C.C., May, T.W., Phillips, R.D., Ruibal, M., Smith, L.M., Peakall, R., 2017. New species
527 of *Tulasnella* associated with terrestrial orchids in Australia. *IMA Fungus* 8, 28-47.

528 Lomsadze, A., Burns, P.D., Borodovsky, M., 2014. Integration of mapped RNA-Seq reads into
529 automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Research* 42,
530 e119-e119.

531 Lowe, T.M., Eddy, S.R., 1997. tRNAscan-SE: A Program for Improved Detection of Transfer
532 RNA Genes in Genomic Sequence. *Nucleic Acids Research* 25, 955-964.

533 Lutzoni, F., Nowak, M.D., Alfaro, M.E., Reeb, V., Miadlikowska, J., Krug, M., Arnold, A.E.,
534 Lewis, L.A., Swofford, D.L., Hibbett, D., Hilu, K., James, T.Y., Quandt, D., Magallón,
535 S., 2018. Contemporaneous radiations of fungi and plants linked to symbiosis. *Nature*
536 *Communications* 9.

537 Martos, F., Munoz, F., Pailler, T., Kottke, I., Gonneau, C., Selosse, M.-A., 2012. The role of
538 epiphytism in architecture and evolutionary constraint within mycorrhizal networks of
539 tropical orchids. *Molecular Ecology* 21, 5098-5109.

540 McCormick, M.K., Taylor, D.L., Whigham, D.F., Burnett, R.K., 2016. Germination patterns in
541 three terrestrial orchids relate to abundance of mycorrhizal fungi. *Journal of Ecology* 104,
542 744-754.

543 Mirarab, S., Reaz, R., Bayzid, M.S., Zimmermann, T., Swenson, M.S., Warnow, T., 2014.
544 ASTRAL: genome-scale coalescent-based species tree estimation. *Bioinformatics* 30,
545 i541-i548.

546 Nagy, L.G., Riley, R., Tritt, A., Adam, C., Daum, C., Floudas, D., Sun, H., Yadav, J.S.,
547 Pangilinan, J., Larsson, K.-H., Matsuura, K., Barry, K., Labutti, K., Kuo, R., Ohm, R.A.,
548 Bhattacharya, S.S., Shirouzu, T., Yoshinaga, Y., Martin, F.M., Grigoriev, I.V., Hibbett,
549 D.S., 2016. Comparative Genomics of Early-Diverging Mushroom-Forming Fungi
550 Provides Insights into the Origins of Lignocellulose Decay Capabilities. *Molecular*
551 *Biology and Evolution* 33, 959-970.

552 Nilsson, R.H., Hyde, K.D., Pawłowska, J., Ryberg, M., Tedersoo, L., Aas, A.B., Alias, S.A.,
553 Alves, A., Anderson, C.L., Antonelli, A., Arnold, A.E., Bahnmann, B., Bahram, M.,
554 Bengtsson-Palme, J., Berlin, A., Branco, S., Chomnunti, P., Dissanayake, A., Drenkhan,
555 R., Friberg, H., Frøslev, T.G., Halwachs, B., Hartmann, M., Henricot, B., Jayawardena,
556 R., Jumpponen, A., Kauserud, H., Koskela, S., Kulik, T., Liimatainen, K., Lindahl, B.D.,
557 Lindner, D., Liu, J.-K., Maharachchikumbura, S., Manamgoda, D., Martinsson, S.,
558 Neves, M.A., Niskanen, T., Nylinder, S., Pereira, O.L., Pinho, D.B., Porter, T.M.,
559 Queloz, V., Riit, T., Sánchez-García, M., De Sousa, F., Stefańczyk, E., Tadych, M.,
560 Takamatsu, S., Tian, Q., Udayanga, D., Unterseher, M., Wang, Z., Wikee, S., Yan, J.,
561 Larsson, E., Larsson, K.-H., Kõljalg, U., Abarenkov, K., 2014. Improving ITS sequence
562 data for identification of plant pathogenic fungi. *67*, 11-19.

563 Nilsson, R.H., Larsson, K.-H., Taylor, A.F S., Bengtsson-Palme, J., Jeppesen, T.S., Schigel, D.,
564 Kennedy, P., Picard, K., Glöckner, F.O., Tedersoo, L., Saar, I., Kõljalg, U., Abarenkov,
565 K., 2019. The UNITE database for molecular identification of fungi: handling dark taxa
566 and parallel taxonomic classifications. *Nucleic Acids Research* 47, D259-D264.

567 Nurk, S., Bankevich, A., Antipov, D., Gurevich, A., Korobeynikov, A., Lapidus, A., Prjibelsky,
568 A., Pyshkin, A., Sirotkin, A., Sirotkin, Y., Stepanauskas, R., McLean, J., Lasken, R.,
569 Clingenpeel, S.R., Woyke, T., Tesler, G., Alekseyev, M.A., Pevzner, P.A., 2013.
570 *Assembling Genomes and Mini-metagenomes from Highly Chimeric Reads*. Springer
571 Berlin Heidelberg, Berlin, Heidelberg, pp. 158-170.

572 Oberwinkler, F., Riess, K., Bauer, R., Kirschner, R., Garnica, S., 2013a. Taxonomic re-
573 evaluation of the *Ceratobasidium-Rhizoctonia* complex and *Rhizoctonia butinii*, a new
574 species attacking spruce. *Mycological Progress* 12, 763-776.

575 Oberwinkler, F., Riess, K., Bauer, R., Selosse, M.-A., Weiß, M., Garnica, S., Zuccaro, A.,
576 2013b. Enigmatic Sebacinaleae. *Mycological Progress* 12, 1-27.

577 Palmer, J., Stajich, J.E., 2018. Funannotate.

578 Peterson, R.L., Massicotte, H.B., 2004. Exploring structural definitions of mycorrhizas, with
579 emphasis on nutrient-exchange interfaces. *Canadian Journal of Botany* 82, 1074-1088.

580 Rasmussen, H.N., Dixon, K.W., Jersakova, J., Tesitelova, T., 2015. Germination and seedling
581 establishment in orchids: a complex of requirements. *Ann Bot* 116, 391-402.

582 Ray, P., Ishiga, T., Decker, S.R., Turner, G.B., Craven, K.D., 2015. A Novel Delivery System
583 for the Root Symbiotic Fungus, *Sebacina vermifera*, and Consequent Biomass
584 Enhancement of Low Lignin COMT Switchgrass Lines. *BioEnergy Research* 8, 922-933.

585 Ruibal, M.P., Peakall, R., Foret, S., Linde, C.C., 2014. Development of Phylogenetic Markers for
586 *Sebacina* (Sebacinaceae) Mycorrhizal Fungi Associated with Australian Orchids.
587 *Applications in Plant Sciences* 2, 1400015.

588 Ruibal, M.P., Peakall, R., Smith, L.M., Linde, C.C., 2013. Phylogenetic and microsatellite
589 markers for *Tulasnella* (Tulasnellaceae) mycorrhizal fungi associated with Australian
590 orchids. *Appl Plant Sci* 1.

591 Ruibal, M.P., Triponez, Y., Smith, L.M., Peakall, R., Linde, C.C., 2017. Population structure of
592 an orchid mycorrhizal fungus with genus-wide specificity. *Scientific Reports* 7.

593 Simao, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V., Zdobnov, E.M., 2015.
594 BUSCO: assessing genome assembly and annotation completeness with single-copy
595 orthologs. *Bioinformatics* 31, 3210-3212.

596 Slater, G.S.C., Birney, E., 2005. Automated generation of heuristics for biological sequence
597 comparison. *BMC Bioinformatics* 6, 31.


- 598 Smith, S.E., Read, D.J., 2008. Mycorrhizal symbiosis. Academic press.
- 599 Spatafora, J.W., Chang, Y., Benny, G.L., Lazarus, K., Smith, M.E., Berbee, M.L., Bonito, G.,
600 Corradi, N., Grigoriev, I., Gryganskyi, A., James, T.Y., O'Donnell, K., Roberson, R.W.,
601 Taylor, T.N., Uehling, J., Vilgalys, R., White, M.M., Stajich, J.E., 2016. A phylum-level
602 phylogenetic classification of zygomycete fungi based on genome-scale data. *Mycologia*
603 108, 1028-1046.
- 604 Stajich, J.E., 2019. Automatic Assembly For the Fungi.
- 605 Stamatakis, A., 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with
606 thousands of taxa and mixed models. *Bioinformatics* 22, 2688-2690.
- 607 Stamatakis, A., Hoover, P., Rougemont, J., 2008. A Rapid Bootstrap Algorithm for the RAxML
608 Web Servers. *Systematic Biology* 57, 758-771.
- 609 Stanke, M., Waack, S., 2003. Gene prediction with a hidden Markov model and a new intron
610 submodel. *Bioinformatics* 19, ii215-ii225.
- 611 Taylor, D.L., McCormick, M.K., 2008. Internal transcribed spacer primers and sequences for
612 improved characterization of basidiomycetous orchid mycorrhizas. *New Phytol* 177,
613 1020-1033.
- 614 Testa, A.C., Hane, J.K., Ellwood, S.R., Oliver, R.P., 2015. CodingQuarry: highly accurate
615 hidden Markov model gene prediction in fungal genomes using RNA-seq transcripts.
616 *BMC Genomics* 16, 170.
- 617 Varma, A., Singh, A., Sudha, Sahay, N.S., Sharma, J., Roy, A., Kumari, M., Rana, D., Thakran,
618 S., Deka, D., Bharti, K., Hurek, T., Blechert, O., Rexer, K.H., Kost, G., Hahn, A., Maier,
619 W., Walter, M., Strack, D., Kranner, I., 2001. *Piriformospora indica*: An Axenically
620 Culturable Mycorrhiza-Like Endosymbiotic Fungus. In: Hock, B. (Ed.), *Fungal*
621 *Associations*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 125-150.
- 622 Veldre, V., Abarenkov, K., Bahram, M., Martos, F., Selosse, M.-A., Tamm, H., Kõljalg, U.,
623 Tedersoo, L., 2013. Evolution of nutritional modes of Ceratobasidiaceae (Cantharellales,
624 Basidiomycota) as revealed from publicly available ITS sequences. *Fungal Ecology* 6,
625 256-268.
- 626 Wang, W., Shi, J., Xie, Q., Jiang, Y., Yu, N., Wang, E., 2017. Nutrient Exchange and Regulation
627 in Arbuscular Mycorrhizal Symbiosis. *Molecular Plant* 10, 1147-1158.
- 628 Weiss, M., Waller, F., Zuccaro, A., Selosse, M.A., 2016. Sebaciniales - one thousand and one
629 interactions with land plants. *New Phytol* 211, 20-40.
- 630 White, T.J., Bruns, T., Lee, S., Taylor, J., 1990. Amplification and direct sequencing of fungal
631 ribosomal RNA genes for phylogenetics. *PCR protocols: a guide to methods and*
632 *applications* 18, 315-322.
- 633 Whitehead, M.R., Catullo, R.A., Ruibal, M., Dixon, K.W., Peakall, R., Linde, C.C., 2017.
634 Evaluating multilocus Bayesian species delimitation for discovery of cryptic mycorrhizal
635 diversity. *Fungal Ecology* 26, 74-84.
- 636 Yoder, J.A., Imfeld, S.M., Heydinger, D.J., Hart, C.E., Collier, M.H., Gribbins, K.M., Zettler,
637 L.W., 2010. Comparative water balance profiles of Orchidaceae seeds for epiphytic and
638 terrestrial taxa endemic to North America. *Plant Ecology* 211, 7-17.
- 639 Zettler, L.W., Corey, L.L., 2018. *Orchid Mycorrhizal Fungi: Isolation and Identification*
640 *Techniques*. *Orchid Propagation: From Laboratories to Greenhouses—Methods and*
641 *Protocols*. Springer, pp. 27-59.
- 642 Zhang, C., Sayyari, E., Mirarab, S., 2017. ASTRAL-III: Increased Scalability and Impacts of
643 Contracting Low Support Branches. Springer International Publishing, Cham, pp. 53-75.

644
645

646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673




Figure 1. Morphological examples of Tulasnella, Ceratobasidium, and Serendipita. One representative from each genus from the Zettler collection. All three isolates started growing on Potato Dextrose Agar on the same day as indicated by the date on the petri dish (25 November 2015). Photographs: Sarah Unruh.



674
675
676
677
678
679
680
681
682
683

Figure 2. Concatenation-based phylogeny of orchid mycorrhizal fungi.


Phylogenetic tree of the orchid mycorrhizal fungi in the Zettler collection with outgroups from the MycoCosm repository (genome.jgi.doe.gov/mycocosm/home). Alignments were made with the Phyling pipeline and the phylogeny was built with RAxML.



684
685
686
687
688
689
690

Figure 3. Quartet-based phylogeny of orchid mycorrhizal fungi.

Phylogenetic tree of the 32 orchid mycorrhizal fungi in the Zettler collection with 16 outgroups from the MycoCosm repository (genome.jgi.doe.gov/mycocosm/home). Alignments were made with the Phyling pipeline the gene trees were produced with RAxML and the tree was inferred using ASTRAL-III. All posterior probabilities are reported on the tree.



- ★ *Campylocentrum pachyrrhizum* ★ *Cypripedium candidum* ★ *Dendrophylax lindenii*
- ★ *Dendrophylax porrectus* ★ *Encyclia tampensis* ★ *Epidendrum amphistomum*
- ★ *Oeceoclades maculata* ★ *Platanthera holochila* ★ *Platanthera integrilabia*
- ★ *Platanthera ladera* ★ *Platanthera leucophaea* ★ *Platanthera paramoena*
- ★ *Polystachya concreta* ★ *Prosthechea cochleata* ★ *Spiranthes brevibrabris* ★ *Spiranthes vernalis*

691
692
693
694
695
696
697
698
699
700
701

Figure 4. Annotated Quartet-based phylogeny.

Phylogenetic tree of the 32 orchid mycorrhizal fungi in the Zettler collection with 18 genomes from the MycoCosm repository (genome.jgi.doe.gov/mycocosm/home). Branches were transformed in FigTree and annotated with colored stars indicating the origin they were isolated from.

Sample ID	Species	Strain	Orchid source	Tissue source	Location
Cerato11750	Ceratobasidium sp	UAMH11750	<i>Dendrophylax lindenii</i> (Lindl.) Benth. ex Rolfe	root	Florida Panther National Wildlife Refuge (NWR)
Cerato370	Ceratobasidium sp	370	<i>Platanthera leucophaea</i> (Nutt.) Lindl.	root	Tuscola Co., MI
Cerato392	Ceratobasidium sp	392	<i>Campylocentrum pachyrrhizum</i> (Rchb.f.) Rolfe	root	Florida Panther National Wildlife Refuge (NWR)
Cerato394	Ceratobasidium sp	394	<i>Dendrophylax lindenii</i> (Lindl.) Benth. ex Rolfe	root	Florida Panther National Wildlife Refuge (NWR)
Cerato395	Ceratobasidium sp	395	<i>Campylocentrum pachyrrhizum</i> (Rchb.f.) Rolfe	root	Florida Panther National Wildlife Refuge (NWR)
Cerato414	Ceratobasidium sp	414	<i>Platanthera lacera</i> (Michx.) G.Don	root	Fayette Co., IL
Cerato423	Ceratobasidium sp	423	<i>Spiranthes vernalis</i> Engelm. & A.Gray	root	Madison Co., IL
Cerato428	Ceratobasidium sp	428	<i>Dendrophylax porrectus</i> (Rchb.f.) Carlswald & Whitten	root	Florida Panther National Wildlife Refuge (NWR)
Serend396	Serendipita sp	396	<i>Prosthechea cochleata</i> (L.) W.E.Higgins	root	Florida Panther National Wildlife Refuge (NWR)
Serend397	Serendipita sp	397	<i>Prosthechea cochleata</i> (L.) W.E.Higgins	root	Florida Panther National Wildlife Refuge (NWR)
Serend398	Serendipita sp	398	<i>Prosthechea cochleata</i> (L.) W.E.Higgins	root	Florida Panther National Wildlife Refuge (NWR)
Serend399	Serendipita sp	399	<i>Encyclia tampensis</i> Small	root	Florida Panther National Wildlife Refuge (NWR)
Serend400	Serendipita sp	400	<i>Encyclia tampensis</i> Small	root	Florida Panther National Wildlife Refuge (NWR)
Serend401	Serendipita sp	401	<i>Epidendrum amphistomum</i> A.Rich	root	Florida Panther National Wildlife Refuge (NWR)
Serend405	Serendipita sp	405	<i>Prosthechea cochleata</i> (L.) W.E.Higgins	root	Florida Panther National Wildlife Refuge (NWR)
Serend407	Serendipita sp	407	<i>Epidendrum amphistomum</i> A.Rich	root	Florida Panther National Wildlife Refuge (NWR)
Serend411	Serendipita sp	411	<i>Prosthechea cochleata</i> (L.) W.E.Higgins	root	Florida Panther National Wildlife Refuge (NWR)
Tulasn330	Tulasnella sp	330	<i>Platanthera holochila</i> (Hillebr.) Kraenzl.	peloton	Molokai, HI
Tulasn331	Tulasnella sp	331	<i>Platanthera holochila</i> (Hillebr.) Kraenzl.	peloton	Molokai, HI
Tulasn332	Tulasnella sp	332	<i>Platanthera holochila</i> (Hillebr.) Kraenzl.	peloton	Molokai, HI
Tulasn403	Tulasnella sp	403	<i>Oeceoclades maculata</i> Lindl.	root	Florida Panther National Wildlife Refuge (NWR)
Tulasn408	Tulasnella sp	408	<i>Polystachya concreta</i> (Jacq.) Garay & H.R.Sweet	root	Florida Panther National Wildlife Refuge (NWR)
Tulasn417	Tulasnella sp	417	<i>Platanthera leucophaea</i> (Nutt.) Lindl.	root	McHenry Co., IL
Tulasn418	Tulasnella sp	418	<i>Platanthera leucophaea</i> (Nutt.) Lindl.	root	McHenry Co., IL
Tulasn419	Tulasnella sp	419	<i>Cypripedium candidum</i> Muhl. ex Willd.	protocorm/seedling	McHenry Co., IL
Tulasn424	Tulasnella sp	424	<i>Platanthera paramoena</i> A.Gray	root	Fayette Co., IL
Tulasn425	Tulasnella sp	425	<i>Platanthera paramoena</i> A.Gray	root	Fayette Co., IL
Tulasn427	Tulasnella sp	427	<i>Dendrophylax porrectus</i> (Rchb.f.) Carlswald & Whitten	root	Florida Panther National Wildlife Refuge (NWR)
Tulasn9824	Tulasnella calospora	UAMH9824	<i>Spiranthes brevilabris</i> Lindl.	root	Levy Co., FL
Tulinq235	Tulasnella inquilina	235	<i>Platanthera integrilabia</i> (Correll) Luer	root	McMinn Co., TN
Tulinq238	Tulasnella inquilina	238	<i>Platanthera integrilabia</i> (Correll) Luer	root	McMinn Co., TN
Tulinq7632	Tulasnella inquilina	UAMH7632	<i>Platanthera integrilabia</i> (Correll) Luer	root	Greenville, SC

Table 1. Description of fungal isolates.

UAMH numbers refer to the repository number for isolates deposited in the UAMH Centre for Global Microfungal Diversity

SampleID	Morphological ID	Top hit UNITE	Top hit GenBank	Primer sequenced	Length edited in base pairs
Cerato11750	<i>Ceratobasidium</i>	Ceratobasidiaceae	Uncultured <i>Ceratobasidium</i> clone LP8-Cer1	ITS1	641
Cerato370	<i>Ceratobasidium</i>	<i>Ceratobasidium</i>	<i>Ceratobasidium</i> UAMH 9847	ITS4	538
Cerato392	<i>Ceratobasidium</i>	Basidiomycota (same as ncbi)	orchid mycorrhizae KH4-8	ITS4	550
Cerato394	<i>Ceratobasidium</i>	<i>Ceratobasidium</i>	<i>Ceratobasidium</i>	ITS1	586
Cerato395	<i>Ceratobasidium</i>	Ceratobasidiaceae	<i>Ceratobasidium</i> sp JTO161	ITS1	548
Cerato414	<i>Ceratobasidium</i>	Ceratobasidiaceae	<i>Ceratobasidium</i> sp	ITS1	100
Cerato423	<i>Ceratobasidium</i>	<i>Ceratobasidium</i>	Uncultured Ceratobasidiaceae clone 207	ITS4	390
Cerato428	<i>Ceratobasidium</i>	Ceratobasidiaceae	<i>Ceratobasidium</i> sp JTO161	ITS1	420
Serend396	<i>Serendipita</i>	Sebacinales (orchid fungus)	uncultured Sebacinales clone	ITS1	189
Serend397	<i>Serendipita</i>	Sebacinales	uncultured Sebacinales clone	NL4	680
Serend398	<i>Serendipita</i>	Sebacinales	uncultured Sebacinales clone	NL4	567
Serend399	<i>Serendipita</i>	Sebacinales	uncultured Sebacinales clone	NL4	740
Serend400	<i>Serendipita</i>	Sebacinales	<i>Serendipita</i> sp MAFF 305831	NL4	780
Serend401	<i>Serendipita</i>	Sebacinales	Uncultured Sebacinales clone LP49-23S	ITS4	614
Serend405	<i>Serendipita</i>	<i>Serendipita</i>	<i>Serendipita</i> sp MAFF 305831	NL4	380
Serend407	<i>Serendipita</i>	Sebacinales	Uncultured <i>Sebacina</i> mycobiont of <i>Riccardia palmata</i>	*	2316
Serend411	<i>Serendipita</i>	Sebacinales	Uncultured Sebacinales clone LP49-23S	ITS3Seb	880

Tulasn330	<i>Tulasnella</i>	Tulasnellaceae	Uncultured Tulasnellaceae isolate 55P-Leu13	ITS4	570
Tulasn331	<i>Tulasnella</i>	Tulasnellaceae	Uncultured Tulasnellaceae	ITS1	620
Tulasn332	<i>Tulasnella</i>	Tulasnellaceae	Uncultured Tulasnellaceae	ITS4-OF	810
Tulasn403	<i>Tulasnella</i>	<i>Tulasnella</i>	<i>Tulasnella sp</i> CH01	ITS1	490
Tulasn408	<i>Tulasnella</i>	Tulasnellaceae	Uncultured Tulasnellaceae clone DOF-YC9	ITS4	622
Tulasn417	<i>Tulasnella</i>	<i>Tulasnella</i>	<i>Tulasnella sp</i> 9 MM-2012	ITS1	870
Tulasn418	<i>Tulasnella</i>	Tulasnellaceae	Uncultured Tulasnellaceae P94	ITS1	350
Tulasn419	<i>Tulasnella</i>	<i>Tulasnella</i>	Tulasnellaceae sp Pch 253	ITS4	368
Tulasn424	<i>Tulasnella</i>	Tulasnellaceae	Tulasnellaceae	ITS4	605
Tulasn425	<i>Tulasnella</i>	<i>Tulasnella</i>	<i>Tulasnella sp</i> 149	ITS1	570
Tulasn427	<i>Tulasnella</i>	Tulasnellaceae	Uncultured <i>Tulasnella</i> clone 998OF	ITS4	380
Tulasn9824	<i>Tulasnella calospora</i>	<i>Tulasnella calospora</i>	<i>Tulasnella calospora</i> isolate Pch-QS-0-1	ITS4-Tul	148
Tulinq235	<i>Epulorhiza inquilina</i>	<i>Tulasnella</i>	<i>Tulasnella sp</i> 3MV-2011 PA 053A	ITS4	650
Tulinq238	<i>Epulorhiza inquilina</i>	Tulasnellaceae	<i>Tulasnella sp</i> 3MV-2011 PA 053A	ITS4	758
Tulinq7632	<i>Epulorhiza inquilina</i>	<i>Tulasnella</i>	<i>Tulasnella sp</i> 3MV-2011 PA 053A	ITS1	790
Isolate 420*	<i>Tulasnella</i>	<i>Phanerochaete australis</i>	<i>Phanerochaete australis</i>	ITS1	350
Isolate 422*	<i>Tulasnella</i>	<i>Trichoderma petersenii</i>	<i>Trichoderma sp</i> isolate ARMI-23	ITS4	390

Table 2. Identifications of fungal isolates based on the internal transcribed spacer (ITS)

SampleID	BioProject or JGI web portal	BioSample
* <i>Ceratobasidium_sp_UAMH11750</i> .Orchid	PRJNA558776	SAMN12498506
<i>Ceratobasidium_sp_370</i> .Orchid	PRJNA557749	SAMN12427929
<i>Ceratobasidium_sp_392</i> .Orchid	PRJNA557750	SAMN12427914
* <i>Ceratobasidium_sp_394</i> .Orchid	PRJNA557751	SAMN12427897
* <i>Ceratobasidium_sp_395</i> .Orchid	PRJNA557752	SAMN12427926
<i>Ceratobasidium_sp_414</i> .Orchid	PRJNA557753	SAMN12427925
* <i>Ceratobasidium_sp_423</i> .Orchid	PRJNA557754	SAMN12427910
<i>Ceratobasidium_sp_428</i> .Orchid	PRJNA557755	SAMN12427923
<i>Serendipita_sp_396</i> .Orchid	PRJNA557757	SAMN12427894
<i>Serendipita_sp_397</i> .Orchid	PRJNA557758	SAMN12427928
<i>Serendipita_sp_398</i> .Orchid	PRJNA557759	SAMN12427900
<i>Serendipita_sp_399</i> .Orchid	PRJNA557760	SAMN12427906
* <i>Serendipita_sp_400</i> .Orchid	PRJNA557761	SAMN12427895
<i>Serendipita_sp_401</i> .Orchid	PRJNA557762	SAMN12427903
* <i>Serendipita_sp_405</i> .Orchid	PRJNA557763	SAMN12427917
* <i>Serendipita_sp_407</i> .Orchid	PRJNA558790	SAMN12498938
* <i>Serendipita_sp_411</i> .Orchid	PRJNA557734	SAMN12427911
<i>Tulasnella_sp_330</i> .Orchid	PRJNA557739	SAMN12427924
<i>Tulasnella_sp_331</i> .Orchid	PRJNA557740	SAMN12427908
* <i>Tulasnella_sp_332</i> .Orchid	PRJNA557741	SAMN12427902
<i>Tulasnella_sp_403</i> .Orchid	PRJNA557742	SAMN12427920
<i>Tulasnella_sp_408</i> .Orchid	PRJNA557743	SAMN12427916
<i>Tulasnella_sp_417</i> .Orchid	PRJNA557744	SAMN12427919
<i>Tulasnella_sp_418</i> .Orchid	PRJNA557745	SAMN12427921
* <i>Tulasnella_sp_419</i> .Orchid	PRJNA557746	SAMN12427922
<i>Tulasnella_sp_424</i> .Orchid	PRJNA557747	SAMN12427899
* <i>Tulasnella_sp_425</i> .Orchid	PRJNA557748	SAMN12427912
* <i>Tulasnella_sp_427</i> .Orchid	PRJNA557733	SAMN12427904
* <i>Tulasnella_calospora_UAMH9824</i> .Orchid	PRJNA558788	SAMN12498837
<i>Tulasnella_inquilina_235</i> .Orchid	PRJNA557736	SAMN12427891

* <i>Tulasnella_inquilina_238</i> .Orchid	PRJNA557737	SAMN12427893
* <i>Tulasnella_inquilina_UAMH7632</i> .Orchid	PRJNA557738	SAMN12427890
<i>Aporpium_caryae_L-13461</i> .	https://mycocosm.jgi.doe.gov/Elmca1	
<i>Auricularia_subglabra_v2.0</i>	https://mycocosm.jgi.doe.gov/Aurde3_1	
<i>Botryobasidium_botryosum_v1.0</i>	https://mycocosm.jgi.doe.gov/Botbo1	
<i>Calocera_cornea_v1.0</i>	https://mycocosm.jgi.doe.gov/Calco1	
<i>Cantharellus_anzutake_C23_v1.0</i>	https://mycocosm.jgi.doe.gov/Cananz1	
<i>Clavulina_sp._PMI_390_v1.0</i>	https://mycocosm.jgi.doe.gov/ClaPMI390	
<i>Exidia_glandulosa_v1.0</i>	https://mycocosm.jgi.doe.gov/Exigl1	
<i>Hydnum_rufescens_UP504_v2.0</i>	https://mycocosm.jgi.doe.gov/Hydru2	
<i>Kockovaella_imperatae_NRRL_Y-17943_v1.0</i>	https://mycocosm.jgi.doe.gov/Kocim1	
<i>Oliveonia_pauxilla_MPI-PUGE-AT-0066_v1.0</i>	https://mycocosm.jgi.doe.gov/Olipa1	
<i>Piriformospora_indica_DSM_11827_from_MPI</i>	https://mycocosm.jgi.doe.gov/Pirin1	
<i>Rhizoctonia_solani_AG-1_IB</i>	https://mycocosm.jgi.doe.gov/Rhiso1	
<i>Sebacina_vermifera_MAFF_305830_v1.0</i>	https://mycocosm.jgi.doe.gov/Sebbe1	
<i>Serendipita_sp._407_v1.0</i>	https://mycocosm.jgi.doe.gov/Serend1	
<i>Thanatephorus_cucumeris_MPI-SDFR-AT-0096_v1.0</i>	https://mycocosm.jgi.doe.gov/Thacu1	
<i>Tulasnella_calospora_AL13_4D_v1.0</i>	https://mycocosm.jgi.doe.gov/Tulca1	

Table 3. List of taxa and data availability.

Asterisks * indicate isolates selected for reference genome sequencing.

SampleID	CONTIG COUNT	TOTAL LENGTH	MIN	MAX	MEDIAN	MEAN	L50	N50	L90	N90
Ceratobasidium_sp_UAMH11750.Orchid	7239	47766782	2000	131072	4320	6598.53	1452	8784	5266	2947
Ceratobasidium_sp_370.Orchid	8028	47284605	2000	145775	3838	5889.96	1613	7304	5991	2715
Ceratobasidium_sp_392.Orchid	8904	52465834	1500	100768	3806	5892.39	1693	8316	6228	2516
Ceratobasidium_sp_394.Orchid	8562	53454716	1500	94250	3864	6243.25	1547	9440	5859	2601
Ceratobasidium_sp_395.Orchid	8769	67010313	1500	125675	4575	7641.73	1478	12298	5716	3127
Ceratobasidium_sp_414.Orchid	4161	50425407	1500	342430	4339	12118.58	349	34639	2143	4179
Ceratobasidium_sp_423.Orchid	15380	66434938	1500	107431	2946	4319.57	3306	5259	11578	2042
Ceratobasidium_sp_428.Orchid	8999	69097995	1500	121965	4552	7678.41	1493	12317	5876	3139
Serendipita_sp_396.Orchid	1431	20638744	2072	267195	9279	14422.6	279	17663	1083	6736
Serendipita_sp_397.Orchid	4253	28775535	1500	270096	3582	6765.94	580	10829	2793	2618
Serendipita_sp_398.Orchid	4302	28851264	1500	267525	3546	6706.48	574	11054	2835	2583
Serendipita_sp_399.Orchid	5013	31004825	1500	127831	3844	6184.88	906	8927	3450	2622
Serendipita_sp_400.Orchid	4392	28560853	1502	143991	3564	6502.93	635	10584	2927	2562
Serendipita_sp_401.Orchid	3823	28571286	1500	662216	3626	7473.52	429	13497	2433	2804
Serendipita_sp_405.Orchid	4254	28724145	1500	280457	3539	6752.27	566	10892	2796	2594
Serendipita_sp_407.Orchid	4028	27230538	1500	297010	3154	6760.31	426	12923	2590	2442
Serendipita_sp_411.Orchid	4211	28400685	1500	296955	3574	6744.4	574	10730	2767	2605
Tulasnella_sp_330.Orchid	1013	42302809	1500	967722	8117	41759.93	66	175815	323	19296
Tulasnella_sp_331.Orchid	3446	44512311	1500	298887	4965	12917.1	311	35916	1764	4762
Tulasnella_sp_332.Orchid	3329	44576393	1501	416704	5260	13390.33	297	36313	1699	5096

Tulasnella_sp_403.Orchid	2963	29964626	1502	501591	4201	10112.93	303	23340	1647	3550
Tulasnella_sp_408.Orchid	10591	63626598	1500	92931	3595	6007.61	1850	9042	7284	2490
Tulasnella_sp_417.Orchid	9866	61487333	1500	224139	3694	6232.25	1709	9469	6703	2528
Tulasnella_sp_418.Orchid	3880	32841821	1501	268652	3496	8464.39	411	18771	2277	2844
Tulasnella_sp_419.Orchid	3865	33665047	1500	229676	3923	8710.23	419	18466	2308	3120
Tulasnella_sp_424.Orchid	781	48431701	1507	1054277	13975	62012.42	71	195410	275	36329
Tulasnella_sp_425.Orchid	769	48399368	1507	1089567	16223	62938.06	74	189857	290	37534
Tulasnella_sp_427.Orchid	6018	39289859	1500	134256	3802	6528.72	956	9789	4051	2663
Tulasnella_calospora_UAMH9824.Orchid	4164	49802335	1500	345740	5668	11960.21	481	25557	2334	4747
Tulasnella_inquilina_235.Orchid	1742	44191844	1503	570014	3948	25368.45	119	102565	508	11931
Tulasnella_inquilina_238.Orchid	2874	45488573	1500	444022	4355	15827.62	240	54071	1242	5477
Tulasnella_inquilina_UAMH7632.Orchid	2898	46174977	1501	520136	4010	15933.39	209	56439	1211	5431

Table 4. Assembly statistics.

SampleID	RNASeq	Gene Count	BUSCO Complete %	BUSCO Single	BUSCO Fragmented	BUSCO Missing	BUSCO # Genes
Ceratobasidium_sp_UAMH11750.Orchid	Cerato379	16971	65.8	60.3	11.6	22.6	1335
Ceratobasidium_sp_370.Orchid	CeratoAll	11343	78.4	77.7	8.5	13.1	1335
Ceratobasidium_sp_392.Orchid	CeratoAll	14816	72.3	66.3	13.4	14.3	1335
Ceratobasidium_sp_394.Orchid	Cerato394	18818	71.1	62.6	13.6	15.3	1335
Ceratobasidium_sp_395.Orchid	Cerato395	19777	71.1	42.9	13.2	15.7	1335
Ceratobasidium_sp_414.Orchid	CeratoAll	12172	96.6	95.5	1.6	1.8	1335
Ceratobasidium_sp_423.Orchid	CeratoAll	13213	77.6	76.6	11.7	10.7	1335
Ceratobasidium_sp_428.Orchid	CeratoAll	25061	71.4	39.4	11.9	16.7	1335
Serendipita_sp_396.Orchid	Serend400	8272	65.5	64.9	5.8	28.7	1335
Serendipita_sp_397.Orchid	Serend400	12078	74.7	73.8	11.9	13.4	1335
Serendipita_sp_398.Orchid	Serend400	12311	74.6	73.6	11.4	14	1335
Serendipita_sp_399.Orchid	Serend400	11252	72.2	64.6	11.6	16.2	1335
Serendipita_sp_400.Orchid	Serend400	12369	72	70.9	11.3	16.7	1335
Serendipita_sp_401.Orchid	Serend400	11951	76.4	75.4	10	13.6	1335
Serendipita_sp_405.Orchid	Serend400	11992	73.3	72.6	10.7	16	1335
Serendipita_sp_407.Orchid	Serend400	11442	69	67.8	13.3	17.7	1335
Serendipita_sp_411.Orchid	Serend400	11996	74.5	73.2	10.6	14.9	1335
Tulasnella_sp_330.Orchid		9146	95.4	94.5	1.8	2.8	1335
Tulasnella_sp_331.Orchid		9039	92.9	91.7	2.8	4.3	1335
Tulasnella_sp_332.Orchid		9025	93	91.7	2.7	4.3	1335

Tulasnella_sp_403.Orchid	Tulinq7632	8272	77.7	77.2	10.3	12	1335
Tulasnella_sp_408.Orchid	Tulinq7632	15407	54.2	48.4	17.5	28.3	1335
Tulasnella_sp_417.Orchid		13362	62.4	56.7	15.3	22.3	1335
Tulasnella_sp_418.Orchid	Tulasn419	12415	86.3	85.5	6.1	7.6	1335
Tulasnella_sp_419.Orchid	Tulasn419	12897	88	87.3	6	6	1335
Tulasnella_sp_424.Orchid		11832	95.6	94.3	1.6	2.8	1335
Tulasnella_sp_425.Orchid		10834	96	94.4	1.4	2.6	1335
Tulasnella_sp_427.Orchid		11876	71.4	62	14.6	14	1335
Tulasnella_calospora_UAMH9824.Orchid	Tulinq7632	12307	88.9	87.3	5.4	5.7	1335
Tulasnella_inquilina_235.Orchid	Tulinq7632	13948	95.4	94.3	1.9	2.7	1335
Tulasnella_inquilina_238.Orchid	Tulinq7632	14664	94.1	92.5	2.6	3.3	1335
Tulasnella_inquilina_UAMH7632.Orchid	Tulinq7632	14741	94.2	92.5	2.3	3.5	1335

Table 5. Annotation and BUSCO completeness metrics.

Taxa without an RNA sequence listed did not sufficiently map to the Tulinq7632 RNA sequences and were annotated without expression data. The colors in the BUSCO complete % column range from blue-green (lowest percentage) to dark red (highest percentage).

Sample ID	Number best hit genes (429 total)
Ceratobasidium_sp_UAMH11750.Orchid	354
Ceratobasidium_sp_370.Orchid	356
Ceratobasidium_sp_392.Orchid	368
Ceratobasidium_sp_394.Orchid	376
Ceratobasidium_sp_395.Orchid	369
Ceratobasidium_sp_414.Orchid	387
Ceratobasidium_sp_423.Orchid	376
Ceratobasidium_sp_428.Orchid	382
Serendipita_sp_396.Orchid	311
Serendipita_sp_397.Orchid	391
Serendipita_sp_398.Orchid	371
Serendipita_sp_399.Orchid	358
Serendipita_sp_400.Orchid	375
Serendipita_sp_401.Orchid	382
Serendipita_sp_405.Orchid	379
Serendipita_sp_407.Orchid	364
Serendipita_sp_411.Orchid	382
Tulasnella_sp_330.Orchid	401
Tulasnella_sp_331.Orchid	401
Tulasnella_sp_332.Orchid	398
Tulasnella_sp_403.Orchid	376
Tulasnella_sp_408.Orchid	291
Tulasnella_sp_417.Orchid	330
Tulasnella_sp_418.Orchid	400
Tulasnella_sp_419.Orchid	408
Tulasnella_sp_424.Orchid	408
Tulasnella_sp_425.Orchid	403
Tulasnella_sp_427.Orchid	376
Tulasnella_calospora_UAMH9824.Orchid	398
Tulasnella_inquilina_235.Orchid	416
Tulasnella_inquilina_238.Orchid	410
Tulasnella_inquilina_UAMH7632.Orchid	417
Aporpium_caryae_L-13461.	423
Auricularia_subglabra_v2.0	425
Botryobasidium_botryosum_v1.0	425
Calocera_cornea_v1.0	410
Cantharellus_anzutake_C23_v1.0	412
Ceratobasidium_sp_AGI_v1.0	422

Clavulina_sp._PMI_390_v1.0	423
Exidia_glandulosa_v1.0	422
Hydnum_rufescens_UP504_v2.0	413
Kockovaella_imperatae_NRRL_Y-17943_v1.0	408
Oliveonia_pauxilla_MPI-PUGE-AT-0066_v1.0	419
Piriformospora_indica_DSM_11827_from_MPI	417
Rhizoctonia_solani_AG-1_IB	410
Sebacina_vermifera_MAFF_305830_v1.0	420
Serendipita_sp._407_v1.0	415
Thanatephorus_cucumeris_MPI-SDFR-AT-0096_v1.0	424
Tulasnella_calospora_AL13_4D_v1.0	405
Tulasnella_calospora_UAMH9824_v1.0	426

Table 6. Matrix Occupany.

Stajich lab CTAB Protocol:

Reagents required:

BUFFER A: 0.35 M sorbitol 0.1 M Tris-HCl, pH 9 5 mM EDTA, pH 8
BUFFER B: 0.2 M Tris-HCl, pH 9 50 mM EDTA, pH 8 2 M NaCl 2% CTAB
BUFFER C: 5% Sarkosyl (N-lauroylsarcosine sodium salt SIGMA L5125)
Potassium Acetate 5M (KAc precipitate polysaccharides) pH 7.5
RNase A (10 mg/ml) Proteinase K (20 mg/ml)
PVP 1 %
(PCI) Phenol:Chloroform:Isoamyl alcohol (25:24:1)
(CI) Chloroform:Isoamyl alcohol (24:1)
Sodium Acetate (NaAc) 3M
Isopropanol 100%
Ethanol 70%

1. Add Lysis Buffer (650 μ L Buffer A, 650 μ L Buffer B, 260 μ L Buffer C, 175 μ L .1% PVP, 10 μ L Proteinase K) to 2 mL microcentrifuge tube, mix, and split equally into two 2 mL tubes.
2. Place in hot plate and heat to 65° C.
3. Grind young fungal tissue in liquid nitrogen, add 50-100 mg of tissue to each tube.
4. Incubate 30 min at 65° mixing by inversion frequently (2-5 min).
5. Add 280 μ L KAc to each tube, mix by inversion, incubate on ice for 5 min.
6. Add 500-700 (the more the better) μ L PCI, mix by inversion (>5 min) or vortex briefly then incubate for 2 min at room temp (RT).
7. Spin at 6,000 g for 10 min
8. Take supernatant, add equal volume CI (usually about 1000ul).
9. Mix by inversion (>5 min) then incubate for 2 min.
10. Spin at 6,000 g for 10 min
11. Take supernatant (usually 700uL):
 - a. RNase treatment (2.5 μ L RNase, 37°, 90-120 min)*
 - b. Optional additional CI washes
12. Add 1/10 vol NaAc, mix, add 1 vol Isopropanol.
13. Incubate at RT 5 min, should start to see lots of DNA threads.
14. Spin at 3,000 g for 2 min, pour out the supernatant.
15. Wash with 1 mL freshly prepared, cold 70% ethanol.
16. Spin at 3,000 g for 2 min, pipette out the EtOH. Remove as much EtOH as possible before drying.
17. Dry pellet at RT for 10-15 min and/or 65° for <2 min to dry any leftover ethanol
 - a. Resuspend in 50-100 μ L TE (adjusted to pH9) at 65° Optional CI wash (add 600-800 TE buffer at 65°, resuspend DNA, add equal volume CI, mix as directed in step 9, carry on protocol from there minus the RNase and CI steps, I usually take 500-600 supernatant if added 800 uL CI).
18. Nanodrop, 260/280 is indicative of nucleic acid and 260/230 indicative of protein
19. Qubit
20. Run on Gel
21. Check ITS and 16S by PCR

Supplemental Figure S1. CTAB DNA extraction protocol from Stajich lab