

SOME NEW ALGORITHMS FOR HIGH-PRECISION COMPUTATION
OF EULER'S CONSTANT

Richard P. Brent and Edwin M. McMillan

January 1979

RECEIVED
LAWRENCE
BERKELEY LABORATORY


MAR 13 1979

LIBRARY AND
DOCUMENTS SECTION

Prepared for the U. S. Department of Energy
under Contract W-7405-ENG-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782*



LBL-8729c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

SOME NEW ALGORITHMS FOR HIGH-PRECISION COMPUTATION
OF EULER'S CONSTANT

by

*Richard P. Brent**

Department of Computer Science
Australian National University
Box 4, P.O., Canberra,
A.C.T. 2600.
AUSTRALIA

and

Edwin M. McMillan

Lawrence Berkeley Laboratory
University of California
Berkeley, California.
U.S.A.

* The work of the first author was supported in part by National Science Foundation grant 1-442427-21164-2 at the University of California, Berkeley. This work was also supported by the U. S. Department of Energy under Contract W-7405-ENG-48.

ABSTRACT:

We describe several new algorithms for the high-precision computation of Euler's constant $\gamma = 0.577\dots$. Using one of the algorithms, which is based on an identity involving Bessel functions, γ has been computed to 30,100 decimal places. By computing their regular continued fractions we show that, if γ or $\exp(\gamma)$ is of the form P/Q for integers P and Q , then $|Q| > 10^{15000}$.

AMS(MOS) Subject Classifications (1970)

Primary 10-04, 10A40; Secondary 10F20, 10F35, 65A05, 68A20.

Key Words and Phrases

Euler's constant, Mascheroni's constant, gamma, Bessel functions, rational approximation, regular continued fractions, multiple-precision arithmetic, Gauss-Kusmin law, arithmetic-geometric mean.

1. INTRODUCTION

Euler's constant γ is defined by

$$(1) \quad \gamma = \lim_{m \rightarrow \infty} (H_m - \ln(m)),$$

where

$$H_m = \sum_{k=1}^m 1/k .$$

Early computations of γ used the Euler-Maclaurin expansion to accelerate convergence of (1): see Brent [7] and Glaisher [12]. Sweeney [16] suggested a method which avoids the need for computation of the Bernoulli numbers which appear in the Euler-Maclaurin expansion, and Brent [7,8] used Sweeney's method to compute γ to 20,700 decimal places. There is some interest in computing γ as accurately as possible because it is not yet known whether γ is rational or irrational (although see Theorem 1 below).

In Section 3 we describe an algorithm which is about twice as fast as Sweeney's. The algorithm depends on some identities, given in Section 2, involving modified Bessel functions. To demonstrate the effectiveness of the algorithm we have used it to compute γ to 30,100 decimal places: see Section 5. Some other algorithms for the high-precision computation of γ are briefly described and compared in Section 4.

2. SOME BESSEL FUNCTION IDENTITIES

The modified Bessel functions $I_\nu(z)$ and $K_0(z)$ are defined by

$$I_\nu(z) = \sum_{k=0}^{\infty} \frac{(z/2)^{\nu+2k}}{k! \Gamma(\nu+k+1)}$$

and

$$K_0(z) = - \partial I_\nu(z) / \partial \nu |_{\nu=0} .$$

It is easy to verify, by carrying out the differentiation indicated above, that

$$(2) \quad \gamma + \ln(z/2) = \frac{S_0(z) - K_0(z)}{I_0(z)} ,$$

where

$$S_0(z) = \sum_{k=0}^{\infty} \frac{(z/2)^{2k}}{(k!)^2} H_k .$$

For real positive z , $K_0(z)$ and $I_0(z)$ have the asymptotic expansions

$$(3) \quad K_0(z) \sim \left(\frac{\pi}{2z} \right)^{1/2} e^{-z} \sum_{k=0}^{\infty} (-1)^k a_k(z)$$

and

$$(4) \quad I_0(z) \sim (2\pi z)^{-1/2} e^z \sum_{k=0}^{\infty} a_k(z) ,$$

where

$$a_k(z) = \frac{1^2 3^2 \dots (2k-1)^2}{k! (8z)^k} = \frac{[(2k)!]^2}{(k!)^3 (32z)^k} .$$

A proof is given in Watson [18, Sec. 7.23].

For real $z \geq 1$, the first terms in the asymptotic expansions (3) and (4) give upper and lower bounds on $K_0(z)$ and $I_0(z)$ respectively:

$$0 < K_0(z) < \left(\frac{\pi}{2z}\right)^{\frac{1}{2}} e^{-z}$$

and

$$I_0(z) > (2\pi z)^{-\frac{1}{2}} e^z.$$

Thus, taking $z = 2n \geq 2$ in (2), we have

$$(5) \quad 0 < U(n)/V(n) - \gamma = K_0(2n)/I_0(2n) < \pi e^{-4n},$$

where

$$(6) \quad U(n) = S_0(2n) - I_0(2n) \ln(n) = \sum_{k=0}^{\infty} \left(\frac{n^k}{k!}\right)^2 (H_k - \ln(n))$$

and

$$(7) \quad V(n) = I_0(2n) = \sum_{k=0}^{\infty} \left(\frac{n^k}{k!}\right)^2.$$

In the following section we describe an algorithm (B1) for computing γ using (5) to (7). It is interesting to note that the relations (2) to (4) were essentially given by Riemann [14] in 1855, but the possibility of using them to compute Euler's constant appears to have been overlooked.

3. THE ALGORITHM B1

Suppose we wish to evaluate γ to d decimal places.

If we choose

$$(8) \quad n = \lfloor c + \frac{1}{4} \ln(10)d \rfloor$$

for some suitable constant c then, from (5),

$$|\gamma - U(n)/V(n)| < \pi e^{4-4c} 10^{-d}.$$

Thus, we need only evaluate $U(n)$ and $V(n)$ sufficiently accurately, and then perform one high-precision division, to evaluate γ to the required accuracy.

Let

$$\begin{aligned} A_k &= \left(\frac{n^k}{k!}\right)^2 (H_k - \ln(n)), & U_k &= \sum_{j=0}^k A_j, \\ B_k &= \left(\frac{n^k}{k!}\right)^2, & V_k &= \sum_{j=0}^k B_j. \end{aligned}$$

Then

$$(9) \quad A_0 = -\ln(n), \quad B_0 = 1, \quad U_0 = A_0, \quad V_0 = 1,$$

and for $k = 1, 2, \dots$ we have

$$(10) \quad \left\{ \begin{aligned} B_k &= B_{k-1} n^2/k^2, \\ A_k &= (A_{k-1} n^2/k + B_k)/k, \\ U_k &= U_{k-1} + A_k, \\ V_k &= V_{k-1} + B_k. \end{aligned} \right.$$

For algorithm B1, n is chosen according to (8), and working precision equivalent to slightly more than d (floating) decimal places is used. $\ln(n)$ is computed, e.g. by the $O(d^2)$ method of [9], and A_0, B_0, U_0 and V_0 are initialized as in (9). The iteration (10) is terminated when, to the working precision, $U_k = U_{k-1}$ and $V_k = V_{k-1}$. The storage required is $O(d)$ as B_k can overwrite B_{k-1} , etc.

For $j \geq 0$, let α_j be the real positive root of

$$(11) \quad \alpha_j \ln \alpha_j - \alpha_j = j.$$

Thus $\alpha_0 = e \approx 2.78$, $\alpha_1 \approx 3.59$, $\alpha_2 \approx 4.32$, $\alpha_3 \approx 4.97$, etc.

Using Stirling's approximation, we see that the number of iterations of (10) required is

$$(12) \quad \begin{aligned} K &= \alpha_1 n + O(\ln(n)) \\ &= \frac{1}{4} \alpha_1 \ln(10) d + O(\ln(d)) \approx 2.07d. \end{aligned}$$

In analysing the time required by algorithm B1 and other algorithms described in Section 4, we make the following simplifying assumptions.

- (a) Only the time required for the inner loop(s) is considered (The computation of $\ln(n)$ is common to all the algorithms considered, so the time required for this is neglected. The final division of U_K by V_K takes time $O(d^2)$ if done as in [9], but the constant factor is relatively small, and $o(d^2)$ methods exist [5].)

- (b) Multiplication or division of a multiple-precision number (e.g. A_{k-1}) by a small integer (e.g. n^2 or k) takes time d units. In the analysis (though not in the implementation of the algorithm) the possibility of reducing the working precision (e.g. for A_k and B_k when $k \approx K$) is neglected. Considering this possibility complicates the analysis but is unlikely to alter the ranking of the algorithms discussed below.
- (c) Addition of two multiple-precision numbers takes time d units. (The constant is assumed to be the same as for (b). Again, this is unlikely to change the ranking of the algorithms).

Using these assumptions, each iteration of (10) requires time $8d$ (for 3 multiple-precision additions, 2 multiplications and 3 divisions by small integers). Thus, from (12), the time required by method B1 is about $2\alpha_1 \ln(10)d^2 \approx 16.5d^2$.

It is important to note that we avoid keeping H_k or $(H_k - \ln(n))$ as a multiple-precision number and multiplying by B_k in the inner loop. This would lead to a method with time $\Omega(d^3)$ if the classical multiplication algorithm were used as in [9]. The idea of using the Bessel function identities to compute γ was suggested by the second author, and the $O(d^2)$ implementation was discovered by the first author.

If terms in the sum (15) are grouped as in [5] and the Schönhage-Strassen fast multiplication algorithm [15] is used, it is possible to compute γ with error bounded by 10^{-d} in time $O\{d[\ln(d)]^3 \ln[\ln(d)]\}$, asymptotically faster than any of the $\Omega(d^2)$ algorithms considered here. However, such "fast" algorithms are very difficult to implement and are

slower than algorithm B1 unless d is very large. Thus, we do not consider them further.

4. RELATED ALGORITHMS

In this section we briefly describe and compare several closely related algorithms for the computation of γ .

4.1 Algorithm B2

From (2) we have

$$\gamma + \ln(n) = \frac{S_0(2n) - K_0(2n)}{I_0(2n)}$$

and, from (3),

$$(13) \quad K_0(2n) = \frac{1}{2}(\pi/n)^{\frac{1}{2}} e^{-2n} \sum_{k=0}^{4n} (-1)^k a_k(2n) + O(e^{-6n}/n)$$

Thus, we can find γ with error $O(n^{-\frac{1}{2}} e^{-8n})$ if $K_0(2n)$ is approximated using (13). If e^{2n} is computed using the Taylor series, and the time required to compute $(\pi/n)^{\frac{1}{2}}$ is neglected, the time required by this method (B2) is about

$$[(8\alpha_3 + 3\alpha_1 + 6)\ln(10)/8]d^2 \approx 16.3 d^2,$$

not appreciably less than for the simpler method B1.

4.2 Algorithm B3

To avoid the computation of $(\pi/n)^{\frac{1}{2}} e^{-2n}$ in (13), we may use the asymptotic series [1, eqn. (9.7.5)]

$$(14) \quad I_0(2n)K_0(2n) \sim \frac{1}{4n} \sum_{k=0}^{k'} \frac{[(2k)!]^3}{(k!)^4 (16n)^{2k}}$$

with $k' \leq 2n$. Empirical evidence suggests that the relative error in (14) with $k' = 2n$ is $O(n^{-1/2}e^{-4n})$, but we have not been able to prove this. Assuming this error bound, the time required with $k' = 2n$ is about

$$(\alpha_3 + 3/8) \ln(10)d^2 \approx 12.3 d^2 .$$

This is less than the estimate $16.5d^2$ for algorithm B1, but we preferred to use B1 because of its simplicity and the difficulty in rigorously bounding the error in (14) .

4.3 Exponential Integral Methods

Several algorithms are based on the identity

$$\gamma + \ln(n) = Q(n) - R(n) ,$$

where

$$(15) \quad Q(n) = \int_0^n \left(\frac{1 - e^{-x}}{x} \right) dx = \sum_{k=1}^{\infty} \frac{n^k (-1)^{k-1}}{k!k}$$

and

$$R(n) = \int_n^{\infty} \frac{e^{-x}}{x} dx = O(e^{-n}/n) .$$

Beyer and Waterman [3, 4] took $n \approx \ln(10)d$, worked to precision equivalent to $2d$ decimal places to compensate for cancellation in the sum for $Q(n)$, and neglected $R(n)$. The time required for this method is about $6\alpha_1 \ln(10)d^2 \approx 49.6d^2$, or three times that for method B1.

Following the suggestion of Sweeney [16], Brent [7, 8] took $n \approx \frac{1}{2} \ln(10)d$, summed the series for $Q(n)$ using the equivalent of $3d/2$ decimal places, and approximated $R(n)$ by its asymptotic expansion

$$(16) \quad R(n) = \frac{e^{-n}}{n} \sum_{k=0}^{n-2} \frac{k!}{(-n)^k} + O(e^{-2n}/n) .$$

Assuming the power series is used to compute e^n , the time required for this method is about

$$\frac{3}{4} (3\alpha_2 + \alpha_0 + 1) \ln(10)d^2 \approx 28.8d^2 ,$$

about twice as much as for method B1. Actual running times confirm this ratio.

Using the identity

$$e^n Q(n) = \sum_{k=0}^{\infty} H_k \frac{n^k}{k!} ,$$

we can evaluate $Q(n)$ by computing $\sum H_k n^k/k!$ and $\sum n^k/k!$ by recurrences similar to (10). Because all the terms in the two sums are positive, there is no need to increase the working precision to much more than the equivalent of d decimal places. If $n \approx \ln(10)d$, and $R(n)$ is neglected, the time required is about $7\alpha_0 \ln(10)d^2 \approx 43.8d^2$, slightly less than for Beyer and Waterman's method. If the asymptotic series (16) is used for $R(n)$, the time required is about

$$\frac{1}{4}(14\alpha_1 + 3)\ln(10)d^2 \approx 30.7d^2 ,$$

slightly greater than for Sweeney's method.

Instead of using the asymptotic expansion (16) for $R(n)$, we could use Euler's continued fraction [17, pg. 350]

$$e^n R(n) = 1/n + 1/1 + 1/n + 2/1 + 2/n + 3/1 + 3/n + \dots$$

and the forward or backward recurrence relations. This has the advantage that $R(n)$ can be evaluated as accurately as desired, whereas with the asymptotic expansion (16) the error is $\Omega(e^{-2n}/n)$. The choice of the optimal n and the optimal number of terms in the continued fraction (evaluated by the backward recurrence relations) gives a method competitive with algorithm B1, but much more complicated.

5. Computational Results

γ was computed to more than 30,100 decimal places using method B1 and a multiple-precision arithmetic package [9] on a Univac 1100/42. Three independent computations were performed, with $n = 17,332$ (using base 10,000 and 7,527 digits), $n = 17,357$ (base 65,535 and 6,260 digits), and $n = 17,387$ (base 65,536 and 6,271 digits). All three agreed to 30,100 decimal places, and the last two agreed to 30,141 decimal places. The computer time required for each computation was about 20 hours, much the same as for the 20,700 decimal place computation [7] using Sweeney's method on the same machine.

We also computed $G = \exp(\gamma)$ to more than 30,100 decimal places using the exponential routine in Brent's package [9] (with base 65,536 and 6,260 digits), and verified it by computing $\ln(G)$ by the Gauss-Salamin algorithm [6] (with base 10,000 and 7,550 digits). The rounded 30,100D values of γ and G are given in [10].

The first 29,200 partial quotients in the regular continued fractions for γ and G were computed and verified as in [7,8]. Statistics on the distributions of the first 29,000 partial quotients are given in Table 1, with notation as in [7, Table 2]. A chi-squared test did not show any

Table 1

Distribution of first 29,000 partial quotients for γ and G

n	number of $q_i(\gamma) = n$	number of $q_i(G) = n$	expected number
1	12112	11992	12036.1
2	4809	4875	4927.8
3	2791	2760	2700.2
4	1727	1757	1707.9
5	1181	1168	1178.6
6	867	848	862.7
7	642	716	658.9
8	497	520	519.7
9	420	417	420.5
10	346	335	347.2
11-20	1624	1729	1694.1
21-50	1148	1103	1133.9
51-100	411	390	400.2
101-1000	378	349	370.4
>1000	47	41	41.8

significant difference (at the 5% level) between the actual distributions and the distribution predicted by the Gauss-Kusmin theorem [13]. A table of the first 29,000 partial quotients for γ and G is given in [11].

From the continued fractions for γ and G we can improve the Theorem of [7, Sec. 7], where the lower bound on $|Q|$ was $10^{10,000}$.

Theorem 1

If γ or $G = P/Q$ for integers P and Q, then $|Q| > 10^{15,000}$.

Acknowledgement

This work was initiated while the first author was visiting the Department of Electrical Engineering and Computer Sciences at the University of California, Berkeley. The computations were performed at the Australian National University.

References

1. M. ABRAMOWITZ & I.A. STEGUN, Handbook of Mathematical Functions with Formulas Graphs and Mathematical Tables, National Bureau of Standards, Washington, D.C., 1964.
2. M. BEELER, R.W. GOSPER & R. SCHROEPPPEL, "Hakmem," Memo No.239, M.I.T. Artificial Intelligence Lab., Cambridge, Mass., 1972, pp. 70-71.
3. W.A. BEYER & M.S. WATERMAN, "Error analysis of a computation of Euler's constant," Math. Comp., v. 28, 1974, pp. 599-604. MR 49 #6555.
4. W.A. BEYER & M.S. WATERMAN, "Decimals and partial quotients of Euler's constant and $\ln 2$," UMT 19, Math. Comp., v. 28, 1974, p.667.
Errata: Math. Comp. TE 549, v. 32, 1978, pp. 317-318.
5. R.P. BRENT, "The complexity of multiple-precision arithmetic," Complexity of Computational Problem Solving (R.S. Anderssen and R.P. Brent, Editors), Univ. of Queensland Press, Brisbane, 1976, pp. 126-165.
6. R.P. BRENT, "Multiple-precision zero-finding methods and the complexity of elementary function evaluation," Analytic Computational Complexity (J.F. Traub, Editor), Academic Press, New York, 1976, pp. 151-176. MR 52 #15938, 54 #11843.
7. R.P. BRENT, "Computation of the regular continued fraction for Euler's constant", Math. Comp., v. 31, 1977, pp. 771-777. MR55 #9490.
8. R.P. BRENT, " γ and $\exp(\gamma)$ to 20700D and their regular continued fractions to 20000 partial quotients", UMT 1, Math. Comp., v. 32, 1978, p. 311.
9. R.P. BRENT, "A Fortran multiple-precision arithmetic package," ACM Trans. Math. Software, v. 4, 1978, pp. 57-70.
10. R.P. BRENT, "Euler's constant and its exponential to 30,100 decimals", Computing Research Group, Australian National University, Sept. 1978. Submitted to Math. Comp. UMT file.

11. R.P. BRENT & E.M. McMILLAN, "The first 29,000 partial quotients in the regular continued fractions for Euler's constant and its exponential", submitted to Math. Comp. UMT file.
12. J.W.L. GLAISHER, "History of Euler's constant," Messenger of Math., v. 1, 1872, pp. 25-30.
13. A. Ya. KHINTCHINE (A. Ja. HINČIN), Continued Fractions, 3rd ed., (English transl. by P. Wynn), Noordhoff, Groningen, 1963. MR 28 #5038.
14. G.F.B. RIEMANN, "Zur Theorie der Nobili'schen Farbenringe", Poggendorff's Annalen der Physik und Chemie, Bd. 95, 1855, pp. 130-139. (Reprinted in Bernhard Riemann's Gesammelte Mathematische Werke und Wissenschaftlicher Nachlass, B.G. Teubner, Leipzig, 1876, pp. 54-61.)
15. A. SCHÖNHAGE & V. STRASSEN, "Schnelle Multiplikation grosser Zahlen", Computing, v. 7, 1971, pp. 281-292.
16. D.W. SWEENEY, "On the computation of Euler's constant," Math. Comp., v. 17, 1963, pp. 170-178. MR 28 #3522.
17. H. WALL, Analytic Theory of Continued Fractions, Van Nostrand, New York, 1948.
18. G.N. WATSON, A Treatise on the Theory of Bessel Functions, Cambridge University Press, London, 1944 (second edition).

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720