

UC Davis

UC Davis Previously Published Works

Title

Current Status and Future Strategies for Advancing Functional Circuit Mapping In Vivo.

Permalink

<https://escholarship.org/uc/item/2h91s8n1>

Journal

The Journal of Neuroscience, 43(45)

Authors

Berndt, Andre

Cai, Denise

Cohen, Adam

et al.

Publication Date

2023-11-08

DOI

10.1523/JNEUROSCI.1391-23.2023

Peer reviewed

Symposium

Current Status and Future Strategies for Advancing Functional Circuit Mapping *In Vivo*

Andre Berndt,¹ Denise Cai,² Adam Cohen,³ Barbara Juarez,⁷ Jaume Taura Iglesias,² Hejian Xiong,⁴ Zhenpeng Qin,⁵ Lin Tian,⁶ and Paul A. Slesinger²

¹University of Washington, Seattle, WA 98109, ²Icahn School of Medicine at Mount Sinai, New York, NY 10029, ³Harvard University, Cambridge, MA 02138, ⁴Southern Medical University, Guangzhou, China, ⁵University of Texas–Dallas, Richardson, TX 75080, ⁶University of California–Davis, Davis, CA 95616, and ⁷University of Maryland–Baltimore, Baltimore, MD 21201

The human brain represents one of the most complex biological systems, containing billions of neurons interconnected through trillions of synapses. Inherent to the brain is a biochemical complexity involving ions, signaling molecules, and peptides that regulate neuronal activity and allow for short- and long-term adaptations. Large-scale and noninvasive imaging techniques, such as fMRI and EEG, have highlighted brain regions involved in specific functions and visualized connections between different brain areas. A major shortcoming, however, is the need for more information on specific cell types and neurotransmitters involved, as well as poor spatial and temporal resolution. Recent technologies have been advanced for neuronal circuit mapping and implemented in behaving model organisms to address this. Here, we highlight strategies for targeting specific neuronal subtypes, identifying, and releasing signaling molecules, controlling gene expression, and monitoring neuronal circuits in real-time *in vivo*. Combined, these approaches allow us to establish direct causal links from genes and molecules to the systems level and ultimately to cognitive processes.

Introduction

Recent advances in deconstructing neuronal circuits *in vivo* involve the synergy of four areas: cell-specific expression of proteins using molecular genetics, monitoring neuronal activity, and neuromodulators over time, triggering the release of specific transmitters, and tracking behaviors. Using model organisms, such as rodents, allows one to target distinct cell types with protein-based tools, such as fluorescent proteins and light-activated ion channels. Viruses encoding these tools are injected into the brain, and their expression is restricted to specific cell types under the control of Cre or Flp recombinase in transgenic rodent driver lines (Gong et al., 2007; Weinholtz and Castle, 2021). For example, tracing experiments label distinct neuronal subtypes with fluorescent proteins, allowing axonal projections to be traced by various volumetric imaging techniques (Oh et al., 2014; Glaser et al., 2023). This type of circuit mapping can scale to subcellular levels and reveal intricate anatomic features, such as dendritic and axonal arborization, identifying convergent or divergent connections from one brain region to another.

While these imaging techniques provide details on the connectome, they lack dynamic information on activity or the biological compounds that regulate these signals. Recent advances in sensor engineering and caging of neuromodulators enable us to monitor intracellular and extracellular signaling molecules with high spatial and temporal resolution in head-fixed or even freely

moving animals. For example, genetically encoded fluorescent sensor proteins can monitor calcium, voltage, dynorphin, dopamine, norepinephrine, acetylcholine, orexin, and more (Patriarchi et al., 2018; Adam et al., 2019; Feng et al., 2019; Jing et al., 2020; Abraham et al., 2021; Duffet et al., 2022; Zhang et al., 2023).

Concurrently, several imaging techniques have been developed to monitor these sensors. Miniaturized fluorescence microscopes are head-mounted onto freely moving rodents for *in vivo* detection (Ghosh et al., 2011; Cai et al., 2016). Two-photon microscopy through transcranial windows of head-fixed animals provides subcellular resolution. It can be complemented by Gradient Refractive Index (GRIN) lenses or prisms to image deep brain areas (Levene et al., 2004; Andermann et al., 2013). Fiber photometry detects bulk fluorescence through implanted optical fibers, targeting specific neuronal subpopulations and analytes (e.g., voltage, Ca²⁺, neurotransmitters, neuromodulators) (Gunaydin et al., 2014). These tools allow for a distinct assessment of the roles of cell types, neuronal circuits, neurotransmitters, and neuromodulators in cognitive functions. For example, they revealed the dynamics of dopamine release from the VTA during reward paradigms, the firing-dependent volumetric release of acetylcholine in the entorhinal cortex, and orexin dynamics in somatosensory cortex and basal forebrain during wake-sleep cycles (Patriarchi et al., 2018; Jing et al., 2020; Duffet et al., 2022).

Optogenetic (Yizhar et al., 2011) and chemogenetic (Vardy et al., 2015) actuators, such as channelrhodopsins and DREADDs, link circuits to behaviors by directly manipulating neuronal activity. Membrane potentials can be depolarized or hyperpolarized, and light-activated ion channels can trigger neurotransmitter release. Intracellular signaling cascades can be modulated by designer GPCRs, such as DREADDs, by light-activated adenylyl

Received July 21, 2023; revised Aug. 24, 2023; accepted Aug. 25, 2023.

The authors declare no competing financial interests.

Correspondence should be addressed to Andre Berndt at berndtuw@uw.edu.

<https://doi.org/10.1523/JNEUROSCI.1391-23.2023>

Copyright © 2023 the authors




Figure 1. Comprehensive *in vivo* circuit mapping requires the modulations and monitoring of genetically defined neuron populations. This review highlights recent advances in techniques to study neuronal circuits on the genetic, molecular, and cellular levels. Applying these methods *in vivo* could link circuit functions to cognitive processes and disease mechanisms. Figure created with BioRender.com.

cyclases, such as bPAC (Stierl et al., 2011) or light-activated GPCRs, such as parainopsin (Copits et al., 2021) and OPN3 (Mahn et al., 2021). Combining these techniques allows simultaneous monitoring and manipulation of targeted neuronal circuits. For instance, one can trigger neuromodulator release from a specific cell type using channelrhodopsin and simultaneously monitor the activity of connected downstream neurons using calcium imaging or sensors for neuromodulators. This combined approach enables the reconstruction of detailed cellular and molecular maps of brain circuitry and informs our understanding of the interplay between brain regions during cognitive processes.

The past 15 years saw tremendous progress in functional circuit mapping in model organisms. In the context of fear memory, specific groups of neurons (known as engram cells) in the entorhinal cortex and within various regions of the hippocampus are thought to encode contextual information. In contrast, engram cells in the amygdala primarily encode fear-associated information. A series of mapping experiments revealed that these distinct cell ensembles form a complex network encapsulating the fear memory (Josselyn and Tonegawa, 2020). Similarly, *in vivo* calcium imaging of the CA1 region in the hippocampus of freely moving rats visualized and tracked hundreds of place cells, crucial for spatial navigation and memory, over weeks. This innovative approach revealed that place cells could maintain their location specificity over extended periods, providing insights into hippocampal circuit organization, which is vital for memory, navigation, and learning (Wirtshafter and Disterhoft, 2022).

Other mapping studies have shown the importance of VTA's (ventral tegmental area) dopaminergic inputs to the mPFC (medial prefrontal cortex) in managing social avoidance and stress resilience. Furthermore, efferent connections from mPFC to other regions, such as the BLA (basolateral amygdala), were

critical in reversing stress-induced behavioral deficits and producing rapid antidepressant effects. Manipulation of the mPFC's various circuits influenced anxiety and depression-like behaviors, indicating the central role mPFC plays in regulating affective states (Liu et al., 2021).

Another study monitored thousands of striatal spiny projection neurons of the direct (dSPNs) and indirect (iSPNs) pathways in the basal ganglia. The results showed that L-DOPA can cause imbalances in the basal ganglia when treating Parkinson's because of the differential responses of dSPN and iSPN pathways. Thus, future treatments need to consider the activity rates and the spatial-temporal coordination of neurons within these circuits (Parker et al., 2018).

These examples show that the future of circuit mapping relies on increasingly precise and multifaceted methods matching the complexity of brain circuits. This review highlights recent technological advances in (1) genetically encoded sensors, (2) recording neuronal activity with mini-microscopes, (3) protein integrators of neuronal activity, (4) CRISPR/Cas9 gene editing, and (5) time-locked release of neuropeptides from nanovesicles (Fig. 1). These topics were discussed at a corresponding symposium at the 2023 annual meeting of the Society for Neuroscience.

Next-generation sensor design: overcoming challenges in protein engineering

One key to circuit mapping is monitoring the release of neuromodulators and neuropeptides from genetically defined neuron populations. Fluorescent sensor proteins are ideally suited for this task because they can be engineered to bind specific ligands, and their expression can be spatially restricted by Cre-Lox recombination or cell-specific promoters. Existing sensors for calcium and neurotransmitters have provided new information

on neuronal circuits. However, much work remains to improve the brightness, sensitivity, kinetics, dynamic range, spectral tuning, and specificity of sensor proteins. For example, *in vivo* imaging of individual neurons through transcranial windows or miniaturized microscopes requires highly optimized sensors with large signal amplitudes. For imaging neuronal activity through calcium transients, this threshold was initially reached with the development of GCaMP6 from GCaMP3 (Chen et al., 2013).

In addition, there is a vast array of targets for future sensor development. Here, we consider the family of opioid receptors and peptides. Despite the large body of research on opioid receptor function, there is a limited understanding of the endogenous opioid peptides released in neuronal circuits. Fluorescent opioid sensors could identify the specific types of neurons that release endogenous opioids during pain responses, reward, or stress. One example is the recently developed sensor kLight1.2, based on the κ -opioid receptor. The sensor could detect dynorphin release in the mPFC of mice under morphine withdrawal (Abraham et al., 2021). A similar prototype sensor based on the μ -opioid receptor, however, failed to generate a sufficient signal response, highlighting the need for further optimization (Patriarchi et al., 2018).

Optimizing fluorescent sensors for detecting neuropeptides via mutagenesis has been a significant challenge because of the immense mutational landscape of proteins. Targeting only five residues by randomized mutations can result in 3.2 million variants. This number far exceeds the current throughput of screening approaches for most sensor proteins (usually in the low hundreds). Newly developed high-throughput systems, such as BeadScan (Koveal et al., 2022) and Opto-MASS (Rappleye et al., 2022), and video-based pooled screens (Tian et al., 2023) can functionally test signal amplitudes, kinetics, and ligand selectivity of semi-randomly mutated variants in minutes. For example, Opto-Mass tested 21,000 variants of mLight and identified a significantly improved variant called uMASS, which features improved membrane trafficking as well as 5 times larger response amplitudes toward the synthetic opioid [D-Ala², N-MePhe⁴, Gly-ol]-enkephalin (DAMGO).

uMASS inherits the ligand binding promiscuity of the μ -receptor. Accordingly, it is highly sensitive to Met-enkephalin but also detects endorphin, Leu-enkephalin, and dynorphin with lower affinities. Future opioid sensors for each type would allow us to distinguish neurons producing different opioids. Given that different opioids can have complementary or opposing effects, multiple opioid-specific fluorescent sensors could uncover competitive receptor binding or the co-release of opioids. High-throughput screening platforms, such as Opto-MASS, can be adapted to screen sensor libraries while several competing agonists are applied sequentially. The screen could identify variants with distinct selectivity profiles, such as high Met-enkephalin and low endorphin affinity or vice versa.

These high-throughput approaches create large datasets that link mutations to sensor functions. Implementing machine learning (ML) to these datasets could significantly advance sensor engineering. In one recent example, ~1000 mutations of GCaMP were used to train ML ensembles to predict response amplitudes and kinetics of known GCaMP mutations (Wait et al., 2023). The final ML ensembles incorporated the five amino acid properties of 554, with the highest predictive power for the function of GCaMP variants. The ML model was applied to a library of 1423 previously untested GCaMP mutants. Variants with high fluorescence and kinetic responses were subsequently

tested. Two variants, called ensemble-GCaMPs (eGCaMP⁺ and eGCaMP2⁺), have generated the largest dynamic range of any calcium indicator, while eGCaMP⁺ is also the fastest known variant in these tests (Wait et al., 2023).

These ML models can be applied to map sequence–function relationships in new sensors instead of more complex structure–function relationships. These approaches will become invaluable to significantly accelerate sensor engineering when combined with high-throughput platforms. Such next-generation methods are urgently needed: ~6000 neuropeptides have been discovered in vertebrate and invertebrate organisms, with most functions only weakly defined at the circuit level (Y. Wang et al., 2015). Beyond neurotransmitters and neuromodulators, sensors that detect changes in neurotrophic factors, such as BDNF, could provide insights into neural development, neural plasticity, and neurodegenerative processes. A sensor for cortisol could help us better understand the brain's response to stress. Furthermore, developing sensors to detect signaling molecules involved in immune responses, such as cytokines, could be revolutionary. Traditional (slow) trial-and-error mutagenesis for engineering and optimizing these sensors would require tremendous resources and time commitments. Further advances in protein engineering are critical to overcome these challenges (Fig. 2).

Functional imaging of neuronal circuits in freely moving rodents with miniscopes

While monitoring the release of neuromodulators provides information on when and where these molecules are released, it is also essential to study the effect on the activity of individual neurons. One of the most frequently used fluorescent sensors is GCaMP for measuring changes in intracellular calcium, which occurs downstream of action potentials. Because of the slower time scales of calcium sensor kinetics compared with voltage sensor kinetics, imaging acquisition rates are lower than for voltage indicators and allow lower excitation light power, reducing hardware demands and photobleaching. Combining fluorescent biosensors with miniaturized microscopes using GRIN lenses instead of traditional objective lenses enables probing deeper brain regions, such as the hippocampus, thalamus, striatum, orbitofrontal cortex, and brainstem. Microscope miniaturization can be combined with robust behavioral paradigms to allow the investigation of circuits underlying functions, such as learning and memory (Ghosh et al., 2011; Cai et al., 2016).

The use and development of the open-source UCLA Miniscope is one example of how *in vivo* imaging can be applied to answer complex questions about memory dynamics *in vivo*. For instance, Miniscopes recorded active neurons as mice explored varying contexts (Fig. 3). These tests proved that memories that occur within a day are more likely to share an overlapping neural ensemble, leading to behavioral linking: the recall of one memory likely triggers recall of the other memory that occurred close in time. Additionally, specific memory-linking deficits in aged mice were rescued using artificial ensemble activation via chemogenetic stimulation by DREADDS (Vardy et al., 2015). These experiments provided novel and critical insight into how the brain organizes and integrates different experiences across time. A future question is how memories are encoded across even longer time scales. A recent preprint details how strong aversive experiences drive ensemble reactivation of a neutral memory formed 2 d prior, effectively transferring fear from the aversive to the neutral context (Zaki et al., 2023). Such findings indicate that




Figure 2. Next-generation approaches for engineering genetically encoded fluorescent sensors. Thousands of biological compounds regulate brain activities and cognitive functions. Fluorescent protein sensors for these compounds can be designed following similar design principles. However, optimizing dynamic range, signal sensitivity, and kinetics is hampered by the immense mutational space of each sensor protein. Therefore, further advancements in protein engineering are required by integrating high-throughput functional testing of large variant libraries, data-driven approaches, and AI-supported structure-guided design. Adapted with permission from Rappleye et al. (2022) and Wait et al. (2023).




Figure 3. **A**, Technologies, such as Miniscope calcium imaging, allow real-time observations of neuronal ensembles *in vivo* while mice perform behavioral tasks (**B**). **C**, Computational analysis of the imaging sessions allows to identify the dynamic activity of hundreds of neurons.

memory linking can occur between events rather than only at initial memory encoding, suggesting a mechanism that allows the brain to associate memories across time.

There is a need to increase access to these powerful imaging technologies, as evidenced by the growing number of users involved in the UCLA Miniscope project. To remove barriers to

technology adoption and iteration, all aspects of the Miniscope, from the hardware components to the software used for analysis, were selected based on their widespread availability. The success of the UCLA Miniscope project has prompted the initiation of many others, including NINscope (de Groot et al., 2020), cScope (Scott et al., 2018), MiniFAST (Juneau et al., 2020), Miniscope3D




Figure 4. Integrators and ticktapes transcend the tradeoffs between *in vivo* and *ex vivo* imaging. **A**, Real-time reporters probe neural dynamics but with a limited Field-of-View; *ex vivo* analyses can access the whole brain but provide static snapshots. Integrators and ticktapes can provide dynamic, brain-wide information. **B**, Light-gated integrators convert the cumulative level of an analyte during an illumination epoch into a transcriptional readout. **C**, Schematic operation of a light-gated integrator. Dynamic fluctuations in an analyte (e.g., Ca^{2+}) are converted into transcription factor (TF) release during an illumination epoch. The TF drives expression of a reporter protein. **D**, Example ticktape recorder. A linearly growing scaffold is marked at periodic intervals via alternating injection of two colors of HaloTag-ligand dye. An activity-dependent promoter (e.g., cFos) drives the expression of a GFP-labeled monomer, which introduces green stripes into the fiber. Localization of the green stripes relative to the red/yellow color transitions identifies the timing of neural activity. **D**, Adapted with permission from Lin et al. (2023).

(Yanny et al., 2020), and Mesoscope (Rynes et al., 2021). These collective efforts have engendered innovative approaches that leverage 3D imaging, variable sampling rates, and different FOVs.

Though powerful in providing anatomically precise, circuit-level investigations of brain function, existing tools have limitations. Miniscope designs generally use a one-photon wide field image similar to a tabletop fluorescent microscope and produce images that are projections from the distal tip of the GRIN lens. These one-photon imaging approaches are prone to out-of-focus light from other depth planes scattering and producing noise in the FOV. Image analysis requires spatiotemporal unmixing of the 2D images to differentiate cells and separate signal from noise. While the currently used algorithms can extract calcium activities from individual neurons fairly reliably amid high noise levels, ensuring that the same neuronal populations are recorded across imaging sessions remains challenging. Because of the chromatic aberration of GRIN lenses that are typically used in Miniscope, Miniscopes are typically only able to record one wavelength of light at a time, precluding the possibility of recording multiple fluorophores simultaneously in the same FOV. In contrast, two-photon microscope systems can measure multiple signals simultaneously and do not suffer from contamination by out-of-focus light scattering; however, their requirement for head-fixed preparations hinders use in many behavioral paradigms.

To overcome these limitations, there are continual developments in open-source single-photon and two-photon Miniscopes with multiwavelength recording capabilities (Zong et al., 2021). This allows for simultaneous imaging of two dynamic signals, such as dopamine and calcium, using dLight (Patriarchi et al., 2018) and GCaMP (Chen et al., 2013), or a static signal with a dynamic signal (e.g., tagging cells with constitutively active

indicators, such as tdTomato and monitoring changes with GCaMP). Innovative software, such as CaImAn (Giovannucci et al., 2019) and Minian (Dong et al., 2022), for calcium imaging analysis, have been developed to be combined with programs, such as DeepLabCut (Mathis et al., 2018), relating behavioral responses to neuronal activity. As these approaches allow researchers to modulate and monitor neuronal signaling in behaving animals with high spatial and temporal resolution, it is critical to continue promoting affordable and user-friendly implementation as new technologies develop.

Integrators and ticktapes for imaging brain-wide neuronal activity

In parallel with the development of new sensors and imaging systems is the search for techniques to monitor the activity of an unlimited number of individual neurons that can be recorded simultaneously. With current imaging techniques, light scatter constrains live-tissue imaging to depths of typically <500 nm for two-photon microscopy and <1 mm for three-photon. Efforts to image deeper require the destruction of overlying tissue and sometimes sacrificing single-cell resolution. On the other hand, brain-wide maps of macromolecular signals (e.g., gene expression or protein levels) can be generated in fixed tissues. Still, these maps probe only a single point in time and do not capture physiological signals, such as calcium or neuromodulators (Fig. 4A). Thus, neuroscientists face a trade-off between dynamical and spatial information.

Several recently developed techniques show a path toward transcending this trade-off. The key feature is to create long-lived chemical traces that record transient events within individual cells. The encoding occurs *in vivo*, but the readout is measured

ex vivo. While whole-brain recordings are not yet possible, there are exciting advances in several areas.

First, light-gated integrators store a snapshot of neural activity in a long-lived chemical signal (Fig. 4B). As their name suggests, these tools build up a signal proportional to the cumulative activity during an interval defined by a light pulse. In one technique, a light-gated voltage integrator and a light-gated voltage sample-and-hold reporter store long-term records of electrical activity via the conformation of a microbial rhodopsin (Venkatachalam et al., 2014). Alternatively, a light-gated calcium integrator, CaMPARI, undergoes an irreversible green-to-red transition in the simultaneous presence of calcium and violet light (Fosque et al., 2015). These tools enable *ex vivo* mapping of circuits that had been active during the illumination epoch. However, they do not provide any downstream genetic access to these cells.

The ST-Cal-Light (Lee et al., 2017b; Hyun et al., 2022), FLiCRE (C. K. Kim et al., 2020), and scFLARE2 (Sanchez et al., 2020) light-gated calcium integrators also sense cumulative calcium during an optically gated interval. These tools also provide a transcriptional readout (Fig. 4C). With these tools, a transcription factor is tethered to the cell membrane by a linker cleaved in the simultaneous presence of blue light and calcium. FLiCRE requires three genetic components (two for the integrator and one for the readout). The scFLARE2 integrator has a single sensing component (but still requires another gene as the readout). This approach allows the coupling of the Ca²⁺ activity signal to a wide range of subsequent measurements or perturbations. For instance, an optogenetic actuator or silencer could be expressed to test the causal roles of the tagged neurons. Alternatively, structural markers could be expressed to assess their connectomes or ribosome pulldown tags to measure their ensembles of translated proteins.

SPARK (M. W. Kim et al., 2017) and iTANGO (Lee et al., 2017a) reporters operate on a similar principle to FLiCRE and scFLARE but are sensitive to the activation of GPCRs. These reporters have a modular design so that they can be coupled to various upstream receptors, as well as arbitrary transcriptional readouts. Integrators have been developed for dopamine receptors D1 and D2, the β -adrenergic receptor, and others.

Existing light-gated integrators have two key limitations. First, these tools only provide a single readout. Ideally, one wishes to map changes in signal associated with a particular experimental condition, but the existing tools can be confounded by basal activity unrelated to the experiment. Readouts at multiple times would also permit comparison of patterns of neural activation under different conditions within the same animal. Second, existing integrators require blue or violet light for the optical gate. Because of strong blue light scattering in tissue, this limits the recording depth to ~ 1 mm. In principle, red or infrared-gated integrators could record throughout a mouse brain, with light sources mounted outside the skull.

Several recent efforts have sought to provide multi-time point measures of neural activity and to overcome the depth limitations of light-gated integrators. Protein tickertapes comprise linearly growing protein fibers that incorporate colored bands during epochs of neural activity, which are then read *ex vivo*. Based on linear crystals of the Pak4 kinase, one approach has an absolute time resolution of ~ 30 min, but the crystals deform the cells and thus are incompatible with *in vivo* use (Fig. 3) (Lin et al., 2023). Another approach is based on filaments of *Escherichia coli* isoaspartyl dipeptidase but has a time resolution of days and does not have an absolute time-base (Mohar et al., 2022). In both

cases, expression of the tags is driven by neural activity-dependent genes (e.g., *cfos*), which are downstream of action potentials.

Pulse-chase labeling with HaloTag-ligand dyes also provides a means to capture aspects of neural activity at multiple time points. In one recently demonstrated technique, dubbed DELTA, protein turnover was mapped brain-wide via pulse-chase labeling of HaloTag-protein fusions with multiple colors of brain-permeant HaloTag-ligand dyes. In another technique, dubbed EPSILON, newly surface-exposed AMPA receptors were identified via pulse-chase labeling with multiple colors of membrane-impermeable dyes (D. Kim et al., 2023). EPSILON could identify synapses potentiated during a particular chemically gated window *in vivo*, providing an approach to mapping the physical basis of memory formation.

The field of light- and chemical-gated integrators and tickertapes is still in its infancy. Nonetheless, these tools promise to provide a view that complements existing *in vivo* and *ex vivo* tools for functional circuit mapping, revealing brain-wide maps of multiple chemical signals as a function of time.

CRISPR/Cas9 gene editing approaches to study gene function

While genetically encoded fluorescent sensors can be targeted to specific types of neurons using specific promoters and Cre-lines of mice, it is also essential to determine the causal roles of specific genes in neural circuit dynamics. The basis for the functional, anatomic, and molecular diversity in the brain is fundamentally linked to the gene expression diversity of cells within neuronal circuits. Consequently, gene editing technologies, such as CRISPR-Cas9, which modify gene expression, are critical for complementing the connectome with genetic information (Charpentier and Marraffini, 2014; Heidenreich and Zhang, 2016). Combined with neurophysiological tools and behavioral analyses, gene-editing technologies will provide new insights into the genetic basis of brain function (Savell and Day, 2017).

Gene-modifying approaches, such as knock-in and knock-out transgenic mice, virally mediated gene delivery of RNA interference (RNAi) constructs, targeting of DNA nucleases, including zinc-finger nucleases (ZFNs) and transcription activator effector nucleases (TALENs), have helped to discover new genetic contributions to brain function (Urnov et al., 2005; Limaye et al., 2009; H. Wang et al., 2013). Yet, the high costs, time for generation, temporal limitations, and complexity of some of these methods restrain their application. Recently, using CRISPR technology combined with Cas bacterial enzymes, such as Cas9, has addressed many shortcomings of earlier methods.

CRISPR-Cas9-based gene-editing approaches engineered for eukaryotic systems are based on the guidance of Cas9 nucleases to a DNA target site of interest using single-guide RNAs (sgRNAs) (Jiang and Doudna, 2017; Kalamakis and Platt, 2023). Once the sgRNAs deliver Cas9 to the site of interest in DNA, the DNA is unwound and cleaved near a protospacer adjacent motif (PAM) (Fig. 4A). Cellular mechanisms of either homology-directed repair (HDR) or nonhomologous end joining (NHEJ) are recruited to repair the double-stranded break (DSB) in the DNA (Day, 2019). HDR pathways are usually recruited in dividing cells and use DNA template strands to precisely edit and repair the DSBs. In nondividing cells (like neurons), NHEJ mechanisms are instead usually recruited, resulting in random insertions, deletions, or base-pair substitutions to correct the DSB, thus disrupting gene expression.




Figure 5. Capabilities of the CRISPR/Cas9 toolbox. **A**, Systems using CRISPR/Cas9 components with Cas9's nuclease activity intact will cause DNA cleavage and error-prone NHEJ, most likely used in neurons or precise editing with HDR. **B**, CRISPR/Cas9 systems with inactive nuclease activity, or dCas9, can be fused with effector proteins to further modify gene expression.

In addition, catalytically inactive (“dead”) Cas9 (dCas9) is altered to prevent DNA cleavage while retaining sgRNA-directed guidance. Thus, dCas9 can guide fused effector proteins to specific DNA sequences (Fig. 4B) (Gilbert et al., 2013). These approaches have been further expanded by CRISPR interference (CRISPRi) to repress gene transcription and CRISPR activation (CRISPRa) that can activate transcriptional states (Pickar-Oliver and Gersbach, 2019; Savell et al., 2019; Duke et al., 2020). Recently, CRISPR/Cas9 systems have been used to shuttle other epigenetic effector proteins to alter gene expression (Fig. 5) (Choudhury et al., 2016; Choi et al., 2023).

Applying these CRISPR-Cas9 approaches *in vivo* has expanded our knowledge of genetic contributions to brain function. For example, two adeno-associated virus (AAV)-mediated gene delivery can express the CRISPR/Cas9 system in the adult mouse brain knocking out genes of interest (Swiech et al., 2015). The development of *Cre*-inducible Cas9 knock-in mice further revolutionized genome editing and its application across systems (Platt et al., 2014). It allowed crossing these mice with *Cre*-driver lines to specifically express Cas9 in cell types of interest in the nervous system. Thus, by virally delivering sgRNAs into targeted brain regions, it can induce rapid genome alterations in a cell type-specific manner (Yamaguchi et al., 2018; McQuillan et al., 2022).

In 2020, a conditional, single AAV-based approach was used to express a smaller Cas9 and sgRNAs in adult mammals (Hunker et al., 2020). The resulting gene mutagenesis was as efficient as gene disruption in KO mice. Juarez et al. (2023) recently combined these approaches with *in vivo* opto-electrophysiology, fiber photometry, and two-photon slice imaging to elucidate how ion channels contribute to patterns of VTA physiology and reinforcement learning. Notably, the conditional, single-AAV-based method could be combined with *Cre*-recombinase or *Flp*-recombinase systems. A compelling finding of this study was that neurons in the lateral hypothalamus produce both stimulating neurotensin and inhibitory GABA. Notably, these signals are

used in a coordinated manner to control neurons in the VTA that regulate dopamine release (Soden et al., 2023).

Advances in CRISPR-Cas9 approaches are now opening the door for ways to manipulate the genome of adult animals to better model human conditions (Cox et al., 2015). Methods that shift the DSB repair mechanism from NHEJ to HDR are being refined to introduce single nucleotide polymorphisms (SNPs) in the adult nervous system (Scholefield and Harrison, 2021). This could broaden the application of CRISPR to activate or repress genes of interest and model human disease-associated SNPs in animals. A critical feature for future applications is the temporal control of CRISPR-Cas9 function, which is under development and includes optogenetic or pharmacological induction of gene editing (Savell and Day, 2017; Choi et al., 2023).

Photosensitive nanovesicles: a promising tool for precise neuromodulation

A missing tool in the arsenal of measuring the release of neuromodulators, studying the activity on neurons, and mapping brain-wide circuits is the ability to control the release of a specific neurotransmitter or neuromodulators within a targeted brain region. Recent advances have improved the ability for spatially and temporally precise delivery of neuroactive compounds.

It is now possible to remotely control the release of neuroactive compounds using external stimuli, such as magnetic fields (Rao et al., 2019), ultrasound (Airan et al., 2017; J. B. Wang et al., 2018), or light (Rapp and DeForest, 2021). Among these methods, light-based modulation offers high temporal and spatial resolution. One approach is to cage a desired neuroactive molecule by blocking a group key for its biological activity, where the molecule remains inactive until photo-uncaging (Ellis-Davies, 2007; Taura et al., 2018). While various caged compounds exist, such as glutamate or GABA (Ellis-Davies, 2020), their use is limited by residual activity, low solubility, and uncaging efficiency (Silva et al., 2019), and the risk of off-target effects (Maier et al., 2005;




Figure 6. Photosensitive nanovesicles for precise neuromodulation. **A**, Schematic of neuropeptide transmission measurement by plasmonic gold-coated nanovesicles (Au-nV-SST) and two clusters of SST2 CNiFERS (Cell-based Neurotransmitter Fluorescent Engineered Reporter). **B**, Two-photon fluorescent image of SST2 CNiFERS at 200 μm in the mouse cortex (Ex: 900 nm; Em: 520–560 nm). **C**, Response curves for SST2 CNiFERS when stimulated (STIM) at different regions (core vs satellite 1/sat-1). **D**, Peak response time of SST2 CNiFER implants at defined distances from the core implant. **E**, Schematic of the photoisomerization of azo-PC. **F**, Schematic of the photoswitchable release of SKF-81297 from azosome. **G**, Real-time fluorescent images of cultured primary mouse striatal neurons before and after the irradiation of sequential 365 nm light and 455 nm light. Fluo-4 was used as the Ca^{2+} indicator. Scale bar, 50 μm . **A–C**, Adapted with permission from Xiong et al. (2022). **D–F**, Adapted with permission from Xiong et al. (2023a).

Noguchi et al., 2011). Along the same line, studying the role of neuropeptides (e.g., oxytocin, vasopressin, or somatostatin) in brain circuits and behavior has been particularly challenging because of the lack of specific methods for localized uncaging or release (DeLaney et al., 2018). While several caged neuropeptides have been reported, their application *in vivo* is limited by instability because of peptidase degradation (Ma et al., 2023; Xiong et al., 2023b). An alternative approach involves targeting neurons that endogenously produce these neuropeptides using cre-recombinase and expressing opsins to either opto-stimulate or inhibit the neuron and the subsequent neuropeptide release (Arrigoni and Saper, 2014; Dao et al., 2019). However, a major limitation of this approach is that neuropeptides are often coreleased with classical neurotransmitters (Merighi, 2002; Hökfelt et al., 2003), making it challenging to isolate and study their specific functions.

A new technology based on photosensitive nanovesicles has been developed to address these limitations. Photosensitive

nanovesicles encapsulate neuroactive molecules in self-assembled nanoscale vesicles, such as those made by a bilayer of phospholipids (liposomes) and a light-sensitive component for the photosensitive release. This innovative approach addresses the limitations of traditional pharmacology and provides a versatile platform to decipher the role of neuropeptides and neuromodulators. Specifically, the physical separation of neuroactive molecules from the surrounding brain environment improves *in vivo* stability compared with caged compounds.

Plasmonic gold-coated nanovesicles, in combination with Cell-based Neurotransmitter Fluorescent Engineered Reporter (PACE), have been used to investigate neuropeptide signaling in the mouse neocortex *in vivo* (Fig. 6A,B) (Xiong et al., 2022). Near-infrared light stimulation triggered the release of femtoliters to picoliters of Somatostatin-14 (SST) from nanovesicles in the brain, which was subsequently detected by activation of SST2 Cell-based Neurotransmitter Fluorescent Engineered Reporter (CNiFERS) tuned to nanomolar concentrations of SST. PACE

revealed a reduced yet coordinated SST transmission within 130 μm , with significantly diminished and delayed transmission at greater distances. PACE provides a novel method for examining the scale and timing of neuropeptide volume transmission and signaling within the brain.

Another class of photosensitive nanovesicles involves photo-switchable azobenzenes. Azobenzene groups have gained significant attention in the field of photoswitches (Bahamonde et al., 2014; Cabré et al., 2019; Morstein et al., 2019, 2020, 2022; DiFrancesco et al., 2020; Kellner and Berlin, 2020; Mukhopadhyay et al., 2022). These compounds exhibit reversible *trans-cis* photoisomerization. Recently, researchers incorporated azobenzene into lipids by replacing one of the hydrophobic tails on the phospholipid, leading to the development of photoswitchable lipids, including the photoswitchable phosphatidylcholine derivative azo-PC (Pernpeintner et al., 2017; Urban et al., 2018, 2020; Pritzl et al., 2020, 2022). Azo-PC enables optical control of membrane organization and permeability, providing a versatile framework for delivering neuroactive compounds. Building on the concept of photoswitchable lipids, a new class of photoswitchable nanovesicles termed “azosomes” have been developed. Azosomes consist of liposomes formulated with azo-PC. When the azobenzene undergoes reversible isomerization, it increases the permeability of the membrane and allows the contents to diffuse out (Fig. 6C–E). Computational studies suggest that *trans*-configuration may decrease the thickness of the lipid bilayer, thereby increasing permeability for encapsulated molecules. The ability to switch the release of molecules on and off using short light pulses (<3 s) demonstrates the potential of azosomes for neuromodulation. Recently, azosomes loaded with the D1 agonist SKF-81 297 were shown to modulate striatal neurons *in vitro* on light stimulation (Xiong et al., 2023a).

To explore the utility of azosomes *in vivo*, they are being tested in awake, behaving mice using fiber photometry to control the release of agonists and simultaneously monitor neural activity. For control, calcein-azosomes were infused into one hemisphere and calcein-liposomes (non-photoswitchable) into the opposite hemisphere. A dual optic fiber delivered light stimulation and monitored fluorescence on both hemispheres. The hemisphere implanted with calcein-azosomes (but not calcein-liposomes) exhibited a fluorescence increase on light photo-switching, confirming the photo-release of calcein *in vivo*. To study the photo-release of SKF-81 297 from azosomes *in vivo*, D1-medium spiny neurons in the dorsomedial striatum, a region involved in locomotor activity control, were targeted. While monitoring D1-medium spiny neuron Ca^{2+} dynamics (with DIO GCaMP expressed in a D1-Cre mouse) and the distance traveled, it is possible to simultaneously photo-release an azosome’s cargo, record neuronal activity, and monitor motor behavior. Additionally, we are exploring the use of azosomes to release neuropeptides, expanding the repertoire of neuroactive compounds that can be remotely released.

In summary, photosensitive nanovesicles possess several features that make them promising tools for the controlled release of neuroactive compounds and precise neuromodulation. Photosensitive nanovesicles can be loaded with fluorescent dyes, drugs, or neuropeptides, allowing for versatile applications. Harnessing the ability to remotely and precisely control the release of neuroactive compounds *in vivo* will further elucidate the *in vivo* function of several neuropeptides in the future.

In conclusion, revealing the brain’s intricacy, with its expansive neural network, synaptic connections, and finely-tuned genetic

and biochemical regulations, remains challenging. Details at the cellular and molecular levels and their control of cognitive function remain especially elusive. However, this gap is continuously narrowing thanks to transgenic model organisms and techniques, such as fluorescence imaging, CRISPR/Cas9 gene editing, protein tickertapes, and time-locked release of neuromodulators, allowing real-time *in vivo* functional circuit mapping. Notably, using protein-based tools in genetically defined neuronal subpopulations has illuminated the granular control of circuit function by a diverse array of cell types and neuromodulators. Concurrent advancements in imaging, epitomized by head-mounted Miniscopes, promise deeper insights into previously inaccessible brain regions. New fluorescent sensors, especially for neuropeptides, are of high priority to maximize the synergistic effects of these developments. Emerging high-throughput techniques for sensor optimization, combined with machine learning, are ideally suited to expedite advancements. Additionally, the revolutionary CRISPR/Cas9 gene editing techniques are increasingly used *in vivo* to delineate the genetic underpinnings of brain function, while photosensitive nanovesicles promise increased precision in neuromodulation. One major challenge for the future is overcoming the trade-off between spatial and temporal information. *In vivo* imaging, electrophysiology, and fiber photometry enable dynamic yet spatially restricted monitoring, while *ex vivo* imaging can provide spatially expansive yet primarily static information. Collectively, the emerging tools and methodologies highlighted in this review are elucidating the brain’s functional architecture and outlining future innovations in circuit mapping.

References

- Abraham AD, Casello SM, Schattauer SS, Wong BA, Mizuno GO, Mahe K, Tian L, Land BB, Chavkin C (2021) Release of endogenous dynorphin opioids in the prefrontal cortex disrupts cognition. *Neuropsychopharmacology* 46:2330–2339.
- Adam Y, et al. (2019) Voltage imaging and optogenetics reveal behaviour-dependent changes in hippocampal dynamics. *Nature* 569:413–417.
- Airan RD, Meyer RA, Ellens NP, Rhodes KR, Farahani K, Pomper MG, Kadam SD, Green JJ (2017) Noninvasive targeted transcranial neuromodulation via focused ultrasound gated drug release from nanoemulsions. *Nano Lett* 17:652–659.
- Andermann ML, Gilfoy NB, Goldey GJ, Sachdev RN, Wölfel M, McCormick DA, Reid RC, Levene MJ (2013) Chronic cellular imaging of entire cortical columns in awake mice using microprisms. *Neuron* 80:900–913.
- Arrigoni E, Saper CB (2014) What optogenetic stimulation is telling us (and failing to tell us) about fast neurotransmitters and neuromodulators in brain circuits for wake-sleep regulation. *Curr Opin Neurobiol* 29:165–171.
- Bahamonde MI, Taura J, Paoletta S, Gakh AA, Chakraborty S, Hernando J, Fernández-Dueñas V, Jacobson KA, Gorostiza P, Ciruela F (2014) Photomodulation of G protein-coupled adenosine receptors by a novel light-switchable ligand. *Bioconjug Chem* 25:1847–1854.
- Cabré G, Garrido-Charles A, Moreno M, Bosch M, Porta-de-la-Riva M, Krieg M, Gascón-Moya M, Camarero N, Gelabert R, Lluch JM, Busqué F, Hernando J, Gorostiza P, Alibés R (2019) Rationally designed azobenzene photoswitches for efficient two-photon neuronal excitation. *Nat Commun* 10:907.
- Cai DJ, et al. (2016) A shared neural ensemble links distinct contextual memories encoded close in time. *Nature* 534:115–118.
- Charpentier E, Marraffini LA (2014) Harnessing CRISPR-Cas9 immunity for genetic engineering. *Curr Opin Microbiol* 19:114–119.
- Chen TW, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, Schreier ER, Kerr RA, Orger MB, Jayaraman V, Looger LL, Svoboda K, Kim DS (2013) Ultra-sensitive fluorescent proteins for imaging neuronal activity. *Nature* 499:295–300.

- Choi EY, Franco D, Stapf CA, Gordin M, Chow A, Cover KK, Chandra R, Kay Lobo M (2023) Inducible CRISPR epigenome systems mimic cocaine induced bidirectional regulation of Nab2 and Egr3. *J Neurosci* 43:2242–2259.
- Choudhury SR, Cui Y, Lubecka K, Stefanska B, Irudayaraj J (2016) CRISPR-dCas9 mediated TET1 targeting for selective DNA demethylation at BRCA1 promoter. *Oncotarget* 7:46545–46556.
- Copits BA, et al. (2021) A photoswitchable GPCR-based opsin for presynaptic inhibition. *Neuron* 109:1791–1809.e11.
- Cox DB, Platt RJ, Zhang F (2015) Therapeutic genome editing: prospects and challenges. *Nat Med* 21:121–131.
- Dao NC, Brockway DF, Crowley NA (2019) In vitro optogenetic characterization of neuropeptide release from prefrontal cortical somatostatin neurons. *Neuroscience* 419:1–4.
- Day JJ (2019) Genetic and epigenetic editing in nervous system. *Dialogues Clin Neurosci* 21:359–368.
- de Groot A, van den Boom BJ, van Genderen RM, Coppens J, van Veldhuijzen J, Bos J, Hoedemaker H, Negrello M, Willuhn I, De Zeeuw CI, Hoogland TM (2020) NINscope, a versatile miniscope for multi-region circuit investigations. *Elife* 9:e49987.
- DeLaney K, Buchberger AR, Atkinson L, Grönder S, Mousley A, Li L (2018) New techniques, applications and perspectives in neuropeptide research. *J Exp Biol* 221:jeb151167.
- DiFrancesco ML, et al. (2020) Neuronal firing modulation by a membrane-targeted photoswitch. *Nat Nanotechnol* 15:296–306.
- Dong Z, Mau W, Feng Y, Pennington ZT, Chen L, Zaki Y, Rajan K, Shuman T, Aharoni D, Cai DJ (2022) Minian, an open-source miniscope analysis pipeline. *Elife* 11:e70661.
- Duffett L, et al. (2022) A genetically encoded sensor for in vivo imaging of orexin neuropeptides. *Nat Methods* 19:231–241.
- Duke CG, Bach SV, Revanna JS, Sultan FA, Southern NT, Davis MN, Carullo NV, Bauman AJ, Phillips RA, Day JJ (2020) An improved CRISPR/dCas9 interference tool for neuronal gene suppression. *Front Genome Ed* 2:9.
- Ellis-Davies GC (2007) Caged compounds: photorelease technology for control of cellular chemistry and physiology. *Nat Methods* 4:619–628.
- Ellis-Davies GCR (2020) Useful caged compounds for cell physiology. *Acc Chem Res* 53:1593–1604.
- Feng J, Zhang C, Lischinsky JE, Jing M, Zhou J, Wang H, Zhang Y, Dong A, Wu Z, Wu H, Chen W, Zhang P, Zou J, Hires SA, Zhu JJ, Cui G, Lin D, Du J, Li Y (2019) A genetically encoded fluorescent sensor for rapid and specific in vivo detection of norepinephrine. *Neuron* 102:745–761.e8.
- Fosque BF, Sun Y, Dana H, Yang CT, Ohyama T, Tadross MR, Patel R, Zlatich M, Kim DS, Ahrens MB, Jayaraman V, Looger LL, Schreiner ER (2015) Neural circuits. labeling of active neural circuits in vivo with designed calcium integrators. *Science* 347:755–760.
- Ghosh KK, Burns LD, Cocker ED, Nimmerjahn A, Ziv Y, Gamal AE, Schnitzer MJ (2011) Miniaturized integration of a fluorescence microscope. *Nat Methods* 8:871–878.
- Gilbert LA, Larson MH, Morsut L, Liu Z, Brar GA, Torres SE, Stern-Ginossar N, Brandman O, Whitehead EH, Doudna JA, Lim WA, Weissman JS, Qi LS (2013) CRISPR-mediated modular RNA-guided regulation of transcription in eukaryotes. *Cell* 154:442–451.
- Giovannucci A, Friedrich J, Gunn P, Kalfon J, Brown BL, Koay SA, Taxisid J, Najafi F, Gauthier JL, Zhou P, Khakh BS, Tank DW, Chklovskii DB, Pnevmatikakis EA (2019) CalmAn an open source tool for scalable calcium imaging data analysis. *Elife* 8:e38173.
- Glaser A, et al. (2023) Expansion-assisted selective plane illumination microscopy for nanoscale imaging of centimeter-scale tissues. *bioRxiv* 544277. <https://doi.org/10.1101/2023.06.08.544277>.
- Gong S, Doughty M, Harbaugh CR, Cummins A, Hatten ME, Heintz N, Gerfen CR (2007) Targeting Cre recombinase to specific neuron populations with bacterial artificial chromosome constructs. *J Neurosci* 27:9817–9823.
- Gunaydin LA, Grosenick L, Finkelstein JC, Kauvar IV, Fenno LE, Adhikari A, Lammel S, Mirzabekov JJ, Airan RD, Zalocusky KA, Tye KM, Anikeeva P, Malenka RC, Deisseroth K (2014) Natural neural projection dynamics underlying social behavior. *Cell* 157:1535–1551.
- Heidenreich M, Zhang F (2016) Applications of CRISPR-Cas systems in neuroscience. *Nat Rev Neurosci* 17:36–44.
- Hökfelt T, Bartfai T, Bloom F (2003) Neuropeptides: opportunities for drug discovery. *Lancet Neurol* 2:463–472.
- Hunker AC, Soden ME, Kravushkina D, Heymann G, Awatramani R, Zweifel LS (2020) Conditional single vector CRISPR/SaCas9 viruses for efficient mutagenesis in the adult mouse nervous system. *Cell Rep* 30:4303–4316.e6.
- Hyun JH, Nagahama K, Namkung H, Mignocchi N, Roh SE, Hannan P, Krüssel S, Kwak C, McElroy A, Liu B, Cui M, Lee S, Lee D, Hugarin RL, Worley PF, Sawa A, Kwon HB (2022) Tagging active neurons by somatargeted Ca-Light. *Nat Commun* 13:7692.
- Jiang F, Doudna JA (2017) CRISPR-Cas9 structures and mechanisms. *Annu Rev Biophys* 46:505–529.
- Jing M, et al. (2020) An optimized acetylcholine sensor for monitoring in vivo cholinergic activity. *Nat Methods* 17:1139–1146.
- Josselyn SA, Tonegawa S (2020) Memory engrams: recalling the past and imagining the future. *Science* 367:eaa4325.
- Juarez B, Kong MS, Jo YS, Elum JE, Yee JX, Ng-Evans S, Cline M, Hunker AC, Quinlan MA, Baird MA, Elerding AJ, Johnson M, Ban D, Mendez A, Goodwin NL, Soden ME, Zweifel LS (2023) Temporal scaling of dopamine neuron firing and dopamine release by distinct ion channels shape behavior. *Sci Adv* 9:eadg8869.
- Juneau J, Duret G, Chu JP, Rodriguez AV, Morozov S, Aharoni D, Robinson JT, St-Pierre F, Kemere C (2020) MiniFAST: a sensitive and fast miniaturized microscope for in vivo neural recording. *bioRxiv* 367466. <https://doi.org/10.1101/2020.11.03.367466>.
- Kalamakis G, Platt RJ (2023) CRISPR for neuroscientists. *Neuron* 111:2282–2311.
- Kellner S, Berlin S (2020) Two-photon excitation of azobenzene photoswitches for synthetic optogenetics. *Appl Sci* 10:805.
- Kim CK, Sanchez MI, Hoerbel P, Fenno LE, Malenka RC, Deisseroth K, Ting AY (2020) A molecular calcium integrator reveals a striatal cell type driving aversion. *Cell* 183:2003–2019.e16.
- Kim D, Park P, Li X, Campos JD, Tian H, Moulton EM, Grimm JB, Lavis L, Cohen AE (2023) Mapping memories: pulse-chase labeling reveals AMPA receptor dynamics during memory formation. *bioRxiv* 541296. <https://doi.org/10.1101/2023.05.26.541296>.
- Kim MW, Wang W, Sanchez MI, Coukos R, von Zastrow M, Ting AY (2017) Time-gated detection of protein-protein interactions with transcriptional readout. *Elife* 6:e30233.
- Koveal D, Rosen PC, Meyer DJ, Díaz-García CM, Wang Y, Cai LH, Chou PJ, Weitz DA, Yellen G (2022) A high-throughput multiparameter screen for accelerated development and optimization of soluble genetically encoded fluorescent biosensors. *Nat Commun* 13:2919.
- Lee D, Creed M, Jung K, Stefanelli T, Wendler DJ, Oh WC, Mignocchi NL, Lüscher C, Kwon HB (2017a) Temporally precise labeling and control of neuromodulatory circuits in the mammalian brain. *Nat Methods* 14:495–503.
- Lee D, Hyun JH, Jung K, Hannan P, Kwon HB (2017b) A calcium- and light-gated switch to induce gene expression in activated neurons. *Nat Biotechnol* 35:858–863.
- Levene MJ, Dombeck DA, Kasischke KA, Molloy RP, Webb WW (2004) In vivo multiphoton microscopy of deep brain tissue. *J Neurophysiol* 91:1908–1912.
- Limaye A, Hall B, Kulkarnil AB (2009) Manipulation of mouse embryonic stem cells for knockout mouse production. *Curr Protoc Cell Biol Chapter* 19:Unit 19.13.1–19.13.24.
- Lin D, Li X, Moulton E, Park P, Tang B, Shen H, Grimm JB, Falco N, Jia BZ, Baker D, Lavis LD, Cohen AE (2023) Time-tagged ticker tapes for intracellular recordings. *Nat Biotechnol* 41:631–639.
- Liu Q, Zhang Z, Zhang W (2021) Optogenetic dissection of neural circuits underlying stress-induced mood disorders. *Front Psychol* 12:600999.
- Ma X, Johnson DA, He XJ, Layden AE, McClain SP, Yung JC, Rizzo A, Bonaventura J, Banghart MR (2023) In vivo photopharmacology with a caged mu opioid receptor agonist drives rapid changes in behavior. *Nat Methods* 20:682–685.
- Mahn M, et al. (2021) Efficient optogenetic silencing of neurotransmitter release with a mosquito rhodopsin. *Neuron* 109:1621–1635.e5.
- Maier W, Corrie JE, Papageorgiou G, Laube B, Grever C (2005) Comparative analysis of inhibitory effects of caged ligands for the NMDA receptor. *J Neurosci Methods* 142:1–9.
- Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M (2018) DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat Neurosci* 21:1281–1289.
- McQuillan HJ, Clarkson J, Kauff A, Han SY, Yip SH, Cheong I, Porteous R, Heather AK, Herbison AE (2022) Definition of the estrogen negative

- feedback pathway controlling the GnRH pulse generator in female mice. *Nat Commun* 13:7433.
- Merighi A (2002) Costorage and coexistence of neuropeptides in the mammalian CNS. *Prog Neurobiol* 66:161–190.
- Mohar B, Grimm JB, Patel R, Brown TA, Tillberg P, Lavis LD, Spruston N, Svoboda K (2022) Brain-wide measurement of protein turnover with high spatial and temporal resolution. *bioRxiv* 516226. <https://doi.org/10.1101/2022.11.12.516226>.
- Morstein J, Hill RZ, Novak AJ, Feng S, Norman DD, Donthamsetti PC, Frank JA, Harayama T, Williams BM, Parrill AL, Tigyi GJ, Riezman H, Isacoff EY, Bautista DM, Trauner D (2019) Optical control of sphingosine-1-phosphate formation and function. *Nat Chem Biol* 15:623–631.
- Morstein J, Dacheux MA, Norman DD, Shemet A, Donthamsetti PC, Citir M, Frank JA, Schultz C, Isacoff EY, Parrill AL, Tigyi GJ, Trauner D (2020) Optical control of lysophosphatidic acid signaling. *J Am Chem Soc* 142:10612–10616.
- Morstein J, Romano G, Hetzler BE, Plante A, Haake C, Levitz J, Trauner D (2022) Photoswitchable serotonin for optical control of the 5-HT_{2A}. *Angew Chem Int Ed Engl* 61:e202117094.
- Mukhopadhyay TK, Morstein J, Trauner D (2022) Photopharmacological control of cell signaling with photoswitchable lipids. *Curr Opin Pharmacol* 63:102202.
- Noguchi J, Nagaoka A, Watanabe S, Ellis-Davies GC, Kitamura K, Kano M, Matsuzaki M, Kasai H (2011) In vivo two-photon uncaging of glutamate revealing the structure–function relationships of dendritic spines in the neocortex of adult mice. *J Physiol* 589:2447–2457.
- Oh SW, et al. (2014) A mesoscale connectome of the mouse brain. *Nature* 508:207–214.
- Parker JG, Marshall JD, Ahanon B, Wu YW, Kim TH, Grewe BF, Zhang Y, Li JZ, Ding JB, Ehlers MD, Schnitzer MJ (2018) Diametric neural ensemble dynamics in parkinsonian and dyskinetic states. *Nature* 557:177–182.
- Patriarchi T, Cho JR, Merten K, Howe MW, Marley A, Xiong WH, Folk RW, Broussard GJ, Liang R, Jang MJ, Zhong H, Dombeck D, von Zastrow M, Nimmerjahn A, Gradinaru V, Williams JT, Tian L (2018) Ultrafast neuronal imaging of dopamine dynamics with designed genetically encoded sensors. *Science* 360:eaat4422.
- Pernpeintner C, Frank JA, Urban P, Roeske CR, Pritzl SD, Trauner D, Lohmüller T (2017) Light-controlled membrane mechanics and shape transitions of photoswitchable lipid vesicles. *Langmuir* 33:4083–4089.
- Pickar-Oliver A, Gersbach CA (2019) The next generation of CRISPR-Cas technologies and applications. *Nat Rev Mol Cell Biol* 20:490–507.
- Platt RJ, et al. (2014) CRISPR-Cas9 knockin mice for genome editing and cancer modeling. *Cell* 159:440–455.
- Pritzl SD, Urban P, Prasselsperger A, Konrad DB, Frank JA, Trauner D, Lohmüller T (2020) Photolipid bilayer permeability is controlled by transient pore formation. *Langmuir* 36:13509–13515.
- Pritzl SD, Konrad DB, Ober MF, Richter AF, Frank JA, Nickel B, Trauner D, Lohmüller T (2022) Optical membrane control with red light enabled by red-shifted photolipids. *Langmuir* 38:385–393.
- Rao S, Chen R, LaRocca AA, Christiansen MG, Senko AW, Shi CH, Chiang PH, Varnavides G, Xue J, Zhou Y, Park S, Ding R, Moon J, Feng G, Anikeeva P (2019) Remotely controlled chemomagnetic modulation of targeted neural circuits. *Nat Nanotechnol* 14:967–973.
- Rapp TL, DeForest CA (2021) Targeting drug delivery with light: a highly focused approach. *Adv Drug Deliv Rev* 171:94–107.
- Rappleye M, Gordon-Fennel A, Castro D, Matarasso A, Zamorano C, Wait S, Lee J, Siebart J, Suko A, Smith N, Muster J, Matreyek K, Fowler D, Stuber G, Bruchas M, Berndt A (2022) Opto-MASS: a high-throughput engineering platform for genetically encoded fluorescent sensors enabling all optical in vivo detection of monoamines and neuropeptides. *BioRxiv* 494241. <https://doi.org/10.1101/2022.06.01.494241>.
- Rynes ML, Surinach DA, Linn S, Laroque M, Rajendran V, Dominguez J, Hadjstamoulou O, Navabi ZS, Ghanbari L, Johnson GW, Nazari M, Mohajerani MH, Kodandaramaiah SB (2021) Miniaturized head-mounted microscope for whole-cortex mesoscale imaging in freely behaving mice. *Nat Methods* 18:417–425.
- Sanchez MI, Nguyen QA, Wang W, Soltesz I, Ting AY (2020) Transcriptional readout of neuronal activity via an engineered Ca²⁺-activated protease. *Proc Natl Acad Sci USA* 117:33186–33196.
- Savell KE, Day JJ (2017) Focus: genome editing: applications of CRISPR/Cas9 in the mammalian central nervous system. *Yale J Biol Med* 90:567.
- Savell KE, Bach SV, Zipperly ME, Revanna JS, Goska NA, Tuscher JJ, Duke CG, Sultan FA, Burke JN, Williams D, Ianov L, Day JJ (2019) A neuron-optimized CRISPR/dCas9 activation system for robust and specific gene regulation. *eNeuro* 6:ENEURO.0495-18.2019.
- Scholefield J, Harrison PT (2021) Prime editing: an update on the field. *Gene Ther* 28:396–401.
- Scott BB, Thiberge SY, Guo C, Tervo DG, Brody CD, Karpova AY, Tank DW (2018) Imaging cortical dynamics in GCaMP transgenic rats with a head-mounted widefield microscope. *Neuron* 100:1045–1058.e5.
- Silva JM, Silva E, Reis RL (2019) Light-triggered release of photocaged therapeutics: where are we now? *J Control Release* 298:154–176.
- Soden ME, Yee JX, Zweifel LS (2023) Circuit coordination of opposing neuropeptide and neurotransmitter signals. *Nature* 619:332–337.
- Stierl M, Stumpf P, Udvari D, Gueta R, Hagedorn R, Losi A, Gärtner W, Petereit L, Efetova M, Schwarzel M, Oertner TG, Nagel G, Hegemann P (2011) Light modulation of cellular cAMP by a small bacterial photoactivated adenylyl cyclase, bPAC, of the soil bacterium *Beggiatoa*. *J Biol Chem* 286:1181–1188.
- Swiech L, Heidenreich M, Banerjee A, Habib N, Li Y, Trombetta J, Sur M, Zhang F (2015) In vivo interrogation of gene function in the mammalian brain using CRISPR-Cas9. *Nat Biotechnol* 33:102–106.
- Taura J, Nolen EG, Cabré G, Hernando J, Squarzialupi L, López-Cano M, Jacobson KA, Fernández-Dueñas V, Ciruela F (2018) Remote control of movement disorders using a photoactive adenosine A_{2A} receptor antagonist. *J Control Release* 283:135–142.
- Tian H, Davis HC, Wong-Campos JD, Park P, Fan LZ, Gmeiner B, Begum S, Werley CA, Borja GB, Upadhyay H, Shah H, Jacques J, Qi Y, Parot V, Deisseroth K, Cohen AE (2023) Video-based pooled screening yields improved far-red genetically encoded voltage indicators. *Nat Methods* 20:1082–1094.
- Urban P, Pritzl SD, Konrad DB, Frank JA, Pernpeintner C, Roeske CR, Trauner D, Lohmüller T (2018) Light-controlled lipid interaction and membrane organization in photolipid bilayer vesicles. *Langmuir* 34:13368–13374.
- Urban P, Pritzl SD, Ober MF, Dirscherl CF, Pernpeintner C, Konrad DB, Frank JA, Trauner D, Nickel B, Lohmueller T (2020) A lipid photoswitch controls fluidity in supported bilayer membranes. *Langmuir* 36:2629–2634.
- Urnov FD, Miller JC, Lee YL, Beausejour CM, Rock JM, Augustus S, Jamieson AC, Porteus MH, Gregory PD, Holmes MC (2005) Highly efficient endogenous human gene correction using designed zinc-finger nucleases. *Nature* 435:646–651.
- Vardy E, et al. (2015) A New DREADD facilitates the multiplexed chemogenetic interrogation of behavior. *Neuron* 86:936–946.
- Venkatachalam V, Brinks D, Maclaurin D, Hochbaum D, Kralj J, Cohen AE (2014) Flash memory: photochemical imprinting of neuronal action potentials onto a microbial rhodopsin. *J Am Chem Soc* 136:2529–2537.
- Wait SJ, Rappleye M, Lee JD, Smith N, Berndt A (2023) Machine learning ensemble directed engineering of genetically encoded fluorescent calcium indicators. *bioRxiv* 536801. <https://doi.org/10.1101/2023.04.13.536801>.
- Wang H, Yang H, Shivalila CS, Dawlaty MM, Cheng AW, Zhang F, Jaenisch R (2013) One-step generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated genome engineering. *Cell* 153:910–918.
- Wang JB, Aryal M, Zhong Q, Vyas DB, Airan RD (2018) Noninvasive ultrasound drug uncaging maps whole-brain functional networks. *Neuron* 100:728–738.e7.
- Wang Y, Wang M, Yin S, Jang R, Wang J, Xue Z, Xu T (2015) NeuroPep: a comprehensive resource of neuropeptides. *Database (Oxford)* 2015:bav038.
- Weinholtz CA, Castle MJ (2021) Intersectional targeting of defined neural circuits by adeno-associated virus vectors. *J Neurosci Res* 99:981–990.
- Wirtshafter HS, Disterhoft JF (2022) In vivo multi-day calcium imaging of CA1 hippocampus in freely moving rats reveals a high preponderance of place cells with consistent place fields. *J Neurosci* 42:4538–4554.
- Xiong H, Lacin E, Ouyang H, Naik A, Xu X, Xie C, Youn J, Wilson BA, Kumar K, Kern T, Aisenberg E, Kircher D, Li X, Zasadzinski JA, Mateo C, Kleinfeld D, Hrabetova S, Slesinger PA, Qin Z (2022) Probing neuropeptide volume transmission in vivo by simultaneous near-infrared light-

- triggered release and optical sensing. *Angew Chem Int Ed Engl* 61: e202206122.
- Xiong H, Alberto KA, Youn J, Taura J, Morstein J, Li X, Wang Y, Trauner D, Slesinger PA, Nielsen SO, Qin Z (2023a) Optical control of neuronal activities with photoswitchable nanovesicles. *Nano Res* 16:1033–1041.
- Xiong H, Wilson BA, Slesinger PA, Qin Z (2023b) Understanding neuropeptide transmission in the brain by optical uncaging and release. *ACS Chem Neurosci* 14:516–523.
- Yamaguchi H, Hopf FW, Li S, Bin de Lecea L (2018) In vivo cell type-specific CRISPR knockdown of dopamine beta hydroxylase reduces locus coeruleus evoked wakefulness. *Nat Commun* 9:5211.
- Yanny K, Antipa N, Liberti W, Dehaeck S, Monakhova K, Liu FL, Shen K, Ng R, Waller L (2020) Miniscope3D: optimized single-shot miniature 3D fluorescence microscopy. *Light Sci Appl* 9:171.
- Yizhar O, Fenno LE, Davidson TJ, Mogri M, Deisseroth K (2011) Optogenetics in neural systems. *Neuron* 71:9–34.
- Zaki Y, Pennington ZT, Morales-Rodriguez D, Francisco TR, LaBanca AR, Dong Z, Lamsifer S, Segura SC, Chen HT, Christenson Wick Z, Silva AJ, van der Meer M, Shuman T, Fenton A, Rajan K, Cai DJ (2023) Aversive experience drives offline ensemble reactivation to link memories across days. *bioRxiv* 532469. <https://doi.org/10.1101/2023.03.13.532469>.
- Zhang Y, et al. (2023) Fast and sensitive GCaMP calcium indicators for imaging neural populations. *Nature* 615:884–891.
- Zong W, Wu R, Chen S, Wu J, Wang H, Zhao Z, Chen G, Tu R, Wu D, Hu Y, Xu Y, Wang Y, Duan Z, Wu H, Zhang Y, Zhang J, Wang A, Chen L, Cheng H (2021) Miniature two-photon microscopy for enlarged field-of-view, multi-plane and long-term brain imaging. *Nat Methods* 18:46–49.