

Lawrence Berkeley National Laboratory

Recent Work

Title

CHEMISTRY OF THE HYDRATED ELECTRON IN OXYGEN-FREE SOLUTIONS OF AMINO ACIDS, PEPTIDES, AND RELATED COMPOUNDS

Permalink

<https://escholarship.org/uc/item/2hk0m7r1>

Authors

Willix, Rodney L.S.
Garrison, Warren M.

Publication Date

1966-11-01

University of California
Ernest O. Lawrence
Radiation Laboratory

**CHEMISTRY OF THE HYDRATED ELECTRON IN
OXYGEN-FREE SOLUTIONS OF AMINO ACIDS,
PEPTIDES, AND RELATED COMPOUNDS**

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to:
Radiation Research

UCRL-17285
Preprint

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

CHEMISTRY OF THE HYDRATED ELECTRON IN
OXYGEN-FREE SOLUTIONS OF AMINO ACIDS,
PEPTIDES, AND RELATED COMPOUNDS

Rodney L. S. Willix and Warren M. Garrison

November 1966

CHEMISTRY OF THE HYDRATED ELECTRON IN OXYGEN-FREE SOLUTIONS OF
AMINO ACIDS, PEPTIDES, AND RELATED COMPOUNDS¹

Rodney L. S. Willix² and Warren M. Garrison

Lawrence Radiation Laboratory
University of California
Berkeley, California

AEC Contract No. W-7405-eng-48

Copies submitted	3
Manuscript pages	21
Tables	2
Figures	4

Rodney L. S. Willix and Warren M. Garrison, Chemistry of the Hydrated Electron in Oxygen-Free Solutions of Amino Acids, Peptides, and Related Compounds
Radiation Res. ____ pp. ____ , ____ .


ABSTRACT

In extending previous studies on the radiation-induced deamination of the simpler α -amino acids in oxygen-free solution it has been found that reductive deamination by the hydrated electron, e_{aq}^- , is not confined to the free amino-acid configuration. The dipeptides, tripeptides, and other derivatives of glycine for example are found to react with e_{aq}^- (essentially quantitatively) to cleave the N-C bond of the primary amino group. Radiolysis studies of variously substituted amino compounds in solutions containing second solutes which are specific scavengers of H, OH in the one case and of e_{aq}^- in the other indicates that reductive deamination by e_{aq}^- requires an unsaturated function at the carbon position α to the amino group. It is suggested that e_{aq}^- adds to this function and that N-C cleavage then ensues. Observed correlations between rates of reaction of e_{aq}^- and pK of the amino group are in accord with this formulation. Velocity constants for reaction of e_{aq}^- with amino acids and derivatives in both acidic and neutral solution were obtained by the method of competition kinetics.


Hydrated electron, amino acids, reductive deamination, reaction rates

INTRODUCTION


Recent work (1,2) has shown that the hydrated electron, e_{aq}^- , which is formed in the radiation-induced decomposition of water (3-5)




reacts with both the zwitterion and cation forms of the simplest α -amino acids, glycine and alanine, via the reductive deamination reaction





and to a lesser extent through the conversion reaction






In acidic solution, reactions 2a, 2b occur in competition with



The radicals OH and H react with both glycine and alanine through hydrogen abstraction at the α -carbon position



which reactions are followed by the oxidative deamination steps³ (1,2,6)



By introducing second solutes which are specific scavengers for OH and H in the one case and for e_{aq}^- in the other, it has been possible to separately evaluate the oxidative and reductive modes of deamination of the simpler α -amino acids (2).

We report here an extension of this earlier work to include other more complex biochemical substances containing primary and secondary amino functions. Effects of pH and of chemical composition on the various modes of reaction of e_{aq}^- are described. Velocity constants for reaction of e_{aq}^- with these organo-nitrogen compounds are derived from measurements of competition kinetics in two-solute systems.

EXPERIMENTAL

The organo-nitrogen compounds (Nutritional Biochemicals) with the exception of N-ethylacetamide (Eastman) were recrystallized from distilled water a minimum of two times. The N-ethyl acetamide was redistilled in vacuo as was the chloroacetic acid (Eastman). Water from a Barnstead still,

redistilled first from alkaline permanganate and then from sulfuric acid was used in the preparation of the solutions. All other chemicals were of reagent grade. The pH adjustments were made with sulfuric acid or sodium hydroxide.


The pyrex irradiation cells containing 10 ml of solution were degassed on the vacuum line, sealed-off, and irradiated with Co^{60} γ rays at a dose rate of 1.6×10^{18} eV/gm/min as determined with the Fricke dosimeter [$G(\text{Fe}^{+++}) = 15.5$, $\epsilon_{305} = 2180$ at 24°C]. Dosage did not exceed 1.5×10^{19} eV/gm; dose-yield relationships were linear over this range.

The Conway diffusion method⁷ was used to separate ammonia from the irradiated solutions. Diffusates were assayed by means of Nessler reagent. Chloride ion was determined by the method of Luce et al. (8) after Hayon and Allen (9). The appropriate blank and control runs were made. Ammonia and chloride measurements were reproducible to within ~5%.

RESULTS AND DISCUSSION

Reaction Rates

Rate constants for reaction of e_{aq}^- with the substances listed in Table I were derived from an analysis of competitive kinetics involving the organo-nitrogen compound at a fixed concentration, 0.05 M,



and a second solute, chloracetic acid, in increasing concentration over the range $\sim 5 \times 10^{-4}$ M to $\sim 5 \times 10^{-2}$ M. The latter reacts with e_{aq}^- according to the stoichiometry (9,12)





to give chloride ion which product is followed analytically. No assumptions regarding the nature of reaction 9 are involved in the rate studies.

All the compounds studied have pK values such that at pH 7 each solute exists almost exclusively as a single species, i.e., the zwitterion form of the α -amino acids, β -amino acids, and dipeptides, the cation form of the simple amines, and the anion forms of the acetylamino acids and the chloroacetic acid. For simplicity we distinguish these proton-deficient species in terms of $\overline{R_3N}$ and $\overline{RCl}$ as defined by the equilibria

$$(\overline{R_3N})(H^+) / (R_3N) = K_{R_3N}$$

$$(\overline{RCl})(H^+) / (RCl) = K_{RCl}$$

The reactions in neutral solution then are written



to distinguish from the corresponding reaction of the acid forms given by Eqs. 9 and 10. For simple competition in these two solute systems at pH 7 we may derive the expression

$$\frac{1}{G(Cl^-)} = \frac{1}{G_{e_{aq}^-}} + \frac{1}{G_{e_{aq}^-}} \left(\frac{k_9}{k_{10}} \right) \left(\frac{\overline{R_3N}}{\overline{RCl}} \right), \quad (11)$$

where $G(\text{Cl}^-)$ represents the experimentally observed chloride yield, $(\overline{\text{R}_3\text{N}})$ and $(\overline{\text{RCl}})$ the concentrations of the two solute-species in neutral solution, and $\bar{k}_9$ and $\bar{k}_{10}$ the respective velocity constants for their reaction with e_{aq}^- . A plot of the reciprocal of the chloride yield as a function of $(\overline{\text{R}_3\text{N}})/(\overline{\text{RCl}})$ gives a straight line with slope $(1/G_{e_{\text{aq}}^-})(\bar{k}_9/\bar{k}_{10})$, as shown by the typical data of Fig. 1.

The intercept value $1/G(\text{Cl}^-) = 0.36$ gives $G_{e_{\text{aq}}^-} \approx 2.8$, in good agreement with published values (5,10). Taking $\bar{k}_{10} = 1.2 \times 10^9 \text{ M}^{-1} \text{ sec}^{-1}$ as determined by Anbar and Hart (11) in pulse radiolysis studies, we obtain the values of $\bar{k}_9$ given in Table I.

A parallel series of experiments was run at pH 3 to obtain rate constants for reactions of e_{aq}^- with these organo-nitrogen compounds in the protonated form. Results are obtained in terms of the rate constant for reaction of e_{aq}^- with the undissociated chloroacetic acid molecule. Corrections must be included for removal of e_{aq}^- by the proton reaction 3, $k_3 = 2.2 \times 10^{10} \text{ M}^{-1} \text{ sec}^{-1}$, and for the competition by $\overline{\text{R}_3\text{N}}$ and $\overline{\text{RCl}}$ at the concentrations determined by the equilibrium constants $K_{\text{R}_3\text{N}}$, K_{RCl} . It can be shown that Eq. 11 then takes the more complicated form

$$\frac{1}{G(\text{Cl}^-)} = \frac{1}{G_{e_{\text{aq}}^-}} + \frac{1}{G_{e_{\text{aq}}^-}} \left\{ \frac{k_9(\text{H}^+) + \bar{k}_9 K_{\text{R}_3\text{N}} + k_3(\text{H}^+)^2/(\text{R}_3\text{N})}{k_{10}(\text{H}^+) + \bar{k}_{10} K_{\text{RCl}}} \right\} \left(\frac{\text{R}_3\text{N}}{\text{RCl}} \right) \quad (13)$$

Here again a plot of the reciprocal chloride ion yield versus $(\text{R}_3\text{N})/(\text{RCl})$ should give a straight line with intercept equal to $1/G_{e_{\text{aq}}^-}$. From the slopes given by



$$\frac{1}{G_{e_{aq}^-}} \left\{ \frac{k_9(H^+) + \bar{k}_9 K_{R_3N} + k_3(H^+)^2 / (R_3N)}{k_{10}(H^+) + \bar{k}_{10} K_{RCl}} \right\}$$

We may calculate the respective values of k_9 , assuming $k_{10} = 6.6 \times 10^9 \text{ M}^{-1} \text{ sec}^{-1}$ as derived from the work of Hayon and Allen (9). Values of k_9 so obtained are listed in Table I.


Some of the rate constants reported in Table I have also been measured by pulse-radiolysis methods in which e_{aq}^- is followed spectrophotometrically (13,14). The latter values are included in parentheses under $\bar{k}_9$ in Table I, and as can be seen the agreement between the two methods is reasonably good. Rate constants for reaction of e_{aq}^- with the cation forms of the amino acids and simple peptides (k_9) are not available as yet from pulse-radiolysis work. Presumably this is because of the very short life-time of e_{aq}^- in aqueous solution at the low pH values needed to retain an appreciable fraction of these solutes in the cation form. At pH 3, for example, assuming $k_3 = 2 \times 10^{10} \text{ M}^{-1} \text{ sec}^{-1}$ the life-time of e_{aq}^- corresponds to $1/(10^{-3})(2 \times 10^{10}) = 0.5 \times 10^{-7} \text{ sec}$.

Chemical Criteria for Reductive Deamination by e_{aq}^- .

We found in the earlier work (1,2) that the yield for reductive deamination of glycine and alanine via reaction 2a could be separately measured through use of a second solute such as formate ion which reacts rapidly with OH and H



$k_{14} = 10^9 \text{ M}^{-1} \text{ sec}^{-1}$, $k_{15} = 2 \times 10^8 \text{ M}^{-1} \text{ sec}^{-1}$ but is relatively unreactive towards e_{aq}^-



$k_{16} = < 10^5 \text{ M}^{-1} \text{ sec}^{-1}$ (15). The preferential scavenging of OH and H via reactions 14 and 15 blocks the formation of the α -carbon radicals $\text{NH}_3^+\dot{\text{C}}\text{R}_2$ which in the absence of formate yield ammonia and carbonyl via steps 6-8. Hence at sufficiently high formate concentrations the observed $G(\text{NH}_3)$ value provides a direct measure of reaction 2a.

We have found, in applying this technique to other systems, that reductive deamination via reaction 2a is not unique to the free α -amino acid configuration. In evaluating the experimental evidence it is convenient here first to compare the ammonia yields from glycine and from the dipeptide glycylglycine at pH 7 in oxygen-free solution under γ rays. We observe, Fig. 2, that $G(\text{NH}_3)$ from both compounds increase abruptly with solute concentration and approach limiting yields under which conditions we may assume that all of the OH, H, e_{aq}^- formed in the radiation-induced reaction are quantitatively scavenged by the solute. The effects of added formate ion on these maximal ammonia yields from glycine and glycylglycine are shown in Fig. 3. In both cases $G(\text{NH}_3)$ decreases with increasing formate concentrations and then levels off to a limiting value which remains essentially constant at the higher formate concentrations. With glycine, the ammonia yield levels off to give $G(\text{NH}_3) = 1.8$ as a measure of the reductive deamination reaction 2a in this system; the yield for conversion of e_{aq}^- to H through reaction 2b corresponds

to $G_{e_{aq}^-} - 1.8 = 1.0$ where, $G_{e_{aq}^-} = 2.8$, represents the yield for production of e_{aq}^- by γ rays in the radiation-induced step 1 (10). This production of H atoms with $G = 1.0$ accounts for the relatively high hydrogen yields observed in the γ radiolysis of neutron solutions of glycine.⁴ The limiting ammonia yield from glycylglycine in the presence of excess formate ion corresponds to $G(NH_3) = 2.5$ and we conclude in this case that the reaction of e_{aq}^- with the glycylglycine zwitterion via reaction 2a is essentially quantitative. The yield for conversion of e_{aq}^- to H by glycylglycine through reaction 2b is given by $G_{e_{aq}^-}$ to H by glycylglycine through reaction (2b) is given by $G_{e_{aq}^-} - 2.5 \leq 0.3$. Confirmation of the above formulation is found in the data of Fig. 4 which shows the effect of a competing electron scavenger, chloracetate ion (reaction 10) on ammonia yields from oxygen-free 0.25 M glycylglycine. The ammonia yield drops rapidly with increasing chloracetate concentration and the decrease in $G(NH_3)$ is accompanied by a corresponding stoichiometric increase in $G(Cl^-)$.

The formate technique has been used to measure the yield of reductive deamination in the γ radiolysis of a number of amino acids and amino acid derivatives in neutral oxygen-free solution. The results are summarized in Table II. We note first that of the compounds studied, only those with an amino group at the carbon position α to a carboxyl, ester, or peptide linkage undergo reductive deamination as a major reaction. Certainly the implication here is that the unsaturated C=O group which is common to all of the above three types of linkage is somehow involved in the deamination reaction; this aspect of the subject is treated in a later section.


We note also from Table II that it is only with glycine and alanine that we observe a marked decrease in $\Delta G(NH_3)$ on addition of the radical scavenger. As we have noted, the reactions of OH and H radicals with these

simplest amino acids occur almost exclusively at the α -carbon position to yield radicals of the type $\text{NH}_3^+\dot{\text{C}}\text{R}_2$. These α -carbon radicals have the property of disproportioning via reactions 6-8 to yield ammonia and carbonyl. On the other hand, the more complex amino acids, valine, for example (Table II) offer competing loci for OH and H attack at the β , γ , ν etc., positions of the side chain. Radicals formed at these sites have the chemical properties of ordinary aliphatic radicals and undergo simple dimerization without the involvement of nitrogen chemistry. Hence, the quenching of these latter reactions by formate ion does not influence $G(\text{NH}_3)$ to as great an extent. The production of diamino acids through combination of radicals formed by radical attack at β , γ , ν side-chain positions of the more complex α -amino acids has been observed by Kopoldova, Liebster, and Babicky (16).

Similarly, the presence of high concentrations of formate ion has but a relatively small effect on $G(\text{NH}_3)$ from glycylglycine, glycylglycylglycine, and glycine ethyl ester. In these cases, also, additional loci are available for OH attack and the evidence is that such attack at the terminal carbon position to give the $\text{NH}_3^+\dot{\text{C}}\text{R}_2$ type radical is relatively unimportant. We have shown elsewhere (17) that peptides are susceptible to OH attack along the main chain to give radicals of the type $\text{RCONH}\dot{\text{C}}\text{R}_2$ which species undergo simple dimerization⁵ to yield the α,α diamino acid derivatives. The present evidence is that reaction of OH with these simple peptide derivatives of glycine occurs preferentially at the C-H bond α to the peptide nitrogen.


Mechanism of Reaction of e_{aq}^- with Amino Compounds

We conclude from the present study that reductive deamination via reaction 2a is a general and characteristic reaction of compounds containing the grouping




when X represents O^- , OH, OR, NHR, etc.⁶ If there is more than one carbon unit between the amino and carbonyl groups reductive deamination does not occur, β -alanine and ϵ -aminocaproic acid are examples of amino compounds that do not undergo reductive deamination. Now, Riesz and Morris (19) found that the simple aliphatic amine cations, methyl ammonium ion, for example, react with e_{aq}^- exclusively via the conversion reaction 2b to yield H. Such reaction is analogous to the conversion of e_{aq}^- to H by NH_4^+ as observed by Jortner, et al. (20) who also showed that the rates of conversion of e_{aq}^- to H by proton donors correlate to a first approximation with the pK values of the donor acids as implied by the Brönsted general acid catalysis law. (The lower the pK the faster the reaction with e_{aq}^- .) And, Braams (13) more recently has studied, by the method of pulse radiolysis, the rate of disappearance of e_{aq}^- in oxygen-free neutral solutions of a variety of simple amines, β -amino acids, α -amino acids and peptides; he finds in all cases a reasonably good correlation between the dissociation constants of the protonated amino groups and their rate constants for reaction with e_{aq}^- . Braams concludes without specifying the nature of the chemistry involved that the protonated amino group of all of these various classes of amino compounds represents the locus of the reaction with e_{aq}^- . It is likely that such is the case for those compounds


that react with e_{aq}^- via the conversion reaction 2b in accord with the earlier work of Jortner, et al. (20). It is not clear that the same correlation between pK and reaction rate in the reductive deamination reactions of the α -amino acids, and their ester and peptide derivatives necessarily implies that e_{aq}^- is reacting at the locus of the amino group. In fact, the finding that an unsaturated double bond must be present α to the amino group for reductive deamination to occur, suggests the possibility that e_{aq}^- adds via




which step is then followed by the dissociation



or the hydrolysis



The recent finding that the glycine- Cu^{+2} chelate undergoes reductive cleavage of the N-C bond on reaction with e_{aq}^- as given by the overall stoichiometry (21)



is also consistent with the interpretation that e_{aq}^- reacts at the C=O bond of the ligand. The formulation 17, 18 is in accord with the finding that the

rate of reaction of e_{aq}^- with the zwitterion form of the α -amino acids is quite low as compared with its rate of reaction with the cation, ester and peptide forms (Table I) since the double bond character of the C=O group of the carboxylate ion is considerably less than that of the C=O group of the acid, ester, and peptide derivatives. While it is true that the rate of reaction of e_{aq}^- with the isolated peptide linkage, N-ethylacetamide, for example, is low (Table I), the presence of the NH_3^+ group in the α -position induces a strong polarization in the C=O bond (22) which would account for the enhanced reactivity of the α -amino acids towards e_{aq}^- . And, since this polarization is manifested in turn as an increase in the acid strength of the NH_3^+ group (22, 23) a correlation between the pK of the amine group and the rate of reaction of e_{aq}^- via paths 17, 18 would be expected.

REFERENCES

1. W. M. Garrison, Actions of ionizing radiations on nitrogen compounds in aqueous media, Radiation Res. Suppl. 4, 158-174 (1964).
2. B. M. Weeks, S. A. Cole, and W. M. Garrison, Reactions of alanine with the reducing species formed in water radiolysis, J. Phys. Chem. 69, 4131-4137 (1965).
3. G. Czapski and H. A. Schwarz, The nature of the reducing species formed in water radiolysis, J. Phys. Chem. 66, 471-474 (1962).
4. E. J. Hart and J. W. Boag, J. Am. Chem. Soc. 84, 4090 (1962).
5. A. O. Allen, Radiation yields and reactions in dilute inorganic solutions, Radiation Res. Suppl. 4, 54-73 (1964).
6. B. M. Weeks and W. M. Garrison, Radiolysis of aqueous solutions of glycine, Radiation Res. 9, 291-304 (1958).
7. E. J. Conway, Microdiffusion Analysis, (Crosby Lockwood and Son, Ltd., London, 1962) p. 90.
8. E. N. Luce, E. C. Denice, and F. E. Akerlund, Turbidimetric determination of small amounts of chlorides, Ind. Eng. Chem. 15, 365-366 (1943).
9. E. Hayon and A. O. Allen, Evidence for two kinds of "H atoms" in the radiation chemistry of water, J. Phys. Chem. 65, 2181-2185 (1961).
10. E. Hayon, Solute scavenging effects in regions of high radical concentration produced in radiation chemistry, Trans. Faraday Soc. 61, 723-733 (1965).
11. M. Anbar and E. J. Hart, The reaction of haloaliphatic components with the hydrated electron, J. Phys. Chem. 69, 271-274 (1965).
12. E. Hayon and J. Weiss, The actions of ionizing radiations on aqueous solutions of mono and trichloroacetic acids, Proc. 2nd Intern. Conf. Peaceful Uses Atomic Energy, Geneva, 29, 80-85 (1958).

13. R. Braams, Reactions of the hydrated electron with amino acids, peptides, and protein, in Pulse Radiolysis, (M. Ebert, et al., ed.) (Academic Press, New York, 1965) p. 171-180.
14. J. V. Davies, M. Ebert, and A. J. Swallow, Reactions of hydrated electrons with glycine and other amino acids and peptides, in Pulse Radiolysis, (M. Ebert, et al., ed.) (Academic Press, New York, 1965) p. 165-170.
15. E. J. Hart, J. C. Thomas, and S. Gordon, A review of the radiation chemistry of single carbon compounds and some reactions of the hydrated electron in aqueous solution, Radiation Res. Suppl. 4, 74-88 (1964).
16. J. Kopoldova, J. Liebster, and A. Babicky, Degradation of α -aminobutyric acid in aqueous solution, J. Appl. Rad. and Isotopes 13, 617-622 (1962).
17. W. M. Garrison and B. M. Weeks, Radiation chemistry of compounds containing the peptide bond, Radiation Res. 17, 341-352 (1962).
18. P. G. Clay and A. Kabi, The radiation-induced decomposition of N-benzyl-dimethylamine and benzyltrimethylammonium ion in aqueous solution, Chem. and Ind. 226-227 (1965).
19. P. Riesz and T. Morris, The radiolysis of aqueous methylammonium ion, Radiation Res. 26, 1-11 (1965).
20. J. Jortner, M. Ottolenghi, J. Rabani, and G. Stein, Conversion of solvated electrons into hydrogen atoms in the photo- and radiation-chemistry of aqueous solutions, J. Chem. Phys. 37, 2488-2495 (1962).
21. R. L. S. Willix and W. M. Garrison, The effect of cupric ion on the radiation chemistry of aqueous glycine, J. Phys. Chem. 69, 1579-1583 (1965).
22. E. S. Gould, Mechanism and Structure in Organic Chemistry, (Henry Holt and Co., New York, 1959).
23. J. P. Greenstein and M. Winitz, Chemistry of the Amino Acids, (John Wiley and Sons, Inc., New York, 1961).

FOOTNOTES

1. Work performed under the auspices of the U. S. Atomic Energy Commission.
2. Present address: Department of Physiology, University of Western Australia, Perth, Western Australia.
3. Some dimerization of $\text{NH}_3^+\dot{\text{C}}\text{R}_2$ radicals does occur but the yield of the α, α diamino acid is low (see Ref. 6).
4. For γ rays the yields of H and H_2 in the radiation decomposition of water, reaction 1 corresponds to $G_{\text{H}} = 0.45$, $G_{\text{H}_2} = 0.55$. With neutral 1 M glycine under γ rays $G(\text{H}_2) = 2.2$ which value is in good agreement with $G_{\text{H}} + G_{\text{H}_2} + 1.0$ (Ref. 2').
5. The radicals $\text{RCONH}\dot{\text{C}}\text{R}_2$ and $\text{NH}_3^+\dot{\text{C}}\text{R}_2$ (or $\text{NH}_2\dot{\text{C}}\text{R}_2$) differ fundamentally in their chemistry. In oxygen-free solution the former dimerize while the latter disproportionate (see Ref. 17).
6. The recent results of Clay and Kobi (Ref. 18) indicate that e_{aq}^- reacts with benzyldimethylamine cation and with benzyltrimethyl ammonium ion to yield dimethyl amine and trimethylamine respectively. This suggests the reactive grouping in the general case corresponds to $\text{R}_2\text{NH}^+-\text{C}(\text{R}_2)-\text{C} \begin{smallmatrix} \text{X} \\ \text{Y} \end{smallmatrix}$.

Table I

Rate Constants for Reaction of e_{aq}^- with Amino Acids and Derivatives
as Measured by Competition Kinetics.^a

Compound	pH 6.7, $\bar{k}_9$ ($M^{-1} \text{ sec}^{-1}$)	pH 3, k_9 ($M^{-1} \text{ sec}^{-1}$)
glycine	1.4×10^7 (0.9×10^7) ^{b,c}	4.5×10^8
alanine	1.8×10^7 (0.6×10^7) ^{b,c}	8.0×10^8
bis-glycinato-CuII	3.5×10^8	---
glycine ethyl ester	1.0×10^9	---
diglycine	1.0×10^8 (2.5×10^8) ^b	8.9×10^8
triglycine	7.2×10^8 (9×10^8) ^b	3.0×10^9
N-ethylacetamide	1.7×10^7	1.5×10^7
N-acetyl alanine	1.1×10^7	1.2×10^8
ethyl amine		$\sim 10^6$

^aBased on $\bar{k}_{10} = 1.2 \times 10^9 M^{-1} \text{ sec}^{-1}$, $k_{10} = 6.6 \times 10^9 M^{-1} \text{ sec}^{-1}$.

^bPulse radiolysis, Ref. 13.

^cPulse radiolysis, Ref. 14.

Table II

Effect of 0.8 M Sodium Formate on $G(\text{NH}_3)$ from Oxygen-Free
Solutions of Amino Acids and Derivatives at pH 7.

Compound ^a	$G(\text{NH}_3)$	
	Formate-free	1.0 M formate
alanine	4.3	2.5
glycine	4.0	1.8
glycine ethyl ester	2.7	2.6
bis-glycinato-CuII ^b	5.0	3.2
glycylglycine	3.0	2.5
valine ^c	1.8	0.8
ϵ -aminocaproic acid	0.35	<0.35
β -alanine	0.75	0.45


^aAt the minimum concentration required to insure the quantitative scavenging of water decomposition-products (see Fig.).

^bAt pH 8.5.

^cMeasured at only one valine concentration, viz 0.25 M.


FIGURE CAPTIONS

- Fig. 1. Typical plots of the reciprocal chloride-ion yields as a function of $(R_3N)/(RCl)$ or $(\overline{R_3N})/(\overline{RCl})$ for glycine zwitterion (Δ), glycine cation ($\triangle$), glycylglycine zwitterion ($\square$), glycine ethyl ester ($\boxtimes$), ethylamine ($\odot$).
- Fig. 2. Effect of glycine ($\circ$) and glycylglycine ($\odot$) concentrations on ammonia yields from oxygen-free solutions, pH 6.5.
- Fig. 3. Effect of formate concentrations on ammonia yields from 1.0 M glycine ($\circ$), 0.20 M glycylglycine ($\odot$), oxygen-free, pH 6.5.
- Fig. 4. Effect of chloracetate concentration on ammonia ($\odot$) and chloride ion ($\triangle$) yields from 0.20 M glycylglycine, oxygen-free, pH 6.5.




MUB 13909

Fig. 1




MUB13910

Fig. 2



MUB-13911

Fig. 3



MUB 13912

Fig. 4

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

