

UCSF

UC San Francisco Previously Published Works

Title

Evidence for Local Antibiotics in the Prevention of Infection in Orthopaedic Trauma

Permalink

<https://escholarship.org/uc/item/2j4594jg>

Journal

Journal of Clinical Medicine, 11(24)

ISSN

2077-0383

Authors

Flores, Michael J

Brown, Kelsey E

Morshed, Saam

et al.

Publication Date

2022

DOI

10.3390/jcm11247461

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Review

Evidence for Local Antibiotics in the Prevention of Infection in Orthopaedic Trauma

Michael J. Flores ^{*}, Kelsey E. Brown, Saam Morshed and David W. Shearer ^{*}

Institute for Global Orthopaedics and Traumatology, Department of Orthopaedic Surgery,
University of California, San Francisco, CA 94110, USA

^{*} Correspondence: michael.flores@yale.edu (M.J.F.); david.shearer@ucsf.edu (D.W.S.)

Abstract: Prevention of fracture-related infection (FRI) remains a substantial challenge in orthopaedic trauma care. There is evolving evidence to support the use of local antibiotics for both the prevention and treatment of musculoskeletal infection. Local antibiotics can achieve higher local tissue concentrations with a lower risk of systemic complications compared to intravenously administered antibiotics. These antibiotics may be administered in powder or liquid form without carrier, or if sustained release is desired, using a carrier. Polymethylmethacrylate (PMMA), ceramics, and hydrogels are examples of antibiotic carriers. Unlike PMMA, ceramics and hydrogels have the advantage of not requiring a second surgery for removal. The VANCO trial supported the use of powdered vancomycin in high-risk fracture cases for the reduction of Gram-positive infections; although, data is limited. Future studies will evaluate the use of aminoglycoside antibiotics to address Gram-negative infection prevention. While theoretical concerns exist with the use of local antibiotics, available studies suggest local antibiotics are safe with a low-risk of adverse effects.

Keywords: fracture-related infection (FRI); orthopaedic trauma; antibiotics; orthopedics fracture healing; bony callus; bone infection



Citation: Flores, M.J.; Brown, K.E.; Morshed, S.; Shearer, D.W. Evidence for Local Antibiotics in the Prevention of Infection in Orthopaedic Trauma. *J. Clin. Med.* **2022**, *11*, 7461. <https://doi.org/10.3390/jcm11247461>

Academic Editors: Roman Pfeifer and Shunsuke Mori

Received: 18 September 2022

Accepted: 9 December 2022

Published: 16 December 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Injury accounts for over 10% of the global disease burden, with trauma accounting for the majority of injury-related disability worldwide [1]. Additionally, it has been found that the global incidence of fractures and fracture-related disability have increased substantially in recent years [2]. Since high-energy mechanisms often accompany traumatic injury, they are particularly prone to infection due to high rates of open injuries with soft tissue damage and the necessity for operative management. Even after operative debridement, infection rates have been cited anywhere between 4–63% [3–5]. Although less common, closed injuries have also been associated with infection [6]. Surgical site infections (SSIs) can be particularly difficult to treat, since some of the common bacterial strains causing SSI have the ability to create a protective biofilm that shields them from host immunity and systemic antibiotics [6]. These infections can lead to devastating morbidity, including delayed wound healing, fracture nonunion, amputation, and even death [7,8].

Early administration of intravenous antibiotics and thorough debridement remain the gold standard interventions to reduce infection after open fracture [9,10]. Although prophylactic systemic antibiotics have been shown to reduce infection rates, high rates of infection persist despite their use [3,11,12]. In an effort to further reduce the risk of infection, the introduction of local antibiotics into the wound after appropriate irrigation and debridement has been implemented with promising results [11,13–17]. This narrative review outlines the evolving evidence related to the use of local antibiotics in the prevention of infection in the setting of orthopaedic trauma.

2. Methods

Evidence for the current narrative review was based on articles obtained through a MEDLINE search (September 2021) using the key words “Surgical Wound Infection” (MeSH term), “Antibiotic Prophylaxis” (MeSH term), and “Local antibiotic” (text word). As the current study was not a systematic review, recommended articles and viewpoints from experts in the field of orthopaedic trauma were also included.

3. Rationale

The use of local antibiotics during surgical procedures has been ongoing for over a century, stemming from the pioneering work of Joseph Lister [6]. Despite their early introduction in the setting of established infections, systemic antibiotics have become the gold standard for prophylaxis as well as therapeutic antibiotics, with much less written about the use of local antibiotics [6,9,10]. However, renewed attention has been placed on the role of prophylactic local antibiotics in recent years, as high rates of surgical site infections (SSIs) have persisted [7,8].

Local antibiotics have potential advantages in the setting of traumatic injury for a variety of reasons. Vascular injury, local tissue damage, or periosteal stripping may all limit the delivery of intravenous antibiotics to the injury site [11]. In these instances, locally administered antibiotics can be more effective in achieving high concentrations at the site of injury and contamination. Additionally, local antibiotics can be administered at much higher local concentrations with lower systemic levels, decreasing the risk of complications such as nephrotoxicity or ototoxicity [3,6]. These high local concentrations of antibiotics have recently been shown not only to reduce the risk of infection, but also decrease biofilm creation and bacterial resistance [8,11,18]. Finally, the use of local antibiotics in conjunction with systemic antibiotics may result in a synergistic effect to further decrease the risk of SSI [19].

The pharmacokinetics of local antibiotics without a carrier are less studied than antibiotics administered intravenously or with a carrier. Studies of locally-applied vancomycin in spine surgery in both adults and children have shown antibiotic levels 1000-fold higher than MICs for common pathogens that persist for 2–3 days post-operatively with relatively low systemic concentrations [20,21]. However, the pharmacokinetics in trauma populations and the potential impact of impurities in local antibiotics are not well understood.

4. Carriers

Local antibiotics can be delivered in multiple ways including with or without a carrier. A recent review published by the Fracture-Related Infection (FRI) consensus group summarises these strategies [22]. Additionally, an overview of the different carriers discussed in the current review can be found in Table 1. Local antibiotic carriers can be both absorbable and non-absorbable. Common carriers include polymethyl methacrylate (PMMA), biodegradable ceramics such as calcium sulfate (CaSO_4), and hydrogels. Carriers provide a means for burst, tapered, and sustained release of antibiotics [22,23]. However, there is some concern for increased risk of the development of antibiotic resistance with carrier use due to sustained release at subtherapeutic levels [24]. Antibiotic resistance is critical to consider when using antibiotic drugs and carriers, and they should be properly assessed for safety and efficacy through high-level evidence studies and proper regulation prior to their general recommendation and use. Additionally, variations in carrier type and formulation, as well as the antibiotics used, can alter antibiotic elution and dilution profiles and the mechanical properties of their carriers [22]. It should be noted that although PMMA and other carriers are commonly used with local antibiotics, this is not an FDA approved use of these products, and hence this off-label use and associated risks should be discussed with patients prior to their application. It is also noteworthy that due to cost, most carriers are utilised for the treatment rather than prevention of infection.

PMMA is the oldest and most common carrier available. Originally designed as a bone cement, it was quickly adopted as an antibiotic carrier in the 1970's, especially for the

treatment of osteomyelitis due its ability to occupy large amounts of dead space [23,25]. PMMA is a solid rigid polymer that can be formed into beads or a mono-block spacer. During the curing phase, antibiotics can be incorporated into the polymerized matrix [22,23,26] (Figure 1). Additionally, antibiotic-infused PMMA has been used to coat orthopaedic implants [27,28] (Figures 2–4). The polymer porosity, antibiotic thermal stability, antibiotic class, and use of fillers all contribute to the final PMMA-mediated antibiotic strength and timing of antibiotic release [22,29–31]. It is important to recognise that the addition of antibiotics to PMMA leads to a non-linear decrease in strength, which may be relevant if a load-bearing spacer is desired. Although many antibiotics have been used with PMMA, vancomycin, tobramycin, and gentamicin are the most commonly used, and when in combination, have been shown to provide a synergistic effect [22,23]. Many studies have shown that local PMMA-mediated antibiotics decrease infection risk and biofilm formation [16,23,32]. However, because PMMA is not absorbable, it typically requires a second surgical procedure for removal. In addition, although its ability to occupy dead space can be an advantage, in cases without bone loss, there may not be sufficient space to allow for the placement of PMMA. For these reasons, PMMA is more commonly used either as prophylaxis in high-energy open fracture situations with bone or soft-tissue loss or in the treatment of infection in the setting of osteomyelitis and fracture-related infection.



Figure 1. PMMA beads can be created by a hand over suture or wire, with or without the use of prefabricated templates. Beads provide no structural integrity but effectively occupy dead space and provide a higher level of antibiotic elution due to greater surface area than mono-block creations.




Figure 2. A PMMA-coated intramedullary nail can be fabricated using a chest tube or a threaded rod or guidewire. After mixing, the cement is injected into the chest tube and the rod is inserted. The tubing is then removed after the PMMA has hardened using a scalpel.




Figure 3. A clinical case of a 27-year-old woman with infected nonunion of the tibia after open fracture (A). She underwent initial irrigation and debridement with removal of hardware and placement of an antibiotic nail, antibiotic beads, and external fixation (B). The beads are ideal in this scenario for maximal surface area and antibiotic elution. At the same time, the antibiotic nail provides stability and local antibiotic delivery in the intramedullary space. A repeat debridement was performed with the placement of a mono-block spacer in the subsequent procedure (C). The mono-block spacer has the advantage of inducing a membrane for subsequent bone grafting and improved structural integrity. Healing of the fracture is shown after bone grafting with the placement of a new intramedullary nail (D).




Figure 4. Clinical images of the same patient are shown at the time of antibiotic spacer placement (A), spacer removal (B), nail placement (C), and bone grafting (D).

Ceramics have been shown to have similar outcomes to PMMA, with the primary advantage of being absorbable and hence not requiring a secondary surgery for removal [33,34]. Common ceramics are derived from formulations of calcium sulfate or calcium phosphate. Although calcium sulfate resorbs at a much faster rate, the antibiotic elution profiles of both ceramics are similar [22,35]. However, resorption time should be taken into account when using ceramics, as slower resorption times can be useful in persistent infections, but can potentially be a nidus for future infection [35]. Although their use in local antibiotic prophylaxis is relatively new [23], they have been shown to maintain antibiotic concentrations longer than PMMA, leading to decreased infection and biofilm formation [34,36]. Like PMMA, ceramics have been proven effective as an antibiotic carrier in other situations such as in treating osteomyelitis [33,37,38].

Hydrogels are emerging as an alternative antibiotic carrier with many potential advantages, but there have been less clinical studies investigating their efficacy [22]. Hydrogels

are a polymer like PMMA but have the advantage of being absorbable like ceramics [22]. Additionally, they have been shown to not interfere with bone healing [39]. Hydrogels have a shorter release period due to their rapid resorption and lack the same structural integrity as other carriers due to their gel-like consistency [22]. This makes them better suited for prophylactic situations where it is less likely for dead space to be present and longer antibiotic elution periods are not needed. Since hydrogels are highly hydrophilic with a unique scaffold, they can be prepared for various situations and adapt or react to various environmental changes [40]. This makes them especially useful for targeted effects.

Table 1. Common Antibiotic Carrier Characteristics.

Carrier	Materials	Antibiotics	Benefits	Disadvantages	References
PMMA	Polymethylmethacrylate	Heat stable only (Aminoglycosides, Glycopeptides, Tetracyclines, and Quinolones)	Availability, Occupies dead space	Not absorbable, long elution profile	[22,23,26]
Ceramics	Calcium sulfate, calcium phosphate, or a combination	Aminoglycosides Glycopeptides Lipopeptides	Absorbable, faster resorption and elution profiles	Possible toxicity or hypercalcemia (rare), wound drainage, cost	[22,34,35]
Hydrogels	PCLA-PEG-PCLA tri-block, poly(ether ester) SynBiosys, etc.	Variable	Absorbable, fast resorption, variety	Shorter release period, lack structural integrity, cost	[22,39,40]

So-called “naked” local antibiotics, including aqueous or powder formulations, deliver antibiotics without a carrier. Aqueous formulations are one of the earliest described forms of local antibiotics in the literature and are injected into the wound after wound closure; whereas, powdered formulations are placed into the wound prior to closure. These methods are advantageous in that they cost less than other delivery methods; however, their effect is shorter lasting [22]. For example, local vancomycin powder is cited in literature to range between \$2.50/g to \$44/g (€3.22/g to €56.72/g using the average 2017 exchange rate of 1.289 USD) [41,42]. In contrast, one cost-effective analysis from the Netherlands showed the mean cost of PMMA per patient with osteomyelitis to be €365.00, while the use of a sulfur-based ceramic glass was €2007.20 [43]. Another economic analysis in Italy documented the price of PMMA for prosthetic implants to be €480, while an antibiotic hydrogel costs €1170 [44].

There are multiple variables to consider in deciding which antibiotic to pair with the different carriers. It is important to choose an antibiotic that acts against the pathogen(s) identified in the wound culture or can effectively act against a wide range of pathogens. Additionally, care must be taken to ensure the antibiotic has a proper toxicity and hypersensitivity profile at the site and a low rate of resistance from the pathogen of interest [29]. In the case of PMMA, which uses an exothermic reaction to create the polymer, thermal stability of the antibiotic is also necessary. It has been shown that beta-lactam antibiotics are not heat stable and should not be used with PMMA, while common heat stable antibiotics include aminoglycosides, glycopeptides, tetracyclines, and quinolones [22,29]. Although certain ceramic formulations also go through an exothermic reaction during their synthesis, the materials and processes vary. Currently, the antibiotics used in ceramic carriers include aminoglycosides, glycopeptides, and lipopeptides [35]. In contrast, since hydrogels are a water-soluble polymer with various preparations, many different antimicrobial substances have been incorporated into hydrogel polymers including metal nanoparticles and antibiotics [40].

5. Effectiveness in Trauma

There is evolving evidence regarding the general efficacy of prophylactic local antibiotic administration. While there are many observational studies that validate the efficacy of

local antibiotics in the prevention of SSI in high-risk fractures, there is only one published prospective randomised controlled clinical trial (RCT) to evaluate their efficacy [45].

The Local Antibiotic Therapy to Reduce Infection After Operative Treatment of Fractures at High Risk of Infection (VANCO) trial evaluated the efficacy of local vancomycin in preventing surgical site infections after fracture surgery [11]. This multi-centre clinical trial collected data from 34 US trauma centres who participated in the Major Extremity Research Consortium (METRC). The trial included participants aged 18–80 years with closed or open tibial plateau and pilon fractures requiring staged treatment and randomised them to either receive standard of care or the placement of 1 gram of vancomycin locally prior to wound closure. The time-to-event estimates for surgical site deep infection rates at 6 months follow up were 6.4% in the group who received local vancomycin and 9.8% in the control group ($p = 0.06$). In a post-hoc subgroup analysis, the rate of a Gram-positive infection was reduced from 6.8% to 3.3%, ($p = 0.02$) while a Gram-negative infection was equal between the two groups. It is important to note that significance in the VANCO trial was only achieved in this subgroup analysis. Additionally, this subgroup analysis was not pre-specified and hence should be interpreted with caution. However, the findings of this single level 1 trial generally support the use of vancomycin in the reduction of Gram-positive SSI in high-risk fractures.

As a follow up to the VANCO trial, the METRC group is beginning to enrol a new multicentre trial, the Topical Antibiotic Therapy to Reduce Infection After Operative Treatment of Fractures at High Risk of Infection (TOBRA) study. The TOBRA trial will compare the efficacy of combining local vancomycin and tobramycin to local vancomycin alone in the prevention of deep SSI in high-risk fractures [46]. The target enrolment for the study is 1900 participants, and results are anticipated to be available in May 2024 [46].

Lawing et al. described the use of a local gentamicin injection in open fractures to prevent infection and reported their results in an observational study before and after implementation of its use [3]. The study found that the deep and superficial infection rate was reduced from 19.7% to 9.5% ($p = 0.010$). Although the study was not randomised, after adjustment using a regression model, local antibiotics were an independent predictor of lower infection risk [3].

The Local Gentamicin for Open Tibial Fractures in Tanzania (GO-Tibia) trial is an ongoing RCT being conducted at the Muhimbili Orthopaedic Institute in Dar as Salaam, Tanzania in collaboration with the Institute for Global Orthopaedics and Traumatology (IGOT) at the University of California, San Francisco [47]. The GO Tibia Trial is aimed to evaluate the efficacy of locally-injected gentamicin on the risk of fracture-related infection (FRI) after open tibia fractures [47]. In contrast to the VANCO and TOBRA trials, the GO Tibia study is masked with a saline placebo control. A pilot trial was recently completed (publication pending), and enrolment in a larger definitive trial is scheduled to begin in January 2022.

6. Potential Limitations

Although prophylactic antibiotics may reduce the risk of infection [3,11,15,16,48,49], some studies have demonstrated inconclusive results [50,51]. Furthermore, as demonstrated in this review, there are currently few high-level evidence studies on prophylactic antibiotic use in musculoskeletal trauma. In spine literature, there seems to be little added benefit of local vancomycin in low-risk cases where infection rates are already low [50]. Furthermore, high antibiotic concentrations may be toxic to local tissues. One in-vitro study showed that local antibiotics may have osteocyte toxicity [52], but animal studies have not shown an effect on bone healing [53]. Additionally, studies in humans have not shown an association between local antibiotics and nonunion or delayed healing. Another potential limitation to local antibiotic use is the development of antibiotic resistance with prophylactic usage, particularly with second-line antibiotics such as vancomycin. However, local use results in low systemic levels for short durations, hence local use would be anticipated to have lower risk of developing resistance than systemic antibiotics. One theoretical

concern with local vancomycin is that impurities in the formulation may act as a nidus for the development of future infections. Another concern is the selection of bacteria, such as Gram-negative organisms, with local vancomycin. However, a meta-analysis on this topic has shown no evidence of selection of pathogens [17]. The TOBRA study will further address this issue by broadening coverage. Finally, the use of prophylactic local antibiotics is an off-label use, and as such, may cause unforeseen adverse events. Additionally, this unregulated use of local antibiotic prophylaxis leads to a lack of quality control, standardisation, and limited understanding of elution and diffusion characteristics. Although there has been concern for the occurrence of adverse events such as nephrotoxicity, ototoxicity, and hypersensitivity reactions with the use of local antibiotics [54,55], there has been little evidence of these complications [19,56,57]. However, given that the potential limitations and complications of prophylactic local antibiotic use are currently addressed by lower-level evidence studies, the use of prophylactic local antibiotics should be considered with caution until further evidence from randomised controlled trials and regulatory bodies have been obtained.

7. Conclusions

There is evolving evidence supporting the use of prophylactic local antibiotics in the prevention and treatment of infections in orthopaedic trauma. Many studies show promising results of their efficacy with few, if any, reported adverse effects despite theoretical concerns. However, current high-level research is limited, and further well-designed randomised controlled trials are needed before definitive recommendations can be made in regard to their use. Greater awareness and appropriate use of local antibiotics has the potential to reduce the global impact of fracture-related infection.

Author Contributions: Conceptualization, D.W.S. and S.M., Literature Review, M.J.F. and K.E.B., Writing—original draft preparation, M.J.F. and D.W.S., Writing—review and editing, M.J.F., K.E.B., S.M. and D.W.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Roser, M.; Ritchie, H. Burden of Disease. Available online: <https://ourworldindata.org/burden-of-disease> (accessed on 1 April 2022).
2. Wu, A.-M.; Bisignano, C.; James, S.L.; Abady, G.G.; Abedi, A.; Abu-Gharbieh, E.; Alhassan, R.K.; Alipour, V.; Arabloo, J.; Asaad, M.; et al. Global, Regional, and National Burden of Bone Fractures in 204 Countries and Territories, 1990–2019: A Systematic Analysis from the Global Burden of Disease Study 2019. *Lancet Healthy Longev.* **2021**, *2*, e580–e592. [[CrossRef](#)] [[PubMed](#)]
3. Lawing, C.R.; Lin, F.-C.; Dahners, L.E. Local Injection of Aminoglycosides for Prophylaxis Against Infection in Open Fractures. *J. Bone Jt. Surg.* **2015**, *97*, 1844–1851. [[CrossRef](#)] [[PubMed](#)]
4. Schenker, M.L.; Yannascoli, S.; Baldwin, K.D.; Ahn, J.; Mehta, S. Does Timing to Operative Debridement Affect Infectious Complications in Open Long-Bone Fractures? A Systematic Review. *J. Bone Jt. Surg.* **2012**, *94*, 1057–1064. [[CrossRef](#)] [[PubMed](#)]
5. Malhotra, A.K.; Goldberg, S.; Graham, J.; Malhotra, N.R.; Willis, M.C.; Mounasamy, V.; Guilford, K.; Duane, T.M.; Aboutanos, M.B.; Mayglothling, J.; et al. Open Extremity Fractures: Impact of Delay in Operative Debridement and Irrigation. *J. Trauma Acute Care Surg.* **2014**, *76*, 1201–1207. [[CrossRef](#)]
6. Cancienne, J.M.; Burrus, M.T.; Weiss, D.B.; Yarboro, S.R. Applications of Local Antibiotics in Orthopedic Trauma. *Orthop. Clin. N. Am.* **2015**, *46*, 495–510. [[CrossRef](#)]
7. Redfern, J.; Wasilko, S.M.; Groth, M.E.; Mcmillian, W.D.; Bartlett, C.S. Surgical Site Infections in Patients with Type 3 Open Fractures: Comparing Antibiotic Prophylaxis with Cefazolin plus Gentamicin versus Piperacillin/Tazobactam. *J. Orthop. Trauma* **2016**, *30*, 415–419. [[CrossRef](#)]
8. Metsemakers, W.J.; Kuehl, R.; Moriarty, T.F.; Richards, R.G.; Verhofstad, M.H.J.; Borens, O.; Kates, S.; Morgenstern, M. Infection after Fracture Fixation: Current Surgical and Microbiological Concepts. *Injury* **2018**, *49*, 511–522. [[CrossRef](#)]
9. Chang, Y.; Kennedy, S.A.; Bhandari, M.; Lopes, L.C.; Bergamaschi, C.d.C.; Carolina de Oliveira e Silva, M.; Bhatnagar, N.; Mousavi, S.M.; Khurshid, S.; Petrisor, B.; et al. Effects of Antibiotic Prophylaxis in Patients with Open Fracture of the Extremities: A Systematic Review of Randomized Controlled Trials. *JBJS Rev.* **2015**, *3*, e2. [[CrossRef](#)]
10. Barker, F.G. Efficacy of Prophylactic Antibiotic Therapy in Spinal Surgery: A Meta-Analysis. *Neurosurgery* **2002**, *51*, 391–400. [[CrossRef](#)]

11. Major Extremity Trauma Research Consortium (METRC); O'Toole, R.V.; Joshi, M.; Carlini, A.R.; Murray, C.K.; Allen, L.E.; Huang, Y.; Scharfstein, D.O.; O'Hara, N.N.; Gary, J.L.; et al. Effect of Intrawound Vancomycin Powder in Operatively Treated High-Risk Tibia Fractures: A Randomized Clinical Trial. *JAMA Surg.* **2021**, *156*, e207259. [[CrossRef](#)]
12. Harris, A.M.; Althausen, P.L.; Kellam, J.; Bosse, M.J.; Castillo, R. Lower Extremity Assessment Project (LEAP) Study Group Complications Following Limb-Threatening Lower Extremity Trauma. *J. Orthop. Trauma* **2009**, *23*, 1–6. [[CrossRef](#)] [[PubMed](#)]
13. Carver, D.C.; Kuehn, S.B.; Weinlein, J.C. Role of Systemic and Local Antibiotics in the Treatment of Open Fractures. *Orthop. Clin. N. Am.* **2017**, *48*, 137–153. [[CrossRef](#)] [[PubMed](#)]
14. O'Toole, R.V.; Joshi, M.; Carlini, A.R.; Murray, C.K.; Allen, L.E.; Scharfstein, D.O.; Gary, J.L.; Bosse, M.J.; Castillo, R.C.; METRC. Local Antibiotic Therapy to Reduce Infection After Operative Treatment of Fractures at High Risk of Infection: A Multicenter, Randomized, Controlled Trial (VANCO Study). *J. Orthop. Trauma* **2017**, *31*, S18–S24. [[CrossRef](#)] [[PubMed](#)]
15. Morgenstern, M.; Vallejo, A.; McNally, M.A.; Moriarty, T.F.; Ferguson, J.Y.; Nijs, S.; Metsemakers, W.J. The Effect of Local Antibiotic Prophylaxis When Treating Open Limb Fractures: A Systematic Review and Meta-Analysis. *Bone Jt. Res.* **2018**, *7*, 447–456. [[CrossRef](#)]
16. Craig, J.; Fuchs, T.; Jenks, M.; Fleetwood, K.; Franz, D.; Iff, J.; Raschke, M. Systematic Review and Meta-Analysis of the Additional Benefit of Local Prophylactic Antibiotic Therapy for Infection Rates in Open Tibia Fractures Treated with Intramedullary Nailing. *Int. Orthop.* **2014**, *38*, 1025–1030. [[CrossRef](#)]
17. Kim, E.K.; Donnelley, C.A.; Ttee, M.; Roberts, H.J.; Von Kaeppler, E.; Shearer, D.; Morshed, S. Prophylactic Topical Antibiotics in Fracture Repair and Spinal Fusion. *Adv. Orthop.* **2021**, *2021*, 1949877. [[CrossRef](#)]
18. Hospenhal, D.R.; Murray, C.K.; Andersen, R.C.; Blice, J.P.; Calhoun, J.H.; Cancio, L.C.; Chung, K.K.; Conger, N.G.; Crouch, H.K.; D'Avignon, L.C.; et al. Guidelines for the Prevention of Infection after Combat-Related Injuries. *J. Trauma* **2008**, *64*, S211–S220. [[CrossRef](#)]
19. Buttaro, M.A.; Gimenez, M.I.; Greco, G.; Barcan, L.; Piccaluga, F. High Active Local Levels of Vancomycin without Nephrotoxicity Released from Impacted Bone Allografts in 20 Revision Hip Arthroplasties. *Acta Orthop.* **2005**, *76*, 336–340. [[CrossRef](#)]
20. Huddleston, P. Intrawound Application of Vancomycin for Prophylaxis in Instrumented Thoracolumbar Fusions: Efficacy, Drug Levels, and Patient Outcomes. *Yearb. Orthop.* **2012**, *2012*, 326–328. [[CrossRef](#)]
21. Armaghani, S.J.; Menge, T.J.; Lovejoy, S.A.; Mencio, G.A.; Martus, J.E. Safety of Topical Vancomycin for Pediatric Spinal Deformity: Nontoxic Serum Levels with Supratherapeutic Drain Levels. *Spine* **2014**, *39*, 1683–1687. [[CrossRef](#)]
22. Metsemakers, W.-J.; Fragomen, A.T.; Moriarty, T.F.; Morgenstern, M.; Egol, K.A.; Zalavras, C.; Obrebsky, W.T.; Raschke, M.; McNally, M.A. Evidence-Based Recommendations for Local Antimicrobial Strategies and Dead Space Management in Fracture-Related Infection. *J. Orthop. Trauma* **2020**, *34*, 18–29. [[CrossRef](#)] [[PubMed](#)]
23. van Vugt, T.A.G.; Arts, J.J.; Geurts, J.A.P. Antibiotic-Loaded Polymethylmethacrylate Beads and Spacers in Treatment of Orthopedic Infections and the Role of Biofilm Formation. *Front. Microbiol.* **2019**, *10*, 1626. [[CrossRef](#)] [[PubMed](#)]
24. Neut, D.; van de Belt, H.; Stokroos, I.; van Horn, J.R.; van der Mei, H.C.; Busscher, H.J. Biomaterial-Associated Infection of Gentamicin-Loaded PMMA Beads in Orthopaedic Revision Surgery. *J. Antimicrob. Chemother.* **2001**, *47*, 885–891. [[CrossRef](#)] [[PubMed](#)]
25. Buchholz, H.; Gartmann, H. Infection Prevention and Surgical Management of Deep Insidious Infection in Total Endoprosthesis. *Chir. Z. Alle Geb. Oper. Medizin* **1972**, *43*, 446–453.
26. Walenkamp, G.H.; Vree, T.B.; van Rens, T.J. Gentamicin-PMMA Beads. Pharmacokinetic and Nephrotoxicological Study. *Clin. Orthop.* **1986**, *205*, 171–183. [[CrossRef](#)]
27. Barger, J.; Fragomen, A.T.; Rozbruch, S.R. Antibiotic-Coated Interlocking Intramedullary Nail for the Treatment of Long-Bone Osteomyelitis. *JBJS Rev.* **2017**, *5*, e5. [[CrossRef](#)]
28. Yu, X.; Wu, H.; Li, J.; Xie, Z. Antibiotic Cement-Coated Locking Plate as a Temporary Internal Fixator for Femoral Osteomyelitis Defects. *Int. Orthop.* **2017**, *41*, 1851–1857. [[CrossRef](#)]
29. Samara, E.; Moriarty, T.; Decosterd, L.; Richards, R.; Gautier, E.; Wahl, P. Antibiotic Stability over Six Weeks in Aqueous Solution at Body Temperature with and without Heat Treatment That Mimics the Curing of Bone Cement. *Bone Jt. Res.* **2017**, *6*, 296–306. [[CrossRef](#)]
30. Greene, N.; Holtom, P.; Warren, C.; Ressler, R.; Shepherd, L.; McPherson, E.; Patzakis, M. In Vitro Elution of Tobramycin and Vancomycin Polymethylmethacrylate Beads and Spacers from Simplex and Palacos. *Am. J. Orthop.* **1998**, *27*, 201–205.
31. van de Belt, H.; Neut, D.; Uges, D.R.; Schenk, W.; van Horn, J.R.; van der Mei, H.C.; Busscher, H.J. Surface Roughness, Porosity and Wettability of Gentamicin-Loaded Bone Cements and Their Antibiotic Release. *Biomaterials* **2000**, *21*, 1981–1987. [[CrossRef](#)]
32. Parvizi, J.; Saleh, K.J.; Ragland, P.S.; Pour, A.E.; Mont, M.A. Efficacy of Antibiotic-Impregnated Cement in Total Hip Replacement. *Acta Orthop.* **2008**, *79*, 335–341. [[CrossRef](#)] [[PubMed](#)]
33. McKee, M.D.; Li-Bland, E.A.; Wild, L.M.; Schemitsch, E.H. A Prospective, Randomized Clinical Trial Comparing an Antibiotic-Impregnated Bioabsorbable Bone Substitute with Standard Antibiotic-Impregnated Cement Beads in the Treatment of Chronic Osteomyelitis and Infected Nonunion. *J. Orthop. Trauma* **2010**, *24*, 483–490. [[CrossRef](#)] [[PubMed](#)]
34. McConoughey, S.J.; Howlin, R.P.; Wiseman, J.; Stoodley, P.; Calhoun, J.H. Comparing PMMA and Calcium Sulfate as Carriers for the Local Delivery of Antibiotics to Infected Surgical Sites. *J. Biomed. Mater. Res. B Appl. Biomater.* **2015**, *103*, 870–877. [[CrossRef](#)] [[PubMed](#)]

35. Ferguson, J.; Diefenbeck, M.; McNally, M. Ceramic Biocomposites as Biodegradable Antibiotic Carriers in the Treatment of Bone Infections. *J. Bone Jt. Infect.* **2017**, *2*, 38–51. [[CrossRef](#)]
36. Howlin, R.P.; Brayford, M.J.; Webb, J.S.; Cooper, J.J.; Aiken, S.S.; Stoodley, P. Antibiotic-Loaded Synthetic Calcium Sulfate Beads for Prevention of Bacterial Colonization and Biofilm Formation in Periprosthetic Infections. *Antimicrob. Agents Chemother.* **2015**, *59*, 111–120. [[CrossRef](#)]
37. McNally, M.A.; Ferguson, J.Y.; Lau, A.C.K.; Diefenbeck, M.; Scarborough, M.; Ramsden, A.J.; Atkins, B.L. Single-Stage Treatment of Chronic Osteomyelitis with a New Absorbable, Gentamicin-Loaded, Calcium Sulphate/Hydroxyapatite Biocomposite: A Prospective Series of 100 Cases. *Bone Jt. J.* **2016**, *98-B*, 1289–1296. [[CrossRef](#)]
38. Ferguson, J.Y.; Dudareva, M.; Riley, N.D.; Stubbs, D.; Atkins, B.L.; McNally, M.A. The Use of a Biodegradable Antibiotic-Loaded Calcium Sulphate Carrier Containing Tobramycin for the Treatment of Chronic Osteomyelitis: A Series of 195 Cases. *Bone Jt. J.* **2014**, *96-B*, 829–836. [[CrossRef](#)]
39. Boot, W.; Gawlitta, D.; Nikkels, P.G.J.; Pouran, B.; van Rijen, M.H.P.; Dhert, W.J.A.; Vogely, H.C. Hyaluronic Acid-Based Hydrogel Coating Does Not Affect Bone Apposition at the Implant Surface in a Rabbit Model. *Clin. Orthop. Relat. Res.* **2017**, *475*, 1911–1919. [[CrossRef](#)]
40. Yang, K.; Han, Q.; Chen, B.; Zheng, Y.; Zhang, K.; Li, Q.; Wang, J. Antimicrobial Hydrogels: Promising Materials for Medical Application. *Int. J. Nanomed.* **2018**, *13*, 2217–2263. [[CrossRef](#)]
41. Hatch, M.D.; Daniels, S.D.; Glerum, K.M.; Higgins, L.D. The Cost Effectiveness of Vancomycin for Preventing Infections after Shoulder Arthroplasty: A Break-Even Analysis. *J. Shoulder Elb. Surg.* **2017**, *26*, 472–477. [[CrossRef](#)]
42. British Pound to US Dollar Spot Exchange Rates for 2017. Available online: <https://www.exchangerates.org.uk/GBP-USD-spot-exchange-rates-history-2017.html> (accessed on 5 January 2022).
43. Geurts, J.; van Vugt, T.; Thijssen, E.; Arts, J.J. Cost-Effectiveness Study of One-Stage Treatment of Chronic Osteomyelitis with Bioactive Glass S53P4. *Materials* **2019**, *12*, 3209. [[CrossRef](#)] [[PubMed](#)]
44. Trentinaglia, M.T.; Van Der Straeten, C.; Morelli, I.; Logoluso, N.; Drago, L.; Romanò, C.L. Economic Evaluation of Antibacterial Coatings on Healthcare Costs in First Year Following Total Joint Arthroplasty. *J. Arthroplast.* **2018**, *33*, 1656–1662. [[CrossRef](#)] [[PubMed](#)]
45. Chiang, H.-Y.; Herwaldt, L.A.; Blevins, A.E.; Cho, E.; Schweizer, M.L. Effectiveness of Local Vancomycin Powder to Decrease Surgical Site Infections: A Meta-Analysis. *Spine J.* **2014**, *14*, 397–407. [[CrossRef](#)]
46. Major Extremity Trauma Research Consortium. Topical Antibiotic Therapy to Reduce Infection after Operative Treatment of Fractures at High Risk of Infection: TOBRA—A Multicenter Randomized Controlled Trial. Available online: <https://clinicaltrials.gov/ct2/show/NCT04597008> (accessed on 8 December 2022).
47. von Kaepler, E.P.; Donnelley, C.; Ali, S.H.; Roberts, H.J.; Ibrahim, J.M.; Wu, H.-H.; Eliezer, E.N.; Porco, T.C.; Haonga, B.T.; Morshed, S.; et al. A Study Protocol for a Pilot Masked, Randomized Controlled Trial Evaluating Locally-Applied Gentamicin versus Saline in Open Tibia Fractures (PGO-Tibia) in Dar Es Salaam, Tanzania. *Pilot Feasibility Stud.* **2021**, *7*, 47. [[CrossRef](#)] [[PubMed](#)]
48. Dodson, V.; Majmundar, N.; Swantic, V.; Assina, R. The Effect of Prophylactic Vancomycin Powder on Infections Following Spinal Surgeries: A Systematic Review. *Neurosurg. Focus* **2019**, *46*, E11. [[CrossRef](#)] [[PubMed](#)]
49. Lemans, J.V.C.; Wijdicks, S.P.J.; Boot, W.; Govaert, G.A.M.; Houwert, R.M.; Öner, F.C.; Kruijt, M.C. Intrawound Treatment for Prevention of Surgical Site Infections in Instrumented Spinal Surgery: A Systematic Comparative Effectiveness Review and Meta-Analysis. *Glob. Spine J.* **2019**, *9*, 219–230. [[CrossRef](#)]
50. Tubaki, V.R.; Rajasekaran, S.; Shetty, A.P. Effects of Using Intravenous Antibiotic Only versus Local Intrawound Vancomycin Antibiotic Powder Application in Addition to Intravenous Antibiotics on Postoperative Infection in Spine Surgery in 907 Patients. *Spine* **2013**, *38*, 2149–2155. [[CrossRef](#)]
51. Martin, J.R.; Adogwa, O.; Brown, C.R.; Bagley, C.A.; Richardson, W.J.; Lad, S.P.; Kuchibhatla, M.; Gottfried, O.N. Experience with Intrawound Vancomycin Powder for Spinal Deformity Surgery. *Spine* **2014**, *39*, 177–184. [[CrossRef](#)]
52. Rathbone, C.R.; Cross, J.D.; Brown, K.V.; Murray, C.K.; Wenke, J.C. Effect of Various Concentrations of Antibiotics on Osteogenic Cell Viability and Activity. *J. Orthop. Res.* **2011**, *29*, 1070–1074. [[CrossRef](#)]
53. Lindsey, R.W.; Probe, R.; Miclau, T.; Alexander, J.W.; Perren, S.M. The Effects of Antibiotic-Impregnated Autogeneic Cancellous Bone Graft on Bone Healing. *Clin. Orthop. Relat. Res.* **1993**, *291*, 303–312. [[CrossRef](#)]
54. Dovas, S.; Liakopoulos, V.; Papatheodorou, L.; Chronopoulou, I.; Papavasiliou, V.; Atmatzidis, E.; Giannopoulou, M.; Eleftheriadis, T.; Simopoulou, T.; Karachalios, T. Acute Renal Failure after Antibiotic-Impregnated Bone Cement Treatment of an Infected Total Knee Arthroplasty. *Clin. Nephrol.* **2008**, *69*, 207–212. [[CrossRef](#)] [[PubMed](#)]
55. Edelstein, A.I.; Okroj, K.T.; Rogers, T.; Della Valle, C.J.; Sporer, S.M. Nephrotoxicity after the Treatment of Periprosthetic Joint Infection with Antibiotic-Loaded Cement Spacers. *J. Arthroplast.* **2018**, *33*, 2225–2229. [[CrossRef](#)] [[PubMed](#)]
56. O’Toole, R.V.; Degani, Y.; Carlini, A.R.; Castillo, R.C.; O’Hara, N.N.; Joshi, M.; Metrc, A. Systemic Absorption and Nephrotoxicity Associated with Topical Vancomycin Powder for Fracture Surgery. *J. Orthop. Trauma* **2021**, *35*, 29–34. [[CrossRef](#)]
57. Henry, S.L.; Seligson, D.; Mangino, P.; Popham, G.J. Antibiotic-Impregnated Beads. Part I: Bead Implantation versus Systemic Therapy. *Orthop. Rev.* **1991**, *20*, 242–247. [[PubMed](#)]