UC Irvine

Journal of Education and Teaching in Emergency Medicine

Title

Acute Ischemic Stroke

Permalink

<https://escholarship.org/uc/item/2j73c2ck>

Journal

Journal of Education and Teaching in Emergency Medicine, 4(2)

Authors

Sangal, Rohit
Siegel, Gregory
Conlon, Lauren
et al.

Publication Date

2019

DOI

10.5070/M542043474

Copyright Information

Copyright 2019 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

SIMULATION

Acute Ischemic Stroke

Rohit Sangal, MD*, Gregory Siegel, MD*, Lauren Conlon, MD* and Kevin Scott, MD*

*Hospital of the University of Pennsylvania, Department of Emergency Medicine, Philadelphia, PA

Correspondence should be addressed to Rohit Sangal, MD at rohit.sangal@uphs.upenn.edu

Submitted: October 6, 2018; Accepted: January 9, 2019; Electronically Published: April 15, 2019; <https://doi.org/10.21980/J8R04X>

Copyright: © 2019 Sangal, et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

ABSTRACT:

Audience: Emergency medicine residents. This simulation was designed for emergency medicine interns to teach the basics of stroke evaluation and care.

Introduction: A cerebrovascular accident (CVA) is an emergent condition that requires rapid diagnosis and treatment. Each year approximately 795,000 people suffer a CVA, leading to over 600,000 emergency department visits for acute ischemic stroke.^{1,2} With such a high incidence of CVAs, there is an increasing demand on emergency medicine (EM) physicians to meet strict metrics for CVA recognition and care. Data have shown that quick and organized stroke care consistent with national guidelines leads to decreased door-to-needle time, lower in-hospital mortality, lower rates of intracranial hemorrhage, and increased percentage of patients discharged home.¹

Emergency medicine interns come from different backgrounds and, although they possess the appropriate medical knowledge, they often lack insight into the hospital protocols and national guidelines that allow them to care for these patients efficiently and effectively. It is important to introduce CVA management early in the education process for early learners to have confidence in managing this disease process. We aim to use simulation to standardize the care and knowledge of CVAs at our institution.

Objectives: By the end of this simulation session, learners will be able to: 1) recognize a CVA using the National Institutes of Health Stroke Scale (NIHSS), 2) understand and properly utilize the NIHSS, 3) list appropriate imaging and laboratory orders for a CVA work-up, 4) determine appropriate subspecialty consultation, 5) discuss common stroke syndromes and associated cerebral locations, 6) review indications and contraindications for tissue plasminogen activator (tPA), 7) review hospital specific stroke protocol.

Method: We recommend use of a standardized patient capable of replicating the details of the neurological exam described. An oral boards format or high-fidelity simulator can also be used if a standardized patient is not available. Debriefing was left to the facilitator who was advised to give open ended questions to gauge reactions and assess understanding.

Topics: Medical simulation, acute cerebrovascular accident, tissue plasminogen activator, neurology, emergency medicine.

USER GUIDE

List of Resources:

Abstract	1
User Guide	2
Instructor Materials	4
Operator Materials	13
Debriefing and Evaluation Pearls	15
Simulation Assessment	21

Learner Audience:

Medical Students, Interns, Junior Residents, Senior Residents

Time Required for Implementation:

Instructor Preparation: 20-30 minutes

Time for case: 15-20 minutes

Time for debriefing: 10-30 minutes

Recommended Number of Learners per Instructor:

3

Topics:

Medical simulation, acute cerebrovascular accident, tissue plasminogen activator, neurology, emergency medicine.

Objectives:

By the end of this simulation session, the learner will be able to:

1. Recognize a CVA using the National Institutes of Health Stroke Scale (NIHSS).
2. Understand and properly utilize the NIHSS.
3. List appropriate imaging and laboratory orders for a CVA work-up.
4. Determine appropriate subspecialty consultation.
5. Discuss common stroke syndromes and associated cerebral locations.
6. Review indications and contraindications for tPA.
7. Review hospital specific stroke protocol.

Linked objectives and methods:

EM interns at University of Pennsylvania participate in a two-week intern orientation “boot camp” to aide in their transition to the role of a physician caring for patients in a new environment. Orientation courses such as these have been shown to improve clinical confidence and skills and to smooth transitions between medical student and doctor. Simulation is widely used within graduate medical education programs and some data in the surgical literature suggests that simulation can help interns achieve skills comparable to that of second or even third year residents.^{6,7} Furthermore, emerging research has shown that simulation training centered on acute stroke

management leads to improved metrics and increased learner confidence.^{3,8}

The case is centered on a new patient, just brought in by medics, displaying symptoms concerning for an acute stroke. The learner is expected to quickly recognize and assess this pathology using the NIHSS (objectives 1 and 2). From there appropriate labs and imaging should be immediately ordered to further work-up the suspected pathology (objective 3). Simultaneously, the learner is expected to acquire subspecialty consultation by means of a “stroke alert,” “code stroke,” or equivalent (objective 4); however, little assistance is provided by this consultant so as to allow the learner to maintain control over the simulation. Once a computed tomography (CT) of the head is obtained and interpreted, the learner should state that tPA is indicated and should order it only after ruling out contraindications (objective 6). The simulation ends when the learner admits the patient to an intensive care unit (ICU). The debrief will consist of a thorough review of the case, pertinent imaging and laboratory findings, use of the NIHSS, localizing symptoms to area of the brain, indications and contraindications for tPA, and relevant institutional protocols (Objectives 1-8).

Recommended pre-reading for instructor:

- The instructor should read any materials on acute cerebrovascular accidents with a focus on acute ischemic strokes and review the National Institutes of Health Stroke Scale as well as contraindications for tPA.
- Additionally, instructors should review their own hospital stroke protocol and provide a worksheet or slide deck reviewing these protocols.

Results and tips for successful implementation:

While this case was piloted with high-fidelity simulation, we recommend use of a standardized patient to allow for more realistic demonstration of neurologic deficits. All standard items were available including code medications, defibrillator, intravenous (IV) line supplies and IV fluids. Learners were able to access laboratory tests and imaging as requested. No specific moulage is required. Supply lists are provided as well for set up.

This simulation was taught to twelve EM interns during their orientation week prior to starting their rotations in the hospital. The simulation was run four times and learners were allowed to divide themselves into groups of three. We recommend the operator of the simulator have basic knowledge of the mannequin, as needed, including changing vital signs and communication through the mannequin. This role can be done by anyone with a script to the case. Additionally, we oriented learners to the simulation center and high-fidelity mannequins

USER GUIDE

prior to beginning the simulation. An EM faculty member was assigned to this case to evaluate the team and troubleshoot any unexpected issues. This case can be modified for senior residents by introducing subtle CVA exam findings and relative contraindications for tPA. Given limited roles and decision makers, groups larger than three are not recommended for this case.

The faculty member in the room evaluated each team based on predetermined critical actions. Debriefing was done with an instructor guide that outlined objectives of the case as well as any notes taken by the operator. Learners evaluated the case at the end with an evaluation form. Of the 12 participants, 12 (100%) completed the survey. Surveys were anonymous and the study was approved by the Institutional Research Board. All 12 participants agreed or strongly agreed the simulation was realistic and useful. 11 (92%) agreed or strongly agreed they have confidence activating the stroke protocol at this institution. Selected comments are below: ⁵

Positive

- "I liked that it was realistic and helped solidify prior knowledge and new info about pathways."
- "Basic stroke eval and treatment of the patient was great to get a bit more comfortable."
- "Very applicable for new ED interns."

Of the surveys commenting on areas for improvement, two indicated that "more subtle signs" or "increased difficulty" of the case was desired. With regards to critical actions, the most commonly missed actions were ordering a fingerstick blood glucose or tPA.

References/suggestions for further reading:

1. Benjamin EJ, Virani SS, Callaway CW, et al. Heart disease and stroke statistics-2018 update: a report from the American Heart Association. *Circulation*. 2018;137(12):e67-e492. doi: 10.1161/CIR.0000000000000558.
2. Stuntz M, Busko K, Irshad S, Paige T, Razhkova V, Coan T. Nationwide trends of clinical characteristics and economic burden of emergency department visits due to acute ischemic stroke. *Open Access Emerg Med*. 2017;9:89-96. doi: 10.2147/OAEM.S146654.
3. Capampangan D, Hentz J, Hoerth M. Simulation based training for acute stroke alerts. *Neurology*. 2012;78:P07.228.
4. Cohen ER, Barsuk JH, Moazed F, et al. Making July safer: simulation-based mastery learning during intern boot camp. *Acad Med*. 2013;88(2):233-239. doi: 10.1097/ACM.0b013e31827bfc0a.
5. Sinha SN, Page W. Interns' Day in Surgery: improving intern performance through a simulation-based course for final year medical students. *ANZ J Surg*. 2015;85(1-2):27-32. doi: 10.1111/ans.12665.
6. Chipman JG, Schmitz CC. Using objective structured assessment of technical skills to evaluate a basic skills simulation curriculum for first-year surgical residents. *J Am Coll Surg*. 2009;209(3):364-370.e2. doi: 10.1016/j.jamcollsurg.2009.05.005.
7. Marshall RL, Gorman PJ, Verne D, et al. Practical training for postgraduate year 1 surgery residents. *Am J Surg*. 2000;179(3): 194-196.
8. Mehta T, Strauss S, Beland D, Fortunato G, Staff I, Lee N. Stroke simulation improves acute stroke management: a systems-based practice experience. *J Grad Med Educ*. 2018;10(1): 57-62. doi: 10.4300/JGME-D-17-00167.1.
9. Kwah LK, Diong J. National Institutes of Health Stroke Scale (NIHSS). *J Physiother*. 2014; 60(1):61. doi: 10.1016/j.jphys.2013.12.012.
10. Kothari RU, Pancioli A, Liu T, Brott T, Broderick J. Cincinnati Prehospital Stroke Scale: reproducibility and validity. *Ann Emerg Med*. 1999;33(4): 373-378.
11. Go S, Worman DJ. Stroke Syndromes. In: Tintinalli JE, Stapczynski JS, Ma OJ, Yealy DM, Meckler GD, Cline DM, eds. *Tintinalli's Emergency Medicine: A Comprehensive Study Guide*. 8th ed. New York, NY: McGraw-Hill; 2016:1128-1129.
12. Demaerschalk BM, Kleindorfer DO, Adeoye OM, et al. Scientific rationale for the inclusion and exclusion criteria for intravenous alteplase in acute ischemic stroke: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2016;47(2):581-641. doi: 10.1161/STR.0000000000000086.
13. National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. Tissue plasminogen activator for acute ischemic stroke. *N Engl J Med*. 1995;333(24):1581-1587.
14. Hacke W, Kaste M, Bluhmki E, et al. Thrombolysis with alteplase 3 to 4.5 hours after acute ischemic stroke. *N Engl J Med*. 2008;359(13): 1317-1329. doi: 10.1056/NEJMoa0804656.
15. Yaghi S, Willey JZ, Cucchiara B, et al. Treatment and outcome of hemorrhagic transformation after intravenous alteplase in acute ischemic stroke: a scientific statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2017;48(12):e343-e361. doi: 10.1161/STR.0000000000000152.
16. Lip GY, Nieuwlaar R, Pisters R, Lane DA, Crijns HJ. Refining clinical risk stratification for predicting stroke and thromboembolism in atrial fibrillation using a novel risk factor-based approach: the euro heart survey on atrial fibrillation. *Chest*. 2010;137(2): 263-272. doi: 10.1378/chest.09-1584.

INSTRUCTOR MATERIALS

Case Title: Acute Ischemic Stroke

Case Description & Diagnosis (short synopsis): A 65-year-old male with a history notable for diabetes, hypertension, hyperlipidemia presents via emergency medical services (EMS) for “not acting right.” Paramedics state the patient has been slurring his speech for about 15 minutes prior to their arrival. The learner should obtain a fingerstick blood glucose and activate a stroke alert. The learner should obtain a non-contrast computed tomography (CT) of the head and begin assessing for contraindications for tPA. The blood pressure will need to be acutely lowered and learners should emergently administer tPA.

Equipment or Props Needed:

IV Supplies:

- Angiocatheters
- IV tubing
- Saline

Airway Management:

- Non-rebreather mask
- Nasal cannula
- Laryngoscope
- Bag-valve mask
- Endotracheal tube and Stylet
- Orogastric/nasogastric tube

Medications:

- Etomidate
- Succinylcholine
- Epinephrine
- Atropine
- Labetelol
- Nicardipine
- Tissue plasminogen activator

Sim Exam:

- Plain clothed mannequin (presenting via EMS)

Miscellaneous:

- Gloves
- Alcohol/chloraprep
- Cardiac Monitor

INSTRUCTOR MATERIALS

Confederates needed:

The simulation operator can be utilized as a pharmacist or other relevant consultant via phone. With multidisciplinary learner teams, a nurse can be used to help with IV-line placement and tPA monitoring.

Stimulus Inventory:

- #1 Non-contrast head computed tomography
- #2 Electrocardiogram (ECG)
- #3 Complete blood count (CBC)
- #4 Basic metabolic panel (BMP)
- #5 Fingerstick blood glucose
- #6 Urine drug screen
- #7 Coagulation panel

Background and brief information: The 65-year-old male patient is brought via EMS to a tertiary medical center with reports of not acting like himself.

Initial presentation: The patient is brought in via EMS with no family present. He appears as an appropriately dressed middle aged male in moderate distress.

How the scenario unfolds: On arrival to the emergency room, learners should begin by assessing the patient's airway, breathing and circulation while performing a brief history leading to a focused physical exam. They should recognize the patient is exhibiting symptoms of a CVA and immediately obtain a fingerstick blood glucose which will be normal. They should immediately activate a stroke alert. If learners do not obtain a fingerstick blood glucose, pharmacy will ultimately refuse to dispense tPA until all contraindications are ruled out. They should obtain a non-contrast head CT and laboratory tests and work through indications and contraindications for tPA. Neurology will respond to the stroke code but will tell the learners they should review the indications and contraindications with the patient and order it if indicated. Learners should recognize the patient's significant hypertension and administer medications to make patient eligible for tPA. If blood pressure control is not done promptly, pharmacy will refuse to dispense tPA until it is corrected.

INSTRUCTOR MATERIALS

Critical actions:

1. Obtains a basic history of symptoms.
2. Obtains past medical history.
3. Perform NIHSS and conduct a thorough physical exam.
4. Obtain fingerstick blood glucose prior to stroke activation.
5. Activate hospital-specific stroke protocol (this should vary based on your hospital protocol so that learners can understand the system they work in).
6. Obtains and correctly interprets head CT (does not need to identify the dense middle cerebral artery (MCA) sign, but does need to state no bleeding).
7. Order appropriate labs (CBC, CMP, coagulation studies, Type and screen).
8. Correctly addresses blood pressure (before pharmacy prompting) prior to tPA.
9. Order tPA (thrombectomy also acceptable) and recognize reason for indication (ie <4.5 hours, not on anticoagulation, no contraindications).
10. Admits patient to the ICU for close monitoring.

INSTRUCTOR MATERIALS

Case title: Acute Ischemic Stroke

Chief Complaint: “Not acting right”

Vitals: Heart Rate (HR) 112 Blood Pressure (BP) 210/125 Respiratory Rate (RR) 18
Temperature (T) 98.0°F Oxygen Saturation (O₂Sat) 98% on room air

General Appearance: Elderly male, appropriately dressed.

Primary Survey:

- **Airway:** Patent.
- **Breathing:** Clear to auscultations bilaterally.
- **Circulation:** 2+ pulses throughout.

History:

- **History of present illness:**
 - **Paramedic history**

The paramedics were called by patient’s wife for patient “not acting right.” Approximately 15 minutes prior to arrival the wife called after the patient was found to be slurring his speech. As a note, the patient had just taken his afternoon meds, which include oxycodone, and he thinks it is related to taking a few extras like he normally does, but his wife wanted to make sure he is alright.
 - **Patient history**

“I’m fine, I think it’s my pain meds, I don’t think it’s anything (slurred speech).” If asked, he took an extra one oxycodone approximately one hour ago and slurred speech began approximately 30 minutes ago. Patient states he had a similar episode about one week ago that resolved after 5-10 minutes, but at that time he had not recently taken any pain medications. He did not tell his wife because he did not want her to worry. If prompted, he states his right arm is feeling heavy, and he had trouble walking in the house because his right leg feels like “it had rocks in it.” Patient is right-hand dominant.
 - If asked: No history of intracranial bleeding, no prior history of stroke. No history of gastrointestinal bleeding or recent major surgery. No oral anticoagulants.

INSTRUCTOR MATERIALS

- **Past medical history:** Type 2 diabetes, hypertension, hyperlipidemia, tobacco abuse, and chronic back pain.
- **Past surgical history:** Lumbar L3-L5 laminectomy 10 years ago.
- **Patient's medications:** Metformin, metoprolol, atorvastatin, baby aspirin, oxycodone for chronic back pain.
- **Allergies:** No known drug allergies.
- **Social history:** 1 pack per day smoker.
- **Family history:** Diabetes mellitus, hypertension, hyperlipidemia, coronary artery disease.

Secondary Survey/Physical Examination:

- **General appearance:** Patient is an elderly appearing male. Patient is awake, alert, and oriented. Comfortable without distress.
- **HEENT:**
 - **Head:** Normocephalic, atraumatic.
 - **Eyes:** Pupils equally round and reactive. Extraocular movements intact.
 - **Ears:** Tympanic membrane intact.
 - **Throat:** Oropharynx clear.
- **Neck:** Within normal limits.
- **Heart:** Irregularly irregular. No murmurs/rubs/gallops.
- **Lungs:** Clear to auscultation bilaterally. No wheezes/rhonchi/rales.
- **Abdominal/GI:** Soft non-tender. Normal bowel sounds. No rebound. No organomegaly.
- **Genitourinary:** Within normal limits.
- **Rectal:** Within normal limits.
- **Extremities:** No contractures, non-tender. No swelling or edema.
- **Back:** Within normal limits.
- **Neuro:** Alert and oriented to person, place, self and situation. Normal mental status. Slurred speech. Right sided flattening of nasolabial fold. Unable to assess dysmetria due to weakness. Right sided pronator drift. 3/5 right upper extremity, 4/5 right lower extremity strength. 5/5 strength throughout on left side.
 - **NIHSS**
 - **Level of Consciousness (LOC):** Alert (0)
 - **LOC Questions:** Able to state month and age (0)
 - **LOC Commands:** Follows commands (0)
 - **Best Gaze:** Normal (0)

INSTRUCTOR MATERIALS

- **Visual:** No vision loss (0)
- **Facial Palsy:** Minor paralysis - right (1)
- **Motor Arm:** Some effort against gravity – right (2), Normal – left (0)
- **Motor Leg:** Drift – right (1), Normal -left (0)
- **Limb Ataxia:** Absent/Unable to complete due to weakness (0)
- **Sensory:** Mild-to-moderate sensory loss (1)
- **Best Language:** No aphasia (0) (able to repeat phrase, name object, and identify what is happening in the picture (if presented))
- **Dysarthria:** Mild-to-moderate dysarthria (1)
- **Extinction and Inattention:** No abnormality (0)
- **Total:** 6
- **Skin:** Normal color and turgor. No diaphoresis. No skin rash.
- **Lymph:** Within normal limits.
- **Psych:** Within normal limits.

INSTRUCTOR MATERIALS

Results:

Non-contrast head CT

Image Source: Dixon A. Hyperdense MCA sign. In: Radiopaedia. rID:

<https://radiopaedia.org/blog/hyperdense-mca-sign>. CC BY-NC-SA 3.0.

INSTRUCTOR MATERIALS

Electrocardiogram

CardioNetworks. Electrocardiogram (ECG) 12 Lead ECG. In: Wikimedia Commons.

[https://commons.wikimedia.org/wiki/File:Afib_ecg_\(CardioNetworks_ECGpedia\).jpg](https://commons.wikimedia.org/wiki/File:Afib_ecg_(CardioNetworks_ECGpedia).jpg).

Published October 7, 2007. CC BY-SA 3.0.

INSTRUCTOR MATERIALS

Complete blood count (CBC)

White blood count (WBC)	11.8 x1000/mm ³ (H)
Hemoglobin (Hgb)	11.8 g/dL
Hematocrit (HCT)	35%
Platelet (Plt)	225 x1000/mm ³

Basic metabolic panel (BMP)

Sodium	133 mEq/L
Chloride	114 mEq/L
Potassium	4.2 mEq/L
Bicarbonate (HCO ₃)	23 mEq/L
Blood Urea Nitrogen (BUN)	16 mg/dL
Creatine (Cr)	1.2 mg/dL
Glucose	49 mg/dL

Fingerstick blood glucose 45 mg/dL

Urine drug screen Pending

Coagulation panel

Prothombin Time (PT)	11.3 seconds
Partial Thromboplastin Time (PTT)	32.6 seconds
International Normalized Ratio (INR)	1.2

OPERATOR MATERIALS

SIMULATION EVENTS TABLE:

Minute (state)	Participant action/ trigger	Patient status (simulator response) & operator prompts	Monitor display (vital signs)
0:00 (Baseline)	Paramedic report. Learner should obtain focused history of present illness, past medical history and brief physical exam.	Patient can answer questions as needed (through operator if running as simulation).	T 98.0°F HR 112 BP 210/125 RR 18 O2 98%
2:00-4:00	Obtain fingerstick blood glucose.	If glucose not obtained, pharmacy will refuse to dispense tPA when asked for.	T 98.0°F HR 98 BP 210/125 RR 18 O2 98%
4:00-6:00	Recognize likely diagnosis of CVA and obtain stat hematocrit. Activate hospital stroke protocol.	If stroke team is not paged or learner does not recognize CVA, the patient will develop more profound weakness and become aphasic. If learners still do not recognize stroke with worsening symptoms then nursing can prompt “do you want to activate a stroke alert? Do you want to get a head CT?”	T 98.0°F HR 98 BP 210/125 RR 18 O2 98%
4:00-6:00 (Cont.)	Obtain and interpret head CT. Obtain appropriate labs (CBC, CMP, coagulation studies, Type and screen) and ECG.	If head CT not obtained, nurse will ask if a head CT is needed, or pharmacy will refuse to dispense tPA. If labs not obtained, pharmacy will refuse to dispense tPA. If ECG not obtained, the nurse will ask what the patient’s rhythm is.	T 98.0°F HR 98 BP 210/125 RR 18 O2 98%
6:00-8:00	Perform NIHSS.	If it is not performed, nursing can ask what the stroke score is. If learners ask for a neurology consult or for help identifying type/location of stroke, they will be told they are unavailable.	T 98.0°F HR 98 BP 210/125 RR 18 O2 98%

OPERATOR MATERIALS

Minute (state)	Participant action/ trigger	Patient status (simulator response) & operator prompts	Monitor display (vital signs)
		If learners have not interpreted the head CT, the nurse should ask what they saw on the head CT.	
6:00-8:00	Recognize patient's elevated blood pressure and need for control (both for potential tPA or management of hemorrhagic CVA).	If elevated blood pressure is recognized at this point or earlier, a single dose of any IV antihypertensive agent will control the blood pressure.	T 98.0°F HR 85 BP 180/100 RR 18 O2 98%
8:00-10:00	Decide to give tPA (thrombectomy also acceptable). Must give some reasoning (ie. <4.5 hrs, not on anticoagulation, no contraindications)	<p>If blood pressure is not controlled and labs are not sent, pharmacy will prompt learners to control BP with IV anti-hypertensives and order INR.</p> <p>If learners are addressing blood pressure for the first time at this step, it will be refractory to a single agent and require an infusion of IV antihypertensive medications.</p> <p>If learners have not stated why they are giving tPA, pharmacy will ask what is the indication for it and ask if contraindications have been reviewed until the learner does so.</p>	<p>With BP control: T 98.0°F HR 85 BP 180/100 RR 18 O2 98%</p> <p>Without BP control: T 98.0°F HR 99 BP 220/130 RR 18 O2 98%</p>
Case Completion	Disposition to the ICU.	Stroke team consultant can prompt learners if no disposition chosen.	T 98.0°F HR 85 BP 175/80 RR 18 O2 98%

Diagnosis:

Acute ischemic stroke

Disposition:

ICU

DEBRIEFING AND EVALUATION PEARLS

Acute Ischemic Stroke

Cerebrovascular Accident (CVA): Sudden neurologic dysfunction due to a vascular cause.

National Institutes of Health Stroke Scale (NIHSS): Developed in 1989, this is a scoring system based on 11 items to assess degree of impairment for patients demonstrating clinical concern for CVA⁹. Each item is scored from zero to four where zero is normal and higher numbers represent more impairment. We recommend instructors print out their institutional NIHSS scoring sheet so that learners can be familiar with the document they will use regularly.

Cincinnati Stroke Scale: An abbreviated scale based on the NIH stroke scale for rapid recognition of possible CVA. It has a high sensitivity and specificity when performed by a physician.¹⁰

1. Facial droop – Do both sides of the face move equally when the patient smiles?
2. Arm drift – Do both hands stay still when the patient holds them out in front of them with their eyes closed?
3. Speech – Does the patient have slurred speech or use inappropriate words?

There are two types of CVAs

- Ischemic strokes make up 90% and can be subclassified (90%)
 - *Cardioembolic (25%):* Atrial fibrillation and valvular heart disease.
 - *Large vessel disease (25%):* Large vessel occlusions often secondary to high cholesterol.
 - *Small vessel disease (25%):* Small vessel occlusions often secondary to hypertension or diabetes.
 - *Other (25%):* Strokes due to arterial dissection or blood coagulation abnormality.
- Hemorrhagic strokes constitute the remaining 10%
 - The rupture of small penetrating arteries causes bleeding into the brain parenchyma and as the bleeding continues, the surrounding brain tissue is displaced and compressed.
 - Overall the most common cause is related to hypertension and in the elderly the most common cause is cerebral amyloid angiopathy.
 - Other causes to consider include acute vestibular syndrome, aneurysms and coagulopathies (congenital or acquired).

DEBRIEFING AND EVALUATION PEARLS

SPECIFIC ISCHEMIC STROKE SYNDROMES¹¹

1. Anterior cerebral artery occlusions are uncommon but usually present with contralateral lower extremity sensory and motor deficits while sparing the hands and face.
 - a. Left sided lesions present with akinetic mutism and transcortical motor aphasia.
 - b. Right sided lesions present with confusion and motor hemineglect.
2. Middle cerebral artery occlusions are most common and present with contralateral hemiparesis including the face and sensory loss.
 - a. Left hemisphere is dominant in most patients and if involved will present with receptive or expressive aphasia.
 - b. Patients with nondominant hemisphere lesion have inattention and neglect.
3. Posterior Cerebral Artery (Including Basilar Artery) occlusions can be challenging to identify as signs are subtle but commonly include ataxia and vertigo.
 - a. Patients have crossed symptoms where the ipsilateral cranial nerve and contralateral extremity motor weakness is present.
 - b. Basilar artery occlusion can rarely cause locked-in syndrome, where there is complete paralysis except with blinking and upward gaze.
4. Cerebellar Artery occlusions have nonspecific symptoms including vertigo, ataxia, and headache. CT scan only is not sensitive enough because the posterior fossa bone produces artifact so MRI must be obtained for definitive diagnosis.
5. Lacunar infarcts involve small penetrating arteries and present with pure motor or sensory deficits. Small Penetrating Arteries (Lacunar infarction). The elderly and those with chronic uncontrolled hypertension are at highest risk.

DEBRIEFING AND EVALUATION PEARLS

INDICATIONS AND CONTRAINDICATIONS FOR ADMINISTRATION OF IV tPA¹²

These represent criteria from the American Heart Association (AHA) and American Stroke Association (ASA).¹² Please refer to your institution-specific protocol for any additional criteria.

1. Indications

- (a) Age ≥ 18 .
- (b) Time of symptom onset < 4.5 hours.
- (c) Diagnosis of ischemic stroke causing measurable neurological deficit.
- (d) The neurological signs should not be clearing spontaneously.
- (e) The neurological signs should not be minor and isolated.
- (f) Caution should be exercised in treating a patient with major deficits.

2. Absolute Contraindications

- (a) History or evidence of intracranial hemorrhage.
- (b) Clinical presentation suggestive of subarachnoid hemorrhage.
- (c) Known intracranial arteriovenous malformation, neoplasm or aneurysm.
- (d) Systolic blood pressure exceeding 185 mmHg or diastolic blood pressure exceeding 110 mmHg.
- (e) Platelet count below 100,000/ μ L.
- (f) Prothrombin time (PT) above 15 or INR above 1.7 or current use of direct oral anticoagulant (DOAC).
- (g) Active internal bleeding or acute trauma (fracture).
- (h) Neurosurgery, head trauma or stroke in the previous 3 months.
- (i) Abnormal blood glucose (<50 or >400).

3. Relative Contraindications

- (a) Minor or rapidly improving symptoms.
- (b) Major surgery or non-serious head trauma in past 14 days.
- (c) History of GI bleed or urinary bleed in past 21 days.
- (d) Seizure with postictal residual neurologic impairment.
- (e) Recent arterial puncture at noncompressible site.
- (f) Recent lumbar puncture.
- (g) Suspected acute pericarditis.
- (h) Myocardial infarction (MI) in the previous 3 months.
- (i) Pregnancy.

DEBRIEFING AND EVALUATION PEARLS

MAJOR RESEARCH STUDIES SUPPORTING tPA ADMINISTRATION

NINDS II (National Institute of Neurologic Disorders and Stroke) ¹³ was a randomized control trial of 312 patients who were given alteplase within three hours of symptom onset.

Results

- Compared to the placebo group, tPA treated patients were at least 30% more likely to have better than minimal disability at three months on the assessment scales.
- Within 36 hours of stroke, 6.4% of tPA given patients had symptomatic intracerebral hemorrhage compared to 0.6% in placebo group ($P < 0.001$).
- There was no significant difference in mortality between control and treatment groups.

ECASS III (European Cooperative Acute Stroke Study III)¹⁴ was published in 2008 by Hacke and colleagues on the results of a randomized control trial of 821 patients given alteplase up to 4.5 hours after symptom onset.

Results

- Patients receiving tPA had favorable outcome 52.4% vs 45.2% in the placebo group. ($p = 0.04$).
- While any intracranial hemorrhage was higher in the tPA group compared to placebo group, there was no difference in mortality between the two groups.
- This study prompted the AHA/ASA guidelines to extend the time window for tPA administration from 3 to 4.5 hours.

While eligibility criteria were similar, this study helps guide additional exclusion criteria to consider including patients older than 80 years, any use of oral anticoagulants, baseline NIHSS greater than 25, and patients with a history of stroke or diabetes.

Some Considerations Regarding tPA

The incidence of symptomatic intracerebral hemorrhage (sICH) post tPA ranges 2% to 7%.

- In the aforementioned NINDS trial, sICH with tPA versus placebo was 6.4% and 0.6%, respectively ($p < 0.01$). This was similar in the ECASS III trial.
- Selection criteria for tPA is very important because the risk of sICH increases with age, larger strokes, and diabetes.¹⁵

ACEP Clinical Guidelines (2015)

DEBRIEFING AND EVALUATION PEARLS

- *Level B Recommendation (moderate clinical certainty) that IV tPA may be given to select patients with ischemic stroke within 3 hours of onset but to consider the risk of symptomatic intracerebral hemorrhage. This medication should only be administered if safety mechanisms are in place.*
- *Level C Recommendation (Expert Consensus) that shared decision making should occur prior to the decision to use IV tPA*

Atrial Fibrillation: Irregular and often rapid heart rhythm where the atria have disorganized electrical activity and do not contract properly.

There are many causes of atrial fibrillation including heart disease, chronic hypertension, infections, thyroid disease, ingestions, and electrolyte abnormalities. The disorganized activity of the atria predispose patients to ischemic stroke from embolism.¹⁶

CHA₂DS₂-VASc Score is a tool to quantify the risk of stroke in patients with atrial fibrillation. This patient's CHA₂DS₂-VASc score is 5 which confers an annual stroke risk of 6.7%.

		Points	Our Patient
C	Congestive Heart Failure	1	0
H	Hypertension > 140/90 or on treatment	1	1
A	Age>75	2	0
D	Diabetes Mellitus	1	1
S	Prior stroke, Transient Ischemic Attack	2	2
V	Vascular disease	1	0
A	Age 65-74	1	1
Sc	Sex Category (Female)	1	0
Total			5

DEBRIEFING AND EVALUATION PEARLS

Other debriefing points:

Discuss how learners felt communication was among team members, especially if there was confusion regarding the diagnosis.

Why is a STAT Head CT important in the care of these patients?

If EMS was not asked for additional history (as opposed to just chief complaint) ask learners, “Were there any other sources of history you could have solicited?”

To generate discussion on atrial fibrillation, a transition question can be used such as, “What do you think was the cause of this patient’s stroke?”

How can we reduce the risk of stroke in patients with atrial fibrillation?

Instructor should prepare debriefing on their individual hospital’s stroke protocol as well.

Wrap Up:

We recommend learners read more on cerebrovascular accidents in available emergency medicine textbooks and the Emergency Neurologic Life Support guidelines on stroke.

Additionally, the American Heart Association provides pocket cards on this disease process (similar to Advanced Cardiovascular Life Support).

SIMULATION ASSESSMENT

Acute Ischemic Stroke

Learner: _____

Assessment Timeline

This timeline is to help observers assess their learners. It allows observer to make notes on when learners performed various tasks, which can help guide debriefing discussion.

Critical Actions:

1. Obtains a basic history of symptoms.2. Obtains past medical history.3. Performs NIHSS and conducts a thorough physical exam.4. Obtains fingerstick blood glucose prior to stroke activation.5. Activates hospital-specific stroke protocol (this should vary based on your hospital protocol so that learners can understand the system they work in).6. Obtains and correctly interprets head CT (does not need to identify the dense MCA sign, but does need to state no bleeding).7. Orders appropriate labs (CBC, CMP, coagulation studies, Type and screen).8. Correctly addresses blood pressure (before pharmacy prompting) prior to tPA.9. Orders tPA (thrombectomy also acceptable) and recognizes reason for indication (ie <4.5 hours, not on anticoagulation, no contraindications).10. Admits patient to the ICU for close monitoring.	0:00

SIMULATION ASSESSMENT

Acute Ischemic Stroke

Learner: _____

Critical Actions:

- ☐ Obtains a basic history of symptoms.
- ☐ Obtains past medical history.
- ☐ Performs NIHSS and conducts a thorough physical exam.
- ☐ Obtains fingerstick blood glucose prior to stroke activation.
- ☐ Activates hospital-specific stroke protocol (this should vary based on your hospital protocol so that learners can understand the system they work in).
- ☐ Obtains and correctly interprets head CT (does not need to identify the dense MCA sign, but does need to state no bleeding).
- ☐ Orders appropriate labs (CBC, CMP, coagulation studies, Type and screen).
- ☐ Correctly addresses blood pressure (before pharmacy prompting) prior to tPA.
- ☐ Orders tPA (thrombectomy also acceptable) and recognizes reason for indication (ie <4.5 hours, not on anticoagulation, no contraindications).
- ☐ Admits patient to the ICU for close monitoring.

Summative and formative comments:

SIMULATION ASSESSMENT

Acute Ischemic Stroke

Learner: _____

Milestones assessment:

	Milestone	Did not achieve level 1	Level 1	Level 2	Level 3
1	Emergency Stabilization (PC1)	☐ Did not achieve Level 1	☐ Recognizes abnormal vital signs	☐ Recognizes an unstable patient, requiring intervention Performs primary assessment Discerns data to formulate a diagnostic impression/plan	☐ Manages and prioritizes critical actions in a critically ill patient Reassesses after implementing a stabilizing intervention
2	Performance of focused history and physical (PC2)	☐ Did not achieve Level 1	☐ Performs a reliable, comprehensive history and physical exam	☐ Performs and communicates a focused history and physical exam based on chief complaint and urgent issues	☐ Prioritizes essential components of history and physical exam given dynamic circumstances
3	Diagnostic studies (PC3)	☐ Did not achieve Level 1	☐ Determines the necessity of diagnostic studies	☐ Orders appropriate diagnostic studies. Performs appropriate bedside diagnostic studies/procedures	☐ Prioritizes essential testing Interprets results of diagnostic studies Reviews risks, benefits, contraindications, and alternatives to a diagnostic study or procedure
4	Diagnosis (PC4)	☐ Did not achieve Level 1	☐ Considers a list of potential diagnoses	☐ Considers an appropriate list of potential diagnosis May or may not make correct diagnosis	☐ Makes the appropriate diagnosis Considers other potential diagnoses, avoiding premature closure

SIMULATION ASSESSMENT

Acute Ischemic Stroke

Learner: _____

	Milestone	Did not achieve level 1	Level 1	Level 2	Level 3
5	Pharmacotherapy (PC5)	☐ Did not achieve Level 1	☐ Asks patient for drug allergies	☐ Selects an medication for therapeutic intervention, consider potential adverse effects	☐ Selects the most appropriate medication and understands mechanism of action, effect, and potential side effects Considers and recognizes drug-drug interactions
6	Observation and reassessment (PC6)	☐ Did not achieve Level 1	☐ Reevaluates patient at least one time during case	☐ Reevaluates patient after most therapeutic interventions	☐ Consistently evaluates the effectiveness of therapies at appropriate intervals
7	Disposition (PC7)	☐ Did not achieve Level 1	☐ Appropriately selects whether to admit or discharge the patient	☐ Appropriately selects whether to admit or discharge Involves the expertise of some of the appropriate specialists	☐ Educates the patient appropriately about their disposition Assigns patient to an appropriate level of care (ICU/Tele/Floor) Involves expertise of all appropriate specialists
9	General Approach to Procedures (PC9)	☐ Did not achieve Level 1	☐ Identifies pertinent anatomy and physiology for a procedure Uses appropriate Universal Precautions	☐ Obtains informed consent Knows indications, contraindications, anatomic landmarks, equipment, anesthetic and procedural technique, and potential complications for common ED procedures	☐ Determines a back-up strategy if initial attempts are unsuccessful Correctly interprets results of diagnostic procedure

SIMULATION ASSESSMENT

Acute Ischemic Stroke

Learner: _____

	Milestone	Did not achieve level 1	Level 1	Level 2	Level 3
20	Professional Values (PROF1)	☐ Did not achieve Level 1	☐ Demonstrates caring, honest behavior	☐ Exhibits compassion, respect, sensitivity and responsiveness	☐ Develops alternative care plans when patients' personal beliefs and decisions preclude standard care
22	Patient centered communication (ICS1)	☐ Did not achieve level 1	☐ Establishes rapport and demonstrates empathy to patient (and family) Listens effectively	☐ Elicits patient's reason for seeking health care	☐ Manages patient expectations in a manner that minimizes potential for stress, conflict, and misunderstanding. Effectively communicates with vulnerable populations, (at risk patients and families)
23	Team management (ICS2)	☐ Did not achieve level 1	☐ Recognizes other members of the patient care team during case (nurse, techs)	☐ Communicates pertinent information to other healthcare colleagues	☐ Communicates a clear, succinct, and appropriate handoff with specialists and other colleagues Communicates effectively with ancillary staff