Lawrence Berkeley National Laboratory

Recent Work

Title

PRECISE MEASUREMENT OF THE ENERGY OP THE RAY FROM K*| DECAY

Permalink


https://escholarship.org/uc/item/2kg3r908

Authors

Smith, Alan R. Miller, Francis M.

Publication Date

1966-09-02



DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory Berkeley, California

AEC Contract No. W-7405-eng-48

PRECISE MEASUREMENT OF THE ENERGY OF THE $_{\gamma}$ RAY FROM K^{40} DECAY

Alan R. Smith and Francis M. Miller

September 2, 1966

Precise Measurement of the Energy of the γ Ray from K^{40} Decay

Alan R. Smith and Francis M. Miller

Lawrence Radiation Laboratory
University of California
Berkeley, California

September 2, 1966.

We have used a Ge(Li) semiconductor γ -ray spectrometer with a 1600-channel analyzer to obtain a precise measurement of the energy of the γ ray which accompanies one branch of K⁴⁰ decay. We find the value to be: 1459.6 keV with an estimated error of 0.2 keV.

Energy calibration of the system in the region of interest was determined precisely using known γ -ray energies of 1293.3 keV and 1507.5 keV from In 116m and 1172.8 keV and 1332.5 keV from Co 60 .

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

