UC San Diego

UC San Diego Previously Published Works

Title

Committee Opinion No. 703 Summary: Asymptomatic Microscopic Hematuria in Women.

Permalink

<https://escholarship.org/uc/item/2kz209vf>

Journal

Obstetrics and gynecology, 129(6)

ISSN

0029-7844

Authors

Nager, Charles W
Sung, Vivian W
Whiteside, James L
et al.

Publication Date

2017-06-01

DOI

10.1097/aog.0000000000002110

Peer reviewed

COMMITTEE OPINION SUMMARY

Number 703 • June 2017

For a comprehensive overview of these recommendations, the full-text version of this Committee Opinion is available at <http://dx.doi.org/10.1097/AOG.0000000000002059>.

Scan this QR code
with your smartphone
to view the full-text
version of this
Committee Opinion.

Committee on Gynecologic Practice American Urogynecologic Society

This Committee Opinion was developed by the American College of Obstetricians and Gynecologists' Committee on Gynecologic Practice and the American Urogynecologic Society in collaboration with committee members Charles W. Nager, MD, Vivian W. Sung, MD, and James L. Whiteside, MD.

This document reflects emerging clinical and scientific advances as of the date issued and is subject to change. The information should not be construed as dictating an exclusive course of treatment or procedure to be followed.

Asymptomatic Microscopic Hematuria in Women

ABSTRACT: Asymptomatic microscopic hematuria is an important clinical sign of urinary tract malignancy. Asymptomatic microscopic hematuria has been variably defined over the years. In addition, the evidence primarily is based on data from male patients. However, whether the patient is a man or a woman influences the differential diagnosis of asymptomatic microscopic hematuria, and the risk of urinary tract malignancy (bladder, ureter, and kidney) is significantly less in women than in men. Among women, being older than 60 years, having a history of smoking, and having gross hematuria are the strongest predictors of urologic cancer. In low-risk, never-smoking women younger than 50 years without gross hematuria and with fewer than 25 red blood cells per high-power field, the risk of urinary tract malignancy is less than or equal to 0.5%. Furthermore, the evaluation may result in more harm than benefit and is unlikely to be cost effective. Thus, data support changing current hematuria recommendations in this low-risk group. The American College of Obstetricians and Gynecologists and the American Urogynecologic Society encourage organizations producing future guidelines on the evaluation of microscopic hematuria to perform sex-specific analysis of the data and produce practical sex-specific recommendations. In the meantime, the American College of Obstetricians and Gynecologists and the American Urogynecologic Society recommend that asymptomatic, low-risk, never-smoking women aged 35–50 years undergo evaluation only if they have more than 25 red blood cells per high-power field.

Recommendations and Conclusions

The American College of Obstetricians and Gynecologists (the College) and the American Urogynecologic Society (AUGS) make the following recommendations and conclusions:

- Urinalysis is a commonly performed test and microscopic hematuria is a common finding.
- Renal cancer and bladder cancer are more common in men than women.
- In low-risk, never-smoking women younger than 50 years without gross hematuria and with fewer than 25 red blood cells per high-power field, the risk of urinary tract malignancy is less than or equal to 0.5%.
- The College and AUGS encourage organizations producing future guidelines on the evaluation of microscopic hematuria to perform sex-specific analysis of the data and produce practical sex-specific recommendations.

- The College and AUGS recommend that asymptomatic, low-risk, never-smoking women aged 35–50 years undergo evaluation only if they have more than 25 red blood cells per high-power field.

Full-text document published concurrently in the June 2017 issue of *Female Pelvic Medicine & Reconstructive Surgery*.

Copyright June 2017 by the American College of Obstetricians and Gynecologists. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, posted on the Internet, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without prior written permission from the publisher.

Requests for authorization to make photocopies should be directed to Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400.

The American College of Obstetricians and Gynecologists
409 12th Street, SW, PO Box 96920, Washington, DC 20090-6920

Official Citation

Asymptomatic microscopic hematuria in women. Committee Opinion No. 703. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2017;129:e168–72.