

# Lawrence Berkeley National Laboratory

## LBL Publications

### Title

EVIDENCE AGAINST  $A_x$  PRODUCTION IN HIGH ENERGY  $K+p$  INTERACTIONS

### Permalink

<https://escholarship.org/uc/item/2mk0h502>

### Authors

Rabin, Monroe S.  
Barbaro-Galtieri, Angela  
Derenzo, Stephen E.  
et al.

### Publication Date

1969-10-01

Submitted to Physical Review Letters

UCRL-19362  
Preprint

cy 2

RECEIVED  
LAWRENCE  
RADIATION LABORATORY

DEC 9 1969

LIBRARY AND  
DOCUMENTS SECTION

EVIDENCE AGAINST  $A_1$  PRODUCTION IN  
HIGH ENERGY  $K^+p$  INTERACTIONS

Monroe S. Rabin, Angela Barbaro-Galtieri, Stephen E. Derenzo,  
Stanley M. Flatté, Jerome H. Friedman, Margaret A. Garnjost,  
Frank T. Solmitz, Norman M. Uyeda, and Roland Windmolders†

October 1969

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. División, Ext. 5545*

LAWRENCE RADIATION LABORATORY  
UNIVERSITY of CALIFORNIA BERKELEY

UCRL-19362

cy 2

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

EVIDENCE AGAINST  $A_1$  PRODUCTION IN  
HIGH ENERGY  $K^+p$  INTERACTIONS\*

Monroe S. Rabin, Angela Barbaro-Galtieri, Stephen E. Derenzo,  
Stanley M. Flatté, Jerome H. Friedman, Margaret A. Garnjost,  
Frank T. Solmitz, Norman M. Uyeda, and Roland Windmolders†

Lawrence Radiation Laboratory  
University of California  
Berkeley, California 94720

October 1969

Abstract

The production of the  $A_1$  has been reported at 12.8 GeV/c in the reaction  $K^+p \rightarrow Kp(3\pi)$ . It has also been reported at 9.0 GeV/c in the reactions  $K^+p \rightarrow Kp(3\pi)$  and  $K^+p \rightarrow Kp(4\pi)$ . At 12.0 GeV/c, with five times the data of the 12.8 GeV/c experiment and three times the data of the 9.0 GeV/c experiment we see no evidence for  $A_1$  production in any of these reactions.

The  $A_1$  enhancement<sup>1</sup> has been seen mainly in the reaction  $\pi^\pm p \rightarrow A_1^\pm p$ ,  $A_1 \rightarrow \rho^0 \pi^\pm$ ,  $\rho^0 \rightarrow \pi^+ \pi^-$ , where its interpretation as a resonance has been questioned because Deck or other diffraction processes may be present.<sup>2</sup> In  $K^+p$  interactions, where  $A_1$  simulating effects of this type are probably not so prominent, the observation of the  $A_1$  would greatly favor the resonance interpretation independently of the concept of duality.<sup>2</sup>

Recently, observations of the  $A_1$  have been reported in the reactions

$$K^+p \rightarrow K^+p\pi^+\pi^-(\pi^0) \quad (\text{Ref. 3}) \quad (1)$$



in bubble chamber experiments at 12.8 (Ref. 3) and 9.0 (Ref. 4) GeV/c (the brackets indicate a particle not detected). In our experiment at 12.0 GeV/c, we have studied these reactions with a path length corresponding to about 35 events/microbarn -- at least three times as great as in either of the other experiments.<sup>5</sup>

All our events were measured on a Spiral Reader, which, in addition to the coordinate measurements, gave track ionization density information useful in selecting among possible kinematic hypotheses. For each kinematic fit a bubble density chi-square ( $\chi_B^2$ ) was calculated, and fits to reactions (1), (2b), and (3), which have one constraint, were considered only if their  $\chi_B^2$  was no more than three greater than the best  $\chi_B^2$  for all hypotheses tried, including missing-mass calculations.<sup>6</sup> The most probable reaction was defined to be that reaction with the lowest sum of kinematic  $\chi^2$  and bubble density  $\chi_B^2$ .

#### Reaction (1)

To make comparisons with other bubble chamber experiments more meaningful we feel that it is necessary to state the criteria used in selecting our sample of events for the study of reaction (1). We required that:

- (a) There be no four-constraint fit to the event with a confidence level greater than  $10^{-5}$ . (b) Reaction (1) be the most probable reaction and have a kinematic confidence level of at least  $10^{-3}$ . (c) In case of a final-state ionization ambiguity between the identities of the  $K^+$  and the proton, the proton be taken to be that track with the lower momentum.<sup>7</sup> (d) The square of the

four-momentum transferred between the target and final state proton,  $|t_{pp}|$ , be less than  $1.0 \text{ GeV}^2$ .

The main ambiguities resulting from this selection involved the identities of the  $\pi^0$  and  $K^0$ . Some events of reaction (2b) were included in the sample of events of reaction (1), whereas some events of reaction (1) were excluded from our sample and called events of reaction (2b). In addition, the identities of the  $\pi^+$  and  $K^+$  were interchanged in some of the events in our sample of reaction (1). By taking events of reaction (2a), ignoring the existence of the visible  $K^0$  decay, and refitting them kinematically, we were able to estimate that 8% of the events in our sample of reaction (1) came from reaction (2b). By passing Monte Carlo generated events<sup>8</sup> through our reconstruction and kinematic fitting programs, we found that for 10 to 20% of the events of reaction (1) the wrong permutation of the identities of the final state  $\pi^+$  and  $K^+$  was used.<sup>9</sup> We have examined the effects of these ambiguities and believe that they do not qualitatively affect the shapes of any of the distributions in our data.

The evidence presented for the production of the  $A_1$  in reaction (1) in the University of Rochester experiment at  $12.8 \text{ GeV}/c$  (Ref. 3) (hereafter referred to as UR) was based upon three observations: (i) the observation of a peak in the three-pion mass spectrum between 950 and 1150 MeV; (ii) the enhancement of this peak when events were selected from the periphery of the three-pion Dalitz plot with the requirement that at least one  $(2\pi)^\pm$  mass combination be consistent with the decay of a  $\rho^\pm$  meson; (iii) the presence of a  $\rho^\pm \pi^\mp$  signal and absence of a  $\rho^0 \pi^0$  signal in the vicinity of the  $A_1$  and no  $\rho$  signal of any kind in the sum of two control regions, above and below the  $A_1$ . We consider each of these three topics in turn.

The three-pion mass spectrum for the events passing our selection criteria is shown in Fig. 1a. We observe a broad plateau extending from 1.0 to 1.3 GeV, but no narrow peak at 1.0 GeV corresponding to that seen by UR.<sup>10</sup> Our results may be quantitatively compared with those of UR [Fig. 1(b) of Ref. 3;  $|t_{pp}| \leq 1.0 \text{ GeV}^2$ ] by minimizing the  $\chi^2$  sum

$$\chi^2 \equiv \sum_i \frac{(A_i R - B_i)^2}{A_i R^2 + B_i},$$

where  $A_i$  is the number of events in the  $i$ th bin of the  $M(3\pi)$  histogram of UR, and  $B_i$  is the number of events in the corresponding bin of the  $M(3\pi)$  histogram for this experiment. The parameter  $R$  is the ratio of the total numbers of events in the two histograms, and is varied to minimize  $\chi^2$ . For the region  $0.85 \leq M(3\pi) \leq 2.0 \text{ GeV}$  we obtain  $R = 0.21$  and  $\chi^2 = 61.7$  for 22 degrees of freedom -- a confidence level of  $10^{-5}$ .<sup>11</sup> The disagreement comes from two regions:  $0.9 \leq M(3\pi) \leq 1.1 \text{ GeV}$  and  $1.65 \leq M(3\pi) \leq 2.0 \text{ GeV}$ . The discrepancy at higher mass is most likely the result of a scanning bias by UR against higher-momentum protons.<sup>12</sup>

To better approximate the conditions of UR, Fig. 1b shows the events of Fig. 1a with the restriction that  $|t_{pp}| \leq 0.3 \text{ GeV}^2$ . The dots in Fig. 1b represent the data of UR multiplied by 2.6. The agreement between the two sets of data in the region  $1.3 \leq M(3\pi) \leq 2.0 \text{ GeV}$  is striking. In the  $\omega$  region, the disagreement could be a result of the better mass resolution of this experiment, and beyond  $M(3\pi) = 2.0 \text{ GeV}$  it is because we excluded all our events with  $|t_{pp}| > 0.3 \text{ GeV}^2$  from Fig. 1b. For the region  $0.85 \leq M(3\pi) \leq 2.0 \text{ GeV}$  we obtain  $R = 2.60$  and  $\chi^2 = 23.1$  for 22 degrees of freedom -- a confidence level of 0.40. Even in the small region around the  $A_1$ ,  $0.9 \leq M(3\pi) \leq 1.2 \text{ GeV}$ , we have

$R = 2.55$  and  $\chi^2 = 7.18$  for five degrees of freedom -- a confidence level of 0.24.

This means that the UR data are in statistical agreement with an experimental distribution containing more than two and a half times the data with no peak at the  $A_1$  mass. Note that our sample of events with  $|t_{pp}| \leq 1.0 \text{ GeV}^2$  (Fig. 1a) contains more than five times the number of events in the UR sample and also shows no peak at the mass of the  $A_1$ .

Selection by UR of events from the periphery of the  $3\pi$  Dalitz plot<sup>13</sup> resulted in greatly enhanced production of  $(\rho^\pm \pi^\mp)$  final states in the vicinity of the  $A_1$ . This was taken to be further evidence for  $A_1$  production, although it was stated that the enhancement might be due in part to a kinematic effect. Figure 2a shows our three-pion mass spectrum for the outer region of the Dalitz plot ( $\lambda \leq 0.006$ ),<sup>13</sup> and Fig. 2b shows those events of Fig. 2a consistent with the decay of at least one charged  $\rho$  ( $0.67 \leq M(\pi^\pm \pi^0) \leq 0.86 \text{ GeV}$ ). The enhancement at 1.0 GeV is striking and looks very similar to the one obtained by UR. We will now show that this enhancement can be understood as being entirely due to kinematics and therefore no resonance interpretation is necessary.<sup>14</sup> Figure 2c shows a normalized Dalitz plot on which the locations of the contour  $\lambda = 0.006$  and the  $\rho^\pm$  bands for  $M(3\pi) = 1.0 \text{ GeV}$  are drawn. As  $M(3\pi)$  increases, the  $\rho^\pm$  bands decrease in width and move across the plot in the directions shown by the arrows. We define the rho probability distribution (RPD) as the fraction of the outer Dalitz plot ( $\lambda \leq 0.006$ ) occupied by the  $\rho^\pm$  bands [ i. e., for  $M(3\pi) = 1 \text{ GeV}$  the value of the RPD is the fraction of the outer Dalitz plot in Fig. 2c occupied by the hatched area]. Figure 2d shows the RPD as a function of  $M(3\pi)$ . A prominent enhancement is seen in the region around 1 GeV. Thus, even a flat three-pion spectrum always appears to have a  $(\rho\pi)$  enhancement in the outer Dalitz plot at about 1 GeV. The result of folding



the RPD into the events in Fig. 2a is shown by the dashed lines in Fig. 2b. The shape of the experimental enhancement is reproduced closely; the absolute normalization is low by only about 15%. It should be kept in mind that the dashed lines are the result of assuming a uniform Dalitz plot population (i. e., no  $\rho$  production), and so agreement to within 15% is remarkable.

We now consider the two-pion spectrum as a function of three-pion mass for events with  $|t_{pp}| \leq 1 \text{ GeV}^2$ . Following UR, we define regions below, in, and above the  $A_1$  as 840 to 940 MeV, 940 to 1140 MeV, and 1140 to 1240 MeV, respectively. Figures 3a and 3b show the  $(\pi^+\pi^-)$  and  $(\pi^\pm\pi^0)$  signals for events in Fig. 1a occurring in the lower region. No evidence for the presence of the  $\rho$  is seen or expected because of lack of phase space. Figures 3c-f show these distributions for events in and above the  $A_1$  region. The existence of the  $\rho$  signal is apparent in the upper control region, and its strength is comparable to that in the  $A_1$  region.<sup>15</sup> The ratio of charged to neutral  $\rho$  appears to be the same above the  $A_1$  region as in the  $A_1$  region, making it unnecessary to invoke the presence of the  $A_1$  to explain the data.

#### Reactions (2a), (2b), and (3)

Evidence for the production of the  $A_1$  in reactions (2a), (2b), and (3) is even weaker than it is in the UR data for reaction (1). UR has 381 examples of reaction (2a), and the 9.0-GeV/c  $K^+p$  experiment<sup>4</sup> has 456 events in reaction (2a), 1475 in (2b), and 1141 in (3).

In our experiment, examples of reactions (2b) and (3) were chosen from the four-pronged and V-four-pronged events, respectively, by criteria similar to those used in the study of reaction (1). In addition, for reaction (2b), we required that the missing mass squared be negative when the track fitted as a  $\pi^+$ , and with the greater momentum in the laboratory frame, was interpreted

as a  $K^+$ . This removed background produced by the reaction  $K^+ p \rightarrow K^+ p \pi^+ \pi^- (n\pi^0)$ , for  $n > 1$ , but did not change the shape of the distribution of interest, namely  $M(\pi^+ \pi^+ \pi^-)$ . Reaction (2a) comes from seven-constraint fits to the V-four-pronged events and is unambiguous.

Imposing the requirement that  $|t_{pp}| \leq 1.0 \text{ GeV}^2$  leaves us with 1454 examples of reaction (2a), 5431 examples of reaction (2b), and 2647 examples of reaction (3). The relevant three-pion mass spectra are shown in Fig. 4. There is no evidence of  $A_1$  production in these data, especially after we have removed from reaction (3) events where one of the  $\pi^+$  is consistent with coming from the decay  $K^{*+}(890) \rightarrow K^0 \pi^+$ , a competing reaction (see hatched graphs in Fig. 4).

In summary, we conclude that there is evidence in our experiment against the production of the  $A_1$  in reaction (1) with a cross section greater than  $5 \mu\text{b}$ , and no convincing evidence for its production in the UR data, where a production cross section of  $40 \mu\text{b}$  was quoted. There is also no evidence in our experiment for  $A_1$  production in reactions (2a), (2b), or (3) with a cross section greater than  $5 \mu\text{b}$ .

We thank Joseph J. Murray for his work in beam design and construction. We gratefully acknowledge the assistance of the staff of the Stanford Linear Accelerator and the 82-Inch Bubble Chamber in obtaining the data for this experiment. We also acknowledge the help of the Lawrence Radiation Laboratory Group A Scanning and Measuring Group in data reduction and we are grateful to Gerald R. Lynch for stimulating discussions.

where  $\vec{P}_+$  and  $\vec{P}_-$  are the three-momenta of the  $\pi^+$  and the  $\pi^-$ , respectively, in the  $3\pi$  c.m.) We define this contour using the parameter  $\lambda = \left| \frac{\vec{P}_+ \times \vec{P}_-}{Q^2 + 2Qm} \right|^2$ , where  $Q$  is the total kinetic energy of the three pions in their c.m. and  $m$  is the sum of the masses of the three pions. This definition of  $\lambda$  is an energy-invariant generalization of the definition given in Ref. 3. The value of  $\lambda$  used is 0.006.

14. The technique we use is similar to that used in studying the kinematical origin of the H meson. See, for example, A. H. Rosenfeld, N. Barash-Schmidt, A. Barbaro-Galtieri, L. R. Price, P. Söding, C. G. Wohl, M. Roos, and W. J. Willis, *Rev. Mod. Phys.* **40**, 77 (1968); S. -Y. Fung, W. Jackson, R. T. Pu, D. Brown, and G. Gidal *Phys. Rev. Letters* **21**, 47 (1968).
15. Adding the control regions together causes the  $\rho$  signal to be obscured. This explains UR's observation of the absence of the  $\rho$  signal when they summed their control regions.

Figure Legends

- Fig. 1. Three-pion mass spectrum for events in reaction (1) with  
 (a)  $|t_{pp}| \leq 1.0 \text{ GeV}^2$ , (b)  $|t_{pp}| \leq 0.3 \text{ GeV}^2$ . The dots represent the data of Fig. 1b of Ref. 3 multiplied by 2.6.
- Fig. 2. (a) Three-pion mass spectrum for those events of Fig. 1b with  $\lambda \leq 0.006$ .  
 (b) Three-pion mass spectrum for those events of Fig. 2a in which at least one  $(\pi^\pm \pi^0)$  combination has a mass between 0.67 and 0.86 GeV. The dashed lines are a prediction assuming a completely uniform Dalitz plot (see text).  
 (c) Normalized Dalitz plot showing the contour  $\lambda = 0.006$  and the position of the  $\rho^\pm$  bands ( $0.67 \leq M(\pi^\pm \pi^0) \leq 0.86 \text{ GeV}$ ) for  $M(3\pi) = 1 \text{ GeV}$ . The arrows show the directions in which the respective  $\rho$  bands move for increasing  $M(3\pi)$ .  
 (d) The  $\rho$  probability distribution (RPD) -- the fraction of the periphery of the Dalitz plot ( $\lambda \leq 0.006$ ) occupied by the  $\rho^\pm$  bands as a function of  $M(3\pi)$ .
- Fig. 3. (a), (c), (e) The  $(\pi^+ \pi^-)$  mass distribution for events of Fig. 1a below (900 events), in (2559 events), and above the  $A_1$  region (1411 events), respectively.  
 (b), (d), (f) The sum of the  $(\pi^+ \pi^0)$  and  $(\pi^- \pi^0)$  mass distributions for events of Fig. 1a below, in, and above the  $A_1$  region, respectively (2 combinations per event).

Fig. 4. Three pion mass spectra for events with  $|t_{pp}| \leq 1.0 \text{ GeV.}^2$


The hatched events are those with a competing  $K^{*+}(890)$  removed.

(a)  $M(\pi^+ \pi^+ \pi^-)$  from reaction (2a).

(b)  $M(\pi^+ \pi^+ \pi^-)$  from reaction (3).


(c)  $M(\pi^+ \pi^- \pi^0)$  from reaction (3), two combinations per event.

(d)  $M(\pi^+ \pi^+ \pi^-)$  from reaction (2b).




XBL6910 - 6068

Fig. 1



XBL6910-6070


Fig. 2



XBL6910-6067

Fig. 3





XBL6910-6069

Fig. 4

LEGAL NOTICE

*This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:*

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or*
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.*

*As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.*

TECHNICAL INFORMATION DIVISION  
LAWRENCE RADIATION LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720