

Lawrence Berkeley National Laboratory

LBL Publications

Title

Nuclear Quadrupole Ratio of Bromine Isotopes in Molecular LiBr

Permalink

<https://escholarship.org/uc/item/2mw9x432>

Authors

Hebert, A J

Street, K, Jr.

Publication Date

1968-08-01

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

cy. 2

University of California

Ernest O. Lawrence
Radiation Laboratory

RECEIVED
LAWRENCE
RADIATION LABORATORY

OCT 23 1968

LIBRARY AND
DOCUMENTS SECTION

NUCLEAR QUADRUPOLE RATIO OF BROMINE ISOTOPES
IN MOLECULAR LiBr

A. J. Hebert and K. Street, Jr.

August 7, 1968

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

UCRL-18353 Rev.
cy. 2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

To be submitted to Physical Review

UCRL-18353 Rev.

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

NUCLEAR QUADRUPOLE RATIO OF BROMINE ISOTOPES IN MOLECULAR LiBr*

A. J. Hebert and K. Street, Jr.

August 7, 1968

NUCLEAR QUADRUPOLE RATIO OF BROMINE ISOTOPES IN MOLECULAR LiBr*

A. J. Hebert and K. Street, Jr.

Lawrence Radiation Laboratory and Department of Chemistry,
University of California,
Berkeley, California

(Received 12 August 1968)

The Molecular Beam Electric Resonance method has been used to determine the ^{79}Br to ^{81}Br nuclear quadrupole moment ratio in molecular LiBr. The observed ratio of 1.197056(6) agrees with one obtained elsewhere for atomic bromine, thus indicating no observable nuclear polarizability effect.

generator at a 19 Hz rate across each line in synchronization with the channel advance in a modified Northern multichannel analyzer. The detected "flop-in" transitions were all $J = 1, m_J = \pm 1$ to $J = 1, m_J = 0$ for the three lowest vibrational levels of each isotope. Peak signal intensities were better than 10^4 transitions per channel above background for all observed lines with no observable baseline slope. Typical run times were from one to three hours (approximately 70 to 210 thousand sweeps) for each line with signal to noise ratios of 100 or better. Line position reproducibility was better than 1 part in 10^6 of frequency for runs made on different days. This corresponds to an uncertainty of less than 15 Hz at 15 MHz, compared to the previous several hundred Hz. All observed lines occurred between 8 and 20 MHz.

Previous strong field electric dipole moment measurements¹ were checked and found to be accurate. Measurements performed in this laboratory by F. W. Breivogel, Jr. on the $^7\text{LiBr}$ isotopes⁴ allowed hyperfine corrections for the ^6Li quadrupole and spin-rotation interactions. Second order quadrupole, Stark-quadrupole, and magnetic dipole spin-spin interaction effects were included in treating the data. Calculations indicate that a large electron coupled spin-spin interaction, (i.e. much greater than the calculated dipole spin-spin interaction of approximately 90 cps), would have a negligible effect on the values reported here.

Calculated hyperfine interaction constants are given in Table I. The present results for each of the three vibrational states measured are in agreement with our previously reported values. The present bromine nuclear quadrupole moment ratios are lower than those previously reported by almost

3 times the probable error. This discrepancy arises primarily from the manner in which the data were treated. In the previous report the quadrupole coupling constants for each isotopic species were separately extrapolated to $v = -1/2$ and the ratio then taken. The probable error was assigned assuming that the errors would be random. The new measurements indicate that there was a non-random trend in the data, which was accentuated by extrapolation to the equilibrium internuclear distance. In our opinion, this non-random trend was produced by a combination of decreasing signal to noise ratio for higher vibrational states, and a previous lack of knowledge about the magnitude of the ${}^6\text{Li}$ quadrupole interaction.

In looking for a nuclear polarizability effect a much better method of treating the data is to simply interpolate the ${}^{81}\text{Br}$ coupling constant for each vibrational state the very short distance in v necessary to achieve the same energy as the corresponding ${}^{79}\text{Br}$ vibrational state. This small correction is much less sensitive to uncertainties in $(eqQ)_{\text{I}}$ and $(eqQ)_{\text{II}}$. The bromine nuclear quadrupole moment ratios calculated in this manner are given in Table II. Treating the previous data in the same manner gives an average ${}^{79}\text{Br}$ to ${}^{81}\text{Br}$ nuclear quadrupole ratio of 1.19712(8) for the weak field data, and an average of 1.19706(8) for the three states observed at intermediate electric field strength.

The best present ${}^{79}\text{Br}$ to ${}^{81}\text{Br}$ nuclear quadrupole ratio of 1.197056(6) for LiBr (calculated from the $v = 0$ vibrational states corrected to a common energy), is in excellent agreement with the value 1.1970568(15) observed by Brown and King for atomic Br, thus indicating no observable nuclear polarizability effect.

Table I. Hyperfine Interaction Constants of LiBr ($\times \frac{1}{h}$ K Hz).*

Vibrational State	${}^6\text{Li } {}^{79}\text{Br}$		${}^6\text{Li } {}^{81}\text{Br}$	
	Present	Previous ^a	Present	Previous ^a
0	38,463.92(30)	$\begin{cases} 38,463(5) \\ 38,464(5) \end{cases}$	32,131.06(30)	$\begin{cases} 32,127(5) \\ 32,130(5) \end{cases}$
1	41,307.01(40)	$\begin{cases} 41,304(5) \\ 41,306(5) \end{cases}$	34,504.11(40)	$\begin{cases} 34,500(5) \\ 34,504(5) \end{cases}$
2	44,069.71(50)	$\begin{cases} 44,067(5) \\ 44,070(5) \end{cases}$	36,810.41(50)	$\begin{cases} 36,805(5) \\ 36,810(5) \end{cases}$
3		46,751(5)		39,046(5)
4		49,355(5)		41,222(5)
5		51,883(5)		
0	4.3(2)		4.3(2)	
1	4.3(4)		4.3(4)	
2	4.3(6)		4.3(6)	
0	9.064(70)	9.2(2)	9.793(70)	9.9(2)
1	9.024(80)	9.2(2)	9.727(80)	9.9(2)
2	9.017(90)	9.2(2)	9.690(90)	9.9(2)
0	0.22(13)		0.22(13)	
1	0.22(13)		0.22(13)	
2	0.22(13)		0.22(13)	
	$(\text{eqQ})_v = (\text{eqQ})_e + (\text{eqQ})_I \left(v + \frac{1}{2}\right) + (\text{eqQ})_{II} \left(v + \frac{1}{2}\right)^2$			
	37,012.233(850)	37,015(3)	30919.501(850)	30,916(2)
	2,923.474(900)	2,918(2)	2,439.798(900)	2,438(2)
	-40.194(350)	-39(2)	-33.374(350)	-33(2)

* All values were calculated using $h = 6.6253 \times 10^{-27}$ erg-sec and $c = 2.997925 \times 10^{10}$ cm/sec.

a. Ref. 1. Where two values are given the second is from data taken at intermediate fields.

b. Isotope corrected values calculated from data in Ref. 4.

Table II. Nuclear Quadrupole Moment Ratios Q_{79}/Q_{81}

V	$\frac{(eqQ)_{79}}{(eqQ)_{81}}$ (Uncorrected)	$\frac{(eqQ)_{79}}{(eqQ)_{81}}$ (Corrected to a common vibrational energy)
0	1.197095(6)	1.197056(6)
1	1.197162(9)	1.197055(9)
2	1.197208(12)	1.197046(12)

$$[(eqQ)_e]_{79_{Br}} / [(eqQ)_e]_{81_{Br}} = 1.197051(12)$$

$$\text{Atomic Bromine Ratio}^3 \frac{Q_{79}}{Q_{81}} = 1.1970568(15)$$

REFERENCES

*This work was supported under the auspices of the U. S. Atomic Energy Commission.

1. A. J. Hebert, F. W. Breivogel, Jr. and K. Street, Jr., J. Chem. Phys. 41, 2368 (1964), and 47, 2202 (1967).
2. P. A. Bonczyk and V. W. Hughes, Phys. Rev. 161, 15 (1967).
3. H. H. Brown and J. G. King, Phys. Rev. 142, 53 (1966).
4. F. W. Breivogel, Jr., Ph.D. Thesis, University of California, Lawrence Radiation Laboratory Report, UCRL-11665, September (1964).

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.