UCLA

UCLA Previously Published Works

Title

Adenosine receptor ligation tips the uveitogenic Th1 and Th17 balance towards the latter in experimental autoimmune uveitis-induced mouse.

Permalink

<https://escholarship.org/uc/item/2n1247fp>

Authors

Sun, Deming
Ko, Minhee
Shao, Hui
et al.

Publication Date

2021

DOI

10.1016/j.crimmu.2021.07.001

Peer reviewed

Adenosine receptor ligation tips the uveitogenic Th1 and Th17 balance towards the latter in experimental autoimmune uveitis-induced mouse

Deming Sun^{a,*}, Minhee Ko^a, Hui Shao^b, Henry J. Kaplan^c

^a Doheny Eye Institute and Department of Ophthalmology, David Geffen School of Medicine at UCLA, Los Angeles, CA, 90033, United States

^b Department of Ophthalmology and Visual Sciences, Kentucky Lions Eye Center, University of Louisville, Louisville, KY, 40202, United States

^c Saint Louis University (SLU) Eye Institute, SLU School of Medicine, Saint Louis, MO, 63104, United States

ARTICLE INFO

Keywords:

Adenosine

Adenosine deaminase

Autoimmune response

$\gamma\delta$ T cells

Th17

ABSTRACT

Various pathological conditions are accompanied by release of adenosine triphosphate (ATP) from the intracellular to the extracellular compartment, where it degrades into adenosine and modulates immune responses. Previous studies concluded that both ATP and its degradation product adenosine are important immunoregulatory molecules; ATP acted as a danger signal that promotes immune responses, but adenosine's effect was inhibitory. We show that adenosine receptor ligation plays an important role in balancing Th1 and Th17 pathogenic T cell responses in experimental autoimmune uveitis (EAU). While its effect on Th1 responses is inhibitory, its effect on Th17 responses is enhancing, thereby impacting the balance between Th1 and Th17 responses. Mechanistic studies showed that this effect is mediated via several immune cells, among which $\gamma\delta$ T cell activation and dendritic cell differentiation are prominent; adenosine- and $\gamma\delta$ -mediated immunoregulation synergistically impact each other's effect. Adenosine receptor ligation augments the activation of $\gamma\delta$ T cells, which is an important promoter for Th17 responses and has a strong effect on dendritic cell (DC) differentiation, tipping the balance from generation of DCs that stimulate Th1 responses to those that stimulate Th17 responses. The knowledge acquired in this study should improve our understanding of the immune-regulatory effect of extracellular ATP-adenosine metabolism and improve treatment for autoimmune diseases caused by both Th1- and Th17-type pathogenic T cells.

1. Introduction

The purinergic system is an evolutionally selected system modulating immune responses (Haskó et al., 2008; Junger, 2011). Under physiological conditions, adenosine triphosphate (ATP) is contained exclusively within cells; however, almost all types of mammalian cells are able to release ATP during tissue damage and inflammation (Junger, 2011). Upon entering the extracellular space, ATP is hydrolyzed into adenosine diphosphate, adenosine-5'-monophosphate, and finally, adenosine in a stepwise manner by ectonucleotidases, including CD73 and CD39 (Fredholm et al., 2011; Haskó et al., 2008; Jacobson and Gao, 2006; Sauer et al., 2012; Yegutkin, 2008). Previous studies have shown that while ATP acts like an endogenously generated Toll-like receptor (TLR) ligand capable of augmenting immune responses (Beigi et al., 2003; Canaday et al., 2002; Hanley et al., 2004; Wilkin et al., 2001), the ATP metabolite adenosine is profoundly anti-inflammatory (Antonoli et al., 2013; Eltzschig and Carmeliet, 2011; Haskó et al., 2009;

Naganuma et al., 2006; Ohta and Sitkovsky, 2001; Zarek et al., 2008). An increase in extracellular adenosine reduces the local inflammatory response, while removal of endogenous adenosine aggravates tissue dysfunction elicited by injury (Grenz et al., 2011). Binding of adenosine to its receptors modulates various pathophysiological responses, including immune responses (Fredholm et al., 2011; Haskó et al., 2008; Jacobson and Gao, 2006; Sauer et al., 2012). The discovery of the regulatory effect of adenosine on inflammation and immune responses has led to attempts to treat immune dysfunctions by targeting adenosine receptor (AR) signaling (Haskó et al., 2008; Jacobson and Gao, 2006). Targeting of ARs and adenosine generation has been successful in treating cancer and neurological diseases (Cronstein et al., 1991; Jacobson and Gao, 2006; Ramlackhansingh et al., 2011).

The extrapolation of adenosine as inhibitory was mostly obtained from studies of Th1-type (interferon (IFN)- γ -producing cells) immune responses, since Th17 responses were discovered only recently. Given the available knowledge that both Th1 and Th17 pathogenic T cells

* Corresponding author. Department of Ophthalmology, University of California Los Angeles, Los Angeles, CA90033, USA.

E-mail addresses: Dsun@doheny.org (D. Sun), mko@doheny.org (M. Ko).

<https://doi.org/10.1016/j.crimmu.2021.07.001>

Received 19 April 2021; Received in revised form 19 July 2021; Accepted 22 July 2021

Available online 26 July 2021

2590-2555/© 2021 University of California, Los Angeles. Published by Elsevier B.V. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

contribute to the pathogenesis of autoimmune diseases (Bettelli et al., 2006; Cua et al., 2003; Dong, 2006), determination of whether adenosine has a similar effect on Th1 and Th17 pathogenic T cell responses is important. In this study we show that the effect of AR ligation on Th17 responses is fundamentally different than its effect on Th1 responses; while it inhibits Th1 responses, it enhances Th17 responses. Mechanistic studies showed that the enhancing effect of adenosine on Th17 responses is accomplished via a sum of effects on various other cellular responses important for T cell activation, including $\alpha\beta$ T cells, $\gamma\delta$ T cells, DCs and regulatory T cells. Adenosine is an important co-stimulating molecule for $\gamma\delta$ T cell activation, and augmented $\gamma\delta$ T cell activation leads to high Th17 responses (Cui et al., 2009; Liang et al., 2013b, 2016a; Nian et al., 2010). We also show that adenosine exposed DCs showed a greater stimulating effect on $\gamma\delta$ T cell activation. Thus, reciprocal interaction between $\gamma\delta$ T cells and DCs leads to enhanced Th17 responses. Adenosine and $\gamma\delta$ -based treatments should be more successful if the mechanisms by which they affect Th1 and Th17 responses are better understood.

2. Materials and methods

2.1. Animals and reagents

All animal studies conformed to the Association for Research in Vision and Ophthalmology statement on the use of animals in Ophthalmic and Vision Research. Institutional approval by Institutional Animal Care and Use Committee (IACUC) of Doheny Eye Institute, University of California Los Angeles was obtained, and institutional guidelines regarding animal experimentation were followed.

Female C57BL/6 (B6) and TCR- $\delta^{-/-}$ mice on the B6 background were purchased from Jackson Laboratory (Bar Harbor, ME, USA). A2AR $^{-/-}$ mice (Chen et al., 1999) were a gift from Dr. Jiang-Fan Chen (Boston University School of Medicine, Boston, MA, USA). They were housed and maintained in the animal facilities of the University of California Los Angeles. Recombinant murine IL-1 β , IL-7, and IL-23 were purchased from R & D Systems (Minneapolis, MN, USA). Fluorescein isothiocyanate (FITC)-, phycoerythrin (PE)-, or allophycocyanin-conjugated antibodies (Abs) against mouse CD4 (GK1.5), $\alpha\beta$ T cell receptor (TCR) (H57-597), or $\gamma\delta$ TCR (GL3) and their isotype control antibodies were purchased from Biolegend (San Diego, CA, USA). (PE)-conjugated anti-mouse IFN- γ (XMG1.2) and IL-17 (TC11-18H10.1) monoclonal antibody was purchased from Santa Cruz Biotechnology (Dallas, TX, USA). The non-selective AR agonist 5'-N-ethylcarboxamidoadenosine (NECA) (Mahamed et al., 2015), selective A2AR agonist 2-p-(2-carboxyethyl) phenethylamino-5'-N-ethylcarboxamidoadenosine (CGS21680), selective A1R agonist (CCPA), A2BR agonist BAY60-6538, A2AR antagonist (SCH 58261) (Feoktistov and Biaggioni, 1997; Zocchi et al., 1996), and erythro-9-(2-hydroxy-3-nonyl) (EHNA, an inhibitor of adenosine deaminase [ADA]) were purchased from Sigma-Aldrich and were dissolved as a 1 mM stock solution in DMSO and diluted 1/10000 in culture medium before use. Toll-like receptor ligands lipopolysaccharide (LPS) and Pam3csk4 (Pam3) and ADA polyclonal antibody were purchased from Invivogen (San Diego, CA, USA).

2.2. Immunization and EAU induction

EAU was induced in B6 mice by subcutaneous injection of 200 μ l of emulsion containing 200 μ g of human interphotoreceptor retinoid-binding protein (IRBP)₁₋₂₀ (Sigma-Aldrich) in complete Freund's adjuvant (CFA; Difco, Detroit, MI, USA) at six spots at the tail base and on the flank and intraperitoneal injection with 300 ng of pertussis toxin.

2.3. T cell preparations

$\alpha\beta$ T cells were purified from B6 mice immunized with IRBP₁₋₂₀, as

described previously (Cui et al., 2009; Liang et al., 2013b; Nian et al., 2010), while $\gamma\delta$ T cells were purified from immunized and control (naïve) B6 mice. Nylon wool-enriched splenic T cells from naïve or immunized mice were incubated sequentially for 10 min at 4 °C with FITC-conjugated anti-mouse $\gamma\delta$ TCR or $\alpha\beta$ TCR Abs and 15 min at 4 °C with anti-FITC Microbeads (Miltenyi Biotec GmbH, Bergisch Gladbach, Germany), then separated into bound and non-bound fractions on an autoMACS™ separator column (Miltenyi Biotec GmbH). The purity of the isolated cells, determined by flow cytometric analysis using PE-conjugated Abs against $\alpha\beta$ or $\gamma\delta$ T cells, was >95 %.

2.4. Prepare $\gamma\delta$ T cells

Non-activated and activated $\gamma\delta$ T cells were separated from either naïve B6 mice or IRBP₁₋₂₀-immunized B6 mice (Liang et al., 2013b) (Liang et al., 2016a), respectively, by positive selection using a combination of FITC-conjugated anti-TCR- δ antibody and anti-FITC antibody-coated Microbeads, followed by separation using an auto-MACS.

2.5. Measurement of Th1 and Th17 responses

$\alpha\beta$ T cells (1.8×10^6) were collected from IRBP₁₋₂₀-immunized B6 mice on day 13 post-immunization, based on previous tests showing that highest T cell responses are detected on days 13–15 post immunization. To obtain a sufficient number of cells, we routinely pool the cells obtained from all six mice in the same group, before the T cells are further enriched. The cells were co-cultured for 48 h with irradiated spleen cells (1.5×10^6 /well) as antigen presenting cells (APCs) and IRBP₁₋₂₀ (10 μ g/ml) in a 24-well plate under either Th1 (culture medium supplemented with 10 ng/ml of IL-12) or Th17 polarized conditions (culture medium supplemented with 10 ng/ml of IL-23) (Liang et al., 2013b, 2014a). Cytokine (IFN- γ and IL-17) levels in the serum and 48 h of culture supernatants were measured by ELISA (R&D Systems). The percentage of IFN- γ^+ and IL-17 $^+$ T cells among the responder T cells was determined by intracellular staining 5 days after in vitro stimulation, followed by FACS analysis, as described previously (Liang et al., 2014a).

2.6. Generation of bone marrow dendritic cells

Bone marrow dendritic cells (BMDCs) were generated by incubating bone marrow cells for 5 days in the presence of 10 ng/ml of recombinant murine GM-CSF and IL-4 (R&D Systems), as described previously (Inaba et al., 1992). Cytokine (IL-1 β , IL-6, IL-12 and IL-23) levels in the culture medium were measured by ELISA after BMDCs were treated with AR agonists. To determine antigen-presenting function, BMDCs were incubated in a 24-well plate with responder T cells isolated from immunized B6 mice under Th1- or Th17-polarizing conditions. Forty-eight hours after stimulation, IFN- γ and IL-17 in the culture medium were measured by ELISA. The percentage of IFN- γ^+ and IL-17 $^+$ T cells among the responder T cells was determined by intracellular staining after 5 days of culture as described above.

2.7. Intracellular cytokine flow cytometry

Unfractionated or purified CD3 $^+$ T cells isolated from immunized mice were stimulated in vitro with 50 ng/ml of PMA, 1 μ g/ml of ionomycin and 1 μ g/ml of brefeldin A (Sigma-Aldrich, St. Louis, MO) for 4 h, then washed, fixed, permeabilized overnight with Cytofix/Cytoperm buffer (eBioscience, San Diego, CA). The cells were then intracellularly stained with antibodies against IFN- γ and IL-17 and analyzed on a FACScalibur flow cytometer.

2.8. Carboxyfluorescein succinimidyl ester (CFSE) assay

Purified $\alpha\beta$ T cells from IRBP₁₋₂₀-immunized B6 mice were stained with carboxyfluorescein succinimidyl ester (CFSE, Sigma-Aldrich) as

described previously [34]. Briefly, the cells were washed and suspended as 50×10^6 cells/ml in serum-free RPMI 1640 medium (Corning Cellgro, VA) and incubated at 37°C for 10 min with gentle shaking with a final concentration of $5 \mu\text{M}$ CFSE. The cells were then washed twice with RPMI 1640 medium containing 10 % fetal calf serum (Atlantic Inc. Santa Fe, CA, USA; complete medium), suspended in complete medium, stimulated with immunizing peptide in the presence of irradiated syngeneic spleen cells as antigen-presenting cells (APCs), and analyzed by flow cytometry.

2.9. ELISA measurement of cytokine

Purified $\alpha\beta$ T cells (3×10^4 cells/well; $200 \mu\text{l}$) from the draining lymph nodes and spleens of IRBP₁₋₂₀-immunized B6 mice were cultured in complete medium at 37°C for 48 h in 96-well microtiter plates with irradiated syngeneic spleen APCs (1×10^5) in the presence of $10 \mu\text{g/ml}$ of IRBP₁₋₂₀. A fraction of the culture supernatant was then assayed for IL-17 and IFN- γ , using ELISA kits (R & D Systems).

2.10. Statistical analysis

The results in the figures are representative of one experiment, which was repeated 3–5 times. Data were analyzed using a paired *t*-test. A *P* value < 0.05 was considered a statistically significant difference and was marked with ** when $P < 0.01$.

3. Results

3.1. Adenosine preferentially inhibits Th1 but not Th17 responses

To determine the adenosine effect on Th1 and Th17 responses in EAU, CD3⁺ responder T cells were harvested 13 days post immunization from the spleens and draining lymph nodes of B6 mice immunized with a uveitogenic antigen (IRBP₁₋₂₀). The responder T cells were stimulated in vitro with the immunizing peptide and APCs (irradiated spleen cells), in the absence or presence of a selective A2AR agonist (CGS21680), under culture conditions that favor Th17 or Th1 autoreactive T cell expansion (medium containing 10 ng/ml, respectively, IL-23 or IL-12) (Liang et al.,

Fig. 1. The effect of adenosine receptor agonist on Th17 responses differed from its effect on Th1 responses. A&B) B6 mice were immunized with interphotoreceptor retinoid-binding protein (IRBP)₁₋₂₀/complete Freund's adjuvant (CFA). Thirteen days after immunization, CD3⁺ cells were separated from spleen and draining lymph nodes cells of immunized mice using a MACS column. They were stimulated with the immunizing peptide (IRBP₁₋₂₀) and APCs, in the absence or presence of an A2AR agonist (CGS21680, 250 nM), under Th17 (right panels) or Th1 (left panels) polarized conditions. The numbers of $\alpha\beta\text{TCR}^+$ IL-17⁺ cells were assessed after a 5-day in vitro stimulation by FACS analysis. Data summarized for 4 separate experiments are plotted in (B). Data were analyzed using a paired *t*-test. ***p* < 0.01 ; ns, not significant, *n* = 6 in each group. C) Carboxyfluorescein succinimidyl ester assay for assessing dose-dependent effect (0–200 nM) of A2AR agonist (CGS21680) on Th1 and Th17 response. MACS column-separated CD3⁺ cells of immunized B6 mice were stimulated with the immunizing peptide (IRBP₁₋₂₀) and antigen presenting cells, under Th17 or Th1 polarized conditions, in the presence of indicated doses of CGS21680. The numbers of activated T cells were assessed by FACS analysis after a 5-day in vitro stimulation. The results shown are representative of those from five experiments. D). Calculated inhibition of Th1 and Th17 response by graded doses of CGS21680. The graphs are showing SEM. Data were analyzed using a paired *t*-test. ***p* < 0.01 ; ns, not significant, *n* = 6 in each group.

2013b; Zuo et al., 2012). Th1 and Th17 responses specific for the immunizing antigen were estimated by assessing responding IFN- γ ⁺ and IL-17⁺ T cells after intracellular staining with Fluorescence-labeled anti-IFN- γ or anti-IL-17 antibodies (Fig. 1A&B). The results showed that the number of IFN- γ ⁺ cells in response to CGS21680 decreased significantly, whereas the number of IL-17⁺ T cells remained unchanged. We have previously shown that $\gamma\delta$ T cell was a major contributor to the regulation of Th17 responses. To determine whether adenosine would have similar effect on Th1 and Th17 responses in the absence of $\gamma\delta$ T cells, which play a major role in Th17 responses (Nian et al., 2011; Rajan et al., 2000; Spahn et al., 1999), we prepared responder CD3⁺ T cell from immunized TCR- δ ^{-/-} mice and assessed T cell activation in the presence of varying doses of A2AR agonist using a CFSE assay (Fig. 1C), in which the responder cells were pre-labeled with CFSE before stimulation under polarizing conditions. The results show that in the absence of $\gamma\delta$ T cells both Th1 and Th17 responses are inhibited by A2AR agonist. However, the Th1 responses were readily

inhibited by a very low dose (20 nM) of the A2AR agonist that is inhibitory for Th1 response; but the Th17 responses remained minimally affected unless a very high dose (>200 nM) of the A2AR agonist was tested. Measurement of cytokine production of the responder T cells showed that IFN- γ production was inhibited by a very low dose (20 nM) of A2AR agonist while IL17 production was only inhibited by doses of A2AR agonist that were 10 times higher (Fig. 1D).

3.2. $\gamma\delta$ T cells offset an inhibitory effect of A2AR agonist on Th17 responses

Previous studies showed that $\gamma\delta$ T cells are important enhancers of Th17 responses (Nian et al., 2011; Rajan et al., 2000; Spahn et al., 1999). To determine the mechanism by which A2AR agonist is more inhibitory for Th1 responses than Th17 response, we compared Th17 responses in the presence or absence of $\gamma\delta$ T cells. The CD3⁺ T cells containing $\gamma\delta$ T cells were purified from immunized B6 mice. Those not containing $\gamma\delta$ T

Fig. 2. $\gamma\delta$ T cell offsets the inhibitory effect of A2AR agonist on Th17 responses. A) Intracellular staining of IL-17⁺ T cells among the responder T cells. Responder T cells were separated from either immunized B6 (top panels) TCR- δ ^{-/-} mice without (mid panels) or with (lower panels) 2 % supplemented $\gamma\delta$ T cells. After stimulation with the immunizing peptide interphotoreceptor retinoid-binding protein (IRBP)₁₋₂₀ and antigen presenting cells (APCs), under Th17 polarized conditions. The numbers of $\alpha\beta$ TCR⁺ IL-17⁺ and $\alpha\beta$ TCR⁺ IFN- γ ⁺ cells were assessed by FACS analysis after a 5-day in vitro stimulation. The results shown are representative of those from five experiments. B) Summary data for all 5 replicates of Fig. 2A. Data were analyzed using a paired *t*-test. ***p* < 0.01, *n* = 6 in each group. C) ELISA test assesses IL-17 (upper panels) and IFN- γ production (upper two panels) by B6 (left panels) and TCR- δ ^{-/-} responder T cells (lower two panels) under effect of agonists for specific adenosine receptors A1R (CCPA, 50 nM), A2AR (CGS 21680, 250 nM), A2BR (BAY60-6538, 100 nM), and vehicle control. The graphs show SEM. Data were analyzed using a paired *t*-test. ***p* < 0.01, *n* = 6 in each group. D) Th17 responses of TCR- δ ^{-/-} responder T cells were not enhanced by A2AR^{-/-} $\gamma\delta$ T cells. Responder T cells of TCR- δ ^{-/-} mice were supplemented by 2 % A2AR^{-/-} $\gamma\delta$ T cells (controls of adding A2AR^{+/+} $\gamma\delta$ T cells were shown in Fig. 2A), before stimulation with IRBP₁₋₂₀ and APCs, under Th17 polarized conditions. The numbers of $\alpha\beta$ TCR⁺ IL-17⁺ cells were assessed by FACS analysis after a 5-day in vitro stimulation. E) A summary data for all 3 replicates of Fig. 2D are shown. Data were analyzed using a paired *t*-test. ***p* < 0.01; ns, not significant, *n* = 6 in each group.

cells were prepared from immunized TCR- $\delta^{-/-}$ mice. The T cells were stimulated in vitro with the immunizing peptide and APCs, and the Th1 and Th17 responses were determined by the number of $\alpha\beta\text{TCR}^+\text{IFN-}\gamma^+$ and $\alpha\beta\text{TCR}^+\text{IL-17}^+$ T cells among responder T cells and the amount of IFN- γ and IL-17 produced in culture supernatants by ELISA. The results in Fig. 2A showed that the generation of $\alpha\beta\text{TCR}^+\text{IL-17}^+$ cells from wild-type (WT) B6 responders (Fig. 2A, top panels) was enhanced by the A2AR agonist CGS21680; but cells from TCR- $\delta^{-/-}$ responders (Fig. 2A, second panels) were not enhanced. Moreover, if 2 % of $\gamma\delta$ T cells were added to TCR- $\delta^{-/-}$ responder T cells before in vitro stimulation, their responses were also enhanced (Fig. 2A third panels) suggesting that $\gamma\delta$ T cells in responder T cells counteracted any inhibitory effect of adenosine, leading to greater Th17 responses. Measurement of Th1 responses under Th1-polarized conditions, however, showed $\gamma\delta$ T cells are less effective in Th1 responses (data not shown). Cytokine production tests after in vitro stimulation showed that IL-17 production was inhibited by the A2AR agonist CGS21680 in TCR- $\delta^{-/-}$ CD3 $^+$ responders (third panel

of Fig. 2C) but not in the responder T cells of B6 mouse (first panel of Fig. 2C), because the presence of $\gamma\delta$ T cells among the B6 responders offset the inhibitor effect of CGS21680 (Cui et al., 2009; Liang et al., 2013b, 2016a; Nian et al., 2010). The IFN- γ production of both responders was inhibited regardless of whether $\gamma\delta$ T cells were absent or present, indicating that Th1 inhibition by CGS21680 was not $\gamma\delta$ T cell dependent. Studies comparing the effect of agonists specific for the ARs A1R, A2AR and A2BR showed that agonists for A2BR and A1R ARs were also ineffective in inhibiting IL-17 production (Fig. 2C). Since A2ARs are not strictly expressed on $\gamma\delta$ T cells, we also compared the adenosine effect on Th17 responses of TCR- $\delta^{-/-}$ responder T cells supplemented with A2AR $^{+/+}$ (from B6 mice) or A2AR $^{-/-}$ $\gamma\delta$ T cells (from A2AR $^{-/-}$ mice). The results showed that adenosine was unable to enhance the Th17 responses supplemented with A2AR $^{-/-}$ $\gamma\delta$ T cells (Fig. 2D&E), suggesting that binding of A2ARs to $\gamma\delta$ T cells crucially involved adenosine-enhanced Th17 responses.

Fig. 3. Adenosine augmented the Th17-, but not Th1-, stimulating effect of BMDCs triggered by TLR ligand. A) LPS treated BMDCs acquired an increased stimulating effect on Th1 and Th17 responses. Responder T cells were isolated from immunized B6 mice ($n = 6$). They were stimulated with the immunizing peptide interphotoreceptor retinoid-binding protein (IRBP) $_{1-20}$ and bone marrow dendritic cells (BMDCs), under Th1 (upper panels) or Th17 (lower panels) polarized conditions. Cytokines in the supernatants were assessed by ELISA 48hr after stimulation. B) Dissociated Th1 and Th17 stimulating effect of BMDCs after dual treatment with lipopolysaccharide (LPS) and adenosine. BMDCs were treated with A2AR agonist (250 nM) or A2BR agonist (100 nM) before (left panels) or after (right panels) LPS treatment. After co-culture with responder T cells, IFN- γ and IL-17 amounts in culture supernatants were determined by ELISA. The results show that after LPS treatment, A2BR agonist treatment augmented BMDCs' Th17-stimulating effect, whereas both A2AR and A2BR agonists decreased BMDCs' Th1-stimulating effect. C). IL-12 and IL-23 production by BMDCs after treatment with LPS, with or without AR agonist. BMDCs produce IL-12 and IL-23 only after treatment with LPS. When LPS treated BMDCs were further exposed to the AR agonist, the IL-12 production was declined, whereas the IL-23 production was significantly increased. The results shown are representative of those from five experiments. Data were analyzed using a paired t -test. $^{**}p < 0.01$, $n = 6$ in each group.

3.3. Adenosine augmented the Th17, but not Th1-stimulating effect of BMDCs triggered by a TLR ligand

Dendritic cells are the principal antigen-presenting (AP) cells for initiating immune responses. Previous studies showed that TLR ligands have a profound effect on DC differentiation and maturation (Fedele et al., 2005). Since the level of extracellular adenosine increases greatly during inflammation (Eltzschig et al., 2012; Ohta and Sitkovsky, 2001; Sitkovsky et al., 2004), we questioned whether adenosine and TLR ligands have counteractive or synergistic effects on DC function and Th1 and Th17 responses. To do so, we assessed GM-CSF-cultured BMDCs for an AP effect in Th1 and Th17 responses, before and after exposure to adenosine and/or TLR ligands. The responder T cells were co-cultured with the treated BMDCs at ratio of DC:T = 1:10 in the presence of immunizing antigen and the cytokine production of responder T cells was measured. After BMDCs were treated with LPS only, both IFN- γ and IL-17 production were increased (Fig. 3A). Unexpectedly, when BMDCs were treated with LPS and A2BR agonists IFN- γ and IL-17 production changed in opposite directions; IL-17 increased whereas IFN- γ declined (Fig. 3B). Thus, the Th1 and Th17-stimulating effects of BMDCs were dissociated under a dual effect of TLR ligand and adenosine, tipping the Th1 and Th17 balance towards the latter. We then investigated whether the higher Th17-promoting effect of adenosine was associated with altered cytokine production by BMDCs after exposure to LPS and/or adenosine. Our results showed that BMDCs did not produce the

cytokines tested before the LPS exposure (not shown); treatment with either LPS (TLR4 ligand) or PAM3 (TLR2 ligand) stimulated a low production of all tested cytokines, including IL-12, IL-23, IL-1 β and IL-6. After LPS and adenosine stimulation, IL-12 production declined and IL-23 production further increased, indicating the dissociated Th1 and Th17 responses can be partly attributed to altered cytokine production of BMDCs. Given that IL-23 (Cua et al., 2003; Peng et al., 2007) and IL-1 β (Bettelli et al., 2007; Korn et al., 2007; Veldhoen et al., 2006) have a strong Th17-promoting effect, changes in patterns and amounts of cytokine production by BMDCs after adenosine presumably contributed to enhanced Th17 T cell response (Fig. 3C).

3.4. Adenosine augmented cytokine-mediated $\gamma\delta$ T cell activation

Given our previous findings that $\gamma\delta$ T cell activation was a major contributor to the regulation of Th17 responses, we questioned whether the enhancing effect of adenosine on Th17 responses was due to augmented $\gamma\delta$ T cell activation. As we have previously reported, purified $\gamma\delta$ T cells can be activated by a number of proinflammatory cytokines and that a mixture of IL-1 β , IL-7, and IL-23 has a strong stimulatory effect (Liang et al., 2013b). We used this combination and tested the activation of $\gamma\delta$ T cells by cytokines and in the absence or presence of adenosine. Responder $\gamma\delta$ T cells were prepared from immunized B6 mice using MACS sorter. Fig. 4A shows that cytokines IL-1 β , IL-7, and IL-23 were able to activate IL-17 production of $\gamma\delta$ T cells; furthermore, a

Fig. 4. Adenosine analogue 5'-N-ethyl-carboxamidoadenosine (NECA) augmented cytokine-mediated $\gamma\delta$ T cell activation. A). IL-17 production by $\gamma\delta$ T cells stimulated by cytokines is augmented by the analogue NECA. MACS purified $\gamma\delta$ T cells were isolated from immunized B6 mice. In a 96-well plate, 2×10^5 /well $\gamma\delta$ T cells were exposed to cytokines (IL-1 β +7+23), in the absence or presence of the non-selective receptor ligand NECA or A2AR specific (250 nM) agonists. IL-17 in the cultured cell supernatants were assessed by ELISA. The results shown are summarized from four separate experiments. Data were analyzed using a paired *t*-test. ***p* < 0.01, *n* = 6 in each group. B). In immunized B6 mice administered with A2BR agonist (Bay60-6538), the number of total $\gamma\delta$ T cells among CD3⁺ T cells (upper panels) and the number of activated (CD44⁺) $\gamma\delta$ T cells (lower panels) are increased. B6 mice (*n* = 6) were immunized with interphotoreceptor binding protein (IRBP)₁₋₂₀/complete Freund's adjuvant (CFA) with or without an injection of A2BR agonist (Bay60-6538, 1 mg/ml). Thirteen days post-immunization CD3⁺ cells isolated were assessed for both abundance and activation status of $\gamma\delta$ T cells. The CD3⁺ cells were gated for assessing total $\gamma\delta$ T cells (upper panels) and $\gamma\delta$ TCR⁺ T cells were gated (lower panels) when CD44⁺ $\gamma\delta$ T cells were determined. The results shown in Fig. 4B are from a single experiment. Summarized data of 4 separate experiments are shown in Fig. 3C. C). Data summarized for 4 separate experiments of Fig. 4B are plotted in Fig. 4C. Data were analyzed using a paired *t*-test. ***p* < 0.01, *n* = 6 in each group. D). $\gamma\delta$ activation is compromised if A2ARs on $\gamma\delta$ T cells are disabled. $\gamma\delta$ T cells isolated from

B6 (A2AR^{+/+}) or A2AR^{-/-} mouse (A2AR^{-/-}) were compared for response to cytokines (IL-1 β +7+23) and/or NECA. $\gamma\delta$ activation is assessed by measurement of IL-17 production. Results show that adenosine enhances A2AR^{+/+} but not A2AR^{-/-} $\gamma\delta$ activation. Data were analyzed using a paired *t*-test. ***p* < 0.01, *n* = 4 in each group.

combination of adenosine analogue NECA and the cytokine mixture greatly augmented IL-17 production by $\gamma\delta$ activation, even though neither NECA nor A2AR agonist itself appreciably stimulated $\gamma\delta$ T cells. A similar synergistic effect was seen when $\gamma\delta$ T cells were exposed to a combination of a TLR ligand and NECA (not shown). Assessment of the in vivo effect of adenosine on $\gamma\delta$ T cells showed that B6 mice that received an A2BR agonist (BAY60-6538) injection after immunization had greater numbers of $\gamma\delta$ T cells, among which the $CD44^{\text{high}} \gamma\delta\text{TCR}^+$ cells were more abundant (Fig. 4B&C). To further determine that adenosine is responsible for $\gamma\delta$ T cell activation we also compared the activation of $A2AR^{+/+}$ and $A2AR^{-/-}$ $\gamma\delta$ T cells by these cytokines. Our results showed that after stimulation with the same dose of cytokines, the activation of $A2AR^{-/-}$ $\gamma\delta$ T cells was significantly lower than that of $A2AR^{+/+}$ $\gamma\delta$ T cells because the AR $A2AR^{-/-}$ on $\gamma\delta$ T cells was disabled (Fig. 4D).

3.5. Adenosine augmented the TLR ligand activation of $\gamma\delta$ T cells by BMDCs

An alternative pathway of $\gamma\delta$ T cell activation is stimulation by DCs. To determine whether BMDCs exposed to adenosine acquired an increased ability to stimulate $\gamma\delta$ T cells, GM-CSF cultured BMDCs were co-incubated with MACS-sorted $\gamma\delta$ T cells, after treatment with LPS and/

or NECA, at a ratio of T:DC = 10:1 for two days. The activation of $\gamma\delta$ T cells was assessed by measuring IL-17 production and the numbers of $CD69^+$ $\gamma\delta$ T cells. The results showed that only the only LPS-treated BMDCs could stimulate $\gamma\delta$ T cells to produce IL-17, and BMDCs treated with LPS plus NECA acquired a greater stimulating effect (Fig. 5A). However, BMDCs treated with NECA alone were not stimulatory, indicating that the effect of adenosine on BMDCs is indirect and needed to be synergized with cytokines. Expression of $CD69$ – a cell surface marker identifying activated T cells – showed that only activated $\gamma\delta$ T cells stimulated by LPS-treated BMDCs could augment $\gamma\delta$ activation leading to augmented Th17 responses; furthermore, treatment of BMDCs with LPS plus NECA further augmented the stimulating effect of adenosine (Fig. 5B).

3.6. Inhibition of ADA by an ADA inhibitor augmented the IL17 responses

Endogenously produced adenosine is degraded by ADA. We observed that Toll ligand activated BMDCs expressed increased amounts of ADA (Fig. 6A). To determine whether Th1 and Th17 responses would be affected if ADA function is deactivated, we determined the AP function of BMDCs with or without prior treatment with EHNA – a reversible inhibitor of ADA (North and Cohen, 1978; Ullman et al., 1976). The results showed that inhibition of ADA by EHNA enhanced both the Th17

Fig. 5. Synergistic effect of 5'-N-ethylcarboxamido-adenosine (NECA) and Toll-like receptor (TLR) ligand in bone marrow dendritic cells' (BMDCs) on $\gamma\delta$ -stimulating activity. **A**). BMDC-stimulated $\gamma\delta$ T cells produced increased amounts of IL-17 if the BMDCs were pre-treated with lipopolysaccharide (LPS) and/or NECA. Data summarized for 4 separate experiments are shown. Data were analyzed using a paired *t*-test. $^{**}p < 0.01$, $n = 4$ in each group. **B&C**). BMDC-stimulated $\gamma\delta$ T cells expressed increased levels of $CD69$ after treated with LPS or LPS&NECA. The gated $\gamma\delta\text{TCR}^+$ T cells were further analyzed. One representative experiment from four separate ones is shown. $^{**}p < 0.01$. Summarized data of three separate experiments are shown in Fig. 5C.

Fig. 6. Adenosine deaminase (ADA) inhibitor erythro-9-(2-hydroxy-3-nonyl) adenine (EHNA) inhibited Th1 responses but enhanced the Th17 responses. **A).** Increased numbers of bone marrow dendritic cells (BMDCs) express ADA after Toll-like receptor (TLR) ligand stimulation. BMDCs were incubated with lipopolysaccharide (LPS) (100 ng/ml), Pam3 (2 µg/ml), or LPS + Pam3 for 48h, before they were stained with a polyclonal anti-ADA antibody. **B).** Prior treatment of BMDCs with EHNA augmented Th17 responses. Responder CD3⁺ T cells isolated from immunized B6 mice (n = 6) were stimulated, under Th17 polarized conditions, with BMDCs with (upper panels) or without (lower panels) a prior treatment with EHNA (10 µM). IL-17 Expression of CD3⁺ responder T cells were evaluated by FACS analysis after intracellular stain. Results show that inhibition of ADA by EHNA on BMDCs enhanced Th17 responses. One representative stain assay is shown. **C).** Data summarized for 3 separate experiments are shown. Data were analyzed using a paired *t*-test. ***p* < 0.01; ns, not significant, n = 4 in each group. **D)** BMDCs produced increased amounts of IL-23 after treatment with EHNA. BMDCs were treated with LPS (100 ng/ml), LPS plus 5'-N-ethylcarboxamidoadenosine (NECA) (100 nM), or LPS, NECA and EHNA (10 µM) as indicated. IL-12 and IL-23 amounts in supernatants were tested 48 h after stimulation. Data were analyzed using a paired *t*-test. ***p* < 0.01, n = 4 in each group.

and γδ T cell responses (Fig. 6B). Measurement of cytokine production of BMDCs showed that untreated BMDCs produced neither IL-23 nor IL-12, but the production of these cytokines was induced by LPS. Adenosine analogue NECA inhibited IL-12 production but enhanced IL-23 production. If the BMDCs were simultaneously treated with EHNA, the IL23 production of BMDCs further increased. These results supported the prediction that regulation of endogenously generated adenosine by ADA crucially controls adenosine levels, and thus controls Th17 responses; when ADA is disabled, adenosine will accumulate and Th17 responses will be enhanced.

4. Discussion

During stress and tissue injury, ATP is released from the intracellular compartment into the extracellular space, where it is degraded to adenosine through a cascade of enzymatic reactions. Elevated amounts of adenosine are found in ischemia, inflammation and trauma (Fredholm et al., 2001; Haskó et al., 2008; Idzko et al., 2014; Linden, 2001). Degradation of ATP to adenosine involves ectonucleotidases including CD39 (nucleoside triphosphate diphosphorylase [NTPDase]) and CD73 (5'-ectonucleotidase [Ecto5'NTase]) (Haskó et al., 2009; Yegutkin, 2008). Produced adenosine is degraded by ADA (North and Cohen, 1978; Ullman et al., 1976). Adenosine is an important regulatory molecule since it modulates a wide range of physiological functions (Fredholm et al., 2011) including the immune response (Fredholm et al., 2011; Haskó et al., 2008; Jacobson and Gao, 2006; Sauer et al., 2012) by

acting on many types of immune cells, including T cells (Jin et al., 2010; Lappas et al., 2005), macrophages/DCs (Naganuma et al., 2006; Panther et al., 2001), NK cells (Hoskin et al., 2008), neutrophils (Fredholm et al., 2001), platelets (Varani et al., 1996), and regulatory T cells (Ehrentraut et al., 2012; Naganuma et al., 2006; Zarek et al., 2008).

Four types of ARs have been defined, designated A1R, A2AR, A2BR, and A3R (Haskó et al., 2008; Ohta and Sitkovsky, 2001). The major functional receptor on T cells is A2AR (Ohta and Sitkovsky, 2001; Sitkovsky and Ohta, 2005). Previous studies have demonstrated that adenosine has a direct inhibitory effect on αβ T cells and macrophages/DCs (Erdmann et al., 2005; Huang et al., 1997; Naganuma et al., 2006; Ohta et al., 2006; Panther et al., 2001; Schnurr et al., 2005; Seigny et al., 2007). Treatment with adenosine reduced Th1 responses (Haskó et al., 2000; Panther et al., 2003), and activation of A2AR on T cells inhibited T-cell-mediated cytotoxicity, cytokine production (Ohta et al., 2009) and T-cell proliferation (Deaglio et al., 2007; Zhang et al., 2004) (Zarek et al., 2008). Regulatory T cells exert their suppressive action through the production of adenosine (Borsellino et al., 2007; Deaglio et al., 2007; Kobia et al., 2006). Adenosine inhibits IL-12 production by DCs via which Th1 responses are inhibited (Csóka et al., 2008). Indeed, A2AR^{-/-} mice developed more severe experimental autoimmune encephalomyelitis, and A2AR antagonism protects against experimental autoimmune encephalomyelitis (Mills et al., 2012); treatment with the A2AR agonist resulted in marked decreases in retinal inflammation in diabetic retinopathy (Ibrahim et al., 2011). Our recent study has tested the protective effect of ADA – an enzyme converting

adenosine into functionally inactive molecules (Mandapathil et al., 2010), in mouse EAU (Liang et al., 2016b). We found that ADA treatment suppresses EAU only when administered to recipients 8–14 days postimmunization, or shortly before EAU expression. Also, treatment of recipients with the ADA inhibitor EHNA enhances EAU development (Liang et al., 2016b). A similar “timing effect” has been found when mice are treated with the nonselective AR agonist NECA, which inhibits autoimmune responses when used at an early stage after immunization, but inhibits the response when administration of the same amount of NECA at a late stage (at 8–14 days post immunization, or prior to the disease onset (manuscript in preparation), indicating that the role of AR ligation in autoimmune pathogenesis is affected by environmental factors. Further investigations are required for better success in application.

The extrapolation of adenosine as inhibitory was mostly obtained from studies of Th1 immune responses, since Th17 responses were discovered only recently. Given that both Th1 and Th17 pathogenic T cells contribute to the pathogenesis of autoimmune diseases (Bettelli et al., 2006; Cua et al., 2003; Dong, 2006; Kolls and Linden, 2004; Langrish et al., 2005), and since extracellular concentration of ATP and its metabolites is abundant at inflammatory sites (Haskó and Cronstein, 2004; Sitkovsky et al., 2004; Wilson et al., 2011), determination of whether adenosine has a similar effect on Th1 and Th17 pathogenic T cells is important. Determination of the mechanisms by which Th17 responses differed from Th1 autoreactive T cells in response to adenosine in EAU, a well-established mouse model of uveitis, showed that the effect of adenosine on Th17 responses is enhancing, while the predominant effect of adenosine in Th1 responses is anti-inflammatory (Panther et al., 2003) (Zarek et al., 2008). As a result, adenosine tips the Th1 and Th17 balance toward the latter. The opposite effect of adenosine on Th1 and Th17 responses could certainly offset therapeutic attempts to regulate Th1 pathogenic reactions. As such, clarification of the conflicting effect of adenosine on Th1 and Th17 responses is of major importance.

The promoting effect on Th17 responses of adenosine has been also previously observed (Wilson et al., 2011); however, comparative effects on Th1 and Th17 responses have not been. Here we show that the enhancing effects of adenosine on Th17 responses is accomplished via several pathways, of which $\gamma\delta$ T cell activation is the most important. An important finding in this study is that adenosine inhibits the $\alpha\beta$ T cell responses but enhances $\gamma\delta$ T cell activation and that the enhancing effect of AR ligation on Th17 responses is modulated by $\gamma\delta$ T cells. We have previously shown that activated $\gamma\delta$ T cells acquire a greatly increased ability to enhance Th17 responses (Liang et al., 2016a; Nian et al., 2011). $\gamma\delta$ T cells can be readily activated by a number of proinflammatory cytokines, in the absence of TCR ligation. For example, a mixture of IL-1, IL-7, and IL-23 has a strong stimulatory effect on $\gamma\delta$ T cells (Liang et al., 2013a). When adenosine was added to the cytokine mixture, $\gamma\delta$ T cell-activation was significantly enhanced, even though adenosine itself does not activate $\gamma\delta$ T cells (Fig. 4). In the absence of $\gamma\delta$ T cells, adenosine is inhibitory for both Th1 and Th17 responder T cells; however, when as few as 2 % $\gamma\delta$ T cells were added to responder $\alpha\beta$ T cells, adenosine inhibition of Th17 responses was abolished but the inhibitory effect on Th1 remained. Furthermore, the Th17 enhancing effect of $\gamma\delta$ T cells was abolished when the effect of A2ARs on $\gamma\delta$ T cells was disabled (Fig. 2), suggesting that the effect of adenosine on $\gamma\delta$ T cells plays an important role in the enhanced responses of Th17. Adenosine promotes DC differentiation into a unique subset that strongly stimulates Th17, but not Th1, responses; in addition, it augments the $\gamma\delta$ -stimulating activity of BMDCs, via which Th17 responses are further enhanced (Mills et al., 2012).

In the study of DCs we found that BMDCs have Th1-stimulating activity but very little Th17 stimulating capacity before adenosine treatment. After treatment with TLR ligands, both the Th1 and the Th17 stimulating effects on BMDCs were enhanced. Unexpectedly, when BMDCs were treated with both TLR ligand and adenosine, the Th1 and Th17-stimulating effects of BMDCs were dissociated; while the Th1-

stimulating function declined, Th17 stimulation increased and tipped the Th1/Th17 balance towards the latter. Given that ATP may function as an endogenous TLR ligand (Matzinger, 2002; Ravichandran, 2010; Vitiello et al., 2012), it is likely that the balance of ATP and its degrading adenosine metabolites plays an important role in the T cell response. To investigate the function of ATP degradation and deactivation of adenosine by ADA enzyme we examined whether deactivation of ADA by a specific enzyme (EHNA) would result in excess adenosine and promote cascading Th17 responses. Our results demonstrated that ADA inhibition favors enhanced Th17 responses.

Alternative pathways may have been also involved in adenosine-induced enhancement of T cell responses. As we previously reported, activated $\gamma\delta$ T cells express greatly increased amounts of high-affinity ARs (A2ARs) (Liang et al., 2014b, 2016a), leading to altered adenosine binding by various immune cells (Liang et al., 2014b). The preferential binding of adenosine by $\gamma\delta$ T cells may lead to a re-distribution of adenosine among various immune cells, leading to diminished adenosine binding by $\alpha\beta$ T cells, for example, which will also favor augmented $\alpha\beta$ T cell responses (Liang et al., 2013b, 2014b, 2016a; Nian et al., 2011).

5. Conclusion

A better knowledge and understanding of the functional conversion of adenosine should facilitate adenosine-mediated immunotherapies. The cellular and molecular basis for enhancing and/or inhibiting the effects of ATP/adenosine remain to be further determined and the outcome of such studies should improve currently available therapies, including adenosine- and $\gamma\delta$ T cell-based immunotherapies.

Grant information

This work was supported by U.S. National Institutes of Health, National Eye Institute Grants EY0022403 and EY018827 and by a grant from Research to Prevent Blindness, NYC.

Data sharing statement

The main data supporting the findings of this study are available within the paper.

CRediT authorship contribution statement

Deming Sun: Formal analysis, Writing – original draft. **Minhee Ko:** Formal analysis, performed the experiments and analyzed data. DS and HK wrote the manuscript. All authors contributed to the article and approved the submitted version. **Hui Shao:** designed research. DS and **Henry J. Kaplan:** Writing – original draft.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Antonoli, L., Pacher, P., Vizi, E.S., Haskó, G., 2013. CD39 and CD73 in immunity and inflammation. *Trends Mol. Med.* 19, 355–367. <https://doi.org/10.1016/j.molmed.2013.03.005>.
- Beigi, R.D., Kertesz, S.B., Aquilina, G., Dubyak, G.R., 2003. Oxidized ATP (oATP) attenuates proinflammatory signaling via P2 receptor-independent mechanisms. *Br. J. Pharmacol.* 140, 507–519. <https://doi.org/10.1038/sj.bjp.0705470>.
- Bettelli, E., Carrier, Y., Gao, W., Korn, T., Strom, T.B., Oukka, M., Weiner, H.L., Kuchroo, V.K., 2006. Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature* 441, 235–238. <https://doi.org/10.1038/nature04753>.
- Bettelli, E., Oukka, M., Kuchroo, V.K., 2007. TH-17 cells in the circle of immunity and autoimmunity. *Nat. Immunol.* 8, 345–350.

- Borsellino, G., Kleinewietfeld, M., Di Mitri, D., Sternjak, A., Diamantini, A., Giometto, R., Höpner, S., Centonze, D., Bernardi, G., Dell'Acqua, M.L., et al., 2007. Expression of ectonucleotidase CD39 by Foxp3⁺ Treg cells: hydrolysis of extracellular ATP and immune suppression. *Blood* 110, 1225–1232. <https://doi.org/10.1182/blood-2006-12-064527>.
- Canaday, D.H., Beigi, R., Silver, R.F., Harding, C.V., Boom, W.H., Dubyak, G.R., 2002. ATP and control of intracellular growth of mycobacteria by T cells. *Infect. Immun.* 70, 6456–6459. <https://doi.org/10.1128/iai.70.11.6456-6459.2002>.
- Chen, J.F., Huang, Z., Ma, J., Zhu, J., Moratalla, R., Standaert, D., Moskowitz, M.A., Fink, J.S., Schwarzschild, M.A., 1999. A(2A) adenosine receptor deficiency attenuates brain injury induced by transient focal ischemia in mice. *J. Neurosci.* 19, 9192–9200.
- Cronstein, B.N., Eberle, M.A., Gruber, H.E., Levin, R.I., 1991. Methotrexate inhibits neutrophil function by stimulating adenosine release from connective tissue cells. *Proc. Natl. Acad. Sci. U.S.A.* 88, 2441–2445.
- Csóka, B., Himer, L., Selmeczy, Z., Vizi, E.S., Pacher, P., Ledent, C., Deitch, E.A., Spolarics, Z., Németh, Z.H., Haskó, G., 2008. Adenosine A2A receptor activation inhibits T helper 1 and T helper 2 cell development and effector function. *Faseb. J.* 22, 3491–3499. <https://doi.org/10.1096/fj.08-107458>.
- Cua, D.J., Sherlock, J., Chen, Y., Murphy, C.A., Joyce, B., Seymour, B., Lucian, L., To, W., Kwan, S., Churakova, T., et al., 2003. Interleukin-23 rather than interleukin-12 is the critical cytokine for autoimmune inflammation of the brain. *Nature* 421, 744–748.
- Cui, Y., Shao, H., Lan, C., Nian, H., O'Brien, R.L., Born, W.K., Kaplan, H.J., Sun, D., 2009. Major role of $\gamma\delta$ T cells in the generation of IL-17⁺ uveitogenic T cells. *J. Immunol.* 183, 560–567.
- Deaglio, S., Dwyer, K.M., Gao, W., Friedman, D., Usheva, A., Erat, A., Chen, J.F., Enjyoji, K., Linden, J., Oukka, M., et al., 2007. Adenosine generation catalyzed by CD39 and CD73 expressed on regulatory T cells mediates immune suppression. *J. Exp. Med.* 204, 1257–1265. <https://doi.org/10.1084/jem.20062512>.
- Dong, C., 2006. Diversification of T-helper-cell lineages: finding the family root of IL-17-producing cells. *Nat. Rev. Immunol.* 6, 329–334. <https://doi.org/10.1038/nri1807>.
- Ehrentauf, H., Westrich, J.A., Eltzschig, H.K., Clambey, E.T., 2012. Adora2b adenosine receptor engagement enhances regulatory T cell abundance during endotoxin-induced pulmonary inflammation. *PLoS One* 7, e32416. <https://doi.org/10.1371/journal.pone.0032416>.
- Eltzschig, H.K., Carmeliet, P., 2011. Hypoxia and inflammation. *N. Engl. J. Med.* 364, 656–665. <https://doi.org/10.1056/NEJMr0910283>.
- Eltzschig, H.K., Sitkovsky, M.V., Robson, S.C., 2012. Purinergic signaling during inflammation. *N. Engl. J. Med.* 367, 2322–2333. <https://doi.org/10.1056/NEJMr1205750>.
- Erdmann, A.A., Gao, Z.-G., Jung, U., Foley, J., Borenstein, T., Jacobson, K.A., Fowler, D. H., 2005. Activation of Th1 and Tc1 cell adenosine A2A receptors directly inhibits IL-2 secretion in vitro and IL-2-driven expansion in vivo. *Blood* 105, 4707–4714. <https://doi.org/10.1182/blood-2004-04-1407>.
- Fedeles, G., Stefanelli, P., Spensieri, F., Fazio, C., Mastrantonio, P., Ausiello, C.M., 2005. Bordetella pertussis-infected human monocyte-derived dendritic cells undergo maturation and induce Th1 polarization and interleukin-23 expression. *Infect. Immun.* 73, 1590–1597. <https://doi.org/10.1128/IAI.73.3.1590-1597.2005>.
- Feoktistov, I., Biaggioni, I., 1997. Adenosine A2B receptors. *Pharmacol. Rev.* 49, 381–402.
- Fredholm, B.B., Ap, I., Jacobson, K.A., Klotz, K.N., Linden, J., 2001. International Union of Pharmacology. XXV. Nomenclature and classification of adenosine receptors. *Pharmacol. Rev.* 53, 527–552.
- Fredholm, B.B., IJzerman, A.P., Jacobson, K.A., Linden, J., Müller, C.E., 2011. International union of basic and clinical pharmacology. LXXXI. Nomenclature and classification of adenosine receptors—an update. *Pharmacol. Rev.* 63, 1–34. <https://doi.org/10.1124/pr.110.003285>.
- Grenz, A., Homann, D., Eltzschig, H.K., 2011. Extracellular adenosine: a safety signal that dampens hypoxia-induced inflammation during ischemia. *Antioxidants Redox Signal.* 15, 2221–2234. <https://doi.org/10.1089/ars.2010.3665>.
- Hanley, P.J., Musset, B., Renigunta, V., Limberg, S.H., Dalpke, A.H., Sus, R., Heeg, K.M., Preisig-Müller, R., Daut, J., 2004. Extracellular ATP induces oscillations of intracellular Ca²⁺ and membrane potential and promotes transcription of IL-6 in macrophages. *Proc. Natl. Acad. Sci. U.S.A.* 101, 9479–9484. <https://doi.org/10.1073/pnas.0400733101>.
- Haskó, G., Cronstein, B.N., 2004. Adenosine: an endogenous regulator of innate immunity. *Trends Immunol.* 25, 33–39.
- Haskó, G., Csoka, B., Németh, Z.H., Vizi, E.S., Pacher, P., 2009. A2B adenosine receptors in immunity and inflammation. *Trends Immunol.* 30, 263–270.
- Haskó, G., Linden, J., Cronstein, B., Pacher, P., 2008. Adenosine receptors: therapeutic aspects for inflammatory and immune diseases. *Nat. Rev. Drug Discov.* 7, 759–770.
- Haskó, G.Y., Kuhel, D.G., Chen, J.F., Schwarzschild, M.A., Deitch, E.A., Mabley, J.G., Marton, A.N.I.T., Szabo, C.S.A.B., 2000. Adenosine inhibits IL-12 and TNF- α production via adenosine A2A receptor-dependent and independent mechanisms. *Faseb. J.* 14, 2065–2074.
- Hoskin, D.W., Mader, J.S., Furlong, S.J., Conrad, D.M., Blay, J., 2008. Inhibition of T cell and natural killer cell function by adenosine and its contribution to immune evasion by tumor cells (Review). *Int. J. Oncol.* 32, 527–535.
- Huang, S., Apasov, S., Koshiba, M., Sitkovsky, M., 1997. Role of A2a extracellular adenosine receptor-mediated signaling in adenosine-mediated inhibition of T-cell activation and expansion. *Blood* 90, 1600–1610.
- Ibrahim, A.S., El-shishtawy, M.M., Zhang, W., Caldwell, R.B., Liou, G.I., 2011. A2A adenosine receptor (A2AR) as a therapeutic target in diabetic retinopathy. *Am. J. Pathol.* 178, 2136–2145.
- Idzko, M., Ferrari, D., Eltzschig, H.K., 2014. Nucleotide signalling during inflammation. *Nature* 509, 310–317. <https://doi.org/10.1038/nature13085>.
- Inaba, K., Inaba, M., Romani, N., Aya, H., Deguchi, M., Ikehara, S., Muramatsu, S., Steinman, R.M., 1992. Generation of large numbers of dendritic cells from mouse bone marrow cultures supplemented with granulocyte/macrophage colony-stimulating factor. *J. Exp. Med.* 176, 1693–1702.
- Jacobson, K.A., Gao, Z.-G., 2006. Adenosine receptors as therapeutic targets. *Nat. Rev. Drug Discov.* 5, 247–264.
- Jin, D., Fan, J., Wang, L., Thompson, L.F., Liu, A., Daniel, B.J., Shin, T., Curiel, T.J., Zhang, B., 2010. CD73 on tumor cells impairs antitumor T-cell responses: a novel mechanism of tumor-induced immune suppression. *Cancer Res* 70, 2245–2255. <https://doi.org/10.1158/0008-5472.can-09-3109>.
- Junger, W.G., 2011. Immune cell regulation by autocrine purinergic signalling. *Nat. Rev. Immunol.* 11, 201–212.
- Kobie, J.J., Shah, P.R., Yang, L., Rebhahn, J.A., Fowell, D.J., Mosmann, T.R., 2006. T regulatory and primed uncommitted CD4 T cells express CD73, which suppresses effector CD4 T cells by converting 5'-adenosine monophosphate to adenosine. *J. Immunol.* 177, 6780–6786.
- Kolls, J.K., Linden, A., 2004. Interleukin-17 family members and inflammation. *Immunity* 21, 467–476.
- Korn, T., Bettelli, E., Gao, W., Awasthi, A., Jäger, A., Strom, T.B., Oukka, M., Kuchroo, V. K., 2007. IL-21 initiates an alternative pathway to induce proinflammatory TH17 cells. *Nature* 448, 484–487.
- Langrish, C.L., Chen, Y., Blumenschein, W.M., Mattson, J., Basham, B., Sedgwick, J.D., McClanahan, T., Kastelein, R.A., Cua, D.J., 2005. IL-23 drives a pathogenic T cell population that induces autoimmune inflammation. *J. Exp. Med.* 201, 233–240.
- Lappas, C.M., Sullivan, G.W., Linden, J., 2005. Adenosine A2A agonists in development for the treatment of inflammation. *Expert Opin. Invest. Drugs* 14, 797–806. <https://doi.org/10.1517/13543784.14.7.797>.
- Liang, D., Zuo, A., Shao, H., Born, W.K., O'Brien, R.L., Kaplan, H.J., Sun, D., 2013a. IL-23 receptor expression on gammadelta T cells correlates with their enhancing or suppressive effects on autoreactive T cells in experimental autoimmune uveitis. *J. Immunol.* 191, 1118–1125. <https://doi.org/10.4049/jimmunol.1300626>.
- Liang, D., Zuo, A., Shao, H., Born, W.K., O'Brien, R.L., Kaplan, H.J., Sun, D., 2013b. IL-23 receptor expression on $\gamma\delta$ T cells correlates with their enhancing or suppressive effects on autoreactive T cells in experimental autoimmune uveitis. *J. Immunol.* 191, 1118–1125. <https://doi.org/10.4049/jimmunol.1300626>.
- Liang, D., Zuo, A., Shao, H., Chen, M., Kaplan, H.J., Sun, D., 2014a. Anti-inflammatory or pro-inflammatory effect of an adenosine receptor agonist on the Th17 autoimmune response is inflammatory environment-dependent. *J. Immunol.* 193, 5498–5505. <https://doi.org/10.4049/jimmunol.1401959>.
- Liang, D., Zuo, A., Shao, H., Chen, M., Kaplan, H.J., Sun, D., 2014b. Roles of the adenosine receptor and CD73 in the regulatory effect of $\gamma\delta$ T cells. *PLoS One* 9, e108932. <https://doi.org/10.1371/journal.pone.0108932>.
- Liang, D., Zuo, A., Zhao, R., Shao, H., Born, W.K., O'Brien, R.L., Kaplan, H.J., Sun, D., 2016a. CD73 expressed on $\gamma\delta$ T cells shapes their regulatory effect in experimental autoimmune uveitis. *PLoS One* 11, e0150078. <https://doi.org/10.1371/journal.pone.0150078>.
- Liang, D., Zuo, A., Zhao, R., Shao, H., Kaplan, H.J., Sun, D., 2016b. Regulation of adenosine deaminase on induced mouse experimental autoimmune uveitis. *J. Immunol.* 196, 2646–2654. <https://doi.org/10.4049/jimmunol.1502294>.
- Linden, J., 2001. Molecular approach to adenosine receptors: receptor-mediated mechanisms of tissue protection. *Annu. Rev. Pharmacol. Toxicol.* 41, 775–787. <https://doi.org/10.1146/annurev.pharmtox.41.1.775>.
- Mahamed, D.A., Toussaint, L.E., Bynoe, M.S., 2015. CD73-Generated adenosine is critical for immune regulation during toxoplasma gondii infection. *Infect. Immun.* 83, 721–729. <https://doi.org/10.1128/iai.02536-14>.
- Mandapatil, M., Hilldorfer, B., Szczepanski, M.J., Czystowska, M., Szajnik, M., Ren, J., Lang, S., Jackson, E.K., Gorelik, E., Whiteside, T.L., 2010. Generation and accumulation of immunosuppressive adenosine by human CD4+CD25highFOXP3+ regulatory T cells. *J. Biol. Chem.* 285, 7176–7186. <https://doi.org/10.1074/jbc.M109.047423>.
- Matzinger, P., 2002. The danger model: a renewed sense of self. *Science* 296, 301–305. <https://doi.org/10.1126/science.1071059>.
- Mills, J.H., Kim, D.-G., Krenz, A., Chen, J.-F., Bynoe, M.S., 2012. A2A adenosine receptor signaling in lymphocytes and the central nervous system regulates inflammation during experimental autoimmune encephalomyelitis. *J. Immunol.* 188, 5713–5722. <https://doi.org/10.4049/jimmunol.1200545>.
- Naganuma, M., Wiznerowicz, E.B., Lappas, C.M., Linden, J., Worthington, M.T., Ernst, P. B., 2006. Cutting edge: critical role for A2A adenosine receptors in the T cell-mediated regulation of colitis. *J. Immunol.* 177, 2765–2769. <https://doi.org/10.4049/jimmunol.177.5.2765>.
- Nian, H., Shao, H., O'Brien, R.L., Born, W.K., Kaplan, H.J., Sun, D., 2011. Activated $\gamma\delta$ cells promote the activation of uveitogenic T cells and exacerbate EAU development. *Invest. Ophthalmol. Vis. Sci.* 52, 5920–5927.
- Nian, H., Shao, H., Zhang, G., Born, W.K., O'Brien, R., Kaplan, H.J., Sun, D., 2010. Regulatory effect of $\gamma\delta$ T cells on IL-17⁺ uveitogenic T cells. *Invest. Ophthalmol. Vis. Sci.* 51, 4661–4667.
- North, T.W., Cohen, S.S., 1978. Erythro-9-(2-hydroxy-3-nonyl)adenine as a specific inhibitor of herpes simplex virus replication in the presence and absence of adenosine analogues. *Proc. Natl. Acad. Sci. U.S.A.* 75, 4684–4688.
- Ohta, A., Gorelik, E., Prasad, S.J., Ronchese, F., Lukashev, D., Wong, M.K.K., Huang, X., Caldwell, S., Liu, K., Smith, P., et al., 2006. A2A adenosine receptor protects tumors from antitumor T cells. *Proc. Natl. Acad. Sci. U.S.A.* 103, 13132–13137. <https://doi.org/10.1073/pnas.0605251103>.
- Ohta, A., Ohta, A., Madasu, M., Kini, R., Subramanian, M., Goel, N., Sitkovsky, M., 2009. A2A adenosine receptor may allow expansion of T cells lacking effector functions in

- extracellular adenosine-rich microenvironments. *J. Immunol.* 183, 5487–5493. <https://doi.org/10.4049/jimmunol.0901247>.
- Ohta, A., Sitkovsky, M., 2001. Role of G-protein-coupled adenosine receptors in downregulation of inflammation and protection from tissue damage. *Nature* 414, 916–920. <https://doi.org/10.1038/414916a>.
- Panther, E., Corinti, S., Idzko, M., Herouy, Y., Napp, M., la Sala, A., Girolomoni, G., Norgauer, J., 2003. Adenosine affects expression of membrane molecules, cytokine and chemokine release, and the T-cell stimulatory capacity of human dendritic cells. *Blood* 101, 3985–3990. <https://doi.org/10.1182/blood-2002-07-2113>.
- Panther, E.L., Idzko, M., Herouy, Y., Rheinen, H., Gebrück-Haerter, P.J., Mrowietz, U., Dichmann, S., Norgauer, J., 2001. Expression and function of adenosine receptors in human dendritic cells. *Faseb. J.* 15, 1963–1970.
- Peng, Y., Han, G., Shao, H., Wang, Y., Kaplan, H.J., Sun, D., 2007. Characterization of IL-17+ interphotoreceptor retinoid-binding protein-specific T cells in experimental autoimmune uveitis. *Invest. Ophthalmol. Vis. Sci.* 48, 4153–4161.
- Rajan, A.J., Asensio, V.C., Campbell, I.L., Brosnan, C.F., 2000. Experimental autoimmune encephalomyelitis on the SJL mouse: effect of $\gamma\delta$ T cell depletion on chemokine and chemokine receptor expression in the central nervous system. *J. Immunol.* 164, 2120–2130.
- Ramlackhansingh, A.F., Bose, S.K., Ahmed, I., Turkheimer, F.E., Pavese, N., Brooks, D.J., 2011. Adenosine 2A receptor availability in dyskinetic and nondyskinetic patients with Parkinson disease. *Neurology* 76, 1811–1816. <https://doi.org/10.1212/WNL.0b013e31821ccce4>.
- Ravichandran, K.S., 2010. Find-me and eat-me signals in apoptotic cell clearance: progress and conundrums. *J. Exp. Med.* 207, 1807–1817. <https://doi.org/10.1084/jem.20101157>.
- Sauer, A.V., Brigida, I., Carriglio, N., Aiuti, A., 2012. Autoimmune dysregulation and purine metabolism in adenosine deaminase (ADA)-deficiency. *Front. Immunol.* 3, 265–275. <https://doi.org/10.3389/fimmu.2012.00265>.
- Schnurr, M., Toy, T., Shin, A., Wagner, M., Cebon, J., Maraskovsky, E., 2005. Extracellular nucleotide signaling by P2 receptors inhibits IL-12 and enhances IL-23 expression in human dendritic cells: a novel role for the cAMP pathway. *Blood* 105, 1582–1589.
- Sevigny, C.P., Li, L., Awad, A.S., Huang, L., McDuffie, M., Linden, J., Lobo, P.I., Okusa, M.D., 2007. Activation of adenosine 2A receptors attenuates allograft rejection and alloantigen recognition. *J. Immunol.* 178, 4240–4249.
- Sitkovsky, M.V., Lukashev, D., Apasov, S., Kojima, H., Koshiba, M., Caldwell, C., Ohta, A., Thiel, M., 2004. Physiological control of immune response and inflammatory tissue damage by hypoxia-inducible factors and adenosine A2A receptors. *Annu. Rev. Immunol.* 22, 657–682. <https://doi.org/10.1146/annurev.immunol.22.012703.104731>.
- Sitkovsky, M.V., Ohta, A., 2005. The 'danger' sensors that STOP the immune response: the A2 adenosine receptors? *Trends Immunol.* 26, 299–304.
- Spahn, T.W., Issazadah, S., Salvin, A.J., Weiner, H.L., 1999. Decreased severity of myelin oligodendrocyte glycoprotein peptide 33-35-induced experimental autoimmune encephalomyelitis in mice with a disrupted TCR δ chain gene. *Eur. J. Immunol.* 29, 4060–4071.
- Ullman, B., Cohen, A., Martin Jr., D.W., 1976. Characterization of a cell culture model for the study of adenosine deaminase- and purine nucleoside phosphorylase-deficient immunologic disease. *Cell* 9, 205–211. [https://doi.org/10.1016/0092-8674\(76\)90111-2](https://doi.org/10.1016/0092-8674(76)90111-2).
- Varani, K., Gessi, S., Dalpiaz, A., Borea, P.A., 1996. Pharmacological and biochemical characterization of purified A2a adenosine receptors in human platelet membranes by [³H]-CGS 21680 binding. *Br. J. Pharmacol.* 117, 1693–1701. <https://doi.org/10.1111/j.1476-5381.1996.tb15341.x>.
- Veldhoen, M., Hocking, R.J., Atkins, C.J., Locksley, R.M., Stockinger, B., 2006. TGF β in the context of an inflammatory cytokine milieu supports de novo differentiation of IL-17-producing T cells. *Immunity* 24, 179–189.
- Vitiello, L., Gorini, S., Rosano, G., la Sala, A., 2012. Immunoregulation through extracellular nucleotides. *Blood* 120, 511–518. <https://doi.org/10.1182/blood-2012-01-406496>.
- Wilkin, F., Duhant, X., Bruyns, C., Suarez-Huerta, N., Boeynaems, J.-M., Robaye, B., 2001. The P2Y₁₁ receptor mediates the ATP-induced maturation of human monocyte-derived dendritic cells. *J. Immunol.* 166, 7172–7177. <https://doi.org/10.4049/jimmunol.166.12.7172>.
- Wilson, J.M., Kurtz, C., Black, S.G., Ross, W.G., Alam, M.S., Linden, J., Ernst, P.B., 2011. The A2B adenosine receptor promotes Th17 differentiation via stimulation of dendritic cell IL-6. *J. Immunol.* 186, 6746–6752.
- Yegutkin, G.G., 2008. Nucleotide- and nucleoside-converting ectoenzymes: important modulators of purinergic signalling cascade. *Biochim. Biophys. Acta* 1783, 673–694. <https://doi.org/10.1016/j.bbamcr.2008.01.024>.
- Zarek, P.E., Huang, C.-T., Lutz, E.R., Kowalski, J., Horton, M.R., Linden, J., Drake, C.G., Powell, J.D., 2008. A2A receptor signaling promotes peripheral tolerance by inducing T-cell anergy and the generation of adaptive regulatory T cells. *Blood* 111, 251–259. <https://doi.org/10.1182/blood-2007-03-081646>.
- Zhang, X., Schwartz, J.C., Guo, X., Bhatia, S., Cao, E., Chen, L., Zhang, Z.Y., Edidin, M.A., Nathanson, S.G., Almo, S.C., 2004. Structural and functional analysis of the costimulatory receptor programmed death-1. *Immunity* 20, 337–347.
- Zocchi, C., Ongini, E., Conti, A., Monopoli, A., Negretti, A., Baraldi, P.G., Dionisotti, S., 1996. The non-xanthine heterocyclic compound SCH 58261 is a new potent and selective A2a adenosine receptor antagonist. *J. Pharmacol. Exp. Therapeut.* 276, 398–404.
- Zuo, A., Liang, D., Shao, H., Born, W.K., Kaplan, H.J., Sun, D., 2012. In vivo priming of IL-17⁺ uveitogenic T cells is enhanced by Toll ligand receptor (TLR)2 and TLR4 agonists via $\gamma\delta$ T cell activation. *Mol. Immunol.* 50, 125–133.