

Lawrence Berkeley National Laboratory

Recent Work

Title

INCLUSIVE pO PRODUCTION IN n-p INTERACTIONS AT 205 GeV/c

Permalink

<https://escholarship.org/uc/item/2nx3n53p>

Authors

Winkelman, F.C.

Bingham, H.H.

Chew, D.M.

et al.

Publication Date

1974-11-14

Submitted to
Physical Review Letters

LBL-3390
Preprint *c.d.*

INCLUSIVE ρ^0 PRODUCTION IN
 π^-p INTERACTIONS AT 205 GeV/c*

RECEIVED
LAWRENCE
RADIATION LABORATORY


JAN 6 1975

LIBRARY AND
DOCUMENTS SECTION

F. C. Winkelman, H. H. Bingham, D. M. Chew,
B. Y. Daugas, W. B. Fretter, G. Goldhaber,
W. R. Graves, A. D. Johnson, J. A. Kadyk,
L. Stutte, G. H. Trilling, and G. P. Yost,
D. Bogert, R. Hanft, F. R. Huson,
S. Kahn, D. Ljung, C. Pascaud,
S. Pruss, and W. M. Smart

November 14, 1974

Prepared for the U. S. Atomic Energy Commission
under Contract W-7405-ENG-48



LBL-3390
c.d.

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

INCLUSIVE ρ^0 PRODUCTION IN π^-p INTERACTIONS AT 205 GeV/c*

F. C. Winkelmann, H. H. Bingham, D. M. Chew,[†] B. Y. Daugeras,[‡]
 W. B. Fretter, G. Goldhaber, W. R. Graves, A. D. Johnson,
 J. A. Kadyk, L. Stutte,^{††} G. H. Trilling, and G. P. Yost

Lawrence Berkeley Laboratory and Department of Physics
 University of California, Berkeley, CA 94720

D. Bogert, R. Hanft, F. R. Huson, S. Kahn,
 D. Ljung,^{‡‡} C. Pascaud,[‡] S. Pruss, and W. M. Smart
 Fermi National Accelerator Laboratory, Batavia, IL 60510

November 14, 1974

ABSTRACT

The inclusive ρ^0 production cross section has been measured in the reaction $\pi^-p \rightarrow \pi^+\pi^-X$ at 205 GeV/c. We find $\sigma(\rho^0) = 13.5 \pm 3.4$ mb, with most of the production occurring in the central region. Assuming $\sigma(\rho^+) \approx \sigma(\rho^-) \approx \sigma(\rho^0)$, it is concluded that approximately one-third of the pions at this energy come from ρ -decay.

In contrast to inclusive single-particle production, very little detailed information now exists on inclusive production of meson resonances, particularly above 25 GeV/c.¹ The major reasons for this are the substantial difficulties of separating resonance from background because of small signal-to-noise ratios, especially when high multiplicities are involved, and the fact that many states, such as the ω and ρ^\pm , are difficult to study inclusively because π^0 detection is required. Nevertheless, resonance decays may account for a significant fraction of pion production at high energy. Furthermore, vector mesons may be an important source of the lepton production recently observed² in nucleon-nucleon collisions at Fermilab and the CERN ISR.

We describe in this paper a measurement of the inclusive ρ^0 production cross section in π^-p interactions at 205 GeV/c ($s = 385 \text{ GeV}^2$). The data were

obtained from a 48,000-picture exposure of the Fermi National Accelerator Laboratory 30-in. hydrogen bubble chamber. To improve the $\pi^+\pi^-$ mass resolution, only events occurring in a restricted fiducial volume located in the upstream half of the bubble chamber were considered for the present analysis, thus allowing at least 35 cm in the forward direction for measurement of outgoing tracks. View-to-view track matching was done by hand using bubble patterns when necessary. All of the 2- and 4-prong events, 72% of the 6- and 8-prongs, and 28% of the ≥ 10 -prongs found in the restricted fiducial volume were measured on film-plane digitizers and geometrically reconstructed with TVGP, yielding a total of 2040 inelastic interactions. Additional details on the experimental arrangement, and on the scanning and measuring of events may be found in Ref. 3.

All charged outgoing tracks were taken to be pions except those of momentum less than 1.4 GeV/c, for which pion-proton separation by means of ionization is possible. From a study of K_S^0 production⁴ we estimate that $\sim 4\%$ of the charged tracks are kaons, assuming $\sigma(K_S^0) \approx \sigma(K^+) \approx \sigma(K^-)$. The $K^{*\pm}(892) \rightarrow K_S^0 \pi^\pm$ and $\Delta^0(1232) \rightarrow p\pi^-$ signals in our data are sufficiently small that K^{*0} or Δ^0 contamination in the ρ^0 mass region (from $K^\pm \pi^\mp$ or $p\pi^-$ misidentified as $\pi^+\pi^-$) is negligible.

Figure 1 shows the inclusive mass distribution for all $\pi^+\pi^-$ combinations. This spectrum peaks at approximately 0.46 GeV and shows a distinct shoulder in the ρ^0 region. For comparison, Fig. 1 also gives the corresponding distribution for pions of like charge ($\pi^+\pi^+$ and $\pi^-\pi^-$), which is observed to be structureless.

Figure 2 shows the inclusive $\pi^+\pi^-$ mass distribution for the following three intervals of center-of-mass rapidity, y , of the $\pi^+\pi^-$ system: $-3 < y < -1$ (backward or target fragmentation region), $-1 < y < 1$ (central region), and $1 < y < 3$ (forward or beam fragmentation region).⁵ All regions show a shoulder

at the ρ^0 mass. There is also some indication of f^0 production in the central region.

The inclusive ρ^0 cross section for each region in y was obtained by fitting the $0.48 < M(\pi^+ \pi^-) < 1.12$ GeV mass range to a relativistic P-wave Breit-Wigner (with a fixed mass of 0.765 GeV and a fixed width of 0.150 GeV) plus a second-order polynomial background. A background of this form was found to provide a good description of the inclusive $\pi^\pm \pi^\pm$ distribution, Fig. 1, in the same mass range. A resolution function obtained from the measurement errors on $M(\pi^+ \pi^-)$ was folded into the Breit-Wigner line shape. The average mass resolution is 12 MeV for $-3 < y < -1$, 23 MeV for $-1 < y < 1$, and 42 MeV for $1 < y < 3$. The resulting fits, shown by the smooth curves in Fig. 2, are excellent. The fit χ^2 per degree of freedom and $\sigma(\rho^0)$ for each rapidity interval are given in Table I. The errors given for $\sigma(\rho^0)$ are statistical only.

We note that ρ^0 production is most copious in the central region, and that, at the present level of statistics, the forward and backward ρ^0 cross sections are similar. This behavior is quite different from that observed in $\pi^- p$ interactions at 8 GeV/c⁶ where the ρ^0 's come predominantly from excitation of the incoming pion and are therefore peaked strongly forward in the center of mass.

Summing over the three rapidity intervals, we obtain an overall ρ^0 cross section of 13.5 ± 3.4 mb. For comparison, the inclusive ρ^0 cross section at lower energy is 3.49 ± 0.42 mb in 24 GeV/c pp reactions⁷ and is ~ 3.2 mb in 8 GeV/c $\pi^- p$ reactions.⁶ Thus $\sigma(\rho^0)$ has increased by about a factor of four in $\pi^- p$ reactions between 8 and 205 GeV/c. Over the same momentum range, the average charged multiplicity has increased^{6,3} from 3.4 to 8.0, a factor of 2.4.

The inelastic $\pi^- p$ cross section at 205 GeV/c is 21.01 ± 0.46 mb,³ so that an average of 0.64 ± 0.16 ρ^0 are produced per inelastic interaction. The fraction of pions (charged plus neutral) coming from ρ decay can be estimated

assuming equal cross sections for ρ^+ , ρ^- , and ρ^0 production. Using $\sigma(\rho) \approx 3 \times \sigma(\rho^0) = 40.5 \pm 10.2$ mb and $\sigma(\pi) = 236 \pm 8$ mb⁸ then gives that $(34.4 \pm 8.7)\%$ of all pions are products of ρ decay. The corresponding fractions at lower energy are $\sim 15\%$ in 24 GeV/c pp reactions⁷ and $\sim 24\%$ in 8 GeV/c π^-p reactions.⁶

The cross-section ratios $\sigma(\rho^0)/\sigma(\pi^+)$ and $\sigma(\rho^0)/\sigma(\pi^-)$ are given in Table I for the backward, central, and forward regions of ρ^0 and π^\pm center-of-mass rapidity. Here, the denominator includes all π^+ or π^- ,⁹ including those from ρ^0 decay. The ρ^0/π^+ ratio in the central region is 0.26 ± 0.09 , in agreement with a value of 0.21 obtained from a general peripheral calculation,¹⁰ and with a value of 0.20 obtained from the quark model.¹¹

We have also investigated the dependence of ρ^0 production on overall charged multiplicity (n_{ch}). We find that a substantial fraction of the ρ^0 cross section is associated with high charged multiplicity; for example, $\sigma(\rho^0) = 9.1 \pm 3.4$ mb for $n_{ch} \geq 10$.

Finally, using a $\rho^0 \rightarrow \mu^+\mu^-$ branching fraction¹² of $(6.7 \pm 1.2) \times 10^{-5}$, we find a ratio

$$(\mu^- \text{ from } \rho^0 \text{ decay}) / (\text{all } \pi^-) = (1.1 \pm 0.3) \times 10^{-5},$$

integrated over rapidity and over transverse momentum up to $p_T \approx 1$ GeV/c. However, due to limited statistics, we are unable to measure the $\rho^0/(\text{all } \pi^\pm)$ ratio for higher p_T , and thus no direct comparison is possible with the data of the lepton production experiments of Ref. 2, where $\mu^\pm/(\text{all } \pi^\pm) \approx 10^{-4}$ is observed for $p_T \gtrsim 1$ GeV/c.

Conclusions.—We have measured the cross section for $\pi^-p \rightarrow \rho^0 + \text{anything}$ at 205 GeV/c to be 13.5 ± 3.4 mb and estimate that approximately one-third of the pions at this energy come from ρ decay. Thus ρ production is an important source of pions in πp collisions at high energy and can therefore be expected to have a substantial effect on single-pion inclusive distributions and two-

pion correlations.¹³ With the present statistics we are not able to explore in detail other aspects of inclusive ρ^0 production such as polarization, scaling, and ρ - π or ρ - ρ correlations.

Acknowledgments.—It is a pleasure to acknowledge the effort put forth by the 30-in. bubble chamber staff, the hadron beam group, the accelerator operations personnel, and our scanning and measuring staffs.

Table I. ρ^0 cross sections and ρ^0/π^\pm ratios in 205 GeV/c π^-p interactions.

rapidity interval	χ^2/dof	$\sigma(\rho^0)$ (mb)	ρ^0/π^+	ρ^0/π^-
$-3 < y < -1$	0.86	2.70 ± 1.25	0.17 ± 0.08	0.20 ± 0.09
$-1 < y < 1$	0.82	8.10 ± 2.81	0.26 ± 0.09	0.25 ± 0.09
$1 < y < 3$	1.08	2.66 ± 1.39	0.13 ± 0.07	0.10 ± 0.05

FOOTNOTES AND REFERENCES

*Work supported by the U. S. Atomic Energy Commission, the National Science Foundation, and the French C.N.R.S.

†On leave from Universite de Paris VI, Paris, France.

‡Present address: L.A.L., Orsay, France.

††Now at California Institute of Technology, Pasadena, CA 91109.

‡‡Now at Yale University, New Haven, Connecticut 06520.


1. The status of inclusive meson resonance production has been reviewed by F. C. Winkelmann, Lawrence Berkeley Laboratory Report No. LBL-3045 (1974), and Proceedings of the IVth International Conference on Meson Spectroscopy, Boston, Massachusetts (1974). This report also contains preliminary results of the present work.
2. J. P. Boymond et al., Phys. Rev. Letters 33, 112 (1974); J. A. Appel et al., Phys. Rev. Letters 33, 722 (1974); D. Binting et al., R. Birge et al., and F. W. Büsler et al., in Proceedings of the XVIIth International Conference on High Energy Physics, London, England (1974).
3. D. Bogert et al., Phys. Rev. Letters 31, 1271 (1973).
4. D. Bogert et al., Inclusive γ , K_S^0 , Λ^0 and $\bar{\Lambda}^0$ Production in 205 GeV/c π^-p Interactions, Fermilab Report No. NAL-Conf-74/55 (1974).
5. Only 0.3% of the observed $\pi^+\pi^-$ combinations fall outside of the $-3 < y < 3$ rapidity range.
6. T. Kitagaki et al., An Inclusive Study of ρ^0 Production in π^-p Interactions at 8 GeV/c, submitted to the IVth International Conference on Meson Spectroscopy, Boston, Massachusetts (1974). See also Ref. 1.
7. V. Blobel et al., Phys. Letters 48B, 73 (1974).

8. We have used $\sigma(\pi) = \sigma(\pi^+) + \sigma(\pi^-) + \sigma(\pi^0)$, with $\sigma(\pi^-) = \langle n_{\text{ch}} \rangle \sigma_{\text{inel}}/2 - \sigma(\text{K}^-) = 80.4 \pm 2.0$ mb; $\sigma(\pi^+) = \langle n_{\text{ch}} \rangle \sigma_{\text{inel}}/2 - \sigma(\text{K}^+) - \sigma(\text{p}) = 70.0 \pm 2.0$ mb; and $\sigma(\pi^0) \approx \sigma(\pi^- \text{p} \rightarrow \gamma + \text{anything})/2 = 85.3 \pm 7.6$ mb (Ref. 4). Here, $\langle n_{\text{ch}} \rangle =$ average charged multiplicity $= 7.99 \pm 0.06$, $\sigma_{\text{inel}} = 21.01 \pm 0.46$ mb (Ref. 3), and $\sigma(\text{p}) \approx \sigma_{\text{inel}}/2$; also, we assume $\sigma(\text{K}^+) \approx \sigma(\text{K}^-) \approx \sigma(\text{K}_S^0) = 3.49 \pm 0.60$ mb (Ref. 4) and neglect possible $\bar{\text{p}}$ contamination in the π^- sample.
9. The π^\pm cross sections used have been corrected for K^\pm contamination assuming a rapidity-independent K^\pm/π^\pm ratio of 4%. In addition, $\sigma(\pi^+)$ for $-3 < y < -1$ has been corrected for an estimated 4.1 mb of contamination due to protons (with momentum $\gtrsim 1.4$ GeV/c) which could not be identified by ionization. Proton contamination in the central and forward regions is assumed to be negligible.
10. R. M. Barnett and D. Silverman, The Relative Production of ρ and π in the Central Plateau, University of California, Irvine, Report No. 74-27 (1974).
11. V. V. Anisovich and V. M. Shekhter, Nucl. Physics B55, 455 (1973) and erratum, Nucl. Physics B63, 542 (1973).
12. V. Chaloupka et al., Review of Particle Properties, Phys. Letters 50B, No. 1 (1974).
13. For discussion of the effects of meson resonance production and decay on one- and two-particle inclusive distributions see, for example, A. Bassetto, M. Toller, and L. Sertorio, Nucl. Physics B34, 1 (1971); T. Sasaki and N. Murai, Phys. Rev. D9, 3090 (1974); J. Randa, Inclusive Rho Effects on Two-Pion Production, Texas A and M University preprint (1974); and F. Hayot, Resonances in High Multiplicity Reactions, University of Michigan Report No. UM-HE-74-36 (1974).

FIGURE CAPTIONS


Fig. 1. Inclusive $\pi^+ \pi^-$ and $\pi^+ \pi^+$ mass distributions for $\pi^- p \rightarrow \pi^+ \pi^- X$ at 205 GeV/c. The arrow indicates the expected position of the ρ^0 at $M(\pi^+ \pi^-) = 0.765$ GeV.

Fig. 2. $\pi^+ \pi^-$ mass distribution for three intervals of center-of-mass rapidity, y , of the $\pi^+ \pi^-$ system in $\pi^- p \rightarrow \pi^+ \pi^- X$ at 205 GeV/c. The arrow indicates the expected position of the ρ^0 at $M(\pi^+ \pi^-) = 0.765$ GeV.



XBL7411-4585

Fig. 1



XBL 7411 - 4584

Fig. 2

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720