Lawrence Berkeley National Laboratory

Recent Work

Title

AUTOIONIZATION STRUCTURE OF NITRIC OXIDE (NO) AT THE FIRST IONIZATION LIMIT

Permalink

<https://escholarship.org/uc/item/2p23f5pv>

Author

Miescher, E.

Publication Date

1978

Submitted to Journal of Chemical Physics

RECEIVED

LBL-7348
Preprint

LAWRENCE
BERKELEY LABORATORY

MAR 8 1978

LIBRARY AND
DOCUMENTS SECTION

AUTOIONIZATION STRUCTURE OF NITRIC OXIDE
(NO) AT THE FIRST IONIZATION LIMIT

E. Miescher and Y. T. Lee

January 1978

Prepared for the U. S. Department of Energy
under Contract W-7405-ENG-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5716*

LBL-7348

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Autoionization structure of Nitric Oxide (NO) at the
first ionization limit.

E.Miescher

Institute of Physics

University of Basel

Basel, Switzerland

and

Y.T.Lee

Department of Chemistry and Lawrence Berkeley

Laboratory

University of California

Berkeley, California 94720

Abstract

A new and more detailed interpretation is given to the fine structure in the photoionization curve of cold NO observed by Ng, Mahan and Lee ¹⁾. Resonances are assigned to autoionizing np and nd Rydberg levels which give prominent diffuse absorption bands in a spectrum recorded with the synchrotron radiation continuum. $\Delta v < -1$ transitions in the vibrational autoionization process are substantiated.

I. Introduction

In 1976 Ng, Mahan and Lee ¹⁾ have published a photoionization curve of NO in the wavelength region 1340 - 1190 Å which comprises the first four well known vibrational steps $v = 0, 1, 2, 3$ of the ion NO^+ in its ground state. In addition to the step structure that indicates direct ionization, the curve shows the fine structure, first seen by Killgoar et al. ²⁾, much more pronounced since the measurements were made on NO molecules in a supersonically cooled molecular beam. The resonance structure is obtained with the spectral resolution of about 6 meV corresponding on the wavenumber scale to 50 cm^{-1} .

In the same year Miescher ³⁾ has published an atlas of the absorption spectrum of cold NO with the resolution of about 3 cm^{-1} in the region 1420 - 1250 Å, and a paper ⁴⁾ on a high resolution (0.1 cm^{-1}) study of the absorption structure in the same region. Prior knowledge ⁵⁾ on the nl-Rydberg series of NO was thus considerably extended by these publications.

The present paper proposes a modified and more detailed interpretation of the fine structure in the photoionization efficiency curve and a new analysis of the autoionization process. In addition it reports on some new results concerning the diffuse structure in the

absorption spectrum recorded with the synchrotron radiation continuum as the background.

II. The Rydberg series of NO

Ng et al. have assigned peaks in the photoionization curve to a single Rydberg series using a quantum defect $\delta = +0.87$. There is, however, no such series observed in absorption. The 4 series observed ⁶⁾, each with its vibrational structure, are ns with $\delta_s = +1.2$, np with $\delta_{p\sigma} = +0.68$, $\delta_{p\pi} = +0.74$, nd with $\delta_{d\sigma,\pi} = -0.02$, $\delta_{d\delta} = +0.07$ and nf with a very small quantum defect. ns and nf of these series show sharp rotational levels, whereas np and nd δ show diffuseness at higher n. nd σ,π - X bands are very weak and are not observed in absorption for $n > 4$. Diffuseness of bands located below the ionization limit ($\lambda > 1340 \text{ \AA}$) must be produced by predissociation. Above the limit autoionization is possible as a additional or alternative channel.

The photoionization efficiency curve of Ng et al. is reproduced again in Fig.1. The figure shows now the positions of the diffuse np and nd δ Rydberg levels for $v = 1, 2, 3, 4$ and 5. The majority of these levels is directly observed in the absorption spectrum. Their energies have been tabulated in ref. 3, 5 and 6. Others can safely be calculated. In some measure the degree of diffuseness of the absorption bands has been characterized in ref. 3. The shapes of the diffuse bands vary considerably and are often difficult to decipher from the high resolution pho-

tographic records because of overlapping lines of nf-X bands. A photoelectrically scanned absorption spectrum of NO could recently be obtained with the smooth and steady continuum of the DORIS storage ring ⁷⁾ at Hamburg. In this experiment the absorption cell was again kept cold with liquid nitrogen and the resolution of the spectrometer, by using the 50 μ m slit, was chosen low enough (8.5 cm^{-1}) to smooth away the overlapping sharp rotational lines. This procedure did bring into prominence the diffuse bands, though it does not give the true shapes. Fig.2 is the result, covering the wave-number range of the first two vibrational steps in the ionization curve.

Only a few peaks of the low Rydberg members $n = 4, 5$ and 6 , corresponding to heads of the sharp nf-X and ns-X bands remain visible in the smeared-out spectrum, all the strong peaks are the diffuse nd δ -X and np-X bands. The latter split into p π and p σ , the two components rapidly narrowing with n . Peaks arising from the $3/2$ -component of the $^2\Pi$ ground state are small and are left unassigned. Based on the new spectrum some of the wavenumbers given in Table 1-3 are corrected compared to numbers given in ref. 3. Additional wavenumbers of diffuse bands are entered that had escaped detection by the photographic method.

III. Discussion

Fig. 1 immediately suggests an interpretation of the fine structure in the ionization curve different from the assignments made by Ng et al. The np and nd δ levels fit quite well the peaks when $\Delta v < -1$ transitions in the autoionization process are not a priori excluded as in ref. 1. The Tables 1 - 3 give the differences $i - a$ in cm^{-1} between the ionization peaks and the absorption data. They are well within the limits of the accuracy which, as mentioned above, is 50 cm^{-1} for well defined ionization peaks and to which the rotational spread of the absorption bands of about equal magnitude must be added. In cases where the latter are very diffuse or overlapping each other (6p is overlapping 9p) an additional uncertainty occurs. Many of the ionization peaks are correlated to several nllv Rydberg levels in the tables. The letter s, m or w characterizes the quality of the ionization peak and Δv indicates the vibrational change for the autoionizing transition considered. The $|\Delta v|$ in Table 2 and 3 are minima. $\Delta v < -1$ transitions are numerous and several of them are unambiguous as particularly for 5p, 6p and 7p peaks. Other than in ref. 1 all peaks are taken as corresponding to molecules in the $X^2\Pi_{\frac{1}{2}}$ component of the ground state. There is now no need to have recourse on the $^2\Pi_{\frac{1}{2}}$ component and on non-equilibrium in the cold NO. More-

*) In fact, a recent investigation on free jet expansion of NO at room temperature by Reuss and coworkers concluded that the electronic relaxation of NO from $^2\Pi_{\frac{3}{2}}$ to $^2\Pi_{\frac{1}{2}}$ is as efficient as the translational relaxation during the expansion, and most of NO molecules in free jet beam were in $^2\Pi_{\frac{1}{2}}$ state 8).

over, the evidence for assignments of the peaks to ground state doublets in ref.1 is not convincing, if one follows the individual pairs beyond $n = 9$, and if one has in view the intensity ratio of the spin-doublets.

The principal quantum number n of a level in a Rydberg series can be changed if the quantum defect δ is simultaneously changed by the same number of units. The quantum defect $\delta = 0.87$ used in ref.1 is close to 0.89 what is obtained as the mean of the quantum defects δ_p and $\delta_{d\delta}+1$ with the values quoted in section II. Since the $nd\delta$ peaks in Fig.1 carry the number n smaller by one unit than the $3/2$ peaks in ref.1 and the np peaks the same n as the $1/2$ peaks in ref.1, it is not surprising that a representation of the series with $\delta = 0.87$ was possible.

The main result of the revised interpretation of the ionization curve is that $\Delta v = -1$ autoionizing transitions are not strongly favored, and that the autoionizing Rydberg levels are specified as np and nd levels. Only the small peak at $1263 \text{ \AA}$ is left unassigned. Multiple assignments outweigh the single ones in Tables 1 to 3. A close inspection of the intensities intimates that ionization into multiple channels is real. For example, absorption to $5d\delta, v+1$ is seen in Fig.2 much stronger than the nearby absorption to $7d\delta, v$ and thus most likely it is the $5d\delta$ levels that are responsible for the height of the first peaks appearing at the steps in the ionization

curve in Fig.1. This peak is followed for $v = 1$ in Fig.1 by two barely separated smaller peaks and for $v = 2$ by a single high peak. $9p, v$ probably contributes little, but the strongly absorbing $6p\pi, v+1$ level can explain the difference in behavior because it coincides for $v = 2$ much closer with $8d\delta, v$ than for $v = 1$, see Table 1 and 2. Unobstructed $\Delta v = -2$ peaks are $7p, v$, clearly seen for $v = 2$ and weakly indicated for $v = 3$ and 4.

High absorption peaks occur in Fig.2 for the isolated $5p\pi$ levels. The peak is much broader for $v = 3$ than for $v = 4$ and furthermore $5p\sigma$ peaks are missing. Either observation is in accordance with previous reports ³⁾ where dissociative interaction with the $A'^2\Sigma^+$ continuum was considered as responsible for the anomalies. Fig.1 gives no evidence for autoionization of the $5p$ levels, clearest for the $5p\pi, 3$ level that is located right at a minimum in the ionization curve.

The theory of the autoionization process in molecules other than H_2 has been discussed comprehensively by Berry ⁹⁾ and detailed ab initio computations for the N_2 molecule have been published by Duzy and Berry ¹⁰⁾. No comparable study for the NO molecule is known. The conditions in NO are different owing to the lack of symmetry. As mentioned in section II the rotational structure of the higher np levels in NO is diffuse in opposition to the conditions for N_2 , and diffuseness in absorption can for NO always

result from predissociation because of the low dissociation energy. It has been previously emphasized ¹¹⁾ that the excited states of $^2\Delta$ symmetry are less exposed to predissociation than the $^2\Pi$ states. In emission spectra up to 70500 cm^{-1} only the $^2\Delta - X$ bands are observed above the dissociation energy. None of the bands in the much higher energy range of Fig.2 has been observed in emission, but it is reasonable to assume that the stability of the higher $nd\delta$ states in that region is still preserved and thus can give autoionization a good chance.

That ns and nf rotational levels appear in absorption sharp within the instrumental width of 0.1 cm^{-1} does not exclude by itself the possibility of autoionization of these levels. nf peaks would not be separated from the $nd\delta$ peaks in Fig.1. These levels are excluded because of their weak interaction with the core. The ns levels are excluded because of absence of peaks in Fig.1 at energies where for $n < 9$ they would be sufficiently isolated if they existed.

Acknowledgments.

E.M. would like to express his gratitude to Drs.E.E. Koch, P.Gürtler and V.Saile at the DESY for the permission and help for the use of the synchrotron continuum. Y.T.L. would like to acknowledge the support of U.S. Department of Energy for his part of work.

References

- 1) C.Y.Ng, B.H.Mahan and Y.T.Lee, J.Chem.Phys. 65, 1965 (1976).
- 2) P.C.Killgoar, G.E.Leroi, J.Berkowitz and W.A.Chupka, J.Chem.Phys. 58, 803 (1973).
- 3) E.Miescher and F.Alberti, J.of Phys.and Chem.Ref.Data, 5, 309 (1976).
- 4) E.Miescher, Can.J.of Phys. 54, 2074 (1976).
- 5) E.Miescher, J.Molec.Spectry. 20, 130 (1966).
- 6) E.Miescher and K.P.Huber, Internat.Review of Science, Physical Chemistry Series 2, Vol.3, Spectroscopy, Butterworth London 1976.
- 7) V.Saile, P.Gürtler, E.E.Koch, A.Kozevnikov, M.Skibowski and W.Steinmann, Appl.Optics, 15, 2559 (1976).
- 8) J.Reuss and S.Stolte, Univ. of Nijmegen. Private communication.
- 9) R.S.Berry, J.Chem.Phys. 45, 1228 (1966).
- 10) C.Duzy and R.S.Berry, J.Chem.Phys. 64, 2421 and 2431 (1976).
- 11) Ch.Jungen and E.Miescher, Astrophys. J. 142, 1660 (1965).

Table I. Autoionization peaks (i) in the Rydberg series converging to the limit $v=1$ of $X'\Sigma^+$ of NO^+ . Possible assignments to absorption levels (a) carrying the quantum numbers $n\ell\lambda v$. Δv = vibrational change in autoionizing transition.

		i	a	i-a	n $\ell\lambda$	v	Δv
λ		cm ⁻¹					
1336.67 s	74813	74780	+ 13	7d δ	1	-1	
		74860	- 47	5d δ	2	-2	
1334.17 s	74953	74995*	- 42	8p	1	-1	
1327.67 m	75320	75315	+ 5	8d δ	1	-1	
		75400	- 80	6p π	2	-2	
1324.17 m	75519	75475	+ 44	9p	1	-1	
		75500	+ 19	6p σ	2	-2	
		75620	-101	5p π	3	-3	
1321.17 m	75690	75690	0	9d δ	1	-1	
		75690*	0	4p π	5	-5	
		75760*	- 70	5p σ	3	-3	
		75800	-110	10p	1	-1	
1316.67 m	75949	75800	+149	10p	1	-1	
		75890	+ 59	4p σ	5	-5	
		75960	- 11	10d δ	1	-1	
		76024	- 75	11p	1	-1	
1313.17 w	76152	76150*	+ 2	11d δ	1	-1	
		76200	- 48	12p	1	-1	
		76240	- 88	6d δ	2	-2	
1310.0 m	76336	76300*	+ 36	12d δ	1	-1	
1305.3 m	76611	76590	+ 21	7p π	2	-2	
		76650	- 39	7p σ	2	-2	
1300.5 w	76894	76830	+ 64	4d δ	4	-4	

* calculated.

s, m, w = strong, medium, weak, respectively.

Table II. Same as Table I, but series converging to the limit $v=2$.

		i	a	i-a	n λ	v	Δv
$\tilde{\nu}$		cm^{-1}					
1296.17	s	77150	77085	+ 65	7d δ	2	-1
			77150	0	5d δ	3	-2
1293.67	w	77299	77290	+ 9	8p	2	-1
1287.67	s	77660	77630	+ 30	8d δ	2	-1
			77670	- 10	6p π	3	-2
1285.17	w	77811	77750	+ 61	9p	2	-1
			77760	+ 51	6p σ	3	-2
			77850	- 39	5p π	4	-3
1281.67	m	78023	78000*	+ 23	5p σ	4	-3
			78000	+ 23	9d δ	2	-1
			78090	- 67	10p	2	-1
1277.67	m	78267	78270*	- 3	10d δ	2	-1
			78330*	- 63	11p	2	-1
1275.17	m	78421	78330*	+ 91	11p	2	-1
			78460*	- 39	11d δ	2	-1
			78530	-109	6d δ	3	-2
1268.1	w	78858	78850	+ 8	7p π	3	-2
			78920*	- 62	7p σ	3	-2
1265.5	w	79020	79030*	- 10	4d δ	5	-4
1263.0	w	79177			?		

Table III. Same as Table I, but series converging to the limit $v=3$.

		i	a	i-a	nl λ	v	Δv
λ		cm ⁻¹					
1259.17	m	79417	79375*	+ 42	7d δ	3	-1
			79385	+ 32	5d δ	4	-2
1256.67	m	79575	79580	- 5	8p	3	-1
1250.67	m	79957	79910*	+ 47	8d δ	3	-1
			79930*	+ 27	6p π	4	-2
			80027*	- 70	6p σ	4	-2
1249.17	s	80053	80027*	+ 26	6p σ	4	-2
			80045*	+ 8	5p π	5	-3
			80050*	+ 3	9p	3	-1
			80225*	-172	5p σ	5	-3
1245.17	w	80310	80225*	+ 85	5p σ	5	-3
			80280*	+ 30	9d δ	3	-1
1243.67	w	80407	80280*	+127	9d δ	3	-1
			80395*	+ 12	10p	3	-1
1241.67	w	80537	80543*	- 6	10d δ	3	-1
1240.17	w	80634	80630*	+ 4	11p	3	-1
1238.67 [†]	w	80732	80735*	- 3	11d δ	3	-1
			80780*	- 48	6d δ	4	-2
1236.67	w	80862	80780*	+ 82	6d δ	4	-2
1233.8	w	81050	81100*	- 50	7p π	4	-2
1232.3	w	81149	81170*	- 21	7p σ	4	-2

[†] error in ref.1 corrected.

Figure Captions

Fig.1 The photoionization efficiency curve of NO (Ng, Mahan and Lee ¹). nd = δ -component of the d complex.

Fig.2 Absorption spectrum of cold NO, obtained with the synchrotron continuum of the DORIS storage ring (DESY, Hamburg) with the resolution of 8.5 cm^{-1} . Length of the absorption cell 52 cm, pressure 0.2 torr. nd = δ -component of the d complex.

Fig. 1

Fig. 2

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720