

# Lawrence Berkeley National Laboratory

## Lawrence Berkeley National Laboratory

### **Title**

Problem Proposed for the American Mathematical Monthly

### **Permalink**

<https://escholarship.org/uc/item/2ph3d7rj>

### **Author**

Bailey, David H.

### **Publication Date**

2009-09-01

# Problem Proposed for the *American Mathematical Monthly*

David H. Bailey\*, Jonathan M. Borwein†, Jörg Waldvogel‡

November 14, 2008

*Problem:* Define

$$P(x) := \sum_{k=1}^{\infty} \arctan \left( \frac{x-1}{(k+x+1)\sqrt{k+1} + (k+2)\sqrt{k+x}} \right). \quad (1)$$

- (a) Find explicit, finite-expression evaluations of  $P(n)$  for all integers  $n \geq 0$ .
- (b) Show  $\tau := \lim_{x \rightarrow -1^+} P(x)$  exists, and find an explicit evaluation for  $\tau$ .
- (c) Are there a more general closed forms for  $P$ , say at half-integers?

*Solution.* With the abbreviations

$$r := \sqrt{k+1}, \quad s := \sqrt{k+x}$$

the argument of  $\arctan$  in (1) becomes

$$\frac{s^2 - r^2}{(s^2 + 1)r + (r^2 + 1)s} = \frac{s - r}{r s + 1} = \frac{\frac{1}{r} - \frac{1}{s}}{1 + \frac{1}{r} \frac{1}{s}}.$$

Therefore, by using the addition theorem of the tangent function, the definition (1) may be written in the more convenient form

$$P(x) = \sum_{k=1}^{\infty} \left( \arctan \frac{1}{\sqrt{k+1}} - \arctan \frac{1}{\sqrt{k+x}} \right) \quad (2)$$

\*Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA, [dhbailey@lbl.gov](mailto:dhbailey@lbl.gov). Supported in part by the Director, Office of Computational and Technology Research, Division of Mathematical, Information and Computational Sciences, U.S. Department of Energy, under contract number DE-AC02-05CH11231.

†School of Mathematical And Physical Sciences, University of Newcastle, NSW 2308 Australia, and Faculty of Computer Science, Dalhousie University, Halifax, NS, B3H 2W5, Canada, [jonathan.borwein@newcastle.edu.au](mailto:jonathan.borwein@newcastle.edu.au), [jborwein@cs.dal.ca](mailto:jborwein@cs.dal.ca). Supported in part by NSERC and the Canada Research Chair Programme.

‡Seminar for Applied Mathematics, Department of Mathematics, ETH Zurich, 8092 Zürich, Switzerland, [waldvogel@math.ethz.ch](mailto:waldvogel@math.ethz.ch).




Figure 1: Plot of  $P(x)$

Now define

$$A(x) = \arctan \frac{1}{\sqrt{x+1}}$$

and note that a telescoping sum argument gives

$$P(x) + A(x) = P(x+1). \quad (3)$$

It is easy to see that the series defining  $P(x)$  is absolutely convergent by the Weierstrass M-test, and to verify that  $P(x)$  is increasing for  $x > -1$ , as shown in Figure 1. Thus,  $\tau$  exists.

(a). First observe that since  $P(1) = 0$ , the identity (3) establishes that  $P(0) = -A(0) = -\pi/4$ , which we had computationally observed. By iteratively applying (3) and applying induction, we establish that

$$\begin{aligned} P(2) &= \arctan \frac{1}{\sqrt{2}} \\ P(3) &= \arctan \frac{1}{\sqrt{2}} + \arctan \frac{1}{\sqrt{3}} \\ P(4) &= \arctan \frac{1}{\sqrt{2}} + \arctan \frac{1}{\sqrt{3}} + \arctan \frac{1}{2}, \end{aligned}$$

and indeed by induction we have, for all  $n \geq 2$ ,

$$P(n) = \sum_{k=2}^n \arctan \frac{1}{\sqrt{k}}.$$

(b). We computationally discovered that to 13-digit accuracy  $\tau = \lim_{x \rightarrow -1^+} P(x) = -3\pi/4$ . This can be rigorously established by noting that

$$\lim_{x \rightarrow -1^+} P(x) + \frac{\pi}{2} = \lim_{x \rightarrow -1^+} P(x) + \lim_{x \rightarrow -1^+} A(x) = \lim_{x \rightarrow -1^+} P(x+1) = P(0) = \frac{-\pi}{4}.$$