

UCSF

UC San Francisco Previously Published Works

Title

CHRNA5 Risk Variant Predicts Delayed Smoking Cessation and Earlier Lung Cancer Diagnosis—A Meta-Analysis

Permalink

<https://escholarship.org/uc/item/2pr040p5>

Journal

Journal of the National Cancer Institute, 107(5)

ISSN

0027-8874

Authors

Chen, Li-Shiun
Hung, Rayjean J
Baker, Timothy
et al.

Publication Date

2015-05-01

DOI

10.1093/jnci/djv100

Peer reviewed

REVIEW

CHRNA5 Risk Variant Predicts Delayed Smoking Cessation and Earlier Lung Cancer Diagnosis—A Meta-Analysis

Li-Shiun Chen, Rayjean J. Hung, Timothy Baker, Amy Horton, Rob Culverhouse, Nancy Saccone, Iona Cheng, Bo Deng, Younghun Han, Helen M. Hansen, Janet Horsman, Claire Kim, Sharon Lutz, Albert Rosenberger, Katja K. Aben, Angeline S. Andrew, Naomi Breslau, Shen-Chih Chang, Aida Karina Dieffenbach, Hendrik Dienemann, Brittni Frederiksen, Jiali Han, Dorothy K. Hatsukami, Eric O. Johnson, Mala Pande, Margaret R. Wrensch, John McLaughlin, Vidar Skaug, Henricus F. van der Heijden, Jason Wampfler, Angela Wenzlaff, Penella Woll, Shanbeh Zienolddiny, Heike Bickeböller, Hermann Brenner, Eric J. Duell, Aage Haugen, Joachim Heinrich, John E. Hokanson, David J. Hunter, Lambertus A. Kiemeny, Philip Lazarus, Loic Le Marchand, Geoffrey Liu, Jose Mayordomo, Angela Risch, Ann G. Schwartz, Dawn Teare, Xifeng Wu, John K. Wiencke, Ping Yang, Zuo-Feng Zhang, Margaret R. Spitz, Peter Kraft, Christopher I. Amos, Laura J. Bierut

Affiliations of authors: Department of Psychiatry, Washington University School of Medicine, St. Louis, MO (LSC, AHO, LJB); The Alvin J. Siteman Cancer Center, Washington University School of Medicine, St. Louis, MO (LSC, LJB); Lunenfeld-Tanenbaum Research Institute of Mount Sinai Hospital, Toronto, Canada (RJH, JM); Tobacco Research and Intervention, University of Wisconsin, School of Medicine, Madison, WI (TB); Department of Medicine, Washington University School of Medicine, St. Louis, MO (RC); Department of Genetics, Washington University School of Medicine, St. Louis, MO (NS); Cancer Prevention Institute of California, Fremont, CA (IC); Division of Epidemiology, Department of Health Sciences Research, Mayo Clinic College of Medicine, Rochester, MN (BD, JW, PY); Department of Community and Family Medicine, Geisel School of Medicine at Dartmouth, Hanover, NH (YH, CIA); Department of Neurological Surgery, University of California San Francisco, San Francisco, CA (HMH, MRW, JKW); Department of Oncology, University of Sheffield, Sheffield, UK (JHo, PW); Department of Epidemiology, UCLA Fielding School of Public Health, University of California Los Angeles, Los Angeles, CA (CK, ZFZ); Department of Biostatistics & Informatics, University of Colorado Anschutz Medical Campus, Denver, CO (SL); Department of Genetic Epidemiology, University of Goettingen Medical School, Goettingen, Germany (AR, HBI); Comprehensive Cancer Centre, the Netherlands, Utrecht, Netherlands & Department for Health Evidence, Radboud University Medical Center, Nijmegen, the Netherlands (KKA); Norris Cotton Cancer Center, Geisel School of Medicine at Dartmouth, Lebanon, NH (ASA); Department of Epidemiology, Michigan State University, East Lansing, MI (NB); Division of Clinical Epidemiology and Aging Research German Cancer Research Center, German Cancer research Center, Heidelberg, Germany (AKD, HBr); German Cancer Consortium, Heidelberg, Germany (AKD, HBr); Department of Thoracic Surgery, Thoraxklinik at University Hospital Heidelberg, Heidelberg, Germany (HD); Translational Lung Research Center Heidelberg, Member of the German Center for Lung, Heidelberg, Germany (HD, AR); Department of Epidemiology, University of Colorado Anschutz Medical Campus, Denver, CO (BF, JEH); Program in Molecular and Genetic Epidemiology, Harvard School of Public Health, Boston, MA (JHa, DJH, PK); Department of Epidemiology, Richard M. Fairbanks School of Public Health, Indiana University, Indianapolis, IN (JHa); Department of Psychiatry, University of Minnesota, Minneapolis, MN (DKH); Division of Health, Social and Economic Research, Research Triangle Institute, International, Research Triangle Park, NC (EOJ); Department of Epidemiology, the University of Texas MD Anderson Cancer Center, Houston, TX (MP, XW); Department of Biological and Chemical Work Environment, National Institute of Occupational Health, Oslo, Norway (VS, SZ, Aha); Department for Lung Diseases, Radboud University Medical Centre, Nijmegen, the Netherlands (HFvdH); Department of Oncology, Karmanos Cancer Institute, Wayne State University, Detroit, MI (AW, AGS); Unit of Nutrition, Environment and Cancer, Epidemiology Program, Catalan Institute of Oncology, Barcelona, Spain (EJD); Helmholtz Zentrum Munchen, Munich, Germany (JHe); Department for Health Evidence & Department of Urology, Radboud University Medical Center, Nijmegen, the Netherlands (LAK); Department of Pharmaceutical Sciences, Washington State University College of Pharmacy, Spokane, WA (PL); University of Hawaii Cancer Center, Honolulu, HI (LLM); Princess Margaret Hospital, Toronto, CA (GL); Division of Medical Oncology, Department of Medicine, University of Colorado, Aurora, CO (JM); Division of Epigenomics and Cancer Risk Factors, German Cancer Research Center, Heidelberg, Germany (AR); School of Health and Related Research, University of Sheffield, Sheffield, UK (DT); Baylor College of Medicine, Houston, TX (MRS).

Correspondence to: Li-Shiun Chen, MD, MPH, ScD, Department of Psychiatry (Box 8134), Washington University School of Medicine, 660 S. Euclid Ave., St. Louis, MO 63110 (e-mail: chenli@psychiatry.wustl.edu).

Abstract

Background: Recent meta-analyses show strong evidence of associations among genetic variants in *CHRNA5* on chromosome 15q25, smoking quantity, and lung cancer. This meta-analysis tests whether the *CHRNA5* variant rs16969968 predicts age of smoking cessation and age of lung cancer diagnosis.

Received: August 1, 2014; Revised: February 16, 2015; Accepted: March 9, 2015

© The Author 2015. Published by Oxford University Press. All rights reserved. For Permissions, please e-mail: journals.permissions@oup.com.

Methods: Meta-analyses examined associations between rs16969968, age of quitting smoking, and age of lung cancer diagnosis in 24 studies of European ancestry ($n = 29\,072$). In each dataset, we used Cox regression models to evaluate the association between rs16969968 and the two primary phenotypes (age of smoking cessation among ever smokers and age of lung cancer diagnosis among lung cancer case patients) and the secondary phenotype of smoking duration. Heterogeneity across studies was assessed with the Cochran Q test. All statistical tests were two-sided.

Results: The rs16969968 allele (A) was associated with a lower likelihood of smoking cessation (hazard ratio [HR] = 0.95, 95% confidence interval [CI] = 0.91 to 0.98, $P = .0042$), and the AA genotype was associated with a four-year delay in median age of quitting compared with the GG genotype. Among smokers with lung cancer diagnoses, the rs16969968 genotype (AA) was associated with a four-year earlier median age of diagnosis compared with the low-risk genotype (GG) (HR = 1.08, 95% CI = 1.04 to 1.12, $P = 1.1 \times 10^{-3}$).

Conclusion: These data support the clinical significance of the *CHRNA5* variant rs16969968. It predicts delayed smoking cessation and an earlier age of lung cancer diagnosis in this meta-analysis. Given the existing evidence that this *CHRNA5* variant predicts favorable response to cessation pharmacotherapy, these findings underscore the potential clinical and public health importance of rs16969968 in *CHRNA5* in relation to smoking cessation success and lung cancer risk.

Tobacco smoking is one of the modifiable risk factors that most greatly contribute to cancer and many other severe medical illnesses. Clear evidence demonstrates that genetic variation in the $\alpha 5$ nicotinic cholinergic receptor subunit gene (*CHRNA5*) is associated with heaviness of smoking and nicotine dependence (1,2). A series of meta-analyses based on tens of thousands of subjects of European ancestry confirmed the association with smoking quantity, defined by number of cigarettes smoked per day (3–6), with the most robust associations reported for rs16969968 and rs1051730, two highly correlated genetic variants (5). The nonsynonymous variant rs16969968 results in an amino acid change in the $\alpha 5$ nicotinic cholinergic receptor subunit, which alters nicotinic receptor conductance in vitro (7,8). These *CHRNA5* variants also are consistently associated with smoking-related disorders including lung cancer and chronic obstructive pulmonary disease (COPD) (9–11).

However, an important gap in our understanding of variation in *CHRNA5* is its contribution to smoking cessation. For example, these *CHRNA5* variants are less consistently associated with smoking cessation outcomes. Some studies show an association between the *CHRNA5* region and successful smoking cessation (12–18), finding that the same risk variants that contribute to smoking quantity and nicotine dependence also predict smoking cessation. Other studies, however, fail to confirm this association (19–21). A genome-wide association study (GWAS) of three treatment cohorts did not identify any nicotinic receptor genes as predictors of prospectively measured smoking cessation (21). Another large genome-wide association meta-analysis that strongly supported the association between 15q25.1 and smoking quantity ($P < 10^{-32}$) reported a below GWAS-level association between *CHRNA5* and smoking cessation ($P < 10^{-4}$), defined as the contrast between current vs former smoking (5).

This gap in knowledge of the relation between *CHRNA5*, smoking cessation, and smoking-related disorders inspired this current research. We meta-analyzed results from 24 studies of European ancestry in collaboration with the International Lung Cancer Consortium (ILCCO) and Genetic Associations and Mechanisms in Oncology (GAME-ON) Consortia (Transdisciplinary Research in Cancer of the Lung [TRICL] and Discovery, Biology, and Risk of Inherited Variants in Breast Cancer [DRIVE]). We explored two linked aims: 1) Does variation in *CHRNA5* affect the age that smokers quit smoking successfully? 2) Does variation in *CHRNA5* affect the onset of lung cancer?

Methods

Samples

In order to examine *CHRNA5*, age of smoking cessation, and age of lung cancer diagnoses, we invited studies in the ILCCO, the GAME-ON Consortia TRICL and DRIVE, and the Genetic Epidemiology of COPD (COPDgene) Study. In addition, we included the Collaborative Genetic Study of Nicotine Dependence (COGEN) and three studies from the dbGap database: Atherosclerosis risk in Communities (ARIC), Multi-Ethnic Study of Atherosclerosis (MESA), and Study of Addiction: Genetics and Environment (SAGE). Results from 24 datasets ($n = 29\,072$ unrelated smokers of European ancestry) contributed to the meta-analyses. Informed consent was obtained from participants, and all studies received approval from the appropriate institutional review board. Standardized quality control measures were applied to all studies. Of these datasets, seven were studies ascertained by conditions other than smoking-related diseases, and 17 were studies of smoking-related diseases: lung cancer (15 studies), COPD (one study), and coronary heart disease (one study). These different study designs allowed us to conduct stratified meta-analysis in patients ascertained for smoking-related diseases and patients ascertained for other reasons. To be included in analyses, each patient was required to have reported smoking cigarettes in his/her lifetime. Additional details for each dataset are provided in [Supplementary Table 1](#) and [Supplementary Methods](#) (available online). [Supplementary Table 2](#) shows the sample size, demographics, cigarettes per day, percentage of smokers who have quit smoking, and minor allele frequency for rs16969968 for each dataset.

Phenotypes

The primary phenotypes were: 1) time (in years) from birth to age of smoking cessation, defined as self-reported age of stopping smoking among ever smokers, and 2) time (in years) from birth to age of lung cancer diagnosis among lung cancer cases. A secondary phenotype defined as the duration from age of smoking onset (not available in all studies) to age of smoking cessation was examined in a subset of studies. Smoking quantity when subjects smoked regularly was assessed with cigarettes smoked per day (CPD), defined as a four-level ordered trait ($CPD \leq 10$; $11 \leq CPD \leq 20$; $21 \leq CPD \leq 30$; $CPD \geq 31$, coded as 0, 1, 2, 3, respectively).

Variants for Analyses

Because of its biological significance, we targeted the *CHRNA5* variant rs16969968 for association testing. The variant rs16969968 was available in all datasets except ARIC, MESA, and the German study. To examine ARIC and MESA, we imputed the rs16969968 genotype based on 1000 Genomes (<http://www.1000genomes.org/>) for analyses. In addition, we used proxy variants (rs951266, $r^2 = 0.98$, in ARIC, and rs9788721, $r^2 = 0.83$, in MESA, r^2 estimated based on 1000 Genomes for the EUR samples including CEU, GBR, TSI, IBS, FIN) (22,23) for analyses and reached similar results. In the German study, imputed genotype was not available and we used a proxy variant (rs1051730, $r^2 = 0.99$ estimated based on 1000 Genomes for the EUR samples) for analyses. In addition, we conducted the meta-analyses with and without the German Studies and reached similar results.

Statistical Analyses and Meta-Analyses

In each dataset, we used Cox regression models to evaluate the association between rs16969968 and the two primary phenotypes (age of smoking cessation among ever smokers, and age of lung cancer diagnosis among lung cancer case patients), and the secondary phenotype of smoking duration. The assumption of proportionality was examined by several different approaches (plotting the log(-log(survival)) against log(time) for different groups, ASSESS statistics, and testing the interactions between the covariate and time). There was a trend of decreasing genetic effects in the very old age group when the cancer incidence decreases. When truncating after age 82.5 (1% of the sample), the proportionality assumption was verified while all results were similar with/without the truncation. Age as a continuous variable and sex were included as covariates. Additional covariates included cigarettes smoked per day (a continuous variable). Genotypes were coded additively as the number of minor alleles (A), where the reference allele was defined as the major allele (G) in the European ancestry population (24). Consistency of allelic coding was confirmed by comparing allele labels and frequencies across datasets (Supplementary Table 2, available online).

Standardized scripts were used for analyses of all participating datasets at each individual research center. For the ILCCO studies, analyses were performed at Washington University as individual level data was provided for data pooling. Results were returned to Washington University for quality checks and meta-analyses. Individual SNP analyses were performed using SAS (SAS Institute, Cary, NC).

We used PLINK to perform variance-based meta-analyses (25). The R package, rmeta, was used to confirm results and generate meta-analysis plots (26). The heterogeneity across studies was assessed with the Cochran Q test and reported for each analysis. There was evidence of heterogeneity across datasets for analyses of age of smoking cessation, but not for the analyses of age at lung cancer diagnosis. This heterogeneity may be because of the varying study designs and ascertainment strategies. We report results from random effects models for all meta-analyses. All statistical tests were two-sided,

Results

The meta-analysis included studies not ascertained for smoking-related-disorders, case patients with smoking-related disorders (lung cancer, COPD, coronary heart disease), and matched control patients without these disorders. This design allowed us to conduct meta-analyses of the genetic associations of

rs16969968, stratified by ascertainment of individuals with and without smoking-related disorders that may alter smoking cessation rates. This comparison is of interest because illness could affect smoking behaviors. First, we confirmed a consistent association between *CHRNA5* rs16969968 and heaviness of smoking in subjects ascertained with smoking-related disease (meta-analysis $\beta = 0.10$, 95% confidence interval [CI] = 0.07 to 0.13, $P = 3.5 \times 10^{-10}$) (Supplementary Figure 1, available online), as well as in subjects ascertained without such disease (meta-analysis $\beta = 0.10$, 95% CI = 0.06 to 0.14, $P = 1.1 \times 10^{-6}$).

The *CHRNA5* rs16969968 Risk Allele and Delayed Smoking Cessation in Smokers Without Smoking-Related Disorders

Among smokers in the studies not ascertained for smoking-related illnesses, the rs16969968 risk allele (A) was associated with a lower likelihood of smoking cessation, adjusted for age and sex. The meta-analysis across seven datasets resulted in a hazard ratio (HR) of 0.95 (95% CI = 0.91 to 0.98, $P = .0042$, $P_{\text{heterogeneity}} = .32$). In addition, we examined this association in subjects who were control patients in studies ascertained for smoking-related illnesses. We saw a similar association in control participants in the studies of lung cancer with significant heterogeneity across these studies (HR = 0.94, 95% CI = 0.87 to 1.02, $P = .15$, $P_{\text{heterogeneity}} = .027$). Non-statistically significant associations were found for control patients in the COPD and coronary heart disease studies. Figure 1 provides a forest plot summary for rs16969968 and age of quitting. The summary median age of smoking cessation was 56 years for individuals with the high-risk genotype (AA), which was a four-year delay compared with individuals with the low-risk genotype (GG), who had a summary median quit age of 52 years. At age 50 years, 42% of individuals with the AA genotype, compared with 48% of individuals with the GG genotype, successfully quit smoking. In addition, rs16969968 was associated with delayed cessation ($n = 13\ 258$, HR = 0.90, 95% CI = 0.88 to 0.93, $P = 1.93 \times 10^{-5}$) in a pooled analysis using available individual-level data adjusted for age, sex, and study. These analyses were repeated with smoking duration as the outcome, and similar results were seen.

To further understand the genetic association between rs16969968 and smoking cessation, we meta-analyzed results from models adjusting for smoking quantity (the four-level cigarettes smoked per day variable). In smokers not ascertained for smoking-related disorders, smoking a higher number of cigarettes per day was associated with a later quitting age (meta-analysis random effect HR = 0.87, 95% CI = 0.76 to 0.99, $P = .037$) with significant heterogeneity ($I^2 = 38.3$, $P_{\text{heterogeneity}} = .033$). Rs16969968 showed a weakened, non-statistically significant association with age of quitting after adjusting for smoking quantity (HR = 0.98, 95% CI = 0.96 to 1.01, $P = .25$).

The *CHRNA5* rs16969968 Risk Allele Predicts Earlier Age of Lung Cancer Diagnosis

The rs16969968 risk allele (A) was associated with earlier age of lung cancer diagnosis, adjusted for sex. The overall meta-analysis across all 15 datasets of lung cancer case patients gave a hazard ratio of 1.08 (95% CI = 1.04 to 1.12, $P = 1.1 \times 10^{-5}$, $P_{\text{heterogeneity}} = .53$) (Figure 2). The summary median age at lung cancer diagnosis was 61 years for individuals with the high-risk genotype (AA), which was a four-year earlier age at onset compared with individuals with the low-risk genotype (GG), who had a summary median




Figure 1. Effect of *CHRNA5* rs16969968 on delayed smoking cessation among subjects without smoking-related disorders. Smoking-related disorders were lung cancer, chronic obstructive pulmonary disorder, and coronary heart disease. All models adjusted for age, sex, and rs16969968. Smoking effect models are shown, $P_{\text{heterogeneity}} = .32$ for group A, .027 for group B, and .0058 when all groups combined. * Individual-level data for these marked studies were available and used for the pooled analysis ($n = 13\ 285$). All statistical tests were two-sided. CHD = coronary heart disease; COPD = chronic obstructive pulmonary disorder.

age of diagnosis of 65 years. In addition, in a pooled analysis using available individual level data adjusted for sex and study, rs16969968 was associated with earlier diagnoses of lung cancer ($n = 7074$, HR = 1.11, 95% CI = 1.07 to 1.15, $P = 2.49 \times 10^{-9}$).

The genetic effect of *CHRNA5* on age at onset of lung cancer remained after adjusting for smoking quantity (four levels of cigarettes smoked per day). Smoking a higher number of cigarettes per day was associated with an earlier age of lung cancer diagnosis (meta-analysis HR = 1.07, 95% CI = 1.02 to 1.13, $P = .0072$), and rs16969968 (A) remained a statistically significant predictor of an earlier age of lung cancer diagnosis (HR = 1.08, 95% CI = 1.04 to 1.11, $P = 3.1 \times 10^{-5}$).

Predictors for Smoking Cessation Change Among Patients With Smoking-Related Disorders

In contrast to what was found in patients not ascertained for smoking-related disease, neither rs16969968 nor heaviness of smoking predicted smoking cessation in patients ascertained with lung cancer, COPD, or coronary heart disease (rs16969968 association with age of quitting: meta-analysis HR = 0.98, 95% CI = 0.91 to 1.05, $P = .50$) (Supplementary Figure 2, available online) (cigarettes per day association with age of quitting: meta-analysis HR = 0.995, 95% CI = 0.90 to 1.11, $P = .93$) (Supplementary Figure 3, available online). In our secondary analyses using meta-regression, we found a statistically significant interaction

in that smoking heaviness was associated with delayed quitting only in patients not ascertained for smoking-related disease, but no association was seen in patients with smoking-related disease ($F = 148.23$, $df = 1$, $P < 1.0 \times 10^{-5}$). The difference in association of *CHRNA5* and delayed quitting between patients with and without smoking-related disease was non-statistically significant ($F = 1.70$, $df = 1$, $P = .21$).

Discussion

This research shows a complex relation amongst smoking quantity, smoking cessation, genetic variation, and lung cancer risk. We found that genetic variation in *CHRNA5*, which strongly predicts risk for nicotine dependence (3–6), lung cancer (9,11), and COPD (10), also predicts delayed smoking cessation. The high-risk rs16969968 A allele is common in individuals of European ancestry (42.5%, dbSNP CEU). Having the high-risk genotype (AA, 18% of European ancestry) was associated with a four-year delay in median age of quitting smoking, compared with the low-risk genotype (GG, 33% of European ancestry) amongst individuals without diagnosed smoking-related disease. This delay in quitting results in a longer exposure to carcinogens in cigarettes, which predicts an increased risk for cancer, pulmonary, and cardiovascular disorders (27,28).

Importantly, the *CHRNA5* variant rs16969968 also predicts an earlier diagnosis of lung cancer, manifested as a median age of




Figure 2. Effect of *CHRNA5* rs16969968 on diagnosis age among smokers with lung cancer. All models adjusted for sex and rs16969968. Random effect model is shown, $P_{\text{heterogeneity}} = .53$. * Individual-level data for all studies are available and used for the pooled analysis ($n = 7074$). All statistical tests were two-sided.

lung cancer diagnosis that is four years earlier in smokers with the high-risk genotype (AA) (61 years of age), compared with those with the low-risk genotype (GG) (65 years of age), an association that remains after adjusting for smoking quantity. Lung cancer is the most common cancer in the United States and worldwide and responsible for 13% of all lung cancer incidence and 27% of all cancer deaths (29,30). The survival rate with a lung cancer diagnosis is low—50% die within a year of diagnosis and the five-year survival rate is 16.6% (31). Most cases (90%) of lung cancer are attributable to smoking. The acceleration of lung cancer diagnosis in those with AA genotypes (18% of ever smokers with lung cancer) by four years compared with those with GG genotypes (33%) can be clinically significant given the high mortality following a diagnosis of lung cancer. This finding is consistent with the previous reports of *CHRNA5* and earlier lung cancer diagnosis in a single study (by five years) (32), and a meta-analysis of five studies (by 1.1 years, $n = 3898$) by Lips et al. (33). The present study is a larger meta-analysis (15 studies, $n = 7074$), one overlapping study (Toronto study, $n = 270$) with the report by Lips et al.

The mechanisms underlying *CHRNA5*, age of smoking cessation, and age of lung cancer diagnosis are complex, and we attempted to summarize our results in Figure 3. The mechanisms through which *CHRNA5* accelerates the age of onset of lung cancer likely involve multiple pathways, including an increased number of cigarettes smoked (3–6), deeper inhalation of cigarettes leading to higher carcinogen exposure (34,35), and a delay in smoking cessation. This combination of risk factors could mediate the genetic effect, so that this one genetic variant, rs16969968, is then associated with earlier diagnosis of lung cancer by four years.

Evidence of the association between *CHRNA5* and smoking cessation has primarily been reported in smoking cessation trials (12,14,15,17,18) and less commonly in general population studies (5,18). However, our findings indicate that association between *CHRNA5* and smoking cessation may be generalizable to the general population of smokers. Examining age of quitting smoking, we confirmed the genetic association between *CHRNA5* and smoking cessation in smokers not ascertained with smoking-related disorders, a step forward in exploring the public health significance of this genetic marker.

These findings can help us understand discrepancies in existing reports of this genetic locus and smoking cessation. Some studies show a relation between *CHRNA5* and smoking cessation (12–16), whereas other studies do not (19–21). We hypothesize that the expression of this genetic risk on cessation varies with several factors, such as developing a smoking-related disease, use of cessation pharmacotherapy (18), and environmental influences on smoking cessation such as partner smoking (36). The effect of this genetic locus is seen most clearly in a sample where there is no smoking-related disorder, pharmacotherapy use is rare, and strong environmental push for smoking cessation is present. In addition, we posit that *CHRNA5* risk alleles dispose individuals to early development of smoking-related disease and that disease development encourages earlier quitting than would otherwise occur. This is why this genetic risk is more likely to be positively related to quitting age amongst individuals without smoking-related disease vs those with such disease; amongst the latter group, the effect of *CHRNA5* to delay quitting is offset by its effect of accelerating disease development, which in turn promotes quitting. Similarly, use of cessation pharmacotherapy may reduce the relation of *CHRNA5* with quitting age because pharmacotherapy appears to blunt this genetic risk (18). The heterogeneity among studies in this meta-analysis indicates the genetic risk association varies with potential moderators for the genetic risk such as the impact of disease/symptom, cessation, medication use, and environmental risks.

These results should be interpreted in the context of multiple limitations. First, smoking cessation in the samples was self-reported and not assessed using biochemical confirmation. However, research shows that self-report is a valid indicator of current smoking, especially when there are no strong incentives to deceive (37). Second, we examined age of smoking cessation as a primary phenotype. There is no evidence of association between rs16969968 and age of onset of smoking, and there is limited variation in age of onset for smoking (38,39). We also analyzed duration of smoking as the secondary outcome for a subset of studies with available information and found similar results. Third, we could not test the effect of cessation treatment on age of quitting, as this information was not available. We assume the use of medication is not common because the majority of smokers quit without medication in the general population (40). Fourth,




Figure 3. Summary diagram of *CHRNA5* rs16969968 genotype, age of smoking cessation, and age of lung cancer diagnosis. * Associations are supported by existing evidence (4–7,19,23). ^a No longer statistically significant after adjusting for heavy smoking. ^b Remains statistically significant after adjusting for heavy smoking.

this work analyzed only one genetic variant, and it is clear that multiple genes or variants contribute to smoking cessation (41). Fifth, we found the genetic effect of *CHRNA5* on earlier age of lung cancer diagnosis after adjusting for the effect of heavy smoking while there is potential residual confounding because of the imprecise nature of self-reported smoking quantity information. Finally, this study included only subjects of European ancestry.

Despite these limitations, this meta-analysis clarifies the clinical and public health significance of a robust common genetic marker identified in genetic studies of nicotine dependence. We demonstrate that the *CHRNA5* variant rs16969968 is a marker of delayed smoking cessation. The median age of smoking cessation for those with the high-risk variants (AA) at rs16969968 is 56 years vs 52 years for those with the low-risk variants (GG). Similarly, those with the high-risk variants have a four-year-earlier age of lung cancer diagnosis (61 years) compared with those with the low-risk variants (65 years). Recent research has revealed that two prominent smoking-related risk factors for lung cancer are heavy smoking quantity and late age of quitting smoking (28), and rs16969968 influences both risk factors. Moreover, the current research suggests the boundary conditions affecting the relation between rs16969968 and cessation, suggesting that development of smoking-related disease obscures the manifestation of this genetic risk. Early detection of lung cancer is important (42), and these results suggest the potential use of a common genetic variant such as *CHRNA5* rs16969968 as a susceptibility marker for earlier lung cancer diagnosis.

Funding

International Lung Cancer Consortium (ILCCO): The data management of ILCCO is supported by Cancer Care Ontario Research Chair awarded to R. Hung and National Institutes of Health (NIH) U19 CA148127.

Transdisciplinary Research in Cancer of the Lung (TRICL): The TRICL study was supported by a grant from the National Institutes of Health (U19CA148127). The Toronto study was also supported by Canadian Cancer Society Research Institute (no. 020214), Ontario Institute of Cancer and Cancer Care Ontario Chair Award to RH. Sample collection for the Heidelberg

lung cancer study was in part supported by grant 70–2919 from the Deutsche Krebshilfe. The work was additionally supported by a Helmholtz-DAAD fellowship (A/07/97379) to MNT and by the National Institute of Health (USA) (U19CA148127). The KORA Surveys were financed by the GSF, which is funded by the German Federal Ministry of Education, Science, Research and Technology and the State of Bavaria. The LUNG Cancer in the Young study (LUCY) was funded in part by the National Genome Research Network (NGFN), the DFG (BI 576/2-1; BI 576/2-2), the Helmholtzgemeinschaft (HGF) and the Federal office for Radiation Protection (BfS: STSch4454). Genotyping was performed in the Genome Analysis Center (GAC) of the Helmholtz Zentrum Muenchen. Funding for the MD Anderson Cancer Study was provided by NIH grants (P50 CA70907, R01CA121197, R01 CA127219, U19CA148127, R01CA55769) and CPRIT grant (RP100443). The Harvard Lung Cancer Study was funded by National Institutes of Health (CA074386, CA092824, CA090578). The National Key Basic Research Program Grant (2011CB503805) and the National Natural Science Foundation of China (30730080, 30972541, 30901233, and 30872178).

MD Anderson: National Cancer Institute (NCI) /NIH K07CA160753.

Collaborative Genetic Study of Nicotine Dependence (COGEN): The COGEN contribution was supported by the National Cancer Institute (P01 CA089392), the National Human Genome Research Institute (NHGRI; U01 HG04422-01), and the National Institute on Drug Abuse (NIDA; K02 DA021237). COGEN genotyping was in part performed under NIDA Contract HHSN271200477471C; phenotypic and genotypic data are stored in the NIDA Center for Genetic Studies (NCGS) at <http://zork.wustl.edu/> under NIDA Contract HHSN271200477451C (PIs J Tischfield and J. Rice); genotyping services were also provided by the Center for Inherited Disease Research (CIDR), which is fully funded through a federal contract from the National Institutes of Health to Johns Hopkins University, contract number HHSN268200782096. Financial support for this study was also provided by the National Institutes of Health under award number R01DA036583. Dr. LiShiun Chen was supported by KL2RR024994, K08DA030398 and by the National Institute On Drug Abuse of the National Institutes of Health under award number R01DA038076. The content is solely

the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Dr. Bierut and Dr. Chen were supported by The Alvin J. Siteman Cancer Center (SCC) at Washington University School of Medicine (WUSM) and Barnes-Jewish Hospital (BJH) cancer center support grant P30 CA091842.

COPDgene: The project described was supported by Award Number U01HL089897 and Award Number U01HL089856 from the National Heart, Lung, and Blood Institute. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Heart, Lung, and Blood Institute or the National Institutes of Health.

The COPDgene project is also supported by the COPD Foundation through contributions made to an Industry Advisory Board comprised of AstraZeneca, Boehringer Ingelheim, Novartis, Pfizer, Siemens, and Sunovion.

German Lung Cancer Study: The HGF GWA study was made of three independent investigations: the LUCY-study (Helmholtz ZentrumMuenchen, PIs Wichmann, Bickeböller), the Heidelberg lung cancer case-control study (German Cancer Research Center, PI Risch), and KORA surveys (“Cooperative health research in the Region of Augsburg,” PI Wichmann). LUCY was partly funded by the National Genome Research Network (NGFN), the DFG (BI 576/2-1; BI 576/2-2), the Helmholtzgemeinschaft (HGF), and the Federal office for Radiation Protection (BfS: STSch4454). The Heidelberg sample collection was partly supported by the Deutsche Krebshilfe (70–2919). The KORA platform predominantly is financed by public funds allocated to the Helmholtz ZentrumMünchen by the Federal Ministry of Education and Research and the State of Bavaria.

Germany Saarland ESTHER Study: This study was supported in part by the Baden-Württemberg State Ministry of Science, Research and Arts, by the German Federal Ministry of Education and Research.

Greater Toronto Area Lung Cancer study: This study is supported by Canadian Cancer Society Research Institute (no. 020214).

Hawaii Case-Control Study: This project was supported by Grant ROI-CA-55874 and Contract NOI-CN-05223 from the United States National Cancer Institute and by Grant EDT-78 from the American Cancer Society.

Karmanos Cancer Institute, Wayne State University: The Karmanos Cancer Institute contribution was supported by the National Cancer Institute (R01 CA60691, R01 CA14176, N01 PC35145, P30 CA022453).

Mayo Study, Mayo Clinic, College of Medicine: The contribution was supported by NIH-R01-80127/84354 and Mayo Foundation Fund.

NELCS: The New England Lung Cancer Study was funded by Grant Number P2ORR018787 from the National Center for Research Resources (NCRR), a component of the National Institutes of Health.

Netherlands Radboudumc: The study was funded by an investment grant of Radboud University Medical Center.

Northern California Lung Cancer Study, University of California San Francisco (UCSF): The UCSF contribution was supported by the National Institute of Environmental Health Sciences (R01 ES06717) and the National Cancer Institute (R01 CA 52689 to MW).

Norway: Funding by the Norwegian Cancer Society and the Norwegian Research Council.

Nurses Health Study (NHS-BrCa, NHS-T2D, NHS-CHD): This work was supported by 2P01CA87969-11 (the Nurses Health Study), U19 CA148065-01, and U19 CA148127 (Discovery Biology and Risk of Inherited Variants in Breast Cancer or DRIVE and

Transdisciplinary Research in Cancer of the Lung or TRICL, part of the NCI’s GAME-ON initiative).

The Resource for Lung Cancer in North Trent (ReSoLUCENT) / Sheffield: This study has been supported by Sheffield ECMC (Experimental Cancer Medicine Centre) and Weston Park Hospital Cancer Charity.

Tampa, FL: This study was supported by Public Health Service grants P01-CA68384 and R01-DE13158 from the National Institutes of Health.

UCLA: This study is supported by the Alper Research Program for Environmental Genomics of the UCLA Jonsson Comprehensive Cancer Center and the National Institute of Health (CA90833, DA11386, CA77954, CA09142, CA96134, and ES 011667).

ARIC: The Atherosclerosis Risk in Communities Study is carried out as a collaborative study supported by National Heart, Lung, and Blood Institute contracts (HHSN268201100005C, HHSN268201100006C, HHSN268201100007C, HHSN268201100008C, HHSN268201100009C, HHSN268201100010C, HHSN268201100011C, and HHSN268201100012C). The authors thank the staff and participants of the ARIC study for their important contributions. Funding for GENEVA was provided by National Human Genome Research Institute grant U01HG004402 (E. Boerwinkle).

MESA: MESA and the MESA SHARE project are conducted and supported by the National Heart, Lung, and Blood Institute (NHLBI) in collaboration with MESA investigators. Support for MESA is provided by contracts N01-HC- 95159, N01-HC-95160, N01-HC-95161, N01-HC-95162, N01-HC-95163, N01-HC-95164, N01-HC-95165, N01-HC- 95166, N01-HC-95167, N01-HC-95168, N01-HC-95169, and CTSA UL1-RR-024156. “Funding for SHARe genotyping was provided by NHLBI Contract N02-HL-64278. Genotyping was performed at Affymetrix (Santa Clara, CA) and the Broad Institute of Harvard and MIT (Boston, MA) using the Affymetric Genome-Wide Human SNP Array 6.0.”

Notes

The authors thank the NCI GAME-ON Consortium to support this collaborative project.

International Lung Cancer Consortium (ILCCO): We thank Li Rita Zhang for the data management of ILCCO Data Repository.

Collaborative Genetic Study of Nicotine Dependence (COGEND): We thank the subjects who participated in this study. We wish to thank Hilary Davidson, Sherri Fisher, Tracey Richmond, Nina Smock, and Heidi Kromrei for administrative support and Louis Fox for data analysis. The Collaborative Genetic Study of Nicotine Dependence (COGEND) study investigators are: Laura Bierut (PI), Michael Brent, Naomi Breslau, Robert Culverhouse, Alison Goate, Richard Gruzca, Dorothy Hatsukami, Anthony Hinrichs, Eric Johnson, Sharon Murphy, John Rice, Nancy Saccone, Scott Saccone, Joe Henry Steinbach, Jerry Stitzel, and Jen-Chyong Wang.

Collaborative Study on the Genetics of Alcoholism (COGA): The Collaborative Study on the Genetics of Alcoholism (COGA), Principal Investigators B. Porjesz, V. Hesselbrock, H. Edenberg, L. Bierut, includes ten different centers: University of Connecticut (V. Hesselbrock); Indiana University (H. J. Edenberg, J. Nurnberger Jr., T. Foroud), University of Iowa (S. Kuperman, J. Kramer), SUNY Downstate (B. Porjesz), Washington University in St. Louis (L. Bierut, A. Goate, J. Rice, K. Bucholz), University of California at San Diego (M. Schuckit), Rutgers University (J. Tischfield), Southwest Foundation (L. Almasy), Howard University (R. Taylor), and Virginia Commonwealth University (D. Dick). Other COGA collaborators include: L. Bauer (University of Connecticut); D. Koller, S. O’Connor, L. Wetherill, X. Xuei (Indiana University); Grace Chan

(University of Iowa); N. Manz, M. Rangaswamy (SUNY Downstate); A. Hinrichs, J. Rohrbaugh, J. C. Wang (Washington University in St. Louis); A. Brooks (Rutgers University); and F. Aliev (Virginia Commonwealth University). A. Parsian and M. Reilly are the National Institute on Alcohol Abuse and Alcoholism (NIAAA) Staff Collaborators. We continue to be inspired by our memories of Henri Begleiter and Theodore Reich, founding PI and Co-PI of COGA, and also owe a debt of gratitude to other past organizers of COGA, including Ting-Kai Li, currently a consultant with COGA, P. Michael Conneally, Raymond Crowe, and Wendy Reich for their critical contributions. This national collaborative study is supported by NIH Grant U10AA008401 from the NIAAA and the National Institute on Drug Abuse (NIDA). Funding support for GWAS genotyping, which was performed at the Johns Hopkins University Center for Inherited Disease Research, was provided by the National Institute on Alcohol Abuse and Alcoholism, the NIH GEI (U01HG004438), and the NIH contract "High throughput genotyping for studying the genetic contributions to human disease" (HHSN268200782096C). The authors thank Kim Doheny and Elizabeth Pugh from CIDR and Justin Paschall from the NCBI dbGaP staff for valuable assistance with genotyping and quality control in developing the dataset available at dbGaP.

COPDgene Investigators: *Administrative Core*: James Crapo, MD (PI), Edwin Silverman, MD, PhD (PI), Barry Make, MD, Elizabeth Regan, MD, PhD, Rochelle Lantz, Lori Stepp, Sandra Melanson. *Genetic Analysis Core*: Terri Beaty, PhD, Barbara Klanderma, PhD, Nan Laird, PhD, Christoph Lange, PhD, Michael Cho, MD, Stephanie Santorico, PhD, John Hokanson, MPH, PhD, Dawn DeMeo, MD, MPH, Nadia Hansel, MD, MPH, Craig Hersh, MD, MPH, Peter Castaldi, MD, MSc, Merry-Lynn McDonald, PhD, Jing Zhou, MD, PhD, Manuel Mattheissen, MD, PhD, Emily Wan, MD, Megan Hardin, MD, Jacqueline Hetmanski, MS, Margaret Parker, MS, Tanda Murray, MS. *Imaging Core*: David Lynch, MB, Joyce Schroeder, MD, John Newell Jr., MD, John Reilly, MD, Harvey Coxson, PhD, Philip Judy, PhD, Eric Hoffman, PhD, George Washko, MD, Raul San Jose Estepar, PhD, James Ross, MSc, Mustafa Al Qaisi, MD, Jordan Zach, Alex Kluiber, Jered Sieren, Tanya Mann, Deanna Richert, Alexander McKenzie, Jaleh Akhavan, Douglas Stinson. *PFT QA Core*, LDS Hospital, Salt Lake City, UT: Robert Jensen, PhD. *Biological Repository*, Johns Hopkins University, Baltimore, MD: Homayoon Farzadegan, PhD, Stacey Meyerer, Shivam Chandan, Samantha Bragan. *Data Coordinating Center and Biostatistics*, National Jewish Health, Denver, CO: Douglas Everett, PhD, Andre Williams, PhD, Carla Wilson, MS, Anna Forssen, MS, Amber Powell, Joe Piccoli. *Epidemiology Core*, University of Colorado School of Public Health, Denver, CO: John Hokanson, MPH, PhD, Marci Sontag, PhD, Jennifer Black-Shinn, MPH, Gregory Kinney, MPH, PhD, Sharon Lutz, MPH, PhD.

COPDgene Clinical Centers: Ann Arbor VA: Jeffrey Curtis, MD, Ella Kazerooni, MD. Baylor College of Medicine, Houston, TX: Nicola Hanania, MD, MS, Philip Alapat, MD, Venkata Bandi, MD, Kalpalatha Guntupalli, MD, Elizabeth Guy, MD, Antara Mallampalli, MD, Charles Trinh, MD, Mustafa Atik, MD, Hasan Al-Azzawi, MD, Marc Willis, DO, Susan Pinero, MD, Linda Fahr, MD, Arun Nachiappan, MD, Collin Bray, MD, L. Alexander Frigini, MD, Carlos Farinas, MD, David Katz, MD, Jose Freytes, MD, Anne Marie Marciel, MD. Brigham and Women's Hospital, Boston, MA: Dawn DeMeo, MD, MPH, Craig Hersh, MD, MPH, George Washko, MD, Francine Jacobson, MD, MPH, Hiroto Hatabu, MD, PhD, Peter Clarke, MD, Ritu Gill, MD, Andetta Hunsaker, MD, Beatrice Trotman-Dickenson, MBBS, Rachna Madan, MD. Columbia University, New York, NY: R. Graham Barr, MD, DrPH, Byron Thomashow, MD, John Austin, MD, Belinda D'Souza, MD. Duke University Medical Center, Durham, NC: Neil MacIntyre Jr., MD, Lacey Washington, MD, H. Page McAdams, MD. Fallon Clinic, Worcester, MA: Richard Rosiello, MD, Timothy

Bresnahan, MD, Joseph Bradley, MD, Sharon Kuong, MD, Steven Meller, MD, Suzanne Roland, MD. Health Partners Research Foundation, Minneapolis, MN: Charlene McEvoy, MD, MPH, Joseph Tashjian, MD. Johns Hopkins University, Baltimore, MD: Robert Wise, MD, Nadia Hansel, MD, MPH, Robert Brown, MD, Gregory Diette, MD, Karen Horton, MD. Los Angeles Biomedical Research Institute at Harbor UCLA Medical Center, Los Angeles, CA: Richard Casaburi, MD, Janos Porszasz, MD, PhD, Hans Fischer, MD, PhD, Matt Budoff, MD, Mehdi Rambod, MD. Michael E. DeBakey VAMC, Houston, TX: Amir Sharafkhaneh, MD, Charles Trinh, MD, Hirani Kamal, MD, Moham Darvishi, MD, Marc Willis, DO, Susan Pinero, MD, Linda Fahr, MD, Arun Nachiappan, MD, Collin Bray, MD, L. Alexander Frigini, MD, Carlos Farinas, MD, David Katz, MD, Jose Freytes, MD, Anne Marie Marciel, MD. Minneapolis VA: Dennis Niewoehner, MD, Quentin Anderson, MD, Kathryn Rice, MD, Audrey Caine, MD. Morehouse School of Medicine, Atlanta, GA: Marilyn Foreman, MD, MS, Gloria Westney, MD, MS, Eugene Berkowitz, MD, PhD. National Jewish Health, Denver, CO: Russell Bowler, MD, PhD, David Lynch, MB, Joyce Schroeder, MD, Valerie Hale, MD, John Armstrong, II, MD, Debra Dyer, MD, Jonathan Chung, MD, Christian Cox, MD. Temple University, Philadelphia, PA: Gerard Criner, MD, Victor Kim, MD, Nathaniel Marchetti, DO, Aditi Satti, MD, A. James Mamary, MD, Robert Steiner, MD, Chandra Dass, MD, Libby Cone, MD. University of Alabama, Birmingham, AL: William Bailey, MD, Mark Dransfield, MD, Michael Wells, MD, Surya Bhatt, MD, Hrudaya Nath, MD, Satinder Singh, MD. University of California, San Diego, CA: Joe Ramsdell, MD, Paul Friedman, MD. University of Iowa, Iowa City, IA: Alejandro Cornellas, MD, John Newell Jr., MD, Edwin JR van Beek, MD, PhD. University of Michigan, Ann Arbor, MI: Fernando Martinez, MD, MeiLan Han, MD, Ella Kazerooni, MD. University of Minnesota, Minneapolis, MN: Christine Wendt, MD, Tadashi Allen, MD. University of Pittsburgh, Pittsburgh, PA: Frank Sciruba, MD, Joel Weissfeld, MD, MPH, Carl Fuhrman, MD, Jessica Bon, MD, Danielle Hooper, MD. University of Texas Health Science Center at San Antonio, San Antonio, TX: Antonio Anzueto, MD, Sandra Adams, MD, Carlos Orozco, MD, Mario Ruiz, MD, Amy Mumbower, MD, Ariel Kruger, MD, Carlos Restrepo, MD, Michael Lane, MD.

German Lung Cancer Study: We thank the subjects who participated in any contributing study, the LUCY-consortium (detail in Sauter et al. 2008), WiebkeSauter, Martina Mittelstrass, Vera Zietemann and Prof. H. E. Wichmann, the KORA study group, H. Dienemann, P. Drings, and the staff at the Thoraxklinik Heidelberg.

Greater Toronto Area Study: We thank Dr. Geoffrey Liu's laboratory for the genotyping work related to this manuscript. Nurses Health Study (NHS-BrCa, NHS-T2D, NHS-CHD): We thank Drs. Frank Hu and Eric Rimm, principal investigators of the NHS Type 2 Diabetes and Coronary Heart Disease genome-wide association studies, respectively. We also thank Ms. Hongyan Huang for data analysis.

Tampa, FL: We thank the patients, physicians, and staff at the H. Lee Moffitt Cancer Center for their participation.

UCLA: We thank the study participants for their dedication and commitment and the Alper Research Program for Environmental Genomics of the UCLA Jonsson Comprehensive Cancer Center for their support.

Conflicts of interest: UCSF: none. COGEND: LB Bierut is listed as an inventor on Issued US Patent 8,080,371, "Markers for Addiction" covering the use of certain SNPs in determining the diagnosis, prognosis, and treatment of addiction. COPDgene: No conflicts of interest from the above three authors. German Lung Cancer Study: H. Bickeböller, A. Rosenberger, A. Risch, and J. Heinrich declare no conflict of interest. Germany Saarland Esther Study. We (HB and AKD) declare no conflicts of interest.

Hawaii Study: No conflict of interest to disclose. Netherland Radboudumc: No conflicts of interest. Norway: No conflict of interest to disclose. ReSoLuCENT: No potential conflicts of interest were disclosed. Tampa, FL: No conflicts of interest to disclose. UCLA: No potential conflicts of interest were disclosed. UGSP has no conflicts of interest to disclose.

References

- Bierut LJ, Madden PA, Breslau N, et al. Novel genes identified in a high-density genome wide association study for nicotine dependence. *Hum Mol Genet.* 2007;16(1):24–35.
- Bierut LJ, Stitzel JA, Wang JC, et al. Nicotine Dependence and the $\alpha 5$ - $\alpha 3$ - $\beta 4$ Nicotinic Receptor gene cluster: Variants in the Nicotinic Receptors Alter the Risk for Nicotine Dependence. *American J Psychiatry.* 2008;9(165):1163–1171.
- Liu JZ, Tozzi F, Waterworth DM, et al. Meta-analysis and imputation refines the association of 15q25 with smoking quantity. *Nat Genet.* 2010;42(5):436–440.
- Saccone NL, Culverhouse RC, Schwantes-An TH, et al. Multiple independent loci at chromosome 15q25.1 affect smoking quantity: a meta-analysis and comparison with lung cancer and COPD. *PLoS Genet.* 2010;6(8):1–16.
- TAG. Genome-wide meta-analyses identify multiple loci associated with smoking behavior. *Nat Genet.* 2010;42(5):441–447.
- Thorgeirsson TE, Gudbjartsson DF, Surakka I, et al. Sequence variants at CHRN3-CHRNA6 and CYP2A6 affect smoking behavior. *Nat Genet.* 2010;42(5):448–453.
- Bierut LJ, Stitzel JA, Wang JC, et al. Variants in nicotinic receptors and risk for nicotine dependence. *Am J Psychiatry.* 2008;165(9):1163–1171.
- Kuryatov A, Berrettini W, Lindstrom J. Acetylcholine receptor (AChR) $\alpha 5$ subunit variant associated with risk for nicotine dependence and lung cancer reduces ($\alpha 4\beta 2$) $\alpha 5$ AChR function. *Mol Pharmacol.* 2011;79(1):119–125.
- Amos CI, Wu X, Broderick P, et al. Genome-wide association scan of tag SNPs identifies a susceptibility locus for lung cancer at 15q25.1. *Nat Genet.* 2008;40(5):616–622.
- Pillai SG, Ge D, Zhu G, et al. A genome-wide association study in chronic obstructive pulmonary disease (COPD): identification of two major susceptibility loci. *PLoS Genet.* 2009;5(3):e1000421.
- Thorgeirsson TE, Geller F, Sulem P, et al. A variant associated with nicotine dependence, lung cancer and peripheral arterial disease. *Nature.* 2008;452(3):638–642.
- Baker TB, Weiss RB, Bolt D, et al. Human neuronal acetylcholine receptor A5-A3-B4 haplotypes are associated with multiple nicotine dependence phenotypes. *Nicotine Tob Res.* 2009;11(7):785–796.
- Freathy RM, Ring SM, Shields B, et al. A common genetic variant in the 15q24 nicotinic acetylcholine receptor gene cluster (CHRNA5-CHRNA3-CHRNA4) is associated with a reduced ability of women to quit smoking in pregnancy. *Hum Mol Genet.* 2009;18(15):2922–2927.
- Munafò MR, Johnstone EC, Walther D, Uhl GR, Murphy MF, Aveyard P. CHRNA3 rs1051730 Genotype and Short-Term Smoking Cessation. *Nicotine Tob Res.* 2011;13(10):982–988.
- Sarginson JE, Killen JD, Lazzeroni LC, et al. Markers in the 15q24 nicotinic receptor subunit gene cluster (CHRNA5-A3-B4) predict severity of nicotine addiction and response to smoking cessation therapy. *Am J Med Genet B Neuropsychiatr Genet.* 2011;156B(3):275–284.
- King DP, Paciga S, Pickering E, et al. Smoking cessation pharmacogenetics: analysis of varenicline and bupropion in placebo-controlled clinical trials. *Neuropsychopharmacology.* 2012;37(3):641–650.
- Bergen AW, Javitt HS, Krasnow R, et al. Nicotinic acetylcholine receptor variation and response to smoking cessation therapies. *Pharmacogenet Genomics.* 2013;23(2):94–103.
- Chen LS, Baker TB, Piper ME, et al. Interplay of genetic risk factors (CHRNA5-CHRNA3-CHRNA4) and cessation treatments in smoking cessation success. *Am J Psychiatry.* 2012;169(7):735–742.
- Breitling LP, Twardella D, Hoffmann MM, Witt SH, Treutlein J, Brenner H. Prospective association of dopamine-related polymorphisms with smoking cessation in general care. *Pharmacogenomics.* 2010;11(4):527–536.
- Conti DV, Lee W, Li D, et al. Nicotinic acetylcholine receptor beta2 subunit gene implicated in a systems-based candidate gene study of smoking cessation. *Hum Mol Genet.* 2008;17(18):2834–2848.
- Uhl GR, Liu QR, Drong T, et al. Molecular genetics of successful smoking cessation: convergent genome-wide association study results. *Arch Gen Psychiatry.* 2008;65(6):683–693.
- Altshuler DM, Gibbs RA, Peltonen L, et al. Integrating common and rare genetic variation in diverse human populations. *Nature.* 2010;467(7311):52–58.
- Durbin RM, Abecasis GR, Altshuler DL, et al. A map of human genome variation from population-scale sequencing. *Nature.* 2010;467(7319):1061–1073.
- Sherry ST, Ward MH, Kholodov M, et al. dbSNP: the NCBI database of genetic variation. *Nucleic Acids Res.* 2001;29(1):308–311.
- Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet.* 2007;81(3):559–575.
- Lumley T. *rmeta: Meta-analysis, R package version 2.15.* In.
- Jha P, Ramasundarahettige C, Landsman V, et al. 21st-century hazards of smoking and benefits of cessation in the United States. *N Engl J Med.* 2013;368(4):341–350.
- Thun MJ, Carter BD, Feskanich D, et al. 50-year trends in smoking-related mortality in the United States. *N Engl J Med.* 2013;368(4):351–364.
- Centers for Disease Control and Prevention, National Center for Health Statistics. CDC Wonder Online Database, compiled from Compressed Mortality File 1999–2010 Series 20 No. 2P; 2013.
- American Cancer Society. Cancer Facts and Figures; 2014.
- Howlander N, Noone AM, Krapcho M, et al. Seer Cancer Statistics Review, 1975–2010. In: InstituteNC, editor. Bethesda, MD; 2013.
- Spitz MR, Amos CI, Dong Q, Lin J, Wu X. The CHRNA5-A3 region on chromosome 15q24-25.1 is a risk factor both for nicotine dependence and for lung cancer. *J Natl Cancer Inst.* 2008;100(21):1552–1556.
- Lips EH, Gaborieau V, McKay JD, et al. Association between a 15q25 gene variant, smoking quantity and tobacco-related cancers among 17 000 individuals. *Int J Epidemiol.* 2010;39(2):563–577.
- Bloom AJ, Hartz SM, Baker TB, et al. Beyond cigarettes-per-day: A genome-wide association study of the biomarker carbon monoxide. *Ann Am Thorac Soc.* 2014;11(7):1003–1010.
- Le Marchand L, Derby KS, Murphy SE, et al. Smokers with the CHRNA lung cancer-associated variants are exposed to higher levels of nicotine equivalents and a carcinogenic tobacco-specific nitrosamine. *Cancer Res.* 2008;68(22):9137–9140.
- Chen LS, Baker TB, Piper ME, et al. Interplay of genetic risk (CHRNA5) and environmental risk (partner smoking) on cigarette smoking reduction. *Drug Alcohol Depend.* 2014;143:36–43.
- Subcommittee on Biochemical Verification SRNT. Biochemical verification of tobacco use and cessation. *Nicotine Tob Res.* 2002;4(2):149–159.
- Stephens SH, Hartz SM, Hoft NR, et al. Distinct loci in the CHRNA5/CHRNA3/CHRNA4 gene cluster are associated with onset of regular smoking. *Genet Epidemiol.* 2013;37(8):846–859.
- Hartz SM, Short SE, Saccone NL, et al. Increased genetic vulnerability to smoking at CHRNA5 in early-onset smokers. *Arch Gen Psychiatry.* 2012;69(8):854–860.
- Hughes JR, Keely J, Naud S. Shape of the relapse curve and long-term abstinence among untreated smokers. *Addiction.* 2004;99(1):29–38.
- Uhl GR, Walther D, Musci R, et al. Smoking quit success genotype score predicts quit success and distinct patterns of developmental involvement with common addictive substances. *Mol Psychiatry.* 2014;19(1):50–54.
- U.S. Preventive Services Task Force Recommendation Statement. Screening for Lung Cancer: U.S. Preventive Services Task Force Recommendation Statement; 2013.