

Lawrence Berkeley National Laboratory

LBL Publications

Title

The Equation of Motion of an Electron

Permalink

<https://escholarship.org/uc/item/2q45q1t9>

Authors

Kim, Kwang-Je

Sessler, Andrew M

Publication Date

1998-10-01



ERNEST ORLANDO LAWRENCE BERKELEY NATIONAL LABORATORY


The Equation of Motion of an Electron

Kwang-Je Kim and Andrew M. Sessler

**Accelerator and Fusion
Research Division**

October 1998

Presented at the
*8th Workshop on Advanced
Accelerator Concepts*,
Baltimore, MD,
July 5–11, 1998,
and to be published in
the Proceedings



REFERENCE COPY
Does Not
Circulate

Lawrence Berkeley National Laboratory
Bldg. 50 Library - Ref.

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

LBNL-42186
ANL/APS/LS-270
CBP Note-267
and LS-270

The Equation of Motion of an Electron

Kwang-Je Kim and Andrew M. Sessler

Accelerator and Fusion Research Division
Ernest Orlando Lawrence Berkeley National Laboratory
University of California
Berkeley, California 94720

October 1998

This work was supported by the Director Office of Science, Office of High Energy and Nuclear Physics, Division of High Energy Physics, of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098, and the Director, Office of Science, Office of Basic Energy Sciences, under Contract No. W-31-109-ENG-38.

Presented at the 8th Workshop on Advanced Acceleration Concepts
July 5-11, 1998, Baltimore, MD.

The Equation of Motion of an Electron^{*}

Kwang-Je Kim[†] and Andrew M. Sessler[‡]

[†]*Argonne National Laboratory, Argonne, IL 60439 and The University of Chicago, Chicago, IL 60637*

[‡]*Lawrence Berkeley National Laboratory, Berkeley, CA 94720*

ABSTRACT. We review the current status of understanding of the equation of motion of an electron. Classically, a consistent, linearized theory exists for an electron of finite extent, as long as the size of the electron is larger than the classical electron radius. Nonrelativistic quantum mechanics seems to offer a fine theory even in the point particle limit. Although there is as yet no convincing calculation, it is probable that a quantum electrodynamic result will be at least as well-behaved as is the nonrelativistic quantum mechanical results.

INTRODUCTION

For almost 100 years there has been consideration of the proper equation of motion of an electron. Many fine physicists, beginning with Abraham (1) and Lorentz (2), have worked on this subject, and there are hundreds of papers in the literature. In this paper we review the present state of understanding, with some historical background, giving the major contributions through the years.

In contrast with what most physicists believe, it is seen that the linearized classical theory, when it is applied in the appropriate regime (nonquantum), is in fine shape: it is finite, has no contradiction with relativity, has no run-away solutions, and has no a-causal behavior.

The Abraham-Lorentz equation for a point electron involving a third derivative in time suffers from two major problems: contradiction with relativity and run-away or a-causal behavior. The work of Poincaré— (3) and Dirac (4) solves the problem with relativity. The run-away problem is solved by going to an extended model of an electron described by a difference-differential equation. The equation for the

^{*} Work supported by the U.S. Department of Energy, Office of Basic Energy Sciences, under Contract No. W-31-109-ENG-38 and Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics, under Contract No. DE-AC03-76SF00098.

nonrelativistic case was derived by Sommerfeld (5) and Page (6) and was generalized to the relativistic case by Caldirola (7). The extended model is finite and causal if the electron size a is larger than the classical electron radius $r_e = e^2/mc^2 = 2.7 \times 10^{-13}$ cm.

Classical theory is clearly not appropriate for examining behavior at a distance less than the Compton wavelength of an electron, $\lambda = \hbar/mc = 4 \times 10^{-11}$ cm. The work by Moniz and Sharp (8) indicates that in nonrelativistic quantum mechanics an electron behaves as an extended particle with a size of the Compton wavelength: The equation of motion is finite and causal even in the point particle limit as long as the Compton wavelength is larger than the classical electron radius. Furthermore, the mass correction δm is not only finite but actually vanishes in quantum theory. We present a new quantum mechanical derivation of this interesting result (9). In quantum electrodynamic (QED) analysis, recent work by Low (10) shows that the electron motion is finite in perturbation theory in $\alpha = e^2/\hbar c$. However, a proper QED analysis has not yet been obtained (and maybe, those do *not* even exist).

There are several excellent text books (11-13) and review articles (14-16) on the classical electron theory.

NOTATION

Notations adopted in this paper are as follows:

m_0		bare mass
m		observed mass ($m = m_0 + \delta m$)
c		speed of light
r_e		classical electron radius (e^2/mc^2)
a		radius of extended electron
t_e		$(2/3) r_e/c$
λ		Compton wavelength
\mathbf{x}, \mathbf{x}'		space coordinate (3 vectors)
$\mathbf{y}(t)$		electron coordinate as a function of time (3 vectors)
$\dot{\mathbf{y}}$	=	$d\mathbf{y}/dt, \ddot{\mathbf{y}} = d^2\mathbf{y}/dt^2$, etc.
\mathbf{u}	=	$d\mathbf{y}/dt$ (three-velocity), $\dot{\mathbf{u}} = d\mathbf{u}/dt$, etc.
β	=	$ \mathbf{u} /c, \gamma = 1/\sqrt{1-\beta^2}$
ds	=	cdt/γ (relativistic invariant)
u^α	=	$dx^\alpha/ds = \gamma(1, \mathbf{u}/c)$ (contra-variant four vector)
u_α	=	$\gamma(1, -\mathbf{u}/c)$ (covariant four vector)
\dot{u}^α	=	$du^\alpha/ds, \ddot{u}^\alpha = d^2u^\alpha/d^2s$, etc.

ABRAHAM-LORENTZ AND OTHER CLASSICAL EQUATIONS OF MOTION

We start by deriving various forms of equation of motion of an electron in nonrelativistic classical mechanics. Let the electron trajectory be $y(t)$. The electron is really a charge distribution centered at $y(t)$ represented by

$$ef(\mathbf{x} - \mathbf{y}(t)), \quad (1)$$

where e is the electron charge, and $f(\mathbf{x})$ is a spherically symmetric function, normalized so that

$$\int f(\mathbf{x})d^3x = 1. \quad (2)$$

The nonrelativistic equation of motion is determined by the Lorentz force law:

$$m_0\ddot{\mathbf{y}}(t) = \mathbf{F}_{\text{ext}} + e \int f(\mathbf{x} - \mathbf{y}(t))(\mathbf{E}(\mathbf{x}) + \frac{1}{c}\dot{\mathbf{y}}(t) \times \mathbf{B}(\mathbf{x}))d^3x, \quad (3)$$

where m_0 is the bare mass, the dot represents the time derivative, \mathbf{F}_{ext} is the external force, and \mathbf{E} and \mathbf{B} are, respectively, the electric and magnet field in Gaussian units. The second term in the above equation is the electromagnetic self force.

It is convenient to work with the potentials A and ϕ in Coulomb gauge.¹ The scalar potential is given by

$$\phi(\mathbf{x}, t) = e \int \frac{f(\mathbf{x}' - \mathbf{y}(t))}{|\mathbf{x} - \mathbf{x}'|} d^3x'. \quad (4)$$

The vector potential is obtained by solving the wave equation

$$\left(\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 \right) \mathbf{A} = \frac{4\pi e}{c} \dot{\mathbf{y}}_{\perp}(t) f(\mathbf{x} - \mathbf{y}(t)). \quad (5)$$

Here $\dot{\mathbf{y}}_{\perp}$ is the transverse part of $\dot{\mathbf{y}}$.

It is convenient to work in the \mathbf{k} representation:

¹⁾ Our discussion follows closely the derivation in Low's recent paper (10).

$$[A(x, t), f(x)] = \frac{1}{(2\pi)^3} \int e^{ik \cdot x} [\tilde{A}(k, t), \tilde{f}(k)] d^3k.$$

We solve Eq. (5) with the retarded boundary condition:

$$\tilde{A}(k, t) = 4\pi e \int_{-\infty}^t dt' \frac{\sin kc(t-t')}{k} e^{-ik \cdot y(t')} \tilde{f}(k), \quad (6)$$

where $k = |\mathbf{k}|$.

Next, we compute $\mathbf{E} = -\nabla\phi - \partial\mathbf{A}/(c\partial t)$ and $\mathbf{B} = \nabla \times \mathbf{A}$, and insert the results into Eq. (3). In doing so, we observe that $\dot{\mathbf{y}}_{\perp} = \dot{\mathbf{y}} - \mathbf{k} (\mathbf{k} \cdot \dot{\mathbf{y}})/k^2$ can be replaced by $(2/3) \dot{\mathbf{y}}$ because of the spherical symmetry. The contribution via the scalar potential ϕ vanishes because of the spherical symmetry and because we are using the Coulomb gauge. The self force arising from the electric field is found to be

$$m_0 \ddot{\mathbf{y}}(t) = \mathbf{F}_{\text{ext}}(t) - \frac{2}{3} e^2 \frac{4\pi}{(2\pi)^3} \int d^3k |\tilde{f}(k)|^2 \int_0^{\infty} d\tau Y(t, \tau) \cos k\tau, \quad (7)$$

where

$$Y(t, \tau) = e^{ik \cdot \mathbf{y}(t)} e^{ik \cdot \mathbf{y}(t-\tau)} \dot{\mathbf{y}}(t-\tau). \quad (8)$$

The force arising from the magnetic field is nonlinear in \mathbf{y} (and its derivative), which we neglect. In linear approximation, the exponential factors in Eq. (8) can also be replaced by 1. Therefore,

$$Y(t, \tau) = \dot{\mathbf{y}}(t-\tau). \quad (9)$$

Equations (7) and (9) are the desired expression for the classical equation of motion and can be shown to be identical to the power series expression derived by Lorentz for linearized radiation reaction in the nonrelativistic approximation (2,11).

For the case of a spherical shell of radius a ,

$$f(\mathbf{x}) = \frac{\delta(|\mathbf{x}| - a)}{4\pi a^2}, \quad \tilde{f}(k) = \frac{\sin ka}{ka}. \quad (10)$$

Equations (7) and (9) become

$$m_0 \ddot{y}(t) = F_{\text{ext}}(t) + \frac{e^2}{3a^2 c} [\dot{y}(t - 2a/c) - \dot{y}(t)]. \quad (11)$$

This differential-difference equation will be referred to as the Sommerfeld-Page equation because they derived it (5,6).

Expanding $\dot{y}(t - 2a/c)$ in Eq. (11) in a Taylor series and neglecting terms that vanish as $a \rightarrow 0$,

$$m_0 \ddot{y}(t) = F_{\text{ext}}(t) - \delta m \ddot{y}(t) + \frac{2e^2}{3c^3} \dddot{y}(t), \quad (12)$$

where

$$\delta m = \frac{4}{3} \frac{e^2}{2ac^2}. \quad (13)$$

Equation (12) will be referred to as the Abraham-Lorentz equation in the following. It is the simplest form of the equation of motion, taking into account the electromagnetic self force in a nonrelativistic linear approximation and in the point particle limit.

The second term on the RHS of the Abraham-Lorentz equation can be interpreted as the inertia due to the electromagnetic mass δm . However, it is in contradiction with the notion of relativity because δm is different from the electrostatic mass $e^2/2ac^2$ by a factor of 4/3.

To see the meaning of the third derivative term in the Abraham-Lorentz equation we move the δm term to the LHS, multiply both sides by \dot{y} , and integrate over a finite interval of time. The LHS becomes the increase in the electron's kinetic energy. The RHS is

$$-\frac{2e^2}{3c^3} \int_{t_1}^{t_2} \ddot{y}^2 dt + \frac{2e^2}{3c^3} \dot{y} \cdot \dot{y} \Big|_{t_1}^{t_2}. \quad (14)$$

As long as the second term is negligible, this is the negative of the well-known dipole radiation. The last term in Eq. (12) is therefore reasonable. However, we will see later that this term gives rise to run-away or acausal behavior.

Equation (12) therefore contains two major difficulties: i) a contradiction with relativity, and ii) run-away and preacceleration behavior. We shall see in the following sections how these troubles are avoided. The bottom line is that there is no problem if the shell radius a is larger than the classical electron radius r_e .

RELATIVITY AND POINCARÉ STRESS

Let us first consider the relativistic generalization of Eq. (12). In 1903, before special relativity was established, Abraham derived (1) the following force equation valid for relativistic speed (neglecting terms vanishing in the limit $a \rightarrow 0$):

$$m_0 \frac{d}{dt} (\gamma \mathbf{u}) = \mathbf{F}_{\text{ext}} - \frac{2}{3} \frac{e^2}{ac^2} \frac{d}{dt} (\gamma \mathbf{u}) + \Gamma, \quad (15)$$

where $\mathbf{u} = d\mathbf{y}/dt$ is the three-velocity, $\gamma = 1/\sqrt{1 - \mathbf{u}^2/c^2}$, and

$$\Gamma = \frac{2}{3} \frac{e^2 \gamma^2}{c^3} \left\{ \ddot{\mathbf{u}} + \frac{3\gamma^2}{c^2} (\mathbf{u} \cdot \dot{\mathbf{u}}) \dot{\mathbf{u}} + \frac{\gamma^2}{c^2} \left[(\mathbf{u} \cdot \ddot{\mathbf{u}}) + \frac{3\gamma^2}{c^2} (\mathbf{u} \cdot \dot{\mathbf{u}})^2 \right] \mathbf{u} \right\}. \quad (16)$$

As was shown by von Laue (17), the Γ in the above is $1/\gamma$ times the space part of the four vector:

$$\Gamma^\mu = \frac{2}{3} e^2 (\ddot{u}^\mu + (\dot{u}^\alpha \dot{u}_\alpha) u^\mu). \quad (17)$$

(See the explanation in the "Notation" section.)

The second term in Eq. (15) gives rise to an electromagnetic mass that is $4/3$ times the electrostatic mass, in contradiction with relativity.

Abraham also found the rate of work done on an electron (neglecting terms vanishing in the limit $a \rightarrow 0$):

$$\frac{dE}{dt} = \mathbf{F}_{\text{ext}} \cdot \mathbf{u} - \frac{2}{3} \frac{e^2}{a} \frac{d}{dt} \left(\gamma - \frac{1}{4\gamma} \right) + \frac{2}{3} e^2 \frac{\gamma^4}{c^3} \left[\mathbf{u} \cdot \ddot{\mathbf{u}} + \frac{3\gamma^2}{c^2} (\mathbf{u} \cdot \dot{\mathbf{u}})^2 \right]. \quad (18)$$

Since $E = mc^2\gamma$ from relativity, it follows that $dE/dt = m(d\gamma u/dt) \cdot \mathbf{u}$. However, the force (Eq. (15)) and the power (Eq. (18)) do not satisfy this fundamental conservation law. The bad term is $2/3 e^2/a (d/dt)(1/4\gamma)$ in Eq. (18). Except for that term, Eqs. (15) and (18) would have formed a four vector equation.

Although these equations were derived before relativity was fully established, Abraham used the correct relativistic model of an electron in which the spherical charge distribution in the rest frame is contracted to a spheroid for a moving electron. The offending term in Eq. (18) arises from this relativistic change of the electron's shape. The derivations of Eq. (15) and Eq. (18) are difficult and confirmed by Schott (18) who carried out a very rigorous and complicated calculation.

We therefore have the strange situation that the relativistic equation of motion is in contradiction with relativity!

In the nonrelativistic limit, Eqs. (15) and (18) become, respectively,

$$m_0 \frac{d\mathbf{u}}{dt} = \mathbf{F}_{\text{ext}} - \frac{2}{3} \frac{e^2}{ac^2} \dot{\mathbf{u}} + \frac{2}{3} \frac{e^2}{c^3} \ddot{\mathbf{u}}, \text{ and} \quad (19)$$

$$\frac{dE}{dt} = \mathbf{F}_{\text{ext}} \cdot \mathbf{u} - \frac{5}{6} \frac{e^2}{ac^2} \mathbf{u} \cdot \dot{\mathbf{u}} + \frac{2}{3} \frac{e^2}{c^3} \mathbf{u} \cdot \ddot{\mathbf{u}}. \quad (20)$$

Once again $\mathbf{u} \cdot \text{force}$ (Eq. (19)) does not yield the power (Eq. (20)). (Note that Eq. (19) is identical to Eq. (12).) The problem was solved by Poincaré (3) with nonelectromagnetic stresses and Dirac by invoking covariance (4).

Poincaré, in his paper submitted in 1905 (3) (without knowledge of Einstein's work on special relativity)², observed that a purely electromagnetic model of the electron, such as the charged sphere, is not internally consistent because it will fly apart due to the electrostatic repulsion. To counteract the repulsive force, he imagined that the inside of the sphere provides a uniform negative pressure (or stress) $p = e^2/8\pi a^4$.

To see how this Poincaré stress solves the problem mentioned in the previous section, consider the charged shell in motion. Due to the Lorentz contraction, the sphere becomes an oblate spheroid with the minor axis in the direction of the motion reduced by γ . The work done by the mechanical force is given by the pressure times the volume change. (The pressure is relativistically invariant since the force and the area element transform the same way.) Thus the mechanical system must lose energy at the rate

$$\begin{aligned} \frac{dE}{dt} &= -\frac{e^2}{8\pi a^4} \frac{d}{dt} \left(\frac{4}{3} \frac{\pi a^3}{\gamma} \right) \\ &= -\frac{2}{3} \frac{e^2}{a} \frac{d}{dt} \frac{1}{4\gamma}. \end{aligned} \quad (21)$$

Here the expression $4\pi a^3/3\gamma$ is the volume of the spheroid. When Eq. (21) is added to Eq. (18), the discrepancy between the force and the power equation is removed.

However, the problem of the mass—the fact that the electromagnetic mass is 4/3 times the electrostatic mass—is not solved yet. One way of solving the problem is to say that the bare mass contains a term $-(1/3)$ times electrostatic mass. For a more

² Relative contributions of Poincaré and Einstein to special relativity is a subject of some debate and considerable historical interest. See A.A. Logunov referred to in (3).

formal approach (19), we can introduce a stress tensor representing the Poincaré stress by

$$\theta_{\text{Poincaré}}^{\mu\nu} = \frac{e^2}{8\pi a^3} (g^{\mu\nu} - qu^\mu u^\nu) \Theta(a-r), \quad (22)$$

where $g^{\mu\nu} = (1, -1, -1, -1)$, q is an arbitrary constant, Θ is the step function, and r is the radial coordinate in the rest frame of the shell. The quantity $\theta_{\text{Poincaré}}^{\mu\nu}$ is constructed so that

$$\partial_\mu \theta^{\mu\nu} = \partial_\mu (\theta_{\text{EM}}^{\mu\nu} + \theta_{\text{Poincaré}}^{\mu\nu}) = 0, \quad (23)$$

where $\theta_{\text{EM}}^{\mu\nu}$ is the electromagnetic stress tensor associated with the Coulomb field of a spherical shell. Equation (23) assures that the total momentum

$$p^\mu = \int d^3x \theta^{\mu 0} \quad (24)$$

is a four vector (11). The momentum associated with the Poincaré stress alone is

$$p_{\text{Poincaré}}^\mu = \int a^3 x \theta_{\text{Poincaré}}^{\mu 0} = + \frac{e^2}{2a} \frac{1}{3} \left[-\frac{1}{\gamma} + q\gamma, q\gamma u \right] \quad (25)$$

We need to add $dp_{\text{Poincaré}}^\mu/dt$ to the RHS of Eqs. (18) and (15). Our previous result (Eq. (21)) corresponds to the case $q = 0$. If, on the other hand, we choose $q = 1$, we obtain

$$m_0 c^2 \frac{d}{ds} u^\mu = f_{\text{ext}}^\mu + \Gamma^\mu - \frac{e^2}{2a} \frac{du^\mu}{ds}. \quad (26)$$

We now have an equation of motion of an electron with the observed mass $m = m_0 + \delta m$, $\delta m = e^2/2ac^2$, solving the so-called 4/3 problem.

The above method of solving the 4/3 problem is clearly rather formal and arbitrary. There are also more intuitive approaches, for example by Boyer (20) who notes that the Poincaré stress may not act at the same time in all parts of the moving electron. A review of different solutions of the 4/3 problem is given by Rohrlich (16).

The saga of the 4/3 problem is in some sense a story of how special relativity proved itself as a theory of internal consistency and beauty. The investigation of the electron's equation of motion started while special relativity was still evolving. Therefore, there was doubt whether the electron theory was consistent with relativity, which lingered even after relativity was fully established. Einstein, never doubting relativity, wasted no time in checking covariance of the electron theory. He was too busy working out general relativity!

Before closing this section, we need to mention Dirac's contribution (4). He understood that the problem with the Abraham-Lorentz equation was in trying to approach too near an electron. He therefore devised an ingenious way to avoid the difficulty. Stay a finite distance away from an electron and demand relativistic covariance. He then obtained Eq. (26), which is therefore referred to as the Abraham-Lorentz-Dirac equation.

Actually, Dirac's derivation of Eq. (4) is valid only up to a term of the form dB^μ/ds , where B^μ is a four vector with the restriction $u^\mu > dB_\mu/ds = 0$. The simplest choice, $B^\mu = ku^\mu$, gives rise to a term that can be incorporated into the inertial term ($mc du^\mu/ds$) in Eq. (4). The next order term involving \dot{u}^μ (the dot indicating the derivative with respect to s) can be shown to be of the form $B^\mu = k'((\dot{u}^\alpha \dot{u}_\alpha)^2 u^\mu + 4(\dot{u}^\alpha \ddot{u}_\alpha) \dot{u}^\mu)$. Terms of such a complexity are disposed of by saying that *a simple thing like an electron cannot possibly have such a complication*.

If we are willing to make a few very plausible assumptions, the Abraham-Lorentz-Dirac equation can be derived very easily as follows (16):

$$m_0 c^2 \frac{du^\mu}{ds} = f_{\text{ext}}^\mu + X^\mu, \quad (27)$$

where X^μ includes radiative effects. Both sides of Eq. (27) are orthogonal to u^μ . Thus we may write $X^\mu = (g^{\mu\nu} - u^\mu u^\nu) Y_\nu$. Try Y^μ of the form $a u^\mu + b \dot{u}^\mu + c \ddot{u}^\mu$. The first term does not contribute, and the second term is of the same form as the mass term, which can be incorporated into m . With the third term, it is easy to show that X^μ is proportional to Γ^μ . (Here we are using $\dot{u}^\alpha \dot{u}_\alpha + \ddot{u}_\alpha \cdot u^\alpha = 0$.) The coefficient is determined by demanding consistency with the nonrelativistic Abraham-Lorentz equation, leading to $\Gamma^\mu = X^\mu$. End of the proof.

RUN-AWAY SOLUTIONS, BOUNDARY CONDITIONS, AND INTEGRAL EQUATIONS

The general solution of the Abraham-Lorentz equation is

$$m \ddot{y}(t) = e^{-t/t_e} \left[m \ddot{y}(0) - 1/t_e \int_0^t dt' e^{-t'/t_e} F_{\text{ext}}(t') \right], \quad (28)$$

where

$$t_e = \frac{2 r_e}{3 c} = \frac{2 e^2}{3 mc^3}. \quad (29)$$

The solution in general exhibits exponential growth, i.e., the run-away behavior. Dirac noted that the run-away can be avoided if we choose the initial condition:

$$m \ddot{y}(0) = 1/t_e \int_0^{\infty} dt' e^{-t'/t_e} F_{\text{ext}}(t'). \quad (30)$$

Then

$$m \ddot{y}(t) = \int_0^{\infty} d\alpha e^{-\alpha} F_{\text{ext}}(t + t_e \alpha). \quad (31)$$

This solution now exhibits preacceleration: the particle starts to move before the force is applied and the initial condition depends upon the entire future path.

Nevertheless, it satisfies the Rohrlich criteria of “*the unobservability of very small charges*” (13). That is, as the charge becomes very small we shouldn’t have a solution widely different from that of an uncharged particle.

The reason why the Abraham-Lorentz equation exhibits either run-away or preacceleration is the fact that, as the electron radius a vanishes, the bare mass m_0 becomes negative to keep the observed mass m finite. A particle with a negative mass can clearly supply an infinite amount of energy.

The difficulty can be avoided in an extended electron model. To see this, we return to the Sommerfeld-Page Eq. (11), which can be written in the following form³:

$$\ddot{y}(t) = \frac{F_{\text{ext}}(t)}{m(1 - ct_e/a)} + \left(\frac{c}{2a} \right) \left(\frac{ct_e}{a} \right) \frac{1}{(1 - ct_e/a)} [\dot{y}(t - 2a/c) - \dot{y}(t)]. \quad (32)$$

In the above expression, the combination $m(1 - ct_e/a)$ is simply the bare mass m_0 . Assume that there are no external forces, and there exists a run-away solution of the form $y(t) = y(0) e^{\alpha t}$ with a positive real part of α . From Eq. (32) it is easy to see that this is possible only when $1 - ct_e/a$ is negative. Thus the run away solution is possible if and only if the electron radius a is less than $(2/3) r_e$. For $a > (2/3) r_e$ we get damped oscillatory solutions.

It is sometimes stated that the run-away behavior is due to the infinite energy associated with a point electron. This is false because the run-away occurs for even a finite a as long as it is less than $(2/3) r_e$. Run-away occurs if and only if the bare mass is negative (and, therefore, the Hamiltonian is no longer positive definite).

We can also show that if $a > ct_e$, the motion is causal with no preacceleration. This can be seen most easily if Eq. (32) is turned into an integral equation with a Green’s function:

³) This derivation here is due to Moniz and Sharp (8). See also Pearle (15).

$$\ddot{y}(t) = \int_{-\infty}^{\infty} G(t-t') F_{\text{ext}}(t') dt', \quad (33)$$

where

$$G(t-t') = \frac{1}{2\pi m} \int_{-\infty}^{\infty} \frac{e^{-i\omega(t-t')}}{B} d\omega, \text{ and} \quad (34)$$

$$B = 1 - \frac{i}{\omega} \left(\frac{c}{2a} \right) \left(\frac{ct_e}{a} \right) \left[e^{\frac{i\omega 2a}{c}} - \frac{i\omega 2a}{c} - 1 \right]. \quad (35)$$

It can be shown that all poles of $1/B(\omega)$ occur in the lower half ω -plane if

$$a > \frac{2}{3} r_e. \quad (36)$$

Then the Green's function $G(\tau)$ vanishes for $\tau < 0$. Therefore, the motion is causal. On the other hand, if $a < \frac{2}{3} r_e$, then there is in general acausal behavior.

A relativistic generalization of the Sommerfeld-Page equation was given by Caldirola (7):

$$m_0 c^2 \dot{u}^\mu = f_{\text{ext}}^\mu + \frac{2}{3} \frac{e^2}{a} \frac{1}{2a} [u^\mu (s-2a) - u^\mu (s) u_\alpha (s) u^\alpha (s-2a)]. \quad (37)$$

The Caldirola equation is fine: no run-aways and no causality problems. However, when we take $a \rightarrow 0$, then there are problems.

Yaghjian (14) attempted to produce a causal equation with no run-aways for a point electron. He modified the Abraham-Lorentz Eq. (12) by multiplying the third derivative term by a function $\eta(t)$, which changes smoothly from 0 for $t < 0$ to 1 for $t > 2a/c$ (or by multiplying the Γ^μ term by a similar function $\eta(s)$), arguing that the Taylor series expansion breaks down when the force changes abruptly. However, the solution of the Yaghjian equation still exhibits acausal behavior.

QUANTUM MECHANICAL EQUATIONS

It had been hoped that the difficulty of the classical theory in taking a point particle limit might be solved by quantum mechanics. The observation that the mass renormalization δm is less singular in quantum electrodynamics than in classical theory seemed to reinforce this hope. However, a real quantum mechanical analysis of the electron equation has not been carried out until modern time. The reason behind this

lack of activity appears to be the success of the renormalization theory in quantum electrodynamics, which established that all observable phenomena can be calculated to a finite answer order by order in perturbation theory in terms of the observed mass and charge.

In 1977, Moniz and Sharp presented a very interesting quantum mechanical calculation (8), according to which the equation of motion in nonrelativistic quantum mechanics is finite even in the point particle limit. However, the paper has not received much attention, probably because the derivation was not transparent due to the elaborate manipulations of infinite series.

One of us (KJK) was able to derive the result of Moniz and Sharp via a more understandable method (9), which will be summarized here.

The derivation of the quantum mechanical equation of motion for Heisenberg operators proceeds similar to the derivation in the classical case given in the section "Abraham-Lorentz and Other Classical Equations of Motion." Again neglecting the magnetic term, one arrives at an equation identical in form to Eq. (7), but the operator Y is given by the symmetrized version of the classical expression, Eq. (8), as follows:

$$Y(t, \tau) = \frac{1}{4} \left\{ e^{ik \cdot y_2}, \left\{ e^{-ik \cdot y_1}, \dot{y}_1 \right\}_+ \right\}_+ \\ = \frac{1}{4} \left(e^{ik \cdot y_2} e^{-ik \cdot y_1} \dot{y}_1 + e^{ik \cdot y_1} \dot{y}_1 e^{-ik \cdot y_2} \right) + [k \rightarrow -k]^+ \quad (38)$$

Here $y_2 = y(t)$, $y_1 = y(t-\tau)$, $\{ \}_+$ is the anticommutator, and $[k \rightarrow -k]^+$ indicates terms obtained by changing the sign of k and taking the Hermitian conjugate.

Noting that $\mathbf{p} = \mathbf{P} - eA/c = m_0 \dot{\mathbf{y}}$ is the kinetic momentum operator (\mathbf{P} = canonical momentum), we obtain

$$e^{ik \cdot y_1} \dot{y}_1 e^{-ik \cdot y_1} = \dot{y}_1 - ck\lambda, \quad (39)$$

where $\lambda = \hbar/m_0c$ is the Compton wavelength for the bare mass m_0 . In classical mechanics, we would have replaced the factor $\exp(ik \cdot y_1)$ by 1 since it would at most contribute to nonlinear terms. Such a procedure is not justified in quantum mechanics as is clear from Eq. (39). Similarly, the factor $\exp(ik \cdot y_2) \cdot \exp(-ik \cdot y_1)$ in Eq. (38) cannot simply be replaced by 1. Instead, we proceed as follows⁴:

⁴ Similar reduction of product of exponential operators was used by Baier and Katkov (21) in their calculation of quantum synchrotron radiation. The steps used here follow closely a simplified formulation of the Baier-Katkov reduction by Cahn and Jackson (22).

$$\begin{aligned} \exp(\mathbf{i}\mathbf{k} \cdot \mathbf{y}_2) \exp(-\mathbf{i}\mathbf{k} \cdot \mathbf{y}_1) &= \exp(i\tau H / \hbar) \exp(\mathbf{i}\mathbf{k} \cdot \mathbf{y}_1) \exp(-i\tau H / \hbar) \exp(-\mathbf{i}\mathbf{k} \cdot \mathbf{y}_1) \\ &= \exp(i\tau H(\mathbf{p}) / \hbar) \exp(-i\tau H(\mathbf{p} - \hbar\mathbf{k}) / \hbar) \quad (\tau = t - t') \end{aligned} \quad (40)$$

Here H is the Hamiltonian. Since H is a sum of the kinetic energy $\mathbf{p}^2/2m_0$ and the electromagnetic energy, it follows that

$$H(\mathbf{p} - \hbar\mathbf{k}) = \frac{\hbar^2 \mathbf{k}^2}{2m_0} + H(\mathbf{p}) - \frac{\hbar \mathbf{p} \cdot \mathbf{k}}{m_0}. \quad (41)$$

Now, using the Campbell-Baker-Hausdorff formula (23) to first order in \mathbf{p} , we can show that

$$\exp(\mathbf{i}\mathbf{k} \cdot \mathbf{y}_2) \exp(-\mathbf{i}\mathbf{k} \cdot \mathbf{y}_1) = \exp\left(-\frac{i\tau \hbar \mathbf{k}^2}{2m_0}\right) [1 + \mathbf{i}\mathbf{k} \cdot (\mathbf{y}_2 - \mathbf{y}_1)]. \quad (42)$$

Thus

$$\mathbf{Y} = \frac{1}{2} \left\{ \exp\left(-\frac{i\tau \hbar \mathbf{k}^2}{2m_0}\right) (1 + \mathbf{i}\mathbf{k} \cdot (\mathbf{y}_2 - \mathbf{y}_1)) \left(\dot{\mathbf{y}}_1 - \frac{c\mathbf{k}\lambda}{2} \right) \right\} + [\mathbf{k} \rightarrow -\mathbf{k}]^+. \quad (43)$$

Noting that odd power in \mathbf{k} vanishes after $d^3\mathbf{k}$ -integration and $\langle \mathbf{k}_x^2 \rangle = \frac{1}{3} \mathbf{k}^2$, one obtains (keeping only terms linear in \mathbf{y})

$$\mathbf{Y}(t, \tau) = \dot{\mathbf{y}}(t - \tau) \cos\left(\lambda c \tau \mathbf{k}^2 / 2\right) - \frac{\mathbf{k}^2 c \lambda}{6} (\mathbf{y}(t) - \mathbf{y}(t - \tau)) \sin\left(\lambda c \tau \mathbf{k}^2 / 2\right). \quad (44)$$

The linearized electron's equation of motion in nonrelativistic quantum mechanics is then given by inserting Eq. (44) into Eq. (7). Expanding the operator $\mathbf{y}(t - \tau)$ in Eq. (44) in a Taylor series around $\tau = 0$ and inserting it in Eq. (7), an equation involving a sum of derivatives of $\mathbf{y}(t)$ is obtained. It can be shown that the coefficients of these derivatives are exactly those derived in Moniz and Sharp⁵:

$$m_0 \mathbf{y}(t) = \mathbf{F}_{\text{ext}}(t) - \frac{2}{3} \frac{e^2}{c^2} \sum_{n=0}^{\infty} \left(\frac{(-1)^n A_n}{n! c^n} \right) \frac{d^{n+2} \mathbf{y}(t)}{dt^{n+2}}, \quad (45)$$

⁵ The derivation in the above gives rise to a term proportional to $\dot{\mathbf{y}}(t)$ with negative coefficient. Such a term, not discussed in Moniz and Sharp, would provide further damping of the motion.

where

$$A_n = \left(1 + \frac{\lambda_c}{3(n+2)} \frac{\partial}{\partial \lambda_c} \right) \left(1 + \frac{\lambda}{(n+1)} \frac{\partial}{\partial \lambda} \right) B_n, \text{ and}$$

$$B_n = \sum_{k=0}^{\infty} \frac{n!}{(n+2k)!} \binom{n+1-2k}{2k} \left(\frac{-\lambda_c^2}{4} \right)^k \int dx dx' f(x) |x-x'|^{n-1+2k} (\nabla_{x'}^2)^{2k} f(x'). \quad (46)$$

In particular, the coefficient of $\ddot{y}(t)$, the quantum mechanical self-mass δm , is found to be

$$\delta m = \left(\frac{2e^2}{3c^2} \right) \left(1 + \frac{\lambda}{6} \frac{\partial}{\partial \lambda} \right) \left(1 + \lambda \frac{\partial}{\partial \lambda} \right) \Omega_0. \quad (47)$$

Here

$$\Omega_0 = \frac{2}{\pi} P \int_0^{\infty} \frac{|\tilde{f}(k)|^2}{(1-\lambda^2 k^2/4)} dk, \quad (48)$$

where P denotes the principal value integration. Equation (48) is remarkable in that the self-mass remains finite in the point particle limit $f(k) \rightarrow 1$. *In fact, the self-mass vanishes in this limit!* $\delta m \rightarrow 0$ ($a \rightarrow 0$ with λ fixed in quantum mechanics).

Therefore, λ (bare mass) = λ (observed mass). This is why we used the same notation for these two Compton wavelengths.

In another limit, $\lambda \rightarrow 0$,

$$\Omega_0 = \frac{2}{\pi} P \int_0^{\infty} |\tilde{f}(k)|^2 dk,$$

and the theory reproduces the classical result, as it should.

Whether the quantum mechanical equation of motion exhibits run-away or acausal behavior can be studied by writing Eq. (7) in a Green's function form, similar to that in the section on "Run-away Solutions, Boundary Conditions, and Integral Equations." Such an analysis in the point particle limit was carried out by Moniz and Sharp. It is found that the motion is causal with no run-aways if

$$\alpha = \frac{e^2}{\hbar c} = \frac{r_e}{\lambda} < 1.75. \quad (50)$$

In quantum mechanics, an electron is spread over a Compton wavelength. Thus, the above condition is reasonable in view of the classical causality condition Eq. (36), which can be written as $r_e/a < 1.5$.

Nonrelativistic quantum mechanics is not valid for rapid motion with frequency $\hbar\omega > mc^2$ or if we approach a distance λ near the electron, and a full QED analysis must be performed. The only calculation by QED was reported by Francis Low (13). He has not derived an equation of motion, but he has shown that the motion is finite in each order of α . This is reasonable in view of the well-established renormalization theory that gives a finite answer for any physical process in perturbation expansion in α . However, the perturbation theory is not suitable for taking the classical limit, and it is unclear whether QED can actually produce an equation of motion.

CONCLUDING REMARKS

The impression one gets from reading text books is that the classical electron theory is in trouble due to run-away solutions and acausal behavior. However, we have seen that classical theory is actually fine if the electron is taken to be a spherical object of not too small a radius, greater than the classical electron radius r_e . The restriction is reasonable since r_e is about 100 times smaller than the Compton wavelength λ , and we cannot consider distances less than λ without considering quantum mechanics.

The nonrelativistic quantum theory also looks fine: In fact, the quantum theory is better behaved than the classical theory because it is finite and causal irrespective of the size of the electron, as long as $\alpha = r_e/\lambda < 1.75$. The inequality is certainly satisfied in the real world where $\alpha = 1/137$. The fact that the quantum theory is better behaved is also reasonable because the electron is smeared out due to the uncertainty principle. The quantum theory as reviewed here has the appropriate feature of having the correct classical limit. However, the limit of validity of the nonrelativistic quantum analysis is not really understood. The vanishing of the self-mass δm in the point particle limit must be accordingly interpreted with care.

In fact, we know that the nonrelativistic treatment cannot strictly be valid in the point-particle limit because of vacuum polarization. It is reasonable that a full QED calculation will be at least as well-behaved as is the case in nonrelativistic quantum mechanics. Unfortunately, there exists as yet no real calculation in QED to confirm these conjectures.

Nevertheless, we are entitled to close this paper on a positive note: We started from the Abraham-Lorentz theory of point electron, which was problematic. We saw how the theory became fine by the use of an extended electron. The nonrelativistic quantum mechanics is fine even in the point-particle limit. A proper relativistic quantum electrodynamics calculation has not yet been done, and may not be possible (within QED as a perturbation theory), but should be no more singular than the nonrelativistic theory.

ACKNOWLEDGMENTS

The authors would like to acknowledge communications with Dave Jackson, Francis Low, Robert Mills, Fritz Rohrlich, and David Sharp.

REFERENCES

1. Abraham, M., "Theorie der Elektrizitat," Vol II: Elektromagnetische Theorie der Strahlung, Leipzig, Teubner (1905).
2. Lorentz, H.A., "The Theory of Electrons," Leipzig: Teubner (1909) (2nd edition, 1916).
3. Poincaré, H., "On the dynamics of the electron," *Rendiconti del Circolo Matematico di Pulermo* **21**, 129 (1906); translated by Scientific Translation Service, Ann Arbor, MI. The article is also translated into English with comments in "On the articles by Henri Poincaré," A.A. Logunov (Publishing Dept. of the Joint Institute of Nuclear Research, Dubna, 1995).
4. Dirac, P.A.M. "Classical theory of radiating electrons," *Proc. Roy. Soc. Lond.* **A167**, 148 (1938).
5. Sommerfeld, A., "Simplified deduction of the field and the forces of an electron moving in any given way," *Akad. van Wetensch. te Amsterdam* **13** (1904) (English translation **7**, 346 (1905)).
6. Page, L., *Phys. Rev.* **11**, 376 (1918).
7. Caldirola, P., "A new model of classical electron," *Nuovo Cimento* **3**, Supplemento **2**, 297 (1956).
8. Moniz, E.J. and Sharp, D.H., "Radiation reaction in nonrelativistic quantum electro-dynamics," *Phys. Review D* **15**, 2850 (1977); see also Grotch, H., Kazes, E., Rohrlich, F. and Sharp, D.H., "Internal retardation," *Acta Physica Austriaca* **54**, 31 (1982).
9. Kim, K.-J., to be published.
10. Low, F., "Run-Away Electrons in Relativistic Spin 1/2 Quantum Electrodynamics," Preprint MIT-CTP-2522 (1997).
11. Panofsky, W.K.H., and Phillips, M., *Classical Electricity and Magnetism*, Reading, MA: Addison-Wesley, 1962.
12. Jackson, J.D., *Classical Electrodynamics*, New York: Wiley, 2nd Edition, 1975; 3rd Edition, 1998. Chapter 16 of 3rd Edition contains a review of the equation of motion of an electron, much more extensive than in previous editions.
13. Rohrlich, F., *Classical Charged Particles*, Reading, MA: Addison-Wesley (1965) (2nd edition 1990).
14. Yaghjian, A.D., "Relativistic Dynamics of a Charged Sphere," *Lecture Notes in Physics* **11**, Springer-Verlag, Berlin (1992)
15. Pearle, P., "Classical Electron Models," in *Electromagnetism: Path to Research*, D. Teplitz, Ed., New York: Plenum, 1982.
16. Rohrlich, F. "The Dynamics of a Charged Sphere and the Electron," *Am. J. Phys.* **65**, 105 (1997).
17. Laue, M., "Die Wellenstrahlung einer bewegten Punktladung nach dem Relativitätsprinzip," *Annalen der Physik* **28**, 436 (1909). Also see Podolsky, B. and Kunz, K.S., "Fundamentals of Electrodynamics," New York, Marcel Dekker, sec. 15 (1969).
18. Schott, G.A., "Electromagnetic Radiation," Cambridge University Press (1912).
19. Schwinger, J., "Electromagnetic Mass Revisited," *Foundations of Physics*, **13**, 373 (1983).
20. Boyer, T.H., *Phys. Rev.* **D25**, 3246 (1982).
21. Baier, V.N., and Katkov, V.M., *Sov. Phys. JETP* **26**, 854 (1968).
22. Cahn, R.N., and Jackson, J.D., unpublished LBL note.
23. Dragt, A.J., and Finn, J.M., *J. Math. Phys.* **17**, 2215 (1976).

**ERNEST ORLANDO LAWRENCE BERKELEY NATIONAL LABORATORY
ONE CYCLOTRON ROAD | BERKELEY, CALIFORNIA 94720**