UC Irvine

UC Irvine Previously Published Works

Title

Radiocarbon constraints imply reduced carbon uptake by soils during the 21st century.

Permalink

<https://escholarship.org/uc/item/2r76q5gt>

Journal

Science (New York, N.Y.), 353(6306)

ISSN

0036-8075

Authors

He, Yujie
Trumbore, Susan E
Torn, Margaret S
et al.

Publication Date

2016-09-01

DOI

10.1126/science.aad4273

Peer reviewed

1 **Title: Radiocarbon constraints imply reduced carbon uptake by soils during**
2 **the 21st century**

3 **Authors:** Yujie He ^{1*}, Susan E. Trumbore², Margaret S. Torn³, Jennifer W. Harden⁴, Lydia J. S.
4 Vaughn³, Steven D. Allison^{1,5}, James T. Randerson¹

5
6**Affiliations:**

7¹ Department of Earth System Science, University of California, Irvine, CA, USA

8² Department of Biogeochemical Processes, Max-Planck-Institute for Biogeochemistry, Jena,
9Germany

10³ Earth Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

11⁴ U.S. Geological Survey, Menlo Park, CA, USA

12⁵ Department of Ecology and Evolutionary Biology, University of California Irvine, Irvine, CA,
13USA

14*Correspondence to: yujie.he@uci.edu

15-

16

17**Subheading:** Reduced carbon uptake by soils during the 21st century

18

19**One Sentence Summary:**

20Global radiocarbon observations show that Earth system models, lacking carbon stabilization
21mechanisms, overestimate the ~~21st century~~ soil carbon sink by almost two-fold.

22**Abstract:**

23Soil is the largest terrestrial carbon reservoir and may influence the sign and magnitude of carbon
24cycle-climate feedbacks. ~~Changes in soil carbon—the largest terrestrial carbon reservoir—may~~
25~~influence the sign and magnitude of climate-carbon cycle feedbacks.~~ Many Earth system models
26(ESMs) estimate a significant soil carbon sink by 2100, yet the underlying carbon dynamics
27determining this response have not been systematically tested against observations. ~~Using~~ We
28~~used~~ $\delta^{14}\text{C}$ data from 157 globally distributed soil profiles sampled to 1 m depth, ~~we to~~ show that
29ESMs underestimated d the mean age of soil carbon by more than six-fold (430 ± 50 years vs.
30 3100 ± 1800 years). Consequently, ESMs overestimated d the carbon sequestration potential of soils
31~~21st century soil carbon sequestration~~ by nearly two-fold ($40 \pm 27\%$). These biases/inconsistencies
32~~suggest that ESMs must better represent carbon stabilization processes and the turnover time of~~
33~~slow and passive reservoirs when simulating future atmospheric CO₂ dynamics.~~
34~~To improve simulations of future atmospheric CO₂ and carbon storage, ESMs must better~~
35~~represent stabilization processes and turnover times for soil carbon pools.~~
36**Keywords:** soil carbon, earth system models, carbon-concentration feedback, mean age,
37radiocarbon

38Main Text:

39 Soil carbon is a dynamic reservoir that may increase substantially in size during the 21st
40century, as predicted by Earth system models (ESMs), thereby influencing the sign and
41magnitude of carbon cycle feedbacks under climate change (1-4). Under a high radiative forcing
42scenario (Representative Concentration Pathway 8.5), changes in soil carbon estimated by
43different models vary from a loss of 20 Pg C to a gain of more than 360 Pg C (5). These models
44suggest that the global carbon inventory in mineral soils may increase by 30% or more over a
45timespan of about two centuries. The multi-model mean soil carbon accumulation of 109 Pg C
46(5) represents about one decade of global fossil fuel emissions at current rates and 5% of
47cumulative fossil emissions by 2100 for this scenario (6). This soil carbon sink represents a
48negative feedback on CO₂ emissions, and if robust, would slow the rate of climate change.

49 Still, there are substantial uncertainties in the soil carbon sink projected by ESMs (5).
50Rapid rates of carbon sequestration in ESMs contrast with findings from CO₂ and warming
51experiments (7, 8) as well as multiple theoretical and observational constraints indicating slow
52(millennial) rates of soil organic carbon accrual and turnover (9-14). Model uncertainty—as
53measured by inter-model spread—is high for soil carbon turnover time (τ) and exceeds the
54uncertainty estimated for carbon uptake through gross primary production (GPP) (15, 16).

55 In coupled model simulations, the relative sink strength (i.e. percentage change in soil
56carbon) depends on the responses of net primary production (NPP) and soil carbon dynamics to
57increasing atmospheric CO₂ concentrations and to a lesser extent climate change (5). Elevated
58CO₂ increases photosynthesis and NPP, which results in greater carbon inputs to soil pools with
59decadal or longer residence times. Carbon sequestration in soils reduces the build-up of CO₂ in
60the atmosphere (the carbon-concentration feedback). On the other hand, elevated CO₂ also

61warms the climate, which tends to accelerate soil carbon turnover and reduce carbon storage (the
62carbon-climate feedback) (17, 18). Although these feedbacks oppose one another, the carbon-
63concentration feedback is more than 4 times greater on average than the carbon-climate feedback
64in current ESMs at the global scale (3). Differences in the representation of elevated CO₂ versus
65climate effects on ecosystem processes result in substantial variation in soil carbon sequestration
66estimates (19) (Table S1).

67 Without [a strong carbon-concentration feedbacks](#), ESMs would likely project smaller
68gains or larger losses of soil carbon over the 21st century. Our aim was to constrain the magnitude
69of the soil carbon-concentration feedback with soil radiocarbon observations. Radiocarbon
70content ~~can be used to constrain~~[provides information about](#) soil carbon turnover over centuries to
71millennia based on radioactive decay and over decades based on inputs of ¹⁴C from atmospheric
72weapons testing (“bomb carbon”). Accurate carbon turnover times are important for ESM
73projections because pools with short turnover times rapidly adjust to increasing NPP, whereas
74pools with long turnover times (and by inference low rates of inputs) change only slowly,
75possibly beyond the time horizon of effective climate mitigation efforts. [Therefore inaccuracies](#)
76[in the representation of carbon turnover times will have consequences for the rate and magnitude](#)
77[of the carbon-concentration feedback simulated by ESMs. If ESMs omit soil carbon pools with](#)
78[long turnover times, they could overestimate the carbon-concentration feedback effect on soil](#)
79[carbon storage during the 21st century while underestimating soil carbon storage at steady-state](#)
80[\(after millennia\).](#)

81 Here we used $\Delta^{14}\text{C}$ measurements at 157 sites across multiple biomes (Fig 1, Table S2)
82along with carbon inventory data to constrain soil carbon dynamics in five biogeochemically-
83coupled ESM simulations (esmFixClim1) from the Coupled Model Intercomparison Project

84Phase 5 (CMIP5) (20). In these idealized simulations the atmospheric CO₂ mole fraction starts at
85a preindustrial value of 285 ppm and rises at a rate of 1% yr⁻¹, thus quadrupling over 140 yrs. The
86biogeochemical components of each model experience the increasing trajectory of atmospheric
87CO₂, whereas the atmospheric radiation submodels do not, limiting impacts solely to the direct
88effects of CO₂ on plant physiology and thus enabling diagnosis of carbon-sink sensitivity to
89increasing CO₂.

90 Total initial soil carbon in the ESMs was not significantly different from the total amount
91in the top meter of the Harmonized World Soil Database (HWSD; Fig 2a, b) for 4 of the 5
92models ($p > 0.05$, except CESM $p = 0.03$). Therefore we compared ESM-derived $\Delta^{14}\text{C}$ — to
93observations derived from soil profiles to [a](#) 1 m depth. The carbon and ¹⁴C patterns of the soil
94profiles we used were similar to those reported in a recent synthesis paper (21), and we used
95some of the same profiles in our analysis.

96 Comparing ESM outputs to ¹⁴C observations requires a model analysis approach because
97most ESMs do not yet explicitly simulate $\Delta^{14}\text{C}$ in soils, and no ESMs had reported turnover times
98for soil carbon pools. Therefore we used a reduced complexity (RC) model to approximate soil
99carbon dynamics in each ESM. This approach allowed us to ~~(1)~~ estimate the ¹⁴C ~~ages and~~
100turnover times [and \$\Delta^{14}\text{C}\$](#) associated with the carbon pools in different ESMs (Table S3), ~~(2)~~
101~~compare with observations, and (3) assess the consequences if ESM parameters were aligned~~
102~~with observations.~~ Where possible, we used a three-pool RC model (with fast, slow, and passive
103pools) to simulate carbon and ¹⁴C dynamics. A multi-pool structure is essential because
104radiocarbon observations show that soil carbon fluxes (NPP inputs and heterotrophic respiration)
105exchange mainly with short-lived pools whereas carbon stocks are dominated by long-lived
106pools (12, 18, 22, 23). The three-pool RC model had five parameters representing turnover times

107of fast, slow, and passive pools (τ_{fast} , τ_{slow} , τ_{passive}) and transfer coefficients (r_f , r_s) that regulated
 108carbon flow from the fast to slow, and slow to passive pools (Fig S1). We used a two-pool RC
 109model for GDFL-ESM2M because it represents soil carbon with two pools (24) and for
 110HadGEM2-ES because it reported carbon for two pools (Table S4). The two-pool RC model had
 111three parameters, representing τ_{fast} , τ_{slow} , and r_f (Fig S1). After verifying that the RC model was a
 112good approximation of each ESM based on minimization of root-mean-square error, we used the
 113RC models to simulate $\delta^{14}\text{C}$ values at each grid cell, with observed atmospheric $\delta^{14}\text{C}$ for the
 114past 50 kyr as a boundary condition and, accounting for radioactive decay (see supplementary
 115material).

116 We used an inverse analysis to determine the RC model parameters that were most
 117consistent with our $\delta^{14}\text{C}$ dataset. In the inversion, we ~~held the total carbon mass in the ESM at its~~
 118~~preindustrial value (except in sensitivity analyses where it was matched to HWSD observations);~~
 119~~and~~ adjusted the parameters described above to match both the total carbon and radiocarbon
 120constraints. With these constraints, turnover time and carbon input rate for each pool were
 121coupled such that an increase in turnover time required a compensatory decline in inputs (Fig
 122S2). RC parameters derived from the inversion were subsequently used to assess consequences
 123of ^{14}C constraints for the carbon-concentration feedback.

124 All ESMs projected an increase in soil carbon over 140 yrs with multi-model mean of
 12532.6% (Table 1). This increase was primarily driven by increasing carbon inputs to soil under the
 126quadrupling of CO_2 (Table S3), as temperature increased by only a small amount (mean $\pm$ 1 s.d.
 127is was 0.52 ± 0.68 °C) for this set of biogeochemically-coupled simulations. CESM showed the
 128smallest soil carbon increase (6.3%) primarily because of low litter inputs relative to other ESMs

129(Table S3). For this time period and set of model runs, storage in soil carbon accounted for
13042±17% of the total accumulation of carbon in the terrestrial biosphere.

131 Both two- and three-pool RC models reproduced the global carbon dynamics of the
132original ESMs (Fig S3-S5; Table S5). The τ_{fast} across all RC models was less than 20 yrs, while
133 τ_{slow} varied from 40 to 600 yrs (Fig S6) with a multi-model mean of 212±104 yrs. The mean τ_{passive}
134for the three-pool RC models from CESM, IPSL and MRI was 1185±123 yrs (Table 1, Fig S7).
135Using the RC model parameters estimated at each grid cell within an ESM, we calculated the
136expected $\Delta^{14}\text{C}$. The resulting global average $\Delta^{14}\text{C}$ for 1995 (median sample year of site profiles)
137from the RC models was significantly higher than the mean of the observations (-6.4±64‰ vs.
138-211±156‰) (Fig 2c,d, $p<0.001$). $\Delta^{14}\text{C}$ values from RC models approximating ESMs with
139passive pools were more negative (-53±35‰) but still significantly higher than the observations
140($p<0.001$). Converting these $\Delta^{14}\text{C}$ observations into mean age for the soil profile yielded an
141estimate of 3100±1800 yrs for the observed soil carbon integrated to 1 m and 430±50 yrs for the
142ESMs (Fig 2e,f). These results indicated that the ESMs did not have enough old carbon that had
143experienced significant levels of radioactive decay; concurrently the models assimilated too
144much bomb ^{14}C . Relative to the observations, the ESM-based RCs underestimated the turnover
145time of bulk soil carbon and thus assimilated too much bomb ^{14}C (and/or had too little old soil
146carbon that would be depleted in radiocarbon).

147 ^{14}C -derived mean ages indicates that organic carbon soils is often thousands of years old
148(12-14, 21), which is an order of magnitude older than suggested by ESM turnover parameters.
149This discrepancy is likely a consequence of incomplete representation of key biogeochemical
150processes and difficulties in developing accurate parameterizations for soil carbon at a global
151scale. Most ESMs do not account for stabilization mechanisms whereby mineral interactions and

152 aggregate formation protect soil organic matter from decomposition over centuries to millennia
153 (13, 25-28). Moreover, first-order decay, as represented in ESMs, may not capture the response
154 of mineral-stabilized carbon to changes in soil moisture, temperature, and other conditions (29-
155 31). In addition, some ESM turnover parameters are based on laboratory incubation studies,
156 which are often biased fast compared to *in situ* decomposition rates (32). Finally, this set of
157 ESMs did not explicitly resolve vertical differences in soil organic matter dynamics, which may
158 cause underestimation of turnover times in deep soils with large carbon stocks (21, 25, 33, 34).

159 Because the turnover times derived from ESMs were inconsistent with ^{14}C observations,
160 we optimized the turnover parameters by fitting our RC models to the observations. We could
161 then run the optimized RC models to re-evaluate ~~21st-century~~ soil carbon storage for the transient
162 1% yr⁻¹ simulations. For this inverse approach, we optimized RC model parameters in each grid
163 cell containing an observation site (Fig 2g, 2h, S8, S9). We optimized the τ of the slowest pool
164 and the corresponding transfer coefficient into this pool based on the ^{14}C observations
165 ~~observations~~ while holding soil inputs and τ for the faster pools at their ESM-derived values. The
166 size of the slowest carbon pool was ~~also~~ constrained by optimizing the turnover time and the
167 transfer coefficient together using both ^{14}C and total carbon ~~and ^{14}C~~ . Consequently the optimized
168 RC model had about the same total carbon stock as the original ESM, thereby maintaining
169 consistency with carbon inventory data. This optimization approach yielded τ_{slow} values of
170 3700 ± 2800 yrs for GFDL and 3500 ± 1300 yrs for HadGEM (using two-pool RC models), which
171 were 16-17 times greater than the turnover times derived from the original ESMs.

172 For ESMs that included a passive pool, the optimization process yielded three distinct
173 outcomes. For CESM, which has the largest passive pool (73% of soil carbon), the optimized
174 τ_{passive} was 4500 yrs, which was 3.7 ± 1.5 times greater than τ_{passive} derived from the original model

(Table 1). IPSL has a smaller fraction of passive carbon (46%) and therefore required a greater τ_{passive} (16,500 yrs) to obtain agreement with the observed $\Delta^{14}\text{C}$. For MRI, the passive pool size was too small (only 13% of soil carbon) to bring $\Delta^{14}\text{C}$ into alignment with the profile observations even after parameter optimization (Fig S10, Table S5). To adjust for MRI's potential bias in the passive pool size, we optimized r_f together with τ_{passive} and r_s to allow for simultaneous changes in slow and passive pool sizes. The resulting RC model for MRI was able to match observations (Fig 2 g,h) with a passive pool fraction of 48% (see Methods; Table S5). These results indicated that increasing the size and turnover time of the passive pool in ESMs would improve agreement with ^{14}C -based mean age estimates. In general, increasing the size and turnover time of the passive pool in ESMs would improve agreement with ^{14}C -based age estimates.

Bringing turnover time and carbon transfer parameters into agreement with ^{14}C observations had significant consequences for the magnitude of the carbon-concentration feedback. Using the ^{14}C -based parameters, we conducted global transient simulations with each of the five RC models. These simulations showed that the soil as a whole (specifically the slow and passive pools) stored much less carbon in response to increasing levels of atmospheric CO_2 , primarily as a consequence of reduced flow into the slow or passive pool. The soil carbon sink decreased from $32 \pm 18\%$ to $18 \pm 12\%$ (Table 1), corresponding to an absolute sink reduction of $170 \pm 127 \text{ Pg C}$ (Fig 3). Relative to the ESMs, these simulations showed much less soil carbon accumulation in response to increasing levels of atmospheric CO_2 because of lower inputs to the slow and/or passive pools. The soil carbon sink decreased from $32 \pm 18\%$ to $18 \pm 12\%$ (Table 1), corresponding to an absolute sink reduction of $170 \pm 127 \text{ Pg C}$ (Fig 3). The magnitude of the soil

197sink reduction varied widely across the different models; those with larger and older passive
198fractions at the onset of the transient simulation (Table 1) generally had smaller sink reductions.

199 To assess the robustness of these sink reductions, we conducted a series of sensitivity
200experiments (see supplementary material). We found that the sink reduction imposed by
201constraining the models with ^{14}C observations ~~is~~was robust to (1) turnover times optimized
202specifically for different biomes; (2) spatial variation ~~and magnitude of~~in soil carbon stocks; and
203(3) variations in $\delta^{14}\text{C}$ across measurement sites (Table 2, S6). Sink reductions declined by a
204factor of 2 when the models were fit to an inventory that was 50% larger than the HSWD dataset,
205suggesting that if soil carbon pools were larger in ESMs, ^{14}C -imposed sink reductions would be
206lower (35). Lastly, we used our RC model approach to analyze four fully-coupled ESM runs
207(1pctCO₂) to address potential interactions between the carbon-climate and the carbon-
208concentration feedback. ^{14}C constraints still reduced the sink by at least 40% on average (Fig
209S11, Table S7) in the fully coupled simulations (see supplementary material).

210 We conclude that ~~CMIP5 current~~ESMs underestimated the mean age of soil carbon,
211especially for slow-cycling pools. By adjusting the turnover times of slow and passive pools to
212bring the models into alignment with ^{14}C observations, the potential for future soil carbon
213sequestration declined by $40 \pm 27\%$. If turnover times of slow and passive pools are adjusted to
214bring the ESMs into alignment with ^{14}C observations, the potential for 21st-century soil carbon
215sequestration declines by $40 \pm 27\%$ in the ESMs we evaluated. Although long-lived soil carbon
216pools consistent with old ^{14}C ages ~~imply~~imply increased a similar potential for carbon storage at
217steady state, the timescale required to reach equilibrium is too long to mitigate the potentially
218damaging climate effects of rising CO₂ concentrations during the 21st century (Fig S2). These
219findings emphasize the need to incorporate ^{14}C and other diagnostics into ESM development and

220evaluation. In addition, models require better representation of long-term mechanisms of soil
221carbon stabilization such as organic matter-mineral interactions. Considered together with
222potential nutrient limitation of NPP inputs to soil (36), our analysis suggests that the
223climatecarbon-concentration feedback may be weaker in the 21st century than currently expected
224from ESMs. Therefore a greater fraction of CO₂ emissions than previously thought could remain
225in the atmosphere and contribute to global warming.

226References and Notes:

2271. P. Friedlingstein *et al.*, Uncertainties in CMIP5 ~~Climate-climate~~ [Projections-projections](#) due to
228 ~~Carbon-carbon~~ [Cycle-cycle](#) ~~Feedbacksfeedbacks~~. *Journal of Climate* **27**, 511-526 (2013).
2292. IPCC, *Summary for Policymakers*. In: *Climate Change 2013: The Physical Science Basis*.
230 *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel*
231 *on Climate Change*. T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A.
232 Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.), Ed., (Cambridge, United Kingdom and New
233 York, NY, USA., 2013).
2343. V. K. Arora *et al.*, Carbon-~~Concentration-concentration~~ and ~~Carboncarbon~~-~~Climate-climate~~
235 ~~Feedbacks-feedbacks~~ in CMIP5 ~~Earth-earth~~ [System-system](#) ~~Modelsmodels~~. *Journal of Climate* **26**,
236 5289-5314 (2013).
2374. C. D. Jones, P. M. Cox, C. Huntingford, Climate-carbon cycle feedbacks under stabilization:
238 Uncertainty and observational constraints. *Tellus B* **58**, 603-613 (2006).
2395. K. E. O. Todd-Brown *et al.*, Changes in soil organic carbon storage predicted by ~~Earth-earth~~
240 [system models](#) during the 21st century. *Biogeosciences* **11**, 2341-2356 (2014).
2416. D. P. Van Vuuren *et al.*, The representative concentration pathways: an overview. *Climatic*
242 *Change* **109**, 5-31 (2011).
2437. J. Lichter *et al.*, Soil carbon sequestration in a pine forest after 9 years of atmospheric CO₂
244 enrichment. *Global Change Biology* **14**, 2910-2922 (2008).
2458. W. H. Schlesinger, J. Lichter, Limited carbon storage in soil and litter of experimental forest plots
246 under increased atmospheric CO₂. *Nature* **411**, 466-469 (2001).
2479. J. W. Harden, R. K. Mark, E. T. Sundquist, R. F. Stallard, Dynamics of soil carbon during
248 deglaciation of the Laurentide ice sheet. *Science* **258**, 1921-1924 (1992).
24910. W. H. Schlesinger, Evidence from chronosequence studies for a low carbon-storage potential of
250 soils. *Nature* **348**, 232-234 (1990).
25111. B. O'brien, J. Stout, Movement and turnover of soil organic matter as indicated by carbon isotope
252 measurements. *Soil Biology and Biochemistry* **10**, 309-317 (1978).
25312. J. B. Gaudinski, S. E. Trumbore, E. A. Davidson, S. H. Zheng, Soil carbon cycling in a temperate
254 forest: radiocarbon-based estimates of residence times, sequestration rates and partitioning of
255 fluxes. *Biogeochemistry* **51**, 33-69 (2000).
25613. M. S. Torn, S. E. Trumbore, O. A. Chadwick, P. M. Vitousek, D. M. Hendricks, Mineral control of
257 soil organic carbon storage and turnover. *Nature* **389**, 170-173 (1997).
25814. S. E. Trumbore, J. W. Harden, Accumulation and turnover of carbon in organic and mineral soils
259 of the BOREAS northern study area. ~~Journal of: Geophys~~~~sical: Research:~~ **102**, 28817-28830
260 (1997).
26115. N. Carvalhais *et al.*, Global covariation of carbon turnover times with climate in terrestrial
262 ecosystems. *Nature* **514**, 213-217 (2014).
26316. A. D. Friend *et al.*, Carbon residence time dominates uncertainty in terrestrial vegetation
264 responses to future climate and atmospheric CO₂. *Proceedings of the National Academy of*
265 *Sciences* **111**, 3280-3285 (2014).
26617. X. Wang *et al.*, Soil respiration under climate warming: differential response of heterotrophic and
267 autotrophic respiration. *Global Change Biology* **20**, 3229-3237 (2014).
26818. W. Knorr, I. C. Prentice, J. I. House, E. A. Holland, Long-term sensitivity of soil carbon turnover
269 to warming. *Nature* **433**, 298-301 (2005).
27019. C. D. Koven *et al.*, Controls on terrestrial carbon feedbacks by productivity vs. turnover in the
271 CMIP5 Earth System Models. *Biogeosciences-Discuss.* **12**, ~~5757-5804~~~~5211-5228~~ (2015).
27220. K. E. Taylor, R. J. Stouffer, G. A. Meehl, An overview of CMIP5 and the experiment design.
273 *Bulletin of the American Meteorological Society* **93**, 485-498 (2012).

27421. J. A. Mathieu, C. Hatté, J. Balesdent, É. Parent, Deep soil carbon dynamics are driven more by
275 soil type than by climate: a worldwide meta-analysis of radiocarbon profiles. *Global Change*
276 *Biology* **21**, 4278-4292 (2015).
27722. S. Trumbore, Age of ~~Soil-soil Organic-organic Matter-matter~~ and ~~Soil-soil Respirationrespiration~~:
278 ~~Radiocarbon-radiocarbon Constraints-constraints~~ on ~~Belowground-belowground Carbon-carbon~~
279 ~~Dynamicsdynamics~~. *Ecological Applications* **10**, 399-411 (2000).
28023. W. Feng et al., Methodological uncertainty in estimating carbon turnover times of soil fractions.
281 *Soil Biology and Biochemistry* **100**, 118-124 (2016).
28224. E. Shevliakova et al., Carbon cycling under 300 years of land use change: Importance of the
283 secondary vegetation sink. *Global Biogeochemical Cycles* **23**, (2009).
28425. M. W. Schmidt et al., Persistence of soil organic matter as an ecosystem property. *Nature* **478**, 49-
285 56 (2011).
28626. M. v. Lützow et al., Stabilization of organic matter in temperate soils: mechanisms and their
287 relevance under different soil conditions—a review. *European Journal of Soil Science* **57**, 426-445
288 (2006).
28927. J. A. Dungait, D. W. Hopkins, A. S. Gregory, A. P. Whitmore, Soil organic matter turnover is
290 governed by accessibility not recalcitrance. *Global Change Biology* **18**, 1781-1796 (2012).
29128. M. Kleber, R. Mikutta, M. S. Torn, R. Jahn, Poorly crystalline mineral phases protect organic
292 matter in acid subsoil horizons. *European Journal of Soil Science* **56**, 717-725 (2005).
29329. J. S. Clemente, A. J. Simpson, M. J. Simpson, Association of specific organic matter compounds
294 in size fractions of soils under different environmental controls. *Organic Geochemistry* **42**, 1169-
295 1180 (2011).
29630. C. Koven et al., The effect of vertically resolved soil biogeochemistry and alternate soil C and N
297 models on C dynamics of CLM4. *Biogeosciences* **10**, 7109-7131 (2013).
29831. B. Ahrens, M. C. Braakhekke, G. Guggenberger, M. Schrumpf, M. Reichstein, Contribution of
299 sorption, DOC transport and microbial interactions to the ¹⁴C age of a soil organic carbon profile:
300 Insights from a calibrated process model. *Soil Biology and Biochemistry* **88**, 390-402 (2015).
30132. G. B. Bonan, M. D. Hartman, W. J. Parton, W. R. Wieder, Evaluating litter decomposition in earth
302 system models with long-term litterbag experiments: an example using the Community Land
303 Model version 4 (CLM4). *Global Change Biology* **19**, 957-974 (2013).
30433. C. R. Lawrence, J. W. Harden, X. Xu, M. S. Schulz, S. E. Trumbore, Long-term controls on soil
305 organic carbon with depth and time: A case study from the Cowlitz River Chronosequence, WA
306 USA. *Geoderma* **247**, 73-87 (2015).
30734. C. Rumpel, I. Kögel-Knabner, Deep soil organic matter—a key but poorly understood component
308 of terrestrial C cycle. *Plant and Soil* **338**, 143-158 (2011).
30935. Y. Luo et al., Toward more realistic projections of soil carbon dynamics by Earth system models.
310 *Global Biogeochemical Cycles* **30**, 40-56 (2016).
31136. W. R. Wieder, C. C. Cleveland, W. K. Smith, K. Todd-Brown, Future productivity and carbon
312 storage limited by terrestrial nutrient availability. *Nature Geoscience* **8**, 441-U435 (2015).

313**Acknowledgments:**

314We thank ~~Lesego Khomo and Oliver Chadwick for use of unpublished data and~~ **C. Hatté** for
315sharing her compilation of published ^{14}C profiles. We received funding support from the Climate
316and Environmental Sciences Division of Biological and Environmental Research (BER) in the
317U.S. Department of Energy Office of Science. This included support from the Regional and
318Global Climate Modeling Program to the Biogeochemical Cycles Feedbacks Science Focus Area
319and several grants from the Terrestrial Ecosystem Science Program (DESC0014374 and DE-
320AC02-05CH11231). This study was supported by the U.S. Department of Energy, Office of
321Science, Office of Biological and Environmental Research, Regional and Global Climate
322Modeling (Biogeochemical Cycles Feedbacks Science Focus Area) and Terrestrial Ecosystem
323Science (DESC0014374 and contract DE-AC02-05CH11231) Programs in the Climate and
324Environmental Sciences Division of Biological and Environmental Research (BER) in the U.S.
325Department of Energy Office of Science. The model simulations analyzed in this study were
326obtained from the Earth System Grid Federation CMIP5 online portal hosted by the Program for
327Climate Model Diagnosis and Intercomparison at Lawrence Livermore National Laboratory
328(<https://pcmdi.llnl.gov/projects/esgf-llnl/>).

329

330

331

332

333

Table 1: Global soil carbon stocks and carbon uptake for CMIP5 models that experienced a quadrupling of atmospheric CO₂ from a preindustrial value of 285 ppm over a period of 140 years.

ESM	Initial SOC (Pg C)	% change in SOC	% change in SOC after ¹⁴ C constraint	¹⁴ C- imposed sink reduction (%)	τ_{slow} (yr) ¹	τ_{passive} (yr)	r_f	r_s	¹⁴ C- imposed correction factors ²			
									τ_{slow}	τ_{passive}	r_f	r_s
CESM1(BGC)	571	6.3	5.1	19	56±16	1310±241	0.06±0.05	0.33±0.05	-	3.7±1.5	-	0.34±0.75
GFDL-ESM2M	1344	26	3.3	87	231±196	-	0.17±0.07	-	16±18	-	0.06±0.14	-
HadGEM2-ES	1028	63	33	46	208±84	-	0.12±0.07	-	17±12	-	0.07±0.32	-
IPSL-CM5A-LR	1340	27	25	5.9	218±82	1181±347	0.06±0.03	0.29±0.07	-	14±8.3	-	0.07±0.14
MRI-ESM1 ³	1403	36	22	40	347±117	1065±257	0.17±0.09	0.10±0.06	-	13±7.2	0.46±0.79	0.34±0.74
Mean⁴	1137±312	32±18	18±12	40±27	212±104	1185±123	0.12±0.06	0.24±0.12	16.5±0.5	10.2±4.6	-	-

¹ τ_{slow} , τ_{passive} denote the turnover time, and r_f , r_s denote the transfer coefficient from the fast to the slow pool, and from the slow to the passive pool respectively. Reported values were estimated as an area-weighted mean and standard deviation of all model grid cells.

² The mean and standard deviation of the ¹⁴C-imposed correction factors were derived from using the ¹⁴C observations at each site in a single optimization, and then averaging these scalar adjustments across the set of 157 optimizations.

³ The ¹⁴C-constrained sink reduction and correction factor for MRI were based on an inverse analysis that changed the pool size of both slow and passive pools. The reported percent change in SOC and sink reduction were derived from transient simulations starting at steady state with the reduced complexity model. See methods in supporting material.

⁴ The multi-model mean and standard deviation were estimated using the mean value from each of the 5 ESMs.

344**Table 2:** Summary of sensitivity experiments.

Experiment	% change in SOC after ^{14}C constraint ¹	^{14}C - imposed sink reduction (%) ¹	Correction factor for turnover time ¹	Correction factor for transfer coefficient ¹
Biome-specific inversions	17±11	43±24	-	-
Match SOC with HWSD at sites ²	18±12	31±40	13±4.5	0.19±0.23
Match SOC with 1.5*HWSD at sites ²	21±12	19±42	11±4.5	0.38±0.39
-1 S.D. of inter- site variation	14±9.9	52±23	-	-
+1 S.D. of inter- site variation	23±16	25±25	-	-

345¹ The mean and standard deviation were estimated from the global mean change of each of the 5
346individual ESMs. The correction factors for the turnover time and transfer coefficients are
347reported for the slowest carbon pool.

348² The correction factors were obtained at each site, and then the mean scalar across all sites was
349applied to the global forward simulation.

350

351**Fig 1:** Location of radiocarbon soil profiles used to constrain ESM soil carbon mean ages and
 352turnover times (N=157). The carbon-weighted $\Delta^{14}\text{C}$ to a depth of 1m is denoted with the color
 353shade of each symbol. A summary of the location, sample year, and reference for each site is
 354provided in Table S2.

355

356**Fig 2:** Soil organic carbon content (a, b) of the original ESMs, $\Delta^{14}\text{C}$ of the reduced complexity
 357model optimized to the original ESMs (c, d), corresponding mean age (e, f), and the $\Delta^{14}\text{C}$ of the
 358 ^{14}C -constrained reduced complexity models (g, h). Left column shows the values of the models
 359sampled at the locations of the individual soil profiles; right column shows the global model
 360distribution. Data from profile sites and the Harmonized World Soil Database represent carbon

361content in the top 1 m of soil; data from ESMs are the total carbon stock. Star denotes the mean;
362the '+' symbol denotes outliers beyond the 25th and 75th percentiles.

364

365**Fig 3:** Absolute change in SOC content from the reduced complexity model fit to the original
 366ESM (bars with white background) and the estimate obtained by applying the ^{14}C constraint to
 367the reduced complexity model (bars with gray background). The estimates on the right side show
 368the total carbon content (sum of fast, slow, and passive) averaged across all the models, before
 369and after applying the radiocarbon constraint.

370**Supplementary Materials:**

371Materials and Methods

372Sensitivity Analysis Results

373Fully-Coupled Simulation Analysis

374Figures S1-S11

375Tables S1-S7

376References 37-1110