

## Pleistocene lagomorphs from Cathedral Cave, Nevada

CHRISTOPHER N. JASS

Department of Geological Sciences, Jackson School of Geosciences, University of Texas at Austin, Austin, TX, USA 78712; and Vertebrate Paleontology Laboratory, University of Texas at Austin, J.J. Pickle Research Campus, 10100 Burnet Road, Building 6, Austin, TX, USA 78758. Current Address: Royal Alberta Museum, 12845 - 102 Ave., Edmonton, AB, Canada T5N 0M6; Chris.Jass@gov.ab.ca

Faunal data from Cathedral Cave, Nevada, provide insight into biotic changes that occurred within the Great Basin prior to the latest Pleistocene. Taxonomic identifications of lagomorphs from Cathedral Cave were made using a morphological approach intended to minimize geographic and temporal assumptions. Although this approach to identification is conservative, the resultant data set is appropriate for inclusion in future analyses of regional biotic change. Lagomorphs recovered from the site include new regional records of two extinct taxa, *Aztlanolagus agilis* and *Brachylagus coloradoensis*. Other lagomorphs from Cathedral Cave include *Brachylagus idahoensis*, *Ochotona* sp., and *Sylvilagus* or *Lepus* sp. The presence of a posterorinternal reentrant fold on the p3 of some specimens of *Ochotona* sp. suggests that the range of variation present in the individual teeth of pikas needs to be described in further detail. In contrast to a previously established hypothesis of increasing enamel complexity in the p4 of *Aztlanolagus agilis*, evaluation of crenulation patterns of *Aztlanolagus agilis* from Cathedral Cave showed no distinct trends.

### INTRODUCTION

A significant body of research on the fossil lagomorphs of the Great Basin is currently available in the literature, primarily as a result of efforts to understand mammalian biogeographic patterns in the region from the late Pleistocene to Recent. Several lagomorphs, including rabbits [*Sylvilagus nuttalli* (Nuttall's cottontail), *Lepus townsendii* (white-tailed jackrabbit)] and pikas [*Ochotona princeps* (North American pika)] were included in Brown's (1978) classic study on montane mammal biogeography in the Great Basin. As a result, subsequent research continued to emphasize the importance of Great Basin lagomorphs in biogeographic studies in the region (Mead et al. 1982, Thompson and Mead 1982, Mead 1987, Grayson 1993, Grayson and Livingston 1993, Hafner 1993, Mead and Spaulding 1995, Mead and Grady 1996, Grayson 2000a, 2000b, 2005, 2006). Here I present a systematic account of the lagomorphs from Cathedral Cave, Nevada, including the first Great Basin records for the extinct species *Aztlanolagus agilis* Russell and Harris and *Brachylagus coloradoensis* Ramos.

Cathedral Cave (CC) is located on a north-facing slope at the mouth of Smith Creek Canyon, Nevada, in the east-central Great Basin (Fig. 1). Preliminary fieldwork in 1989 tested the paleontological significance of the site (Bell 1990). Initial age estimates for portions of the excavated fauna were maximally between  $15,000 \pm 200$  yr and  $24,600 \pm 2100$  yr B.P. and were based on uranium-series analyses of a flowstone fragment attached to a small stalagmite and a whole bone fragment from different arbitrary excavation levels (Bell 1990). Subsequent analyses of the fauna and new research in the same region of the cave indicate an older age for the deposit (Bell 1995, Bell and Barnosky 2000, Bell et al. 2004b, Jass 2005, 2007). The most recent age data, based on uranium-series analyses of fine samples of *in situ* flowstone,

place lower portions of the deposit in a maximum age range of  $146.02 \pm 2.584$  ka to  $151.2 \pm 4.4$  ka (Jass 2007). This age range places the fauna in a chronologically older context relative to other regional faunal assemblages. Specifically, faunal data from Cathedral Cave potentially provides insight into biotic changes that occurred within the Great Basin prior to the latest Pleistocene.




Figure 1. The geographic location of Cathedral Cave, NV.

## MATERIALS AND METHODS

The lagomorphs described here were collected during a new excavation conducted at Cathedral Cave in 2003. The excavation consisted of a 1.5 m x 2 m area of undisturbed sediments that were removed to a maximum depth of 110 cm below an arbitrary datum, or roughly 76 cm below the surface of the deposit. The excavation was stopped where large blocks of roof spall were interpreted as being situated directly on the bedrock floor of the cave. Excavation levels were hand dug using trowels and brushes in arbitrary 5-cm intervals except where natural sedimentary levels could be removed as distinct units. Unique horizontal (e.g., 1N 2E) and vertical (e.g., 45–50 cm below datum) provenience data were recorded for all excavated sediments and fossils.

Screen-washing of sediments was conducted at the Vertebrate Paleontology Laboratory of the Texas Natural Science Center and the Laboratory of Quaternary Paleontology at Northern Arizona University. All sediments were washed through nested pairs of screens (3.175 mm- and 0.7 mm-mesh). Washed sediments were air dried prior to being sorted for fossils. Additional details about the excavation and screen-washing methods employed in this study are presented elsewhere (Jass 2007).

The composite assemblage of 183 lagomorph fossils presented here represents 50 stratigraphic levels within the excavation. Specimens are housed at the Vertebrate Paleontology Laboratory of the Texas Memorial Museum (TMM) under locality number 43693, each with a unique specimen number (e.g., TMM 43693-1).

The majority of lagomorph specimens from Cathedral Cave consist of isolated teeth. Patterns of evolution and taxonomic identification in lagomorphs are usually based at least partially on changes in the enamel pattern of the p3 (e.g., White 1984, 1987, 1991a, 1991b). This study focuses on isolated p3s, dentaries that retain the p3, and a few edentulous dentaries that retain characteristics that allow for reliable taxonomic identification.

The lagomorph p3 is characterized by patterns of enamel-banded dentine that range from simple to complex, and cementum-filled re-entrant folds. The enamel folds in many leporid teeth appear wrinkled or crenulated and the degree of crenulation is sometimes used for identification, particularly for extinct taxa (Ramos 1999a). A binocular microscope with an ocular micrometer was used to examine the specimens and determine maximum occlusal length and width measurements.

## SYSTEMATIC PALEONTOLOGY

The following summary of the lagomorphs from Cathedral Cave, Nevada, is based on results of the 2003 excavation. Taxonomy and classification above the species level follows McKenna and Bell (1997).

ORDER: LAGOMORPHA Brandt, 1855

FAMILY: OCHOTONIDAE Thomas, 1897

GENUS: *Ochotona* Link, 1795


*Ochotona* sp.

**Referred Specimens**—See Appendix 1.

**Description**—The p3s of *Ochotona* sp. from Cathedral Cave exhibit some variation in size (Fig. 2) and are characterized by the presence of at least three enamel reentrant folds (anteroventral, anteroexternal, and posteroexternal; Figs. 3a,b). The folds are simple and lack the crenulated enamel patterns seen in many leporid species. The degree to which enamel reentrant folds are developed varies in some specimens and depends on how far the enamel penetrates from the margin towards the midline of the tooth. Some specimens (e.g., TMM 43693-1857) display weak development (i.e., little or indistinct enamel penetration towards the midline) of the anteroventral reentrant, as in the '*Ochotona* sp. near Trout Cave form' from Porcupine Cave, Colorado (Mead et al. 2004). Several specimens retain at least some development of a fourth, posteroventral reentrant fold. A strongly developed posteroventral reentrant fold, in which there is distinct penetration of enamel towards the midline, is apparent in both occlusal and ventral views of specimen TMM 43693-1856 (Fig. 3c).

*Ochotona* dentaries lacking the p3 were identified on the basis of diminutive size and general shape similarity to those of modern pikas. Whenever possible, the shape of the m3 or m3 alveolus was evaluated because it is triangular in leporids, but anteroposteriorly short in ochotonids (Heaton 1985). Dentaries of *Ochotona* sp. from Cathedral Cave that preserve the m3 or m3 alveolus retain a flattened posterior margin that gives the tooth and alveolus a more rectangular appearance than those of leporids.

**Discussion**—Two species of pika, *O. collaris* and *O. princeps* inhabit North America today. *Ochotona collaris* is geographically restricted to portions of Alaska and northwestern Canada, while *O. princeps* inhabits portions of southwestern Canada and the western United States (Wilson and Ruff



**Figure 2.** Scatter plot of length versus width in specimens of *Ochotona* sp. (n=74) from Cathedral Cave, NV. Measurements were taken to nearest 0.1 mm using an ocular micrometer. See Appendix 1 for individual specimen data.

1999). Fossil specimens, mainly attributed to *O. princeps*, are known from eastern and western portions of the United States and Canada (Mead 1987). Two extinct species of pika are also known from North America. *Ochotona spanglei* occurs in the late Miocene-early Pliocene of Oregon (Shotwell 1956), and the larger *O. whartoni* was found in the Pleistocene of Alaska and Canada (Guthrie and Matthews 1971, Mead and Grady 1996). Other morphological variants are recognized in the fossil record but their taxonomic status is uncertain (Mead et al. 2004).

The average length of the p3 in Cathedral Cave specimens is 1.23 mm ( $n = 75$ ;  $SD = 0.11$ ; range = 1.0–1.5) and the average width is 1.32 mm ( $n = 74$ ;  $SD = 0.16$ ; range = 1.0–1.7). Although there are size differences among individual specimens, the scatterplot of measurements for the p3 (i.e., length versus width; Fig. 2) is not clearly bi- or polymodal. This might suggest the presence of a single species exhibiting intraspecific variation at Cathedral Cave. However, Mead and Grady (1996) presented scatterplots (length versus width) of the m1 and P3 in extant and fossil pikas that show species “clusters”. If this result can be extrapolated to the p3s from Cathedral Cave, several species might be represented by the size range found here.

The amount of variation occurring in quantitative and qualitative morphologic characters of the p3 in fossil and extant pikas needs further study. There is some indication that morphological variants occur in the fossil record (e.g., Mead and Grady 1996, Mead et al. 2004), but the taxonomic level at which that variation is expressed remains uncertain. It is possible that most, if not all, of the specimens from Cathedral Cave are *Ochotona princeps*; however, the presence of a few unique morphological variants suggests that other species might be represented in the fauna. Because our understanding of morphologic variation in the individual teeth of *Ochotona* is rudimentary, assigning the specimens from Cathedral Cave to *O. princeps* would, by default, be

partially based on the assumption that fossil *Ochotona* occurring within or near the modern range of *O. princeps* are that species. As noted elsewhere, assumptions about geography during the identification process may lead to circularity in subsequent hypotheses concerning regional faunal dynamics (e.g., biogeography; Bell et al. 2004a). Therefore, I restricted my identifications to *Ochotona* sp. but provide an illustration of a specimen (TMM 43693-1856) with a morphotype that might be outside of the natural range of variation in the p3 of *O. princeps* (Fig. 3c).

TMM 43693-1856 has a well-developed postero-internal fold that is unique among North American pikas examined in this study. Erbajeva's (1994) review of the phylogeny and evolution of ochotonids includes an illustration of a p3 of *Ochotona antiqua* that exhibits four reentrant folds as in TMM 43693-1856. *Ochotona antiqua* is known only from late Miocene-early Pliocene deposits in the Old World (Erbajeva 1994). I am hesitant to assign TMM 43693-1856 to any particular species prior to a thorough review of variation in the p3 of New and Old World species of *Ochotona*. The occurrence of this p3 morphology in Cathedral Cave specimens could represent a new species or a new record of an Old World species, or it may fall within the natural, but as yet undocumented, range of variation exhibited by extant North American pikas.

FAMILY: LEPORIDAE Fischer de Waldheim, 1817

GENUS: *Aztlanolagus* Russell and Harris, 1986


Type species: *Aztlanolagus agilis* Russell and Harris, 1986

*Aztlanolagus agilis* Russell and Harris, 1986


**Referred Specimens**—See Appendix 1.

**Description**—The p3s of *Aztlanolagus agilis* are characterized by the presence of five reentrant folds (Fig. 4a; Russell and Harris 1986). No extant North American leporid retains this pattern, and among fossil taxa only *Nekrolagus* is similar (Russell and Harris 1986). All specimens identified as *A. agilis* exhibit a strongly developed antero-internal fold and a well-developed postero-internal fold (Fig. 4b) or lake. The presence of a strongly developed antero-internal fold on the p3 distinguishes *A. agilis* from *Nekrolagus*, in which the fold is less developed or absent (Russell and Harris 1986). The degree of crenulation within individual folds varies among individual specimens from Cathedral Cave. Table 1 summarizes the number of major crenulations in the antero-external reentrant, along the posterior wall of the postero-internal reentrant, and along the posterior wall of the postero-external reentrant. For specimens of *A. agilis* from Cathedral Cave, the p3 average length is 2.40 mm ( $n = 25$ ;  $SD = 0.23$ ; range = 1.7–2.8) and average width is 2.29 mm ( $n = 25$ ;  $SD = 0.34$ ; range = 1.2–2.8). These are slightly larger than average measurements recorded by Winkler and Tomida (1988) for other localities in the southwestern United States.

**Discussion**—*Aztlanolagus* is known from Blancan to late Rancholabrean deposits in the western United States and Mexico. Records other than Cathedral Cave come from



**Figure 3.** The p3 of *Ochotona* from Cathedral Cave. **A.** Occlusal pattern and terminology discussed in text; **B.** Left p3 (TMM 43693-1814) of *Ochotona* sp. from Cathedral Cave; **C.** Right p3 of *Ochotona* sp. exhibiting a fourth reentrant fold (TMM 43693-1856). Scale bar equals 1 mm.



**Figure 4.** Leporid teeth from Cathedral Cave. A. Enamel pattern and terminology for the p3; B. Right p3 of *Aztlanolagus agilis* (TMM 43693-1750) from Cathedral Cave; C. Left p3 of *Brachylagus idahoensis* (TMM 43693-1777) from Cathedral Cave; D. Right p3 of *Brachylagus coloradoensis* (TMM 43693-1774) from Cathedral Cave. Scale bars equal 1 mm.

localities within Porcupine Cave, CO (Barnosky and Hopkins 2004, Baxter 2004), and localities in northern Mexico, New Mexico, Arizona, and Texas (Russell and Harris 1986, Winkler and Tomida 1988). Although Winkler and Tomida (1988) tentatively referred all specimens that they examined to *A. agilis*, Baxter (2004) reported Porcupine Cave specimens as *Aztlanolagus* spp. because of variation in tooth size and crenulation patterns.

*Aztlanolagus* might be synonymous with *Pliopentalagus*, an extinct leporid known mostly from China that has five reentrant folds on the p3 (Tomida and Jin 2002, 2004). However, until the argument for synonymy is formalized, *Aztlanolagus* should be retained. Taxonomic issues notwithstanding, the disappearance of *A. agilis* is one of the few known late Quaternary small mammal extinctions in continental North America.

The records of *Aztlanolagus agilis* from Cathedral Cave presented here represent its first known occurrence in the Great Basin. The only other specimen known from the Great Basin is unnumbered in bulk material from Smith Creek Cave, Nevada, housed at the Nevada State Museum (Jass *personal observation*).

GENUS: *Brachylagus* Miller, 1900

Type species: *Lepus idahoensis* Merriam, 1891

*Brachylagus idahoensis* (Merriam, 1891)

**Referred Specimens**—See Appendix 1.

**Description**—Three specimens of *Brachylagus idahoensis* were identified from Cathedral Cave (Fig. 4c). Two of the specimens retain simple anteroexternal reentrants and the trigonid and talonid are either completely (TMM 43693-1777), or nearly (TMM 43693-1776), separated by cementum. The extension of the anteroexternal folds across the tooth is characteristic of *B. idahoensis* (Ramos 1999b). Length and width of the p3s were: (TMM 43693-1777) -  $l=1.8$  mm,  $w=1.6$  mm; (43693-1776) -  $l=1.8$  mm,  $w=1.8$  mm.

A third p3 specimen (TMM 43693-1775) with a damaged occlusal surface that was also identified as *Brachylagus idahoensis* is unique. In ventral view, it has a posteroexternal reentrant that extends across two-thirds of the tooth, but exhibits an enamel lake in the position of a posterointernal reentrant. Because the joining of the posteroexternal reentrant with the enamel lake would result in a morphology more similar to *B. idahoensis* than *B. coloradoensis* (see below), the specimen is identified as the former. Further review of variation in the p3 enamel patterns of *B. idahoensis* is needed.

**Discussion**—*Brachylagus* is distinguished from most extant and fossil North American leporids by the lack of an anterior reentrant on the p3 (Kurtén and Anderson 1980, Ramos 1999a). The extinct genus *Hypolagus* is similar in some qualitative aspects to an extinct form of *Brachylagus* (*B. coloradoensis*) but is larger in size (Ramos 1999b). Until recently, the genus *Brachylagus* was considered monotypic, with *B. idahoensis* as the only recognized species (Ramos 1999b).

Fossils of *Brachylagus idahoensis* are known only from the western United States and most fall within or near the present distribution of the species (Kurtén and Anderson 1980). Exceptions are the records from Isleta Cave No. 2 and Sheep Camp Shelter in New Mexico (Harris 1993). The occurrence of *B. idahoensis* at Cathedral Cave was previously noted by Ramos (1999b).

*Brachylagus coloradoensis* Ramos 1999

**Referred Specimens**—See Appendix 1.

**Description**—The two specimens identified as *Brachylagus coloradoensis* are similar to *B. idahoensis*, but both have a weakly developed posterointernal reentrant and a posteroexternal reentrant that extends only between one-third and two-thirds across the tooth (Fig. 4d). As such, these specimens are consistent with those identified as *B. coloradoensis* from Porcupine Cave (Ramos, 1999b).

**Discussion**—Prior to this report, *B. coloradoensis* was known only from several localities of Irvingtonian or unknown age within Porcupine Cave (Ramos 1999b, Barnosky 2004, Barnosky and Hopkins 2004, Baxter, 2004; age assignments from Barnosky et al. 2004). Its occurrence in Cathedral Cave extends both its geographic and chronologic ranges.

GENUS: *Sylvilagus* Gray 1867 or *Lepus* Linnaeus 1758

*Sylvilagus* or *Lepus* sp.

**Referred Specimens**—See Appendix 1.

**Description**—Specimens identified as *Sylvilagus* or *Lepus* sp. are characterized by the presence of three (anterior, anteroexternal, and posteroexternal) reentrant folds on the p3. The presence of the anterior fold distinguishes these specimens from *Ochotona*, which also typically retains three reentrant folds on the p3. In most specimens the anterior and anteroexternal reentrants are simple with little crenulation, while the posteroexternal reentrant fold may or may not exhibit crenulations. Length and width of the p3s are listed

in Appendix 1 and were graphed on a scatter plot used to evaluate if distinct size groupings exist (Fig. 5).

**Discussion**—*Sylvilagus audubonii*, *S. nuttalli*, *Lepus californicus*, and *L. townsendii* all have modern ranges that include the region surrounding Cathedral Cave. One, all, or none of these taxa may be represented in Cathedral Cave. Previous authors noted that there is considerable difficulty in the identification of *Lepus* versus *Sylvilagus* (e.g., Kurtén and Anderson 1980) and that size overlap occurs (Dalquest and Schultz 1992, Ramos 1999a). Because it would be necessary to invoke either geographic or evolutionary assumptions (i.e., constant size in the past relative to the modern) to separate these genera, no attempt was made to differentiate *Lepus* and *Sylvilagus* from Cathedral Cave (see discussion below).

### DISCUSSION


Bell's (1990) report on lagomorphs from Cathedral Cave included *Ochotona princeps*, *Brachylagus idahoensis* (as *Sylvilagus idahoensis*), *Sylvilagus* sp., and *Lepus* sp. The 2003 excavation resulted in the recognition of a more diverse lagomorph assemblage that includes new records of *B. coloradoensis* and *Aztlanolagus agilis*. The record of *A. agilis* is particularly notable because it consists of a reasonable sample size ( $n=25$ ) that might be used to further evaluate the variation in crenulation patterns of this taxon. Older specimens of *A. agilis* appear to have less-complex crenulation patterns than do more recent specimens (Winkler and Tomida 1988). My attempt to quantitatively evaluate crenulation patterns of the p3 in *A. agilis* from Cathedral Cave by counting the number and distinctness of major crenulations failed to show a pattern of increasing complexity from lower to upper excavated levels (Table 1). This does not negate the possibility of such a pattern over a broader time scale, but does provide a measure by which specimens from other localities might be compared. Nevertheless, the occurrence of *A. agilis* at Cathedral Cave provides an important record

for understanding Pleistocene lagomorph diversity in the Great Basin and western North America.

Fossil lagomorphs have had a prominent role in paleontological research in the Great Basin since the 1970s when Brown (1971, 1978) published his classic research on montane mammal biogeography. Those studies helped initiate a number of research projects on Great Basin lagomorphs (e.g., Grayson 1977, Mead 1987, Hafner 1993, Mead and Spaulding 1995, Osterhadt 1999), and continue to influence research on lagomorphs of the region (e.g., Grayson 2005). Great Basin lagomorph fossils are ideal for such studies because they are common constituents of many Pleistocene and Holocene vertebrate assemblages (e.g., Heaton 1985).

The taxonomic approach taken here is more conservative than those of prior studies of Great Basin lagomorphs so that taxa are identified based exclusively upon morphological criteria and, in turn, excluded from regional-scale biogeographic and evolutionary assumptions. Minimal use of geography as either an implicit or explicit character in the fossil identification process results in identifications that are more appropriate for broad biogeographic analyses (e.g., Bell et al. 2004a). The use of modern geographic range data to restrict possible identifications only to those species currently inhabiting or in close proximity to the Great Basin (e.g., *Ochotona princeps*, *Sylvilagus audubonii*, *S. nuttalli*, *Lepus californicus*, and *L. townsendii*) would foster circularity in biogeographic interpretations of the region. At present, there is not enough primary data on morphologic variation in the p3 to substantiate species-level identifications of pika (*Ochotona* sp.) and some leporids (*Sylvilagus* or *Lepus* sp.). Similarly, the invocation of size alone as a criterion for distinguishing species or genera (e.g., *Lepus* vs. *Sylvilagus*) assumes a pattern of stasis in body size relative to modern populations, where such an assumption may not be appropriate. For example, it is possible that certain species of *Sylvilagus* were larger 130,000 years ago and more comparable in size to extant *Lepus*. In the absence of other independent characters, the size of a particular taxon might simply reflect ecophenotypic variation. The identification of other lagomorph faunas and re-evaluation of previously described faunas under comparable, conservative identification criteria are needed to better understand how this mode of identification influences our understanding of broad biotic change in the Great Basin.

In summary, Cathedral Cave preserves the first Great Basin records of the extinct species *Aztlanolagus agilis* and *Brachylagus coloradoensis*. By virtue of their presence at Cathedral Cave, the records of these taxa in the Great Basin are limited to time periods preceding the last glacial episode, possibly as old as  $146.02 \pm 2.584$  ka to  $151.2 \pm 4.4$  ka. Other records from Cathedral Cave include *Ochotona* sp., *B. idahoensis*, and *Sylvilagus* or *Lepus* sp. Additional work focusing on variation in the p3 of lagomorphs is needed to determine if there are morphologic characters that support species-level identifications for taxa conservatively identified here as *Ochotona* sp. and *Sylvilagus* or *Lepus* sp. Future work



**Figure 5.** Scatter plot of length versus width in specimens identified as *Sylvilagus* or *Lepus* sp. ( $n=42$ ) from Cathedral Cave, NV. Measurements were taken to nearest 0.1 mm using an ocular micrometer. See Appendix 1 for individual specimen data.

**Table 1.** Variation in the number of major crenulations on the anteroexternal reentrant (AER), posterior wall of the posteroexternal reentrant (PW-PIR), and posterior wall of the posteroexternal reentrant (PW-PER) in the p3 of *Aztlanolagus agilis* from Cathedral Cave, Nevada, by excavated level. (?) = minimum number of major crenulations. Dash (-) = data could not be evaluated. Specimen No. = unique specimen number for locality TMM 43693 (i.e., TMM 43693-1748 is the full number for the first specimen listed below).

| Specimen No. | Excavated Level                               | AER | PW-PIR | PW-PER |
|--------------|-----------------------------------------------|-----|--------|--------|
| 1748         | 2N 3E 40-45 cmbd                              | 0   | 1      | 0      |
| 1749         | 2N 3E 40-45 cmbd                              | 0   | 2(?)   | 0      |
| 1750         | 1N 3E Cemented Level                          | 2   | 4      | 2      |
| 1751         | 1N 3E 43.5-55 cmbd; below Cemented Level      | —   | 2      | 2      |
| 1752         | 1N 3E 43.5-55 cmbd; below Cemented Level      | 0   | 4      | 3      |
| 1753         | 2N 2-3E 45-60 cmbd                            | 0   | 2      | 2      |
| 1754         | 2N 2-3E 45-60 cmbd                            | —   | —      | —      |
| 1755         | 2N 2-3E 45-60 cmbd                            | 0   | 2      | 2      |
| 1756         | 1N 2E 55-60 cmbd                              | —   | 2(?)   | 3      |
| 1757         | 1N 2E 60-65 cmbd                              | 1   | 2      | 0      |
| 1758         | 2N 2-3E 60-65 cmbd                            | 2   | 3      | 2      |
| 1759         | 2N 2-3E 60-65 cmbd                            | —   | 3      | 2      |
| 1760         | 1N 2E 65-70 cmbd                              | 0   | 0      | 1      |
| 1761         | 1N 2E 65-70 cmbd                              | 0   | 1      | 0      |
| 1486         | 1N 3E 65-70 cmbd                              | 2   | 4      | 3      |
| 1762         | 2N 2-3E 65-70 cmbd                            | 2   | 2      | 1      |
| 1763         | 2N 2-3E 65-70 cmbd                            | 0   | 4      | 3      |
| 1764         | 2N 2-3E 70-75 cmbd                            | 0   | 2      | 0      |
| 1765         | 2N 2-3E 70-75 cmbd                            | 0   | 2      | 0      |
| 1766         | 1N 2-3E 75-80 cmbd                            | 0   | 1      | 1      |
| 1767         | 2N 2-3E 75-80 cmbd                            | 0   | 2      | 2      |
| 1768         | 2N 2-3E 80-85 cmbd                            | 2   | 4      | —      |
| 1769         | 1N 2-3E 85-90 cmbd                            | 1   | 0      | 1      |
| 1770         | 1-2N 2-3E 87-102 cmbd; below pedasteled rocks | 1   | 1      | 1      |
| 1771         | 2N 2-3E 90-95 cmbd                            | 1   | 2      | 1      |
| 1772         | 2N 2-3E 90-95 cmbd                            | 1   | 0      | 1      |

will examine how data on lagomorphs from Cathedral Cave relate to broad patterns of biotic change in the region (e.g., montane island biogeography).

#### ACKNOWLEDGMENTS

Funding for this project came from the Geology Foundation of the Jackson School of Geosciences at UT-Austin, a Stephen J. Gould Student Research Grant from the Paleontological Society, and a Student Research Award from the Texas Academy of Science. I thank my dissertation committee (C. Bell, T. Rowe, J. Mead, J. Sprinkle, and J. Kappelman) and E. Lundelius for comments on early drafts of this manuscript, and the following people for field and laboratory assistance: C. Bell, J. Mead, S. Swift, M. Osborne, N. Czaplewski, J. Meyers, M. Poteet, A. Mead, D. Balakrishnan, T. Eiting, W. Jass, L. Nairn, L. Naski, P. Owen, J. Pape, D. Sanders, G. Tembe, K. Urban, and M. Wickwire. I am also grateful to K. Finger, B. Kraatz, and an anonymous reviewer for critical reviews.

#### LITERATURE CITED

- Barnosky, A.D. 2004. A summary of fossilized species in Porcupine Cave. Pp. 95–116 *in* A.D. Barnosky (ed.). Biodiversity Response to Climate Change in the Middle Pleistocene: The Porcupine Cave Fauna from Colorado. University of California Press, Berkeley, CA.
- Barnosky, A.D., C.J. Bell, R.G. Raynolds, and L.H. Taylor. 2004. The Pleistocene fossils of Porcupine Cave, Colorado: Spatial distribution and taphonomic overview. Pp. 6–26 *in* A.D. Barnosky (ed.). Biodiversity Response to Climate Change in the Middle Pleistocene: The Porcupine Cave Fauna from Colorado. University of California Press, Berkeley, CA.
- Barnosky, A.D., and S.B. Hopkins. 2004. Identification of miscellaneous mammals from the Pit locality: Including Soricidae, Leporidae, Geomyidae. Pp. 169–171 *in* A.D. Barnosky (ed.). Biodiversity Response to Climate Change in the Middle Pleistocene-The Porcupine Cave fauna from Colorado. University of California Press, Berkeley, CA.
- Baxter, C.N. 2004. Leporidae of the DMNH Velvet Room excava-

- tions and Mark's Sink. Pp. 164–168 in A.D. Barnosky (ed.). Biodiversity Response to Climate Change in the Middle Pleistocene: The Porcupine Cave Fauna from Colorado. University of California Press, Berkeley, CA.
- Bell, C.J. 1990. A Rancholabrean mammalian fauna from Cathedral Cave, White Pine County, Nevada. M.S. thesis. Northern Arizona University, Flagstaff, AZ.
- Bell, C.J. 1995. A middle Pleistocene (Irvingtonian) microtine rodent fauna from White Pine County, Nevada, and its implications for microtine rodent biochronology. *Journal of Vertebrate Paleontology* 15(3, Supplement):18A.
- Bell, C.J., and A.D. Barnosky 2000. The microtine rodents from the Pit Locality in Porcupine Cave, Park County, Colorado. *Annals of Carnegie Museum* 69:93–134.
- Bell, C.J., J.J. Head, and J.I. Mead. 2004a. Synopsis of the herpetofauna from Porcupine Cave. Pp. 117–126 in A.D. Barnosky (ed.). Biodiversity Response to Climate Change in the Middle Pleistocene: The Porcupine Cave Fauna from Colorado. University of California Press, Berkeley, CA.
- Bell, C.J., C.A. Repenning, and A.D. Barnosky. 2004b. Arvicoline rodents from Porcupine Cave—identification, spatial distribution, taxonomic assemblages, and biochronologic significance. Pp. 207–263 in A.D. Barnosky (ed.). Biodiversity Response to Climate Change in the Middle Pleistocene: The Porcupine Cave Fauna from Colorado. University of California Press, Berkeley, CA.
- Brown, J.H. 1971. Mammals on mountaintops: Nonequilibrium insular biogeography. *American Naturalist* 105:467–478.
- Brown, J.H. 1978. The theory of insular biogeography and the distribution of boreal birds and mammals. *Great Basin Naturalist Memoirs* 2:209–227.
- Dalquest, W.W., and G.E. Schultz. 1992. Ice Age Mammals of Northwestern Texas. Midwestern University Press, Wichita Falls, TX. 309 pp.
- Erbajeva, M.A. 1994. Phylogeny and evolution of Ochotonidae with emphasis on Asian Ochotonids. In Y. Tomida, C.-K. Li, and T. Setoguchi (eds.). Rodent and Lagomorph Families of Asian Origins and Diversification. *National Science Museum Monographs* 8:1–13.
- Grayson, D.K. 1977. On the Holocene history of some northern Great Basin lagomorphs. *Journal of Mammalogy* 58:507–513.
- Grayson, D.K. 1993. The Desert's Past: A Natural Prehistory of the Great Basin. Smithsonian Institution Press, Washington, DC. 356 pp.
- Grayson, D.K. 2000a. Mammalian responses to middle Holocene climatic change in the Great Basin of the western United States. *Journal of Biogeography* 27:181–192.
- Grayson, D.K. 2000b. The Homestead Cave mammals. In D.B. Madsen (ed.). Late Quaternary Paleocology in the Bonneville Basin. *Utah Geological Survey Bulletin* 130:67–89.
- Grayson, D.K. 2005. A brief history of Great Basin pikas. *Journal of Biogeography* 32:2101–2111.
- Grayson, D.K. 2006. The Late Quaternary biogeographic histories of some Great Basin mammals (western USA). *Quaternary Science Reviews* 25:2964–2991.
- Grayson, D.K., and S.D. Livingston. 1993. Missing mammals on Great Basin mountains: Holocene extinctions and current knowledge. *Conservation Biology* 7:527–532.
- Guthrie, R.D., and J.V. Matthews, Jr. 1971. The Cape Deceit fauna: Early Pleistocene mammalian assemblage from the Alaskan Arctic. *Quaternary Research* 1:474–510.
- Hafner, D.J. 1993. North American pika (*Ochotona princeps*) as a late Quaternary biogeographic indicator species. *Quaternary Research* 39:373–380.
- Harris, A.H. 1993. Quaternary vertebrates of New Mexico. In S.G. Lucas and J. Zidek (eds.). Vertebrate Paleontology in New Mexico. *New Mexico Museum of Natural History and Science Bulletin* 2:179–197.
- Heaton, T.H. 1985. Quaternary paleontology and paleoecology of Crystal Ball Cave, Millard County, Utah: With emphasis on mammals and description of a new species of fossil skunk. *Great Basin Naturalist* 45:337–390.
- Jass, C.N. 2005. A re-evaluation of the age assignment of Cathedral Cave, Nevada. *Journal of Vertebrate Paleontology* 25(3, Supplement):74A.
- Jass, C.N. 2007. New perspectives on Pleistocene biochronology and biotic change in the east-central Great Basin: An examination of the vertebrate fauna from Cathedral Cave, Nevada. Ph.D. diss. University of Texas at Austin, Austin, TX.
- Kurtén, B., and E. Anderson. 1980. Pleistocene Mammals of North America. Columbia University Press, New York, NY. 442 pp.
- McKenna, M.C., and S.K. Bell. 1997. Classification of Mammals Above the Species Level. Columbia University Press, New York, NY. 631 pp.
- Mead, J.I. 1987. Quaternary records of pika, *Ochotona*, in North America. *Boreas* 16:165–171.
- Mead, J.I., M. Erbajeva, and S.L. Swift. 2004. Middle Pleistocene (Irvingtonian) *Ochotona* (Lagomorpha: Ochotonidae) from Porcupine Cave. Pp. 155–163 in A.D. Barnosky (ed.). Biodiversity Response to Climate Change in the Middle Pleistocene—The Porcupine Cave Fauna from Colorado. University of California Press, Berkeley, CA.
- Mead, J.I., and F. Grady. 1996. *Ochotona* (Lagomorpha) from late Quaternary cave deposits in eastern North America. *Quaternary Research* 45:93–101.
- Mead, J.I., and W.G. Spaulding. 1995. Pika (*Ochotona*) and paleoecological reconstructions of the intermountain west, Nevada and Utah. In D.W. Steadman and J.I. Mead (eds.). Late Quaternary Environments and Deep History: A Tribute to Paul S. Martin. *The Mammoth Site of Hot Springs, South Dakota Scientific Papers* 3:165–186.
- Mead, J.I., R.S. Thompson, and T.R. Van Devender. 1982. Late Wisconsinan and Holocene fauna from Smith Creek Canyon, Snake Range, Nevada. *Transactions of the San Diego Society of Natural History* 20(1):1–26.
- Osterhadt, J. 1999. A review of Quaternary Lagomorpha of the Great Basin and Snake Creek Burial Cave, Nevada. M.S. thesis. Northern Arizona University, Flagstaff, AZ.
- Ramos, C.N. 1999a. Morphometric variation among some leporids (Mammalia: Lagomorpha) of North America. *Proceedings of*

- the Denver Museum of Natural History, Series 3, no. 16:1–12.
- Ramos, C.N. 1999b. An Irvingtonian species of *Brachylagus* (Mammalia: Lagomorpha) from Porcupine Cave, Park County, Colorado. *Great Basin Naturalist* 59:151–159.
- Russell, B.D., and A.H. Harris. 1986. A new leporine (Lagomorpha: Leporidae) from Wisconsinan deposits of the Chihuahuan Desert. *Journal of Mammalogy* 67:632–639.
- Shotwell, J.A. 1956. Hemphillian mammalian assemblage from northeastern Oregon. *Bulletin of the Geological Society of America* 67:717–738.
- Thompson, R.S., and J.I. Mead. 1982. Late Quaternary environments and biogeography in the Great Basin. *Quaternary Research* 17:39–55.
- Tomida, Y., and C. Jin. 2002. Morphological evolution of the genus *Pliopentalagus* based on the fossil material from Anhui Province, China: A preliminary study. In T. Kubodera, M. Higuchi, and R. Miyawaki (eds.). Proceedings of the 3<sup>rd</sup> and 4<sup>th</sup> Symposia on Collection Building and Natural History Studies in Asia and the Pacific Rim. *National Science Museum Monographs* 22:97–107.
- Tomida, Y., and C. Jin. 2004. *Aztlanolagus* (Lagomorpha, Mammalia) revisited: Origin, migration, evolution, and taxonomy. *Journal of Vertebrate Paleontology* 24 (3, Supplement):121A–122A.
- White, J.A. 1984. Late Cenozoic Leporidae (Mammalia, Lagomorpha) from the Anza-Borrego Desert, southern California. In R.M. Mengel (ed.). Papers in Vertebrate Paleontology Honoring Robert Warren Wilson. *Carnegie Museum of Natural History Special Publication* 9:41–57.
- White, J.A. 1987. The Archaeolaginae (Mammalia: Lagomorpha) of North America, excluding *Archaeolagus* and *Panolax*. *Journal of Vertebrate Paleontology* 7:425–450.
- White, J.A. 1991a. North American Leporinae (Mammalia: Lagomorpha) from late Miocene (Clarendonian) to latest Pliocene (Blancan). *Journal of Vertebrate Paleontology* 11:67–89.
- White, J.A. 1991b. A new *Sylvilagus* (Mammalia: Lagomorpha) from the Blancan (Pliocene) and Irvingtonian (Pleistocene) of Florida. *Journal of Vertebrate Paleontology* 11:243–246.
- Wilson, D.E., and S. Ruff, eds. 1999. The Smithsonian Book of North American Mammals. Smithsonian Institution Press, Washington, DC. 750 pp.
- Winkler, A.J., and Y. Tomida. 1988. New records of the small leporid *Aztlanolagus agilis* Russell and Harris (Leporidae: Leporinae). *Southwestern Naturalist* 33:391–396.

## APPENDIX 1

Identified specimens of lagomorphs collected during the 2003 excavation at Cathedral Cave, Nevada, by taxon and provenience. Measurements of length (*l*) and width (*w*) of the third premolar (p3) are presented in millimeters to the nearest 0.1 mm. The column headed by the number symbol (#) contains unique specimen numbers associated with Cathedral Cave locality number TMM 43693 (i.e., TMM 43693-1748 is the full number for the first specimen listed below).

| #    | TAXON                      | SIDE | ELEMENT | <i>l</i> | <i>w</i> | PROVENIENCE                                   |
|------|----------------------------|------|---------|----------|----------|-----------------------------------------------|
| 1748 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.1      | 2.1      | 2N 3E 40-45 cmbd                              |
| 1749 | <i>Aztlanolagus agilis</i> | R    | p3      | 2.4      | 2.7      | 2N 3E 40-45 cmbd                              |
| 1750 | <i>Aztlanolagus agilis</i> | R    | p3      | 2.6      | 2.5      | 1N 3E Cemented Level                          |
| 1751 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.6      | 2.3      | 1N 3E 43.5-55 cmbd; below Cemented Level      |
| 1752 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.6      | 2.4      | 1N 3E 43.5-55 cmbd; below Cemented Level      |
| 1753 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.5      | 2.1      | 2N 2-3E 45-60 cmbd                            |
| 1754 | <i>Aztlanolagus agilis</i> | R    | p3      | —        | —        | 2N 2-3E 45-60 cmbd                            |
| 1755 | <i>Aztlanolagus agilis</i> | R    | p3      | 2.3      | 2.3      | 2N 2-3E 45-60 cmbd                            |
| 1756 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.1      | 1.2      | 1N 2E 55-60 cmbd                              |
| 1757 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.6      | 2.4      | 1N 2E 60-65 cmbd                              |
| 1758 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.6      | 2.5      | 2N 2-3E 60-65 cmbd                            |
| 1759 | <i>Aztlanolagus agilis</i> | R    | p3      | 2.8      | 2.6      | 2N 2-3E 60-65 cmbd                            |
| 1760 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.3      | 2.1      | 1N 2E 65-70 cmbd                              |
| 1761 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.2      | 2.1      | 1N 2E 65-70 cmbd                              |
| 1486 | <i>Aztlanolagus agilis</i> | R    | p3      | 2.5      | 2.8      | 1N 3E 65-70 cmbd                              |
| 1762 | <i>Aztlanolagus agilis</i> | R    | p3      | 2.2      | 2.0      | 2N 2-3E 65-70 cmbd                            |
| 1763 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.5      | 2.3      | 2N 2-3E 65-70 cmbd                            |
| 1764 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.6      | 2.5      | 2N 2-3E 70-75 cmbd                            |
| 1765 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.4      | 2.3      | 2N 2-3E 70-75 cmbd                            |
| 1766 | <i>Aztlanolagus agilis</i> | R    | p3      | 1.7      | 1.7      | 1N 2-3E 75-80 cmbd                            |
| 1767 | <i>Aztlanolagus agilis</i> | R    | p3      | 2.5      | 2.5      | 2N 2-3E 75-80 cmbd                            |
| 1768 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.5      | 2.1      | 2N 2-3E 80-85 cmbd                            |
| 1769 | <i>Aztlanolagus agilis</i> | L    | p3      | 2.2      | 2.1      | 1N 2-3E 85-90 cmbd                            |
| 1770 | <i>Aztlanolagus agilis</i> | R    | p3      | 2.5      | 2.5      | 1-2N 2-3E 87-102 cmbd; below pedestaled rocks |

| #    | TAXON                            | SIDE | ELEMENT                           | <i>l</i> | <i>w</i> | PROVENIENCE                              |
|------|----------------------------------|------|-----------------------------------|----------|----------|------------------------------------------|
| 1771 | <i>Aztlanolagus agilis</i>       | R    | p3                                | 2.5      | 2.6      | 2N 2-3E 90-95 cmbd                       |
| 1772 | <i>Aztlanolagus agilis</i>       | R    | p3                                | 2.3      | 2.5      | 2N 2-3E 90-95 cmbd                       |
| 1773 | <i>Brachylagus coloradoensis</i> | L    | dentary<br>with p3-p4             | 1.6      | 1.6      | 1N 2E 65-70 cmbd                         |
| 1774 | <i>Brachylagus coloradoensis</i> | R    | p3                                | 1.8      | 1.8      | 2N 2-3E 80-85 cmbd                       |
| 1775 | <i>Brachylagus idahoensis</i>    | L    | p3                                | 1.4      | 1.2      | 1N 2E 25-30 cmbd                         |
| 1776 | <i>Brachylagus idahoensis</i>    | R    | p3                                | 1.8      | 1.8      | 2N 2-3E 45-60 cmbd                       |
| 1777 | <i>Brachylagus idahoensis</i>    | L    | p3                                | 1.8      | 1.6      | 2N 2-3E 75-80 cmbd                       |
| 1778 | Leporidae                        | R    | p3                                | —        | —        | 1N 3E 30-35 cmbd                         |
| 1779 | Leporidae                        | R    | p3                                | —        | —        | 2N 2E E Half Cemented Level              |
| 1780 | Leporidae                        | R    | p3                                | 2.0      | —        | 1N 2-3E 70-75 cmbd                       |
| 1781 | <i>Ochotona</i> sp.              | L    | p3                                | 1.2      | 1.2      | 1N 3E 45-70 cmbd; Wall Collapse          |
| 1782 | <i>Ochotona</i> sp.              | R    | p3                                | 1.2      | 1.5      | 1N 3E 45-70 cmbd; Wall Collapse          |
| 1783 | <i>Ochotona</i> sp.              | R    | p3                                | —        | —        | 1N 3E 45-70 cmbd; Wall Collapse          |
| 1784 | <i>Ochotona</i> sp.              | L    | p3                                | 1.1      | 1.2      | 1N 3E 45-70 cmbd; Wall Collapse          |
| 1785 | <i>Ochotona</i> sp.              | R    | p3                                | —        | —        | 1N 2E 30-35 cmbd                         |
| 1786 | <i>Ochotona</i> sp.              | R    | p3                                | 1.2      | 1.3      | 1N 3E 35-40 cmbd                         |
| 1787 | <i>Ochotona</i> sp.              | L    | p3                                | 1.1      | 1.2      | 1N 2E 40-45 cmbd                         |
| 1788 | <i>Ochotona</i> sp.              | L    | p3                                | 1.1      | 1.0      | 1N 2E 40-45 cmbd                         |
| 1789 | <i>Ochotona</i> sp.              | R    | p3                                | 1.3      | 1.5      | 1N 3E 40-45 cmbd                         |
| 1790 | <i>Ochotona</i> sp.              | L    | p3                                | 1.5      | 1.6      | 2N 3E 40-45 cmbd                         |
| 1791 | <i>Ochotona</i> sp.              | R    | p3                                | —        | —        | 2N 3E 40-45 cmbd                         |
| 1792 | <i>Ochotona</i> sp.              | L    | p3                                | 1.4      | 1.6      | 1N 3E Cemented Level                     |
| 1793 | <i>Ochotona</i> sp.              | R    | p3                                | 1.4      | 1.7      | 1N 3E Cemented Level                     |
| 1794 | <i>Ochotona</i> sp.              | R    | p3                                | 1.3      | 1.4      | 1N 3E Cemented Level                     |
| 1795 | <i>Ochotona</i> sp.              | R    | p3                                | 1.3      | 1.3      | 1N 3E Cemented Level                     |
| 1796 | <i>Ochotona</i> sp.              | L    | p3                                | 1.5      | —        | 2N 2E E Half Cemented Level              |
| 1797 | <i>Ochotona</i> sp.              | R    | p3                                | 1.2      | 1.1      | 1N 3E 43.5-55 cmbd; below Cemented Level |
| 1798 | <i>Ochotona</i> sp.              | R    | p3                                | 1.3      | 1.4      | 1N 3E 43.5-55 cmbd; below Cemented Level |
| 1799 | <i>Ochotona</i> sp.              | R    | dentary<br>with m1-m3             | —        | —        | 1N 3E 43.5-55 cmbd; below Cemented Level |
| 457  | <i>Ochotona</i> sp.              | L    | p3                                | 1.0      | 1.2      | 2N 2-3E 45-60 cmbd                       |
| 470  | <i>Ochotona</i> sp.              | R    | p3                                | 1.1      | 1.3      | 2N 2-3E 45-60 cmbd                       |
| 471  | <i>Ochotona</i> sp.              | L    | p3                                | 1.3      | 1.4      | 2N 2-3E 45-60 cmbd                       |
| 1800 | <i>Ochotona</i> sp.              | R    | p3                                | 1.3      | 1.3      | 2N 2-3E 45-60 cmbd                       |
| 1801 | <i>Ochotona</i> sp.              | R    | p3                                | 1.2      | 1.4      | 2N 2-3E 45-60 cmbd                       |
| 1802 | <i>Ochotona</i> sp.              | R    | dentary<br>with p4-m3             | —        | —        | 2N 2-3E 45-60 cmbd                       |
| 1803 | <i>Ochotona</i> sp.              | L    | p3                                | 1.3      | 1.2      | 1N 2E 50-55 cmbd                         |
| 1804 | <i>Ochotona</i> sp.              | L    | p3                                | 1.3      | 1.3      | 1N 2E 55-60 cmbd                         |
| 1805 | <i>Ochotona</i> sp.              | R    | dentary<br>with p4-m1<br>or m1-m2 | —        | —        | 1N 2E 55-60 cmbd                         |
| 1806 | <i>Ochotona</i> sp.              | L    | p3                                | 1.3      | 1.6      | 1N 2E 55-60 cmbd                         |
| 1807 | <i>Ochotona</i> sp.              | R    | p3                                | 1.3      | 1.5      | 1N 2E 55-60 cmbd                         |
| 1808 | <i>Ochotona</i> sp.              | R    | p3                                | 1.2      | 1.4      | 1N 2E 55-60 cmbd                         |
| 1809 | <i>Ochotona</i> sp.              | L    | p3                                | 1.3      | 1.4      | 1N 2E 55-60 cmbd                         |
| 1810 | <i>Ochotona</i> sp.              | R    | p3                                | 1.3      | 1.5      | 1N 3E 55-60 cmbd                         |
| 1811 | <i>Ochotona</i> sp.              | R    | dentary                           | —        | —        | 1N 3E 55-60 cmbd                         |
| 1812 | <i>Ochotona</i> sp.              | R    | dentary<br>with m1-m2             | —        | —        | 1N 2E 60-65 cmbd                         |
| 1813 | <i>Ochotona</i> sp.              | L    | p3                                | 1.3      | 1.6      | 1N 2E 60-65 cmbd                         |
| 1814 | <i>Ochotona</i> sp.              | L    | p3                                | 1.2      | 1.2      | 1N 3E 60-65 cmbd                         |
| 1815 | <i>Ochotona</i> sp.              | L    | p3                                | 1.2      | 1.2      | 1N 3E 60-65 cmbd                         |

| #    | TAXON               | SIDE | ELEMENT               | <i>l</i> | <i>w</i> | PROVENIENCE                                      |
|------|---------------------|------|-----------------------|----------|----------|--------------------------------------------------|
| 1816 | <i>Ochotona</i> sp. | L    | p3                    | 1.2      | 1.1      | 2N 2-3E 60-65 cmbd                               |
| 1817 | <i>Ochotona</i> sp. | R    | p3                    | 1.3      | 1.5      | 2N 2-3E 60-65 cmbd                               |
| 1818 | <i>Ochotona</i> sp. | R    | p3                    | 1.0      | 1.1      | 2N 2-3E 60-65 cmbd                               |
| 1819 | <i>Ochotona</i> sp. | R    | p3                    | 1.3      | 1.4      | 2N 2-3E 60-65 cmbd                               |
| 1820 | <i>Ochotona</i> sp. | R    | p3                    | 1.3      | 1.2      | 1N 2E 65-70 cmbd                                 |
| 1821 | <i>Ochotona</i> sp. | L    | p3                    | 1.3      | 1.5      | 1N 2E 65-70 cmbd                                 |
| 1310 | <i>Ochotona</i> sp. | R    | p3                    | 1.2      | 1.2      | 1N 3E 65-70 cmbd                                 |
| 1311 | <i>Ochotona</i> sp. | R    | p3                    | 1.3      | 1.4      | 1N 3E 65-70 cmbd                                 |
| 1822 | <i>Ochotona</i> sp. | L    | p3                    | 1.3      | 1.3      | 2N 2-3E 65-70 cmbd                               |
| 1823 | <i>Ochotona</i> sp. | R    | p3                    | 1.1      | 1.1      | 2N 2-3E 65-70 cmbd                               |
| 1824 | <i>Ochotona</i> sp. | R    | p3                    | 1.4      | 1.4      | 2N 2-3E 65-70 cmbd                               |
| 1825 | <i>Ochotona</i> sp. | L    | p3                    | 1.5      | 1.3      | 2N 2-3E 65-70 cmbd                               |
| 884  | <i>Ochotona</i> sp. | L    | p3                    | 1.1      | —        | 1N 2-3E 70-75 cmbd                               |
| 1826 | <i>Ochotona</i> sp. | R    | p3                    | 1.3      | 1.5      | 1N 2-3E 70-75 cmbd                               |
| 1827 | <i>Ochotona</i> sp. | R    | p3                    | 1.1      | 1.3      | 2N 2-3E 70-75 cmbd                               |
| 1828 | <i>Ochotona</i> sp. | R    | dentary<br>with p3-p4 | 1.1      | 1.3      | 1N 2-3E 75-80 cmbd                               |
| 1829 | <i>Ochotona</i> sp. | R    | p3                    | 1.3      | 1.3      | 1N 2-3E 75-80 cmbd                               |
| 1830 | <i>Ochotona</i> sp. | L    | p3                    | 1.2      | 1.3      | 1N 2-3E 75-80 cmbd                               |
| 1831 | <i>Ochotona</i> sp. | L    | p3                    | 1.2      | 1.2      | 2N 2-3E 75-80 cmbd                               |
| 1832 | <i>Ochotona</i> sp. | L    | p3                    | 1.2      | 1.2      | 2N 2-3E 75-80 cmbd                               |
| 1833 | <i>Ochotona</i> sp. | L    | p3                    | 1.2      | 1.1      | 2N 2-3E 75-80 cmbd                               |
| 1834 | <i>Ochotona</i> sp. | L    | p3                    | 1.0      | 1.1      | 2N 2-3E 75-80 cmbd                               |
| 1835 | <i>Ochotona</i> sp. | L    | p3                    | 1.1      | 1.0      | 2N 2-3E 75-80 cmbd                               |
| 1836 | <i>Ochotona</i> sp. | L    | p3                    | 1.1      | 1.2      | 2N 2-3E 75-80 cmbd                               |
| 1837 | <i>Ochotona</i> sp. | R    | p3                    | —        | —        | 1N 2-3E 80-85 cmbd                               |
| 1839 | <i>Ochotona</i> sp. | R    | p3                    | 1.3      | 1.3      | 2N 2-3E 80-85 cmbd                               |
| 1840 | <i>Ochotona</i> sp. | R    | p3                    | 1.3      | 1.3      | 2N 2-3E 80-85 cmbd                               |
| 1841 | <i>Ochotona</i> sp. | R    | p3                    | —        | —        | 2N 2-3E 80-85 cmbd                               |
| 1842 | <i>Ochotona</i> sp. | L    | p3                    | 1.3      | 1.6      | 1N 2-3E 85-90 cmbd                               |
| 1843 | <i>Ochotona</i> sp. | L    | p3                    | 1.3      | 1.3      | 1N 2-3E 85-90 cmbd                               |
| 1844 | <i>Ochotona</i> sp. | L    | p3                    | 1.2      | 1.2      | 1N 2-3E 85-90 cmbd                               |
| 1845 | <i>Ochotona</i> sp. | R    | p3                    | 1.1      | 1.2      | 2N 2-3E 85-90 cmbd                               |
| 1846 | <i>Ochotona</i> sp. | L    | p3                    | 1.3      | 1.3      | 2N 2-3E 85-90 cmbd                               |
| 1847 | <i>Ochotona</i> sp. | R    | p3                    | 1.1      | 1.2      | 2N 2-3E 85-90 cmbd                               |
| 1848 | <i>Ochotona</i> sp. | R    | p3                    | —        | —        | 2N 2-3E 85-90 cmbd                               |
| 1849 | <i>Ochotona</i> sp. | R    | p3                    | 1.2      | 1.3      | 2N 2-3E 85-90 cmbd and below                     |
| 1850 | <i>Ochotona</i> sp. | L    | p3                    | 1.0      | 1.0      | 2N 3E 85-110 cmbd; below flowstone               |
| 1851 | <i>Ochotona</i> sp. | L    | p3                    | 1.2      | 1.3      | 2N 3E 85-110 cmbd; below flowstone               |
| 1852 | <i>Ochotona</i> sp. | L    | p3                    | 1.3      | 1.4      | 2N 3E 85-110 cmbd; below flowstone               |
| 1853 | <i>Ochotona</i> sp. | R    | p3                    | 1.2      | 1.3      | 2N 3E 85-110 cmbd; below flowstone               |
| 1854 | <i>Ochotona</i> sp. | L    | p3                    | 1.2      | 1.3      | 1-2N 2-3E 87-102 cmbd; below pedestaled<br>rocks |
| 1855 | <i>Ochotona</i> sp. | R    | dentary<br>with p3-p4 | 1.1      | 1.1      | 1-2N 2-3E 87-102 cmbd; below pedestaled<br>rocks |
| 1856 | <i>Ochotona</i> sp. | R    | p3                    | 1.2      | 1.5      | 2N 2-3E 90-95 cmbd                               |
| 1857 | <i>Ochotona</i> sp. | L    | p3                    | 1.3      | 1.4      | 2N 2-3E 90-95 cmbd                               |
| 1858 | <i>Ochotona</i> sp. | L    | p3                    | 1.3      | 1.4      | 2N 2-3E 90-95 cmbd                               |
| 1859 | <i>Ochotona</i> sp. | L    | p3                    | —        | —        | 1N 2-3E 90 cmbd and below                        |
| 1860 | <i>Ochotona</i> sp. | R    | p3                    | —        | —        | 2N 2-3E 95-102 cmbd                              |
| 1861 | <i>Ochotona</i> sp. | R    | p3                    | 1.4      | 1.5      | 2N 3E 100-105 cmbd                               |
| 1862 | <i>Ochotona</i> sp. | L    | p3                    | 1.4      | 1.4      | 2N 3E 100-105 cmbd                               |
| 1863 | <i>Ochotona</i> sp. | R    | p3                    | 1.3      | 1.4      | 2N 3E 100-105 cmbd                               |
| 1864 | <i>Ochotona</i> sp. | R    | p3                    | 1.3      | 1.6      | 2N 3E 100-105 cmbd                               |

| #    | TAXON                                 | SIDE | ELEMENT               | <i>l</i> | <i>w</i> | PROVENIENCE                              |
|------|---------------------------------------|------|-----------------------|----------|----------|------------------------------------------|
| 1865 | Ochotonidae                           | R    | p3                    | —        | —        | 1N 2E 65-70 cmbd                         |
| 1866 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 2.9      | 3.1      | 1N 3E 25-60 cmbd; Wall Scrapings         |
| 1867 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 3.4      | 3.2      | 1N 3E 45-70 cmbd; Wall Collapse          |
| 1868 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 3.3      | 3.2      | 2N 3E 30-35 cmbd                         |
| 1869 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 1.7      | 1.5      | 2N 3E 40-45 cmbd                         |
| 1870 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 2.4      | 2.4      | 2N 3E 40-45 cmbd                         |
| 1871 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 2.7      | 2.6      | 2N 3E 40-45 cmbd                         |
| 1872 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 2.5      | 1.8      | 1N 3E Cemented Level                     |
| 1873 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 3.2      | 3.2      | 2N 2E E Half Cemented Level              |
| 1874 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | dentary<br>with p3-p4 | 2.4      | 2.2      | 1N 3E 43.5-55 cmbd; below Cemented Level |
| 1875 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 3.1      | 3.0      | 1N 3E 43.5-55 cmbd; below Cemented Level |
| 1876 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 3.1      | 2.9      | 1N 3E 43.5-55 cmbd; below Cemented Level |
| 1877 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 2.1      | 2.0      | 1N 3E 43.5-55 cmbd; below Cemented Level |
| 1878 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 2.3      | 2.2      | 1N 3E 43.5-55 cmbd; below Cemented Level |
| 1879 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 3.2      | 3.0      | 2N 2-3E 45-60 cmbd                       |
| 1880 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 2.5      | 2.2      | 2N 2-3E 45-60 cmbd                       |
| 1881 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 3.4      | 2.6      | 2N 2-3E 45-60 cmbd                       |
| 1882 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 3.9      | 3.5      | 2N 2-3E 45-60 cmbd                       |
| 1883 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 2.4      | 2.4      | 2N 2-3E 45-60 cmbd                       |
| 463  | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 2.0      | 1.7      | 2N 2-3E 45-60 cmbd                       |
| 452  | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | —        | —        | 2N 2-3E 45-60 cmbd                       |
| 480  | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 2.2      | 1.2      | 2N 2-3E 45-60 cmbd                       |
| 1884 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 2.7      | 2.5      | 2N 2-3E 45-60 cmbd                       |
| 1885 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | —        | —        | 2N 2-3E 45-60 cmbd                       |
| 1886 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 3.3      | 3.0      | 2N 2-3E 45-60 cmbd                       |
| 1887 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 3.3      | 2.8      | 1N 2E 50-55 cmbd                         |
| 1888 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 3.2      | 3.0      | 1N 2E 55-60 cmbd                         |
| 1889 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 3.5      | 3.6      | 1N 3E 55-60 cmbd                         |
| 1890 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 2.1      | 2.0      | 1N 2E 60-65 cmbd                         |
| 1891 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | —        | —        | 1N 3E 60-65 cmbd                         |
| 1892 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | dentary<br>with p3-m2 | 2.5      | 2.5      | 1N 3E 60-65 cmbd                         |
| 1893 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 3.0      | 2.7      | 2N 2-3E 60-65 cmbd                       |
| 1894 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | —        | —        | 1N 2E 65-70 cmbd                         |
| 1895 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 3.6      | 3.4      | 1N 2E 65-70 cmbd                         |
| 1896 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 2.5      | 2.2      | 1N 3E 65-70 cmbd                         |
| 1897 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 2.8      | 2.4      | 1N 3E 65-70 cmbd                         |
| 1898 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | —        | —        | 1N 2-3E 70-75 cmbd                       |
| 1899 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 3.3      | 2.8      | 1N 2-3E 70-75 cmbd                       |
| 1900 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 3.3      | 2.9      | 2N 2-3E 70-75 cmbd                       |
| 1901 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 2.9      | 2.5      | 2N 2-3E 70-75 cmbd                       |
| 1902 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 2.9      | 2.8      | 2N 2-3E 70-75 cmbd                       |
| 1903 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | —        | —        | 2N 2-3E 70-75 cmbd                       |
| 1904 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | —        | —        | 2N 2-3E 70-75 cmbd                       |
| 1905 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | —        | 3.5      | 2N 2-3E 70-75 cmbd                       |
| 1906 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 2.3      | 2.0      | 2N 2-3E 70-75 cmbd                       |
| 1907 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 3.0      | 2.6      | 2N 2-3E 75-80 cmbd                       |
| 1908 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 3.0      | 2.8      | 2N 2-3E 75-80 cmbd                       |
| 1909 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 2.9      | 2.9      | 2N 2-3E 75-80 cmbd                       |
| 1910 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | dentary<br>with p3-m2 | 3.6      | 3.4      | 2N 2-3E 75-80 cmbd                       |
| 1911 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 3.7      | —        | 1N 2-3E 80-85 cmbd                       |
| 1912 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 2.7      | 2.6      | 2N 2-3E 80-85 cmbd                       |

| #    | TAXON                                 | SIDE | ELEMENT               | <i>l</i> | <i>w</i> | PROVENIENCE                        |
|------|---------------------------------------|------|-----------------------|----------|----------|------------------------------------|
| 1913 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | —        | —        | 2N 2-3E 80-85 cmbd                 |
| 1914 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | —        | —        | 2N 2-3E 80-85 cmbd                 |
| 1915 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 3.0      | 2.5      | 2N 2-3E 80-85 cmbd                 |
| 1916 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | 3.4      | 3.7      | 2N 2-3E 80-85 cmbd                 |
| 1917 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | R    | p3                    | —        | —        | 2N 2-3E 80-85 cmbd                 |
| 1918 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | P3                    | —        | —        | 2N 2-3E 90-95 cmbd                 |
| 1919 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | dentary<br>with p3-m2 | —        | —        | 2N 2-3E 90-95 cmbd                 |
| 1119 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | 2.9      | 2.7      | 2N 3E 85-110 cmbd; below flowstone |
| 1920 | <i>Sylvilagus</i> or <i>Lepus</i> sp. | L    | p3                    | —        | —        | 2N 3E 100-105 cmbd                 |