UC San Diego

UC San Diego Previously Published Works

Title

First night of CPAP: impact on memory consolidation attention and subjective experience.

Permalink

<https://escholarship.org/uc/item/2s00z9v5>

Journal

Sleep medicine, 16(6)

ISSN

1389-9457

Authors

Djonlagic, Ina
Guo, Mengshuang
Matteis, Paul
et al.

Publication Date

2015-06-01

DOI

10.1016/j.sleep.2015.01.017

Peer reviewed

Original Article

First night of CPAP: impact on memory consolidation attention and subjective experience

Ina Djonlagic^{a,b,c,*}, Mengshuang Guo^{b,c}, Paul Matteis^{b,c}, Andrea Carusona^c,
Robert Stickgold^b, Atul Malhotra^{c,d}

^a Department of Neurology, Beth Israel Deaconess Medical Center, Boston, MA, USA

^b Center for Sleep and Cognition, Department of Psychiatry, Beth Israel Deaconess Medical Center, Boston, MA, USA

^c Division of Sleep Medicine, Sleep Disorders Program Brigham & Women's Hospital and Harvard Medical School, Boston, MA, USA

^d Division of Pulmonary and Critical Care, University of California San Diego, La Jolla, CA, USA

ARTICLE INFO

Article history:

Received 2 September 2014

Received in revised form 23 December 2014

Accepted 23 January 2015

Available online 27 February 2015

Keywords:

Obstructive sleep apnea
Sleep-dependent memory
CPAP
PVT

ABSTRACT

Objective: Neurocognitive deficits are common and serious consequences of obstructive sleep apnea (OSA). Currently, the gold standard treatment is continuous positive air pressure (CPAP) therapy, although the clinical responses to this intervention can be variable. This study examined the effect of one night of CPAP therapy on sleep-dependent memory consolidation, attention, and vigilance as well as subjective experience. **Methods:** Fifteen healthy controls and 29 patients with obstructive sleep apnea of whom 14 underwent a full-night CPAP titration completed the psychomotor vigilance test (PVT) and motor sequence learning task (MST) in the evening and the morning after undergoing overnight polysomnography. All participants also completed subjective evaluations of sleep quality.

Results: Participants with OSA showed significantly less overnight improvement on the MST compared to controls without OSA, independent of whether or not they had received CPAP treatment, while there was no significant difference between the untreated OSA and CPAP-treated patients. Within the OSA group, only those receiving CPAP exhibited faster reaction times on the PVT in the morning. Compared to untreated OSA patients, they also felt subjectively more rested and reported that they slept better.

Conclusion: Our results demonstrate an instant augmentation of subjective experience and, based on PVT results, attention and vigilance after one night of CPAP, but a lack of an effect on offline sleep-dependent motor memory consolidation. This dissociation may be explained by different brain structures underlying these processes, some of which might require longer continued adherence to CPAP to generate an effect.

© 2015 Elsevier B.V. All rights reserved.

1. Introduction

Disorders of breathing during sleep are a major health problem; in particular, obstructive sleep apnea (OSA) has been associated with important health consequences including an increased risk for cardiovascular [1–6] and cerebrovascular disease [7,8], mortality [5,9–14], and cognitive impairment, independent of daytime sleepiness [15–21].

The majority of studies assessing cognition show spared global cognitive functioning and basic language functions [22–24]. By contrast, most OSA patients are impaired on testing of vigilance, that is, tests that require sustained attention for a period of time, a finding which has important implications for the ability to drive and for

general occupational functioning [25–28]. It is therefore not surprising that OSA has been linked to an increased risk of motor vehicle accidents and, thus, if left untreated, imposes a health risk not only to the individual but also to those around them.

Continuous positive airway pressure (CPAP) treatment restores normal respiration during sleep, thereby improving sleep architecture and is considered the first line of treatment for OSA [29,30]. Nightly use of CPAP can markedly reduce sympathetic activity and blood pressure and thus decrease the risk of mortality [31,32]. Studies examining cognitive function pre- and post-CPAP treatment have indicated selective improvement in neuropsychological deficits after CPAP treatment [33–38]. The results have been difficult to interpret due to variations in hypoxia, apnea–hypopnea index (AHI), and cognitive areas tested. Some investigators have also speculated that CPAP only improves deficits related to sleep fragmentation but not due to hypoxemia [38], with which there may be irreversible injury to certain brain regions [39].

Sleep can affect learning and memory in a number of ways. Specific sleep-dependent memory tasks have been used with increasing

* Corresponding author. Department of Neurology, Beth Israel Deaconess Medical Center, 330 Brookline Avenue/KS434, Boston, MA 02115, USA. Tel.: +617 667 5129; fax: +617 667 5216.

E-mail address: jdjonlag@bidmc.harvard.edu (I. Djonlagic).

popularity to assess the dynamic processes during which initial, labile memory representations are converted (either stabilized or enhanced) into more permanent ones. But thus far, few studies have investigated sleep-dependent memory processes in patients with obstructive sleep apnea [40,41]. Given that the precise domains of sleep-dependent memory impairment present in OSA have not been sufficiently studied, important questions remain concerning their reversibility with CPAP treatment.

Several studies have demonstrated that the motor sequence task shows a robust enhancement of performance after a night of sleep, but not after an equivalent time spent awake [42,43]. We recently published a study, in which we demonstrated that memory consolidation for this procedural motor skill task is affected by OSA even in young people with relatively mild disease, and that this effect is not the result of a difference in attention or vigilance [40]. Thus, given the significance of sleep-related processes for improvement on this task, we sought to examine how quickly patients with OSA could recover from the effects of their disease on this form of sleep-dependent memory consolidation.

Patients with OSA often report a major subjective improvement after using CPAP for the first time. The aim of this study was thus to delineate the effect of first-time experience with CPAP on subjective experience, measures of vigilance, and sleep-dependent procedural motor memory consolidation.

2. Methods

The first 44 patients who underwent either a diagnostic polysomnography or a CPAP titration study at Brigham and Women's Hospital Sleep Disorders Laboratory agreed to participate and met inclusion criteria were enrolled in the study. Post polysomnography (PSG), participants were subdivided into three groups: (1) Healthy controls (control group), (2) patients who were found to have obstructive sleep apnea, but did not undergo any treatment intervention (OSA group), and (3) patients with OSA who were CPAP-naïve and underwent a whole night with CPAP (CPAP group). To qualify as healthy controls, participants had to have an AHI <5/h, whereas an AHI of >15/h was required for the OSA group.

2.1. Exclusion criteria

Participants were excluded if they (1) were found to have a periodic limb movement index of >15/h based on their PSG; (2) had another diagnosed sleep or circadian disorder; (3) had a history of alcohol, narcotic, or other drug abuse; (4) had a history of a medical, neurologic, or psychiatric disorder (other than OSA and treated hypertension) that could influence excessive daytime sleepiness; (5) used medications known to have an effect on sleep and daytime vigilance (eg, psychoactive drugs or medications, sedatives, or hypnotics, including selective serotonin re-uptake inhibitors or SSRIs); or (6) were left-handed.

2.2. Study design

All participants performed a 5-min version of the psychomotor vigilance task (PVT) and then trained on the motor sequence task (MST) in the evening between 8 and 9 PM [44,45]. (Fig. 1).

The MST requires repeated typing of a five-digit sequence on a standard computer keyboard (eg, 4-1-3-2-4) with their nondominant (left) hand. The specific sequence to be typed is displayed on the computer screen at all times. Typing is performed in 30-s trials separated by 30-s rest periods. Training and retest each included 12 trials. Following training, participants spent the night in the laboratory and underwent standard sleep recording. The following morning subjects were woken up and then repeated the PVT and were tested on the MST. After a 10-min break, they trained on a new MST sequence. As learning the motor sequence task is sequence specific, there is no transfer of learning to new sequences [46]. This study design allowed us to compare learning in the evening and morning, to confirm that there was no general time-of-day effect on motor sequence performance, and that overnight improvement reflected specific enhancement of performance on the previously learned. MST sequences were counterbalanced for evening and morning training across all participants within groups to control for any order effect. The main performance measure for the MST is the number of correctly typed sequences per 30-s trial, thereby reflecting both speed and accuracy.

The PVT is an objective measurement of sustained attention and reaction time (RT) and has been shown to be sensitive to sleep deprivation, partial sleep loss, and circadian variation in performance efficiency [47]. As such, it allowed us to verify that differences in overnight improvement on the MST were related to differences in overnight memory consolidation, and not to differences in attention. In this task, subjects push a button as fast as they can whenever they see a small (3-mm-high, 4-digit-wide) LED millisecond clock begin counting up from 0000. Pressing the button stops the digital clock, allowing the subject 1.5 s to read the RT. The inter-stimulus interval on the task varies randomly from 2 to 10 s. The duration of the task can be either 5 – as in our case – or 10 min [48]. Participants also completed subjective evaluations of sleep quality in the morning. These questions included two categories: (1) “Would you describe your sleep last night as (a) better, (b) same, (c) worse than usual” and 2) a yes–no question “Do you feel rested this morning?”. In addition, they filled out a questionnaire asking them about their previous typing experience.

2.3. Polysomnography

Standard overnight PSG recording and data interpretation were performed in accordance with the American Academy of Sleep Medicine (AASM) scoring manual [49,50]. This approach included standard electroencephalogram (EEG) leads (F3, F4, C3, C4, O1, and O2), as well as bilateral electrooculogram (EOG), submental electromyogram

Fig. 1. Schematic overview of the visit protocol.

(EMG), bilateral anterior tibialis EMG, and standard electrocardiogram (ECG) electrodes. Nasal/oral airflow (thermistor), nasal pressure (Validyne transducer), chest plus abdominal wall motion (piezo electrodes), and oxygen saturation were also recorded.

All studies were scored by a registered PSG technologist, blinded to subject performance. In particular, hypopneas required a clear (discernable) amplitude reduction of a validated measure of breathing during sleep, and were associated with either an oxygen desaturation of $>3\%$ or an arousal lasting ≥ 10 s.

Arousals were scored visually according to the AASM manual scoring criteria, which require an abrupt shift of EEG frequency including alpha, theta, and/or frequencies >16 Hz (but not spindles) that last at least 3 s preceded by 10 s of stable sleep.

2.4. PSG data analysis

PSG data were preprocessed and analyzed using BrainVision Analyzer 2.0 (BrainProducts, Munich, Germany). Artifacts were automatically detected and removed and EEG data were filtered at 0.5–35 Hz. Artifact rejection was confirmed by visual inspection. Power spectral density ($\mu V^2/Hz$) was calculated by fast Fourier transform (FFT), applying a Hanning window to 5-s epochs of N2 and N3 sleep with 50% overlap.

Based on previous studies, spindle analysis (number and density) was performed from the C3 and C4 location for stage 2 sleep manually and by using a wavelet-based algorithm. Data were preprocessed and analyzed using BrainVision Analyzer 2.0 and MatLab R2009b (The MathWorks, Natick, MA, USA) as described by Wamsley et al. [51].

2.5. Statistical analysis

Variables were compared between healthy controls and OSA patients with and without CPAP. One-way analyses of variance (ANOVAs) were performed to compare demographic and PSG-derived sleep data including sleep spindle analyses. Subjective sleep quality evaluations were analyzed using chi-square tests.

Overnight MST improvement was calculated as percent change in performance speed (correct sequences per 30-s trial) for initial and plateau improvement taking into account the characteristic performance curves over the 12 training trials and 12 test trials [*initial improvement* = percent increase in performance from the last three training trials in the evening to the first three test trials in the morning and *plateau improvement* = percent improvement from the last six training trials in the evening to the last six test trials in the morning].

Percentage values for improvement across the three groups were compared by applying one-way ANOVAs and post-tested with Tukey's Honestly Significant Differences (HSD) tests.

Statistical analysis was performed using JMP Version 10 (SAS Institute Inc., Cary, NC, USA). A p -value of <0.05 was considered significant. Variability is expressed as standard deviation (S.D.).

2.5.1. Sample size justification

For the motor sequence task, published results indicate an effect size of 1.64 [44]. The required sample size to achieve a power of 80% with the alpha-level set to 0.05 is an $n = 14$ for a two-tailed test and $n = 12$ for a single-tailed test. All groups had $n \geq 14$.

2.6. Ethics statement

All participants provided written informed consent. The study was approved by the Partners' Institutional Review Board.

3. Results

3.1. Demographic and sleep variables

Demographic and sleep characteristics are shown in Table 1. Among the three groups, there was no significant difference in age or BMI. In addition, measures of sleep architecture including total sleep time, sleep efficiency, and sleep stage distribution were very similar.

As expected, untreated OSA patients had a significantly higher AHI and arousal index as well as the lowest oxygen nadir compared to healthy controls and OSA patients treated with CPAP.

3.2. Performance measures of the MST

3.2.1. Evening training

Though healthy controls were overall able to type more sequences throughout the 12 evening training trials, this difference was not significant. All three groups showed similar practice-dependent improvement across the 12 evening training trials (percent increase in performance from the first two trials to the last three trials; ANOVA, $p = 0.291$) (Fig. 2).

3.2.2. Overnight improvement

Independent of whether they received treatment, OSA participants (both OSA and CPAP groups) showed significantly less initial overnight improvement on the MST compared to healthy sleepers (OSA $2.0 \pm 13.0\%$ [S.D.], CPAP $1.4\% \pm 12.6\%$, controls $13.4\% \pm 13.5\%$; $F = 4.1$, $df = 2$, $p = 0.02$ [ANOVA across groups]; Fig. 3A). Post-hoc Tukey HSD comparisons revealed $p = 0.05$ for controls compared to OSA patients and $p = 0.04$ for controls compared to the CPAP group. No significant differences were seen between the OSA and CPAP groups ($p = 0.99$). Similar results were found for comparisons of overnight plateau improvement (OSA $8.9\% \pm 7.8\%$, CPAP $8.0\% \pm 9.1\%$, controls $24.0\% \pm 19.3\%$; $F = 6.0$, $df = 2$, $p = 0.005$ Fig. 3B [ANOVA across groups]; Fig. 2B). Here, post-hoc comparisons revealed significant differences between controls and OSA patients ($p = 0.02$) and between controls and the CPAP group ($p = 0.01$), but no significant difference between the OSA and CPAP groups ($p = 0.98$). However, the level of performance at the end of the evening training session (average of last three trials) did not correlate with the amount of overnight improvement (Pearson correlation $r^2 = 0.05$, $p = 0.14$).

Table 1
Group demographics and sleep characteristics.

	CPAP ($n = 14$)	OSA ($n = 15$)	Control ($n = 15$)	P -value
Age (years)	44.4 $\pm$ 11.1	43.3 $\pm$ 13.0	37.3 $\pm$ 10.5	0.19
BMI (kg/m^2)	32.0 $\pm$ 5.4	30.2 $\pm$ 5.5	27.6 $\pm$ 5.8	0.12
TST (min)	332.3 $\pm$ 57.7	308.5 $\pm$ 51.9	345.7 $\pm$ 27.9	0.11
Sleep Efficiency (%)	84.3 $\pm$ 8.5	82.5 $\pm$ 11.5	86.0 $\pm$ 10.7	0.61
Sleep stages:				
Stage N1 (%)	8.0 $\pm$ 4.0	8.6 $\pm$ 4.4	8.3 $\pm$ 4.8	0.66
Stage N2 (%)	60.8 $\pm$ 10.0	63.4 $\pm$ 8.3	58.5 $\pm$ 8.42.2	0.30
Stage N3 (%)	11.6 $\pm$ 13.3	11.9 $\pm$ 10.5	15.2 $\pm$ 11.73.0	0.71
REM (%)	19.7 $\pm$ 3.4	16.1 $\pm$ 7.9	18.0 $\pm$ 7.82.0	0.37
AHI (events/h)	5.9 $\pm$ 4.6	23.9 $\pm$ 10.0 ^a	3.5 $\pm$ 1.30.4	<0.0001
Oxygen nadir (%)	90.7 $\pm$ 3.3	86.4 $\pm$ 4.2 ^a	91.4 $\pm$ 2.6	<0.0001
Arousal index (A/h)	17.2 $\pm$ 5.6	25.6 $\pm$ 12.7 ^a	18.3 $\pm$ 7.3	0.03

Values are presented as mean $\pm$ S.D. P -values represent results of ANOVA across the three groups.

Asterisks represent post-hoc comparisons using Tukey HSD tests: ^a(CPAP/OSA).

^a (Control/OSA).

Fig. 2. Trial-by-trial MST performance across groups.

3.2.3. Morning training

After completing the MST testing in the morning, all participants were trained on a novel sequence in order to control for circadian influences on performance. No significant within-group differences in performance were found between the two training sequences (evening vs. morning; all p 's > 0.28). Thus, circadian influences do not explain the overnight changes in performance between groups.

3.2.4. Sleep spindles

Previous research has shown an association between motor sequence learning and sleep spindles during stage 2 (N2) sleep [52,53]. Based on these reports, sleep spindles were assessed during stage 2 (N2) sleep only. A significant difference among the three groups in total number of sleep spindles as well as in spindle density was observed at both central electrodes (C3, C4). The most notable difference was seen over the C3 electrodes where CPAP-treated OSA patients demonstrated significantly higher total number of sleep spindles ($p = 0.01$ [ANOVA across groups], $p = 0.01$ [CPAP vs. OSA]) and central sleep spindle density ($p = 0.008$ [ANOVA across groups], $p = 0.007$ [CPAP/OSA]) compared to controls and untreated OSA patients. Significant differences were also present over C4 ($p = 0.03$ [ANOVA, C4 total number], $p = 0.02$ [ANOVA, C4 density]). Post-hoc Tukey HSD test demonstrated significant differences again only for CPAP/OSA group comparisons ($p = 0.02$ [C4 total number], $p = 0.01$ [C4 density]) (Table 2).

Fig. 3. MST Overnight Improvement.

Table 2
Sleep spindles.

	CPAP	OSA	Control	P-value
Spindles – C3	435.9 ± 149.4*	318.5 ± 135.8	377.4 ± 75.3	0.01
Spindles – C4	420 ± 162.2*	310.9 ± 123.8	400.7 ± 85.8	0.03
Spindle density C3	2.3 ± 1.0*	1.6 ± 0.5	1.9 ± 0.3	0.01
Spindle density C4	2.2 ± 1.0*	1.6 ± 0.5	2.0 ± 0.4	0.02

Values are presented as mean ± S.D. P-value represents results of ANOVA across the three groups. Asterisks represent post-hoc comparisons using Tukey HSD tests: *(CPAP/OSA).

Fig. 4. Changes in PVT reaction time between evening and morning.

Contrary to previous reports, we did not find any correlation between total number of central spindles or central spindle density and overnight MST improvement for any of the groups.

3.3. Measures of alertness

3.3.1. PVT performance

In the evening, the mean RT for baseline PVT performance was 387.9 ± 121.4 ms (lapses [RT > 500 ms] = 9.0 ± 11.5) for OSA patients without CPAP and 440.0 ± 142.6 ms (lapses = 11.6 ± 13.1) for OSA patients on CPAP, compared to a mean RT of 409.7 ± 125.3 ms (lapses = 8.1 ± 11.9) for controls ($p = 0.56$ [ANOVA across all groups]). The control group showed very similar PVT performance in the morning compared to the evening ($p = 0.88$). Among all the OSA patients, only those who received a full night on CPAP (CPAP group) exhibited faster RT on the PVT in the morning than those who did not (OSA group) ($p = 0.04$ [OSA /CPAP]) (Fig. 4).

3.3.2. Subjective ratings

Compared to untreated OSA patients, more full-night CPAP participants reported both feeling more rested (64% vs. 36%, $p = 0.005$) and having improved sleep quality (38.5% vs. 6.7%, $p = 0.04$) the next morning (Fig. 5 A–D). We did not find any correlations between subjective ratings and PVT performance or overnight MST improvement in any of the three groups.

4. Discussion

Previous studies have indicated selective improvement in neurocognitive deficits after CPAP treatment [37,54]. However, little is known about the effect of CPAP on sleep-dependent memory consolidation.

Our study shows resolution of sleep-disordered breathing during the first night of treatment with CPAP, including oxygen saturations and arousal indices similar to healthy controls. Those OSA patients who experienced their first night on CPAP demonstrated an instant

Fig. 5. Subjective Sleep Ratings.

improvement in attention and vigilance based on PVT results the next morning. They also felt subjectively more rested and reported improved sleep quality. Despite this finding, however, their overnight improvement on the MST remained virtually identical to that seen in untreated OSA patients and distinctly poorer than in healthy controls. Therefore, even though patients who received CPAP reported that they felt they had slept better and felt more rested, they failed to show the normal memory benefits from the night's sleep.

Our results demonstrate that acute recovery after one night of CPAP is limited and selective and confirm that improvement of cognitive deficits caused by OSA depends on multiple likely intertwined mechanisms. Previous studies investigating the effects of CPAP on cognitive recovery have applied CPAP exposure duration from one week [55], to up to 12 months [56]. The study by Bardwell et al. investigated the effectiveness of one week CPAP treatment versus placebo CPAP on a board battery cognitive outcomes and was able to demonstrate that CPAP improved overall cognitive functioning, but not in any specific domain tested. In particular, only one of the 22 neuropsychological tests scores (Digit Vigilance–Time) showed significant changes specific to CPAP treatment. Munoz et al. investigated the long-term use of CPAP and demonstrated that a significant improvement in somnolence and vigilance was already evident after three months of treatment and continued to persist after 12.1 months [56]. On the other hand, their study results also showed that long-term use of CPAP does not normalize the levels of depression and anxiety.

The present study is the first study to investigate the effect of the first night on CPAP and the results provide compelling evidence that improvement in some cognitive areas may already be present after one day of treatment.

While many cognitive areas improve after CPAP, there have also been some reports of persistent deficits after continued use of CPAP, which have included deficits attributable to frontal lobe function

such as executive function, short-term memory, planning abilities, as well as manual dexterity and psychomotor function [38,57].

It is therefore likely that the sensitivity to the effects of CPAP treatment may not only depend on the duration of treatment and nightly compliance, but also depend on the specific cognitive areas tested and the recoverability of their underlying brain structures.

Thus, whether sleep-dependent memory consolidation processes can recover with continued adherence to CPAP remains unknown.

Sleep spindles are presumed to reflect experience-dependent activation patterns as part of the hippocampal–neocortical dialogue involved in the consolidation of new learning, thus facilitating synaptic plasticity and also representing an index of the capacity for learning [58,59]. Few studies have analyzed sleep spindles in patients with sleep-disordered breathing. Our findings confirm a reduction of sleep spindles in OSA patients compared to healthy controls.

By contrast, those in the CPAP group showed spindle numbers significantly higher than the OSA group and even the control group. Without spindle values from pre-CPAP, interpretations of this finding remain speculative. Based on the link between sleep spindles and stability of sleep, one possible explanation might be that the higher number of sleep spindles in the CPAP group could be interpreted as a recovery phenomenon in conjunction with the improvement of the sleep-fragmenting nature of OSA [60,61].

Contrary to the differences in sleep spindles, there were no significant differences in sleep architecture among the three groups. It is unclear why we do not see significantly poorer sleep architecture in the OSA group, despite their significantly higher AHI and arousal index, and lower oxygen nadir. It may simply reflect how well apnea patients can fall back asleep after these arousals. These findings replicate our previous study [40] where the extent of impairment of sleep-dependent consolidation correlated with number of arousals, but not other sleep parameters.

We acknowledge limitations to our study. Despite the limited sample size, we were still well powered for our primary outcomes and do believe that we can draw meaningful conclusions from these data. Second, we studied a relatively young population compared to some prior studies, and our conclusions cannot be extended to older OSA patients. Further research will be required to determine the effects of various covariates such as age and gender on these outcomes. Third, in an effort to dissociate the subjective improvements in symptoms from the objective changes in neurocognitive function, we studied only a single night of CPAP. Thus, we cannot draw conclusions about the effects of longer-term therapy on sleep-dependent memory consolidation or whether the observed abnormalities are in fact irreversible. A related factor is the duration of disease prior to diagnosis; however, this parameter is notoriously difficult to estimate and thus we are unable to determine whether patients with long-standing disease have less improvement in memory consolidation compared to those with more recent onset of disease.

Despite these limitations, we believe that our findings represent a useful addition to the literature, and help fuel additional research in this area.

Conflict of interest

The ICMJE Uniform Disclosure Form for Potential Conflicts of Interest associated with this article can be viewed by clicking on the following link: <http://dx.doi.org/10.1016/j.sleep.2015.01.017>.

Acknowledgments

The authors wish to thank Dr. Erin Wamsley and Dr. Matthew Tucker for providing spindle analysis tools and their helpful advice.

This was not an industry-supported study. The study was funded by NIH grant # K23HL103850 and American Sleep Medicine Foundation grant #54-JF-10.

References

- [1] Grunstein RR, Stenlof K, Hedner J, Sjostrom L. Impact of obstructive sleep apnea and sleepiness on metabolic and cardiovascular risk factors in the Swedish Obese Subjects (SOS) Study. *Int J Obes Relat Metab Disord* 1995;19:410–18.
- [2] Jennum P, Sjol A. Snoring, sleep apnoea and cardiovascular risk factors: the MONICA II Study. *Int J Epidemiol* 1993;22:439–44.
- [3] Newman AB, Enright PL, Manolio TA, Haponik EF, Wahl PW. Sleep disturbance, psychosocial correlates, and cardiovascular disease in 5201 older adults: the Cardiovascular Health Study. *J Am Geriatr Soc* 1997;45:1–7.
- [4] Newman AB, Nieto FJ, Guidry U, et al. Relation of sleep-disordered breathing to cardiovascular disease risk factors: the Sleep Heart Health Study. *Am J Epidemiol* 2001;154:50–9.
- [5] Newman AB, Spiekerman CF, Enright P, et al. Daytime sleepiness predicts mortality and cardiovascular disease in older adults. The Cardiovascular Health Study Research Group. *J Am Geriatr Soc* 2000;48:115–23.
- [6] Shahar E, Whitney CW, Redline S, et al. Sleep-disordered breathing and cardiovascular disease: cross-sectional results of the Sleep Heart Health Study. *Am J Respir Crit Care Med* 2001;163:19–25.
- [7] Yaggi H, Mohsenin V. Obstructive sleep apnoea and stroke. *Lancet Neurol* 2004;3:333–42.
- [8] Yaggi HK, Concato J, Kernan WN, Lichtman JH, Brass LM, Mohsenin V. Obstructive sleep apnea as a risk factor for stroke and death. *N Engl J Med* 2005;353:2034–41.
- [9] Ancoli-Israel S, Kripke DF, Klauber MR, et al. Morbidity, mortality and sleep-disordered breathing in community dwelling elderly. *Sleep* 1996;19:277–82.
- [10] Kripke DF, Garfinkel L, Wingard DL, Klauber MR, Marler MR. Mortality associated with sleep duration and insomnia. *Arch Gen Psychiatry* 2002;59:131–6.
- [11] Manabe K, Matsui T, Yamaya M, et al. Sleep patterns and mortality among elderly patients in a geriatric hospital. *Gerontology* 2000;46:318–22.
- [12] Mant A, King M, Saunders NA, Pond CD, Goode E, Hewitt H. Four-year follow-up of mortality and sleep-related respiratory disturbance in non-demented seniors. *Sleep* 1995;18:433–8.
- [13] Rockwood K, Davis HS, Merry HR, MacKnight C, McDowell I. Sleep disturbances and mortality: results from the Canadian Study of Health and Aging. *J Am Geriatr Soc* 2001;49:639–41.
- [14] Stone KL, Ewing SK, Ancoli-Israel S, et al. Self-reported sleep and nap habits and risk of mortality in a large cohort of older women. *J Am Geriatr Soc* 2009;57:604–11.
- [15] Bonnet MH. Cognitive effects of sleep and sleep fragmentation. *Sleep* 1993;16:S65–7.
- [16] Crenshaw MC, Edinger JD. Slow-wave sleep and waking cognitive performance among older adults with and without insomnia complaints. *Physiol Behav* 1999;66:485–92.
- [17] Dealberto MJ, Pajot N, Courbon D, Alperovitch A. Breathing disorders during sleep and cognitive performance in an older community sample: the EVA Study. *J Am Geriatr Soc* 1996;44:1287–94.
- [18] Edinger JD, Glenn DM, Bastian LA, Marsh GR. Slow-wave sleep and waking cognitive performance II: findings among middle-aged adults with and without insomnia complaints. *Physiol Behav* 2000;70:127–34.
- [19] Engleman HM, Kingshott RN, Martin SE, Douglas NJ. Cognitive function in the sleep apnea/hypopnea syndrome (SAHS). *Sleep* 2000;23(Suppl. 4):S102–8.
- [20] Granat P. Sleep loss and cognitive performance. *J Fam Pract* 1990;30:632.
- [21] McCann UD, Penetar DM, Shaham Y, et al. Sleep deprivation and impaired cognition. Possible role of brain catecholamines. *Biol Psychiatry* 1992;31:1082–97.
- [22] Greenberg GD, Watson RK, Deptula D. Neuropsychological dysfunction in sleep apnea. *Sleep* 1987;10:254–62.
- [23] Lojander J, Kajaste S, Maasilta P, Partinen M. Cognitive function and treatment of obstructive sleep apnea syndrome. *J Sleep Res* 1999;8:71–6.
- [24] Redline S, Strauss ME, Adams N, et al. Neuropsychological function in mild sleep-disordered breathing. *Sleep* 1997;20:160–7.
- [25] Barbe F, Pericas J, Munoz A, Findley L, Anto JM, Agusti AG. Automobile accidents in patients with sleep apnea syndrome. An epidemiological and mechanistic study. *Am J Respir Crit Care Med* 1998;158:18–22.
- [26] Findley L, Unverzagt M, Guchu R, Fabrizio M, Buckner J, Suratt P. Vigilance and automobile accidents in patients with sleep apnea or narcolepsy. *Chest* 1995;108:619–24.
- [27] Findley LJ, Suratt PM, Dinges DF. Time-on-task decrements in “steer clear” performance of patients with sleep apnea and narcolepsy. *Sleep* 1999;22:804–9.
- [28] Bedard MA, Montplaisir J, Richer F, Malo J. Nocturnal hypoxemia as a determinant of vigilance impairment in sleep apnea syndrome. *Chest* 1991;100:367–70.
- [29] Loreda JS, Ancoli-Israel S, Dimsdale JE. Effect of continuous positive airway pressure vs placebo continuous positive airway pressure on sleep quality in obstructive sleep apnea. *Chest* 1999;116:1545–9.
- [30] Giles TL, Lasserson TJ, Smith BH, White J, Wright J, Cates CJ. Continuous positive airways pressure for obstructive sleep apnoea in adults. *Cochrane Database Syst Rev* 2006;(1):CD001106.
- [31] Keenan SP, Burt H, Ryan CF, Fleetham JA. Long-term survival of patients with obstructive sleep apnea treated by uvulopalatopharyngoplasty or nasal CPAP. *Chest* 1994;105:155–9.
- [32] Faccenda JF, Mackay TW, Boon NA, Douglas NJ. Randomized placebo-controlled trial of continuous positive airway pressure on blood pressure in the sleep apnea-hypopnea syndrome. *Am J Respir Crit Care Med* 2001;163:344–8.
- [33] Bedard MA, Montplaisir J, Malo J, Richer F, Rouleau I. Persistent neuropsychological deficits and vigilance impairment in sleep apnea syndrome after treatment with continuous positive airways pressure (CPAP). *J Clin Exp Neuropsychol* 1993;15:330–41.
- [34] Montplaisir J, Bedard MA, Richer F, Rouleau I. Neurobehavioral manifestations in obstructive sleep apnea syndrome before and after treatment with continuous positive airway pressure. *Sleep* 1992;15:S17–19.
- [35] Engleman HM, Martin SE, Deary IJ, Douglas NJ. Effect of CPAP therapy on daytime function in patients with mild sleep apnoea/hypopnoea syndrome. *Thorax* 1997;52:114–19.
- [36] Kotterba S, Rasche K, Widdig W, et al. Neuropsychological investigations and event-related potentials in obstructive sleep apnea syndrome before and during CPAP-therapy. *J Neurol Sci* 1998;159:45–50.
- [37] Matthews EE, Aloia MS. Cognitive recovery following positive airway pressure (PAP) in sleep apnea. *Prog Brain Res* 2011;190:71–88.
- [38] Naegel B, Pepin JL, Levy P, Bonnet C, Pellat J, Feuerstein C. Cognitive executive dysfunction in patients with obstructive sleep apnea syndrome (OSAS) after CPAP treatment. *Sleep* 1998;21:392–7.
- [39] Gozal D. CrossTalk proposal: the intermittent hypoxia attending severe obstructive sleep apnoea does lead to alterations in brain structure and function. *J Physiol* 2013;591:379–81.
- [40] Djonic I, Saboisky J, Carusona A, Stickgold R, Malhotra A. Increased sleep fragmentation leads to impaired off-line consolidation of motor memories in humans. *PLoS ONE* 2012;7:e34106.
- [41] Kloepper C, Riemann D, Nofzinger EA, et al. Memory before and after sleep in patients with moderate obstructive sleep apnea. *J Clin Sleep Med* 2009;5:540–8.
- [42] Huber R, Ghilardi MF, Massimini M, Tononi G. Local sleep and learning. *Nature* 2004;430:78–81.
- [43] Kuriyama K, Stickgold R, Walker MP. Sleep-dependent learning and motor-skill complexity. *Learn Mem* 2004;11:705–13.
- [44] Walker M, Brakefield T, Morgan A, Hobson JA, Stickgold R. Practice with sleep makes perfect: sleep dependent motor skill learning. *Neuron* 2002;35:205–11.
- [45] Karni A, Meyer G, Rey-Hipolito C, et al. The acquisition of skilled motor performance: fast and slow experience-driven changes in primary motor cortex. *Proc Natl Acad Sci U S A* 1998;95:861–8.
- [46] Fischer S, Hallschmid M, Elsner AL, Born J. Sleep forms memory for finger skills. *Proc Natl Acad Sci U S A* 2002;99:11987–91.
- [47] Van Dongen HP, Maislin G, Mullington JM, Dinges DF. The cumulative cost of additional wakefulness: dose-response effects on neurobehavioral functions and sleep physiology from chronic sleep restriction and total sleep deprivation. *Sleep* 2003;26:117–26.
- [48] Roach GD, Dawson D, Lamond N. Can a shorter psychomotor vigilance task be used as a reasonable substitute for the ten-minute psychomotor vigilance task? *Chronobiol Int* 2006;23:1379–87.
- [49] Iber C, Ancoli-Israel S, Al C, Sf Q. The AASM manual for the scoring of sleep and associated events: rules, terminology, and technical specifications. Westchester, Ill: American Academy of Sleep Medicine; 2007.
- [50] Anbar RD, Hehir DA. Hypnosis as a diagnostic modality for vocal cord dysfunction. *Pediatrics* 2000;106:E81.
- [51] Wamsley EJ, Tucker MA, Shinn AK, et al. Reduced sleep spindles and spindle coherence in schizophrenia: mechanisms of impaired memory consolidation? *Biol Psychiatry* 2012;71:154–61.
- [52] Nishida M, Walker MP. Daytime naps, motor memory consolidation and regionally specific sleep spindles. *PLoS ONE* 2007;2:e341.
- [53] Ungerleider LG, Doyon J, Karni A. Imaging brain plasticity during motor skill learning. *Neurobiol Learn Mem* 2002;78:553–64.
- [54] Canessa N, Castronovo V, Cappa SF, et al. Obstructive sleep apnea: brain structural changes and neurocognitive function before and after treatment. *Am J Respir Crit Care Med* 2011;183:1419–26.
- [55] Bardwell WA, Ancoli-Israel S, Berry CC, Dimsdale JE. Neuropsychological effects of one-week continuous positive airway pressure treatment in patients with obstructive sleep apnea: a placebo-controlled study. *Psychosom Med* 2001;63:579–84.
- [56] Munoz A, Mayoralas LR, Barbe F, Pericas J, Agusti AG. Long-term effects of CPAP on daytime functioning in patients with sleep apnoea syndrome. *Eur Respir J* 2000;15:676–81.
- [57] Feuerstein C, Naegel B, Pepin JL, Levy P. Frontal lobe-related cognitive functions in patients with sleep apnea syndrome before and after treatment. *Acta Neurol Belg* 1997;97:96–107.
- [58] Fogel SM, Smith CT. The function of the sleep spindle: a physiological index of intelligence and a mechanism for sleep-dependent memory consolidation. *Neurosci Biobehav Rev* 2011;35:1154–65.
- [59] Wilson MA, McNaughton BL. Reactivation of hippocampal ensemble memories during sleep. *Science* 1994;265:676–9.
- [60] Kim A, Latchoumane C, Lee S, et al. Optogenetically induced sleep spindle rhythms alter sleep architectures in mice. *Proc Natl Acad Sci U S A* 2012;109:20673–8.
- [61] Gottselig JM, Bassetti CL, Achermann P. Power and coherence of sleep spindle frequency activity following hemispheric stroke. *Brain* 2002;125:373–83.