

Zinc concentration affects the functional groups of microbial communities in sugarcane-cultivated soil

Author links open overlay panel [Acacio A. Navarrete^{ab}](#) [Estêvão V. Mellis^c](#) [Arthur Escalas^d](#) [Leandro N. Lemos^a](#) [José Lavres Junior^e](#) [José Antonio Quaggio^b](#) [Jizhong Zhou^c](#) [Siu M. Tsai^a](#)

Show more

<https://doi.org/10.1016/j.agee.2016.12.009> Get rights and content

Highlights

-

[Zinc](#) concentration in soil affects functional gene structure of [microbial community](#).

-

A threshold in gene family abundance was revealed between 5 and 10 kg Zn ha⁻¹.

-

Gene family responses can assist to define potential soil microbial indicators.

Abstract

The addition of zinc (Zn) to the soil has significant gains in sugarcane productivity. The understanding of the consequences of zinc application requires insight into ecological aspects of changes in soil microbial communities. Here, we evaluated the effects of zinc application on richness, diversity, evenness and structure of microbial functional genes in sugarcane-cultivated soil, and response thresholds of abundance for groups of microbial functional genes on different zinc concentrations added to the soil (0, 5, 10 and 20 kg Zn ha⁻¹) in two sugarcane fields in the Northeastern São Paulo state, Brazil. Using a high-density functional gene array termed GeoChip 5.0, which contains 101,796 distinct probes belonging to gene categories involved in zinc transport, secondary metabolism, stress, virulence and nutrient cycling, differences were observed in the functional gene structure of microbial communities and abundance of specific gene families with a threshold between 5 and 10 kg Zn ha⁻¹. However, the richness, diversity and evenness of functional genes did not differ across the zinc gradients in soil. Cluster analysis based on microbial functional subcategories revealed variation in abundances across the gradient of zinc concentrations in soil mainly for virulence, stress, secondary metabolism, and carbon- and phosphorus-cycling-related gene families. A threshold in abundance was observed between 5 and 10 kg Zn ha⁻¹ with high abundance of

microbial gene families associated with zinc transporter proteins, antioxidant enzymes, exoenzymes associated with infection, secretion proteins, virulence regulatory genes, carbon fixation, and phosphorus utilization in soils supplemented with up to 5 kg Zn ha⁻¹. In these same soils, reduced abundance was observed for gene families associated with methanogenesis, metabolism of halogen and carotenoid, and antiphagocytosis compared with soil supplements with 10 or 20 kg Zn ha⁻¹. The results suggest that changes in microbial functional gene structure and abundance could be used to evaluate the impact of soil management practices on soil microbial communities in sugarcane production fields.

- [**Previous article in issue**](#)
- [**Next article in issue**](#)

Keywords

Soil management

Soil ecosystem service

Microbial functional gene

Microbial indicator

1. Introduction

[Zinc](#) is an essential [cofactor](#) required by all living organisms. Zinc is the second most abundant [transition metal](#) in living organisms (after iron), has important functions in [enzymatic activity](#), [transcription factors](#) (which regulate gene expression), and serves as a cofactor in more than 300 proteins ([Palmgren et al., 2008](#)). According to a global study of the [micronutrient](#) status of soils ([Sillanpää, 1990](#)), the zinc content in soils from Brazilian regions are highly variable, covering a wide range of values from high to low, potentially leading to zinc deficiency for important crops cultivated in this area (e.g., [sugarcane](#), coffee, oranges, rice, maize, [sorghum](#), [cassava](#) and rubber). In this sense, several studies have demonstrated significant gains in sugarcane productivity with zinc application ([Cambria et al., 1989](#), [Marinho and Albuquerque, 1981](#), [Korndörfer et al., 1995](#), [Wang et al., 2005](#), [Mellis et al., 2016](#)).

The number of zinc-binding proteins correlates linearly with the total number of proteins encoded by the genome of an organism, but the proportionality constant of Eukaryota (8.8%) is significantly increased compared with that observed in Bacteria and Archaea (from 5% to 6%) due to the larger portfolio of regulatory proteins in Eukaryota ([Andreini et al., 2006](#)). Zinc levels must be tightly regulated; too little zinc does not support [cellular growth](#), whereas too much zinc is cytotoxic ([Koh et al., 1996](#), [Costello et al., 1997](#)). The

free zinc concentration in the [cytoplasm](#) is maintained well below the picomolar range ([Outten and O'Halloran, 2001](#)). It is generally accepted that accumulated [trace metals](#) reduce the soil microbial biomass ([Brookes and McGrath, 1984](#)) and ([Chander et al., 1995](#)) and the activity of various microbial [enzymes](#) ([Kandeler et al., 1996](#)). Several microbial cellular processes, such as synthesis of [secondary metabolites](#) ([Weinberg, 1990](#)), and mechanisms associated with the infection and virulence of microorganisms ([Campoy et al., 2002](#), [Davis et al., 2009](#)) seem to be sensitive to zinc stress ([Kunito et al., 2001](#)) and zinc [homeostasis](#) ([Hantke, 2001](#)) and thus might be useful for evaluation of the toxic effects of zinc on microorganisms. Soil [microbial communities](#) hold a central place in [terrestrial ecosystems](#), performing essential [ecological processes or functions](#), such as [nutrient cycling](#), plant nutrition or disease suppression ([Bardgett et al., 2008](#)). Therefore, it is useful to understand the consequences of zinc application on diversity and structure of microbial functional genes in [soil ecosystems](#).

Different methods and approaches have been used to study the functional diversity of soil microbial community. Biolog EcoPlate, a technique based on levels of activities on various substrates, was used in different studies to evaluate the metabolic profile of soil microbial communities ([Pajak et al., 2016](#), [Tosi et al., 2016](#), [Kelly et al., 1999](#), [Zak et al., 1994](#)) and determine the relative zinc sensitivity on the metabolic activities of the isolated microorganisms ([Lock and Janssen, 2005](#)). A range of novel methods, most of which are based on rRNA and rDNA analyses, have been developed to assess the functional diversity and structure of microbial communities. [Polymerase chain reaction](#) (PCR)-based fingerprinting, such as [terminal restriction fragment length polymorphism](#) (T-RFLP), has been used to analyze changes in diversity and structure of soil microbial community based on specific functional genes ([Bannert et al., 2011](#)). Quantitative real-time [PCR](#) is useful to analyze differences in the abundance of microbial function-specific genes ([Lammel et al., 2015](#)). However, there are some problems and biases in the PCR amplification step; therefore, these methods cannot be used as definite indicators of richness of ecological communities.

Shotgun [metagenomic sequencing](#) and [gene expression](#) array analysis constitute an alternative approach to taxonomic and functional characterization of soil microbial community that avoids these limitations. The GeoChip-based functional gene array method simultaneously monitors the expression levels of thousands of genes ([He et al., 2007](#)). Such information is very useful in functional diversity studies to track highly expressed genes and genes critical in biogeochemical pathways. It is also important to

obtain information on the microbial community response to [environmental changes and factors](#) that regulate diversity ([Cong et al., 2015](#), [Li et al., 2014](#)).

Considering the potential increases in sugarcane productivity provided by zinc application in natural low-fertility soils worldwide, it is necessary to obtain a better understanding of the effects of zinc application on specialized functioning of the soil performed by microbial groups. For this purpose, we applied a functional gene microarray (GeoChip v. 5.0) (*i*) to assess the effects of zinc application on richness, diversity, evenness and structure of microbial functional genes in sugarcane-cultivated soil and (*ii*) to identify the response thresholds in abundance for groups of microbial functional genes to the zinc concentration added to the sugarcane-cultivated soil.

2. Materials and methods

2.1. Field experiment

Regional-scale [field experiments](#) were performed with RB867515 variety of [sugarcane](#) at the two major producing regions of sugarcane in the State of São Paulo, Brazil: Serra Azul (latitude 21°18'41" S and longitude 47°33'59"W) and Assis (latitude 22°39'40" S and longitude 50°23'58" W) municipalities. The climate in the region is humid subtropical. [Oxisol](#) and [Ultisol](#) are the [soil types](#) at the Serra Azul and Assis municipalities, respectively. The field experiments at both municipalities were characterized by [sandy soil](#) with low natural fertility and available [zinc](#) concentrations of less than 0.5 mg dm⁻³ ([Mellis et al., 2016](#)). Four treatments with three replicate plots each were used in a completely randomized design in each location. Each experimental plot (75 m²) consisted of five neighboring rows, each 10 m long and 1.4 m apart. Zinc was applied in the form of zinc sulfate (ZnSO₄) in the furrow at three different doses (5, 10 and 20 kg ha⁻¹) once at the beginning of the experiment on March 2011. The control treatment consisted of soil without application of zinc. In addition, fertilization was performed at sugarcane planting and consisted of 25 kg nitrogen (N) in the form of [urea](#), 150 kg P₂O₅ (triple superphosphate), 100 kg K₂O (potassium oxide) and 3 kg of [sulfur](#) (S) per hectare. Broadcast fertilization was performed six months after planting (September 2011) with 27 kg N and 40.5 kg K₂O per hectare. In addition, 125 kg N and 100 kg K₂O per hectare were applied in three subsequent cycles of sugarcane ratoon.

2.2. Soil sampling

Soil samples were collected from each of three treatment and control soils in both sugarcane field experiments (4 treatments × 3 soil samples per treatment × 2 experimental fields = 24 soil samples) after the third sugarcane ratoon. [Soil](#)

[sampling](#) consisted of five points in the furrow for each experimental plot: one central sampling point and four other sampling points at least 2.8 m apart from the central point and directed towards the four cardinal points. The soil cores were taken from the 0 to 40 cm topsoil layer using a 5-cm diameter aseptic cylindrical core. Each soil sample consisted of a composite of five soil cores collected within each experimental plot. Soil samples for chemical determination were immediately processed. Aliquots of soil samples were transported to the laboratory on ice and stored at $-20\text{ }^{\circ}\text{C}$ until further processing within 72 h after sampling.

2.3. Soil chemical measurements

The chemical measurements of each of the 24 soil samples were determined according to [EMBRAPA \(2011\)](#). Soil pH was measured on a soil/0.01 M CaCl_2 (1:5) suspension. Aluminum (Al), calcium (Ca), and [magnesium](#) (Mg) were extracted with 1 M [potassium chloride](#). Ca and Mg were determined by atomic absorption [spectrometry](#), whereas Al was determined by acid-base [titration](#). Available phosphorous (P) and [potassium](#) (K) were extracted by [ion-exchange resin](#) and determined by [colorimetry](#) and atomic emission spectroscopy, respectively. Combined results were used for calculation of exchangeable bases (SB) as the sum of Ca, Mg, and K; [cation-exchange capacity](#) (CEC) as the sum of Ca, Mg, K, Al, and H; base saturation (V) as the percent relation between SB and [CEC](#); aluminum saturation (m) as the percent relation between exchangeable Al and CEC; and potential [acidity](#) (H + Al) using an equation based on the pH determined in Shoemaker-McLean-Pratt (SMP) buffer solution. Soil zinc [fractionation](#) was performed according to the Standard [U.S. Environmental Protection Agency \(USEPA\), 1998](#). Briefly, the fractionation involved a [sequential extraction](#) procedure with (a) CaCl_2 to extract water-soluble plus exchangeable zinc (Ex-Zn), (b) Na [acetate](#) (pH 5.0) to extract carbonate-bound zinc (CARB-Zn), (c) NaOCl (pH 8.5) to obtain organically bound zinc (OM-Zn), (d) $\text{H}_2\text{C}_2\text{O}_4\text{-C}_2\text{H}_8\text{N}_2\text{O}_4\text{-C}_6\text{H}_8\text{O}_6$ (pH 3.0) to extract [oxide](#) bound zinc (Ox-Zn), and (e) $\text{HNO}_3\text{-HCl}$ to extract residual zinc (RES-Zn). The percent recovery of the sum of the extracted zinc in all fractions (Ex-Zn, CARB-Zn, OM-Zn, and Ox-Zn) plus the residual fraction (RES-Zn) was expressed as total zinc (TOT-Zn).

2.4. Soil DNA extraction, purification and labeling

[DNA](#) was extracted from 250 mg of soil using the PowerLyzer PowerSoil DNA isolation kit (MO BIO Laboratories Inc., Carlsbad, CA, USA) according to the manufacturer's instructions. Three DNA extractions were performed per soil sample, combined and

concentrated with the [Genomic](#) DNA Clean and Concentrator kit (Zymo Research Corporation, Irvine, CA, USA). Concentrated DNA was re-suspended in 20 μl of [PCR](#) grade water, and purity was assessed spectrophotometrically (Nanodrop ND-1000, NanoDrop Technologies, Inc., Wilmington, DE, USA) to determine [absorbance](#) at the following wavelengths: 230, 260, 280 and 320 nm. DNA concentration was measured with the Quant-iT PicoGreen kit (Molecular Probes/Invitrogen, Carlsbad, CA, USA). The DNA extraction from Assis and Serra Azul samples had an average of 26.90 $\text{ng } \mu\text{l}^{-1}$ (± 1.34) and 23.72 $\text{ng } \mu\text{l}^{-1}$ (± 1.61), respectively. All 24 of the DNA samples were stored at $-20\text{ }^{\circ}\text{C}$ until use.

Purified DNA was labeled with the fluorescent dye Cy-3 according to [Wu et al. \(2006\)](#). Briefly, DNA (600 ng) was mixed with random [primers](#) (300 $\text{ng } \mu\text{l}^{-1}$), denatured at $99.9\text{ }^{\circ}\text{C}$ for 5 min and immediately chilled on ice. A solution containing 5 mM dAGC-TP, 2.5 mM dTTP, Klenow fragment (40 U) and Cy-3 dUTP (25 nM) was added to the denatured DNA, and the reaction volume was adjusted to 50 μl with H_2O . The labeling solution was incubated at $37\text{ }^{\circ}\text{C}$ for 6 h followed by 3 min at $95\text{ }^{\circ}\text{C}$. The labeled DNA was purified with the QIAquick Kit (Qiagen, Valencia, CA, USA), and dye incorporation was confirmed with a NanoDrop [spectrophotometer](#). Labeled samples were vacuum-dried and stored at $-20\text{ }^{\circ}\text{C}$ until [hybridization](#).

2.5. GeoChip 5.0 experiment, scanning and image processing

The GeoChip 5.0 was manufactured by Agilent (Agilent Technologies Inc., Santa Clara, CA, USA) in the $4 \times 180\text{ K}$ format. The chip contains 167,044 distinct probes belonging to different key gene categories involved in carbon (e.g., degradation, fixation, methane), nitrogen (e.g., [ammonification](#), [nitrification](#), fixation), sulfur, phosphorus cycling, metal [homeostasis](#) (e.g., zinc transport), [secondary metabolism](#) (e.g., [antibiotic](#) metabolism, pigments), stress responses (e.g., oxidative stress), virulence (e.g., infection), and other microbial genes of known function. Here, gene sequence refers to each unique sequence targeted by GeoChip 5.0, and a gene family comprises all gene sequences assigned with the same name (e.g., named *ZnuC*) and coding for the same class of proteins. The probes were designed to target Bacteria, Archaea, fungi, and viruses (bacteriophage) gene sequences.

Prior to hybridization, labeled DNA was resuspended in 27.5 μl of DNase/RNase-free distilled water and mixed completely with 99.4 μl of hybridization solution master mix containing 2 x Hi-RPM hybridization buffer (Agilent Technologies Inc., Santa Clara, CA, USA), 10 x aCGH blocking agent (Agilent Technologies Inc., Santa Clara, CA, USA), 10% formamide (J.T. Baker, Philipsburg, NJ, USA), 0.05 $\mu\text{g}/\mu\text{l}$ Cot-1 DNA (Agilent,

Technologies Inc., Santa Clara, CA, USA), and 10 pM Cy5-labelled universal standard DNA for normalization ([Liang et al., 2010](#)). The solution was then denatured at 95 °C for 3 min, incubated at 37 °C for 30 min, and hybridized with GeoChip 5.0 arrays. GeoChip hybridization proceeded at 67 °C in the presence of 10% formamide in a microarray hybridization oven (Agilent Technologies Inc., Santa Clara, CA, USA) for 24 h. After hybridization, the slides were washed using Agilent Wash Buffers 1 and 2 following the manufacturer's protocol. Then, the arrays were scanned at a laser power of 100% and a [photomultiplier tube](#) gain of 75% with a NimbleGen MS200 Microarray Scanner (Roche NimbleGen, Madison, WI, USA). The image data were extracted using the Agilent [Feature Extraction](#) program.

2.6. Raw data processing

Spot alignment and signal intensity determination were performed with the Feature Extraction 11.5 (Agilent Technologies Inc., Santa Clara, CA, USA). The raw data were submitted to the Microarray Data Manager at the Institute for [Environmental Genomics](#) website (<http://www.ou.edu/content/ieg/tools/data-analysis-pipeline.html>) using the following parameters: (i) spots with signal intensity lower than 1.3 times background or with a [signal-to-noise ratio](#) (SNR) less than 2.0 were removed as poor-quality spots (SNR = spot signal intensity – background signal average/SD background); (ii) unrepresentative data were removed using group cut 1 (by default) to remove singletons (probes that were positive in only one sample of an experimental group); (iii) normalization was performed by dividing the signal intensity of each spot by the average of the universal standard spots and then by the average of the signal intensities in each sample (i.e., mean-ratio approach).

2.7. Statistical analysis

One-way ANOSIM (analysis of similarity) was performed on Bray-Curtis dissimilarity matrices (incorporating 999 permutations for R statistics) after a process of [standardization](#) of all variables using the L_1 distance to determine the significance of differences across treatments regarding all soil chemical factors and only the zinc concentration in soil. All procedures were computed using PRIMER v6.1.9 (Primer-E Ltd, Plymouth, UK). One-way ANOVA (analysis of variance) with post hoc Tukey's HSD test was used to determine the significance of the differences among treatments for each soil factor within each of the two field experiments. The procedure was performed using the STATISTICA 10 package (StatSoft Inc., Tulsa, OK, USA). Spearman's rank correlation coefficients were calculated to explore the relationship between gene family

abundance and zinc concentration in soil using the R package 'multtest' (version 2.6.0; [R Core Team, 2015](#)) and the false discovery rate controlling procedure ([Benjamini and Hochberg, 1995](#)). The abundance of each gene family was calculated based on the sum of all gene sequences assigned with the same name and coding for the same class of proteins. Ecological metrics, such as gene richness, diversity (Shannon and inverse Simpson indexes) and evenness (Sheldon index), were calculated using the 'diversity' function from the R package 'vegan' ([R Core Team, 2015](#)). A Tukey's test was used to determine the significance of the differences among treatments for each ecological metric within each of the two field experiments. The similarities in gene family composition across the samples were visualized by multidimensional scaling (MDS) ordination and tested by ANOSIM using Bray-Curtis index ([Clarke, 1993](#)) based on the abundance of gene families correlated to the zinc concentration in soil. To determine significant differences between two zinc rate categories defined by MDS, the Statistical Analysis of [Metagenomic](#) Profiles (STAMP) software package was used ([Parks and Beiko, 2010](#)). *P*-values were calculated using the two-sided Welch's *t*-test ([Welch, 1947](#)), whereas confidence intervals were calculated using Welch's inverted test. In addition, correction was performed using Benjamini-Hochberg false discovery rate. The total gene abundance in different functional subcategories was ordered in a [heat map](#). Dissimilarity between samples and gene abundance within functional subcategories was estimated using the 'vegdist' function from the R package 'vegan' and with Bray-Curtis index ([R Core Team, 2015](#)). Clustering was performed using the average linkage algorithm (also called [UPGMA](#) for Unweighted Pair Group Method with Arithmetic Mean) ([R Core Team, 2015](#)).

3. Results

3.1. Content of zinc in the soil

The [zinc](#) concentration in soils revealed the intentional gradient defined by the treatments containing the different zinc concentration, the total content of zinc and zinc fractions that differed significantly among the treatments ([Table 1](#)) (ANOSIM, Assis: $R = 0.989$, $P < 0.001$; Serra Azul: $R = 0.966$, $P < 0.001$). Based on the [soil chemical properties](#) (Supplementary Table S1), the three zinc treatments did not form significantly distinct groups for either [sugarcane field experiment](#) (ANOSIM, Assis: $R = 0.436$, $P < 0.001$; Serra Azul: $R = 0.448$, $P < 0.001$).

Table 1. Total content of [zinc](#) (Zn) and its fractions in sugarcane-cultivated soils under different zinc concentration.

Treatments (Zinc concentration)	Zinc fractions in soil					TOT-Zn (mg dm ⁻³)
	Ex-Zn (mg dm ⁻³)	CARB-Zn (mg dm ⁻³)	OM-Zn (mg dm ⁻³)	Ox-Zn (mg dm ⁻³)	RES-Zn (mg dm ⁻³)	
Assis (kg ha ⁻¹)						
0	0.03 ^a ± 0.004 [§]	0.02a ± 0.004	0.05a ± 0.002	0.03a ± 0.008	0.11a ± 0.011	0.24a ± 0.006
5	1.03b ± 0.002	1.03b ± 0.006	1.08b ± 0.001	1.06b ± 0.002	1.35b ± 0.018	5.57b ± 0.013
10	2.04c ± 0.006	2.03c ± 0.004	2.08c ± 0.004	2.06c ± 0.008	2.65c ± 0.042	10.86c ± 0.032
20	3.04d ± 0.001	3.03d ± 0.004	3.08d ± 0.019	3.06d ± 0.001	3.48d ± 0.067	15.70d ± 0.058
Serra Azul (kg ha ⁻¹)						
0	0.04a ± 0.007	0.05a ± 0.014	0.09a ± 0.007	0.06a ± 0.025	0.14a ± 0.041	0.38a ± 0.093
5	1.03b ± 0.003	1.03b ± 0.010	1.09b ± 0.003	1.07b ± 0.005	1.33b ± 0.029	5.55b ± 0.021
10	2.04c ± 0.001	2.03c ± 0.002	2.08c ± 0.004	2.06c ± 0.007	2.55c ± 0.170	10.76c ± 0.165
20	3.05d ± 0.004	3.02d ± 0.012	3.08d ± 0.028	3.06d ± 0.002	3.45d ± 0.131	15.65d ± 0.100

Ex-Zn, exchangeable zinc; CARB-Zn, carbonate-bound Zn; OM-Zn, organically bound Zn; Ox-Zn, [oxide](#) bound Zn; RES-Zn, residual zinc; TOT-Zn, total zinc.

†

Average for each of three replicates of soil.

‡

Tukey's HSD test followed by Bonferroni correction for multiple comparisons was performed considering all treatments regarding each of two field experiments (Assis and Serra Azul) across replicates of soil for each treatment. Significance level: $P < 0.05$.

§

Standard deviation of the average for each of three replicates of soil.

3.2. Ecological metrics and structure similarity of microbial functional genes

A total of 40,489 out of 101,796 (40%) microbial functional genes targeted by GeoChip 5.0 were detected across all soil samples from Assis and Serra Azul experimental fields. These genes belonged to 525 different gene families and were associated with the following functional categories: zinc transport; carbon-, nitrogen-, phosphorus-, and sulfur-cycling; [secondary metabolism](#); stress and virulence. Overall, 53 of the 525 detected gene families (10%) were significantly correlated with zinc concentration in soil ([Table 2](#)). Gene richness did not differ across the zinc gradient within each of the two experiments ([Table 3](#)). Functional gene diversity based on Shannon and inverse Simpson [diversity indexes](#) and evenness calculated using Sheldon index were also not

significantly different across the zinc gradient within each of the two experiments ([Table 3](#)). Although the richness, diversity and evenness of gene and gene families did not differ between [microbial communities](#) across the zinc gradient in soil, we observed differences in the identity of genes correlated with zinc concentration ([Fig. 1](#)). ANOSIM revealed similar grouping with some differences for control treatment and soil amended with 5 kg Zn ha⁻¹ in both field experiments ([Table 4, Fig. 1](#)). Similar patterns were observed for soils amended with 10 and 20 kg Zn ha⁻¹ ([Table 4, Fig. 1](#)). However, clear difference were observed between both described clusters ($R = 0.987$).

Table 2. Quantity of gene families and gene sequences detected by GeoChip 5.0 and correlated with [zinc](#)(Zn) concentration belonging to gene categories in both [field experiments](#).

Gene categories	Gene families		Gene sequences	
	Detected	Correlated with Zn [†]	Detected	Correlated with Zn [†]
Zinc transport	14	5	2145	972
Carbon cycling	118	15	13145	785
Nitrogen cycling	29	2	1041	379
Phosphorus cycling	6	2	1668	23
Sulfur cycling	26	0	1730	0
Secondary metabolism	52	6	1356	127
Stress	85	8	9259	869
Virulence	195	15	10145	335
Total	525	53	40489	3490

†

Values indicate the number of gene families and gene sequences significantly ($P < 0.05$) correlated with zinc concentration in soil (total content of Zn and its fractions).

Table 3. Ecological metrics based on functional gene sequences and gene families detected by GeoChip 5.0 and correlated with [zinc](#) concentration in sugarcane-cultivated soils.

Assis (Zinc concentration)				Serra Azul (Zinc concentration)			
0 kg ha ⁻¹	5 kg ha ⁻¹	10 kg ha ⁻¹	20 kg ha ⁻¹	0 kg ha ⁻¹	5 kg ha ⁻¹	10 kg ha ⁻¹	20 kg ha ⁻¹
<i>Richness estimates</i>							
Gene sequences detected							
12628 ^a ab ^b ± 169 ^a	12480 ab ± 100	12249 a ± 611	13366 b ± 721	13653 a ± 280	14024 a ± 157	13105 a ± 139	13393 a ± 275
Gene sequences belonging to gene families correlated with zinc concentration							
2632 a ± 40	2577 a ± 10	2547 a ± 131	2783 a ± 159	3007 ab ± 131	3266 b ± 105	2768 a ± 58	2874 ab ± 87


Assis (Zinc concentration)				Serra Azul (Zinc concentration)			
0 kg ha⁻¹	5 kg ha⁻¹	10 kg ha⁻¹	20 kg ha⁻¹	0 kg ha⁻¹	5 kg ha⁻¹	10 kg ha⁻¹	20 kg ha⁻¹
<i>Gene families detected</i>							
501 a ± 3	499 a ± 1	500 a ± 8	511 a ± 9	513 a ± 3	519 a ± 3	506 a ± 3	509 a ± 5
<i>Gene families correlated with zinc concentration</i>							
70 a ± 7	69 a ± 3	64 a ± 6	77 a ± 6	69 a ± 4	79 a ± 12	67 a ± 9	67 a ± 5
<i>Shannon index</i>							
<i>Gene families detected</i>							
4.81 a ± 0.005	4.80 a ± 0.003	4.80 a ± 0.01	4.81 a ± 0.01	4.82 a ± 0.002	4.83 a ± 0.002	4.81 a ± 0.003	4.82 a ± 0.005
<i>Gene families correlated with zinc concentration</i>							
2.97 a ± 0.005	2.98 a ± 0	2.97 a ± 0.01	2.99 a ± 0.01	2.99 a ± 0.01	2.99 a ± 0.005	2.98 a ± 0.01	2.98 a ± 0.005
<i>Inverse Simpson index</i>							
<i>Gene families detected</i>							
41.99 a ± 0.42	41.61 a ± 0.30	41.49 a ± 0.74	42.0 a ± 0.90	43.29 a ± 0.20	43.56 a ± 0.19	42.34 a ± 0.31	43.04 a ± 0.26
<i>Gene families correlated with zinc concentration</i>							
12.76 a ± 0.12	12.76 a ± 0.01	12.59 a ± 0.31	13.14 a ± 0.24	13.09 a ± 0.12	13.29 a ± 0.03	12.86 a ± 0.11	13.17 a ± 0.22
<i>Sheldon index</i>							
<i>Gene families detected</i>							
0.77 a ± 0.0002	0.77 a ± 0.0004	0.77 a ± 0.0005	0.77 a ± 0.0003	0.77 a ± 0.0004	0.77 a ± 0.0003	0.77 a ± 0.0003	0.77 a ± 0.0004
<i>Gene families correlated with zinc concentration</i>							
0.76 a ± 0.0017	0.76 a ± 0.0016	0.76 a ± 0.0046	0.76 a ± 0.0023	0.76 a ± 0.0018	0.76 a ± 0.0003	0.76 a ± 0.0021	0.76 a ± 0.0016

†

Average for each of three replicates of soil.

‡

Samples under different zinc addition were contrasted within each field experiment. Values with the same lower-case letters were not significantly different ($P < 0.05$) based on Tukey's HSD test followed by Bonferroni correction for multiple comparisons.



1. [Download high-res image \(142KB\)](#)
2. [Download full-size image](#)

Fig. 1. Multidimensional scaling of functional gene families significantly correlated with [zinc](#) concentration in sugarcane-cultivated soils. Correlation was determined using Pearson coefficient, and we used a confidence level of $P < 0.05$.

Table 4. Analysis of similarity (ANOSIM). Pairwise comparison between functional gene family structures of [microbial communities](#) in sugarcane-cultivated soils under different [zinc](#) concentrations.

	0 kg Zn ha ⁻¹	5 kg Zn ha ⁻¹	10 kg Zn ha ⁻¹	20 kg Zn ha ⁻¹
0 kg Zn ha ⁻¹	–	0.072	0.793	0.851
5 kg Zn ha ⁻¹	0.005 [*]	–	0.754	0.810
10 kg Zn ha ⁻¹	0.006 [*]	0.007 [*]	–	0.254
20 kg Zn ha ⁻¹	0.005 [*]	0.005 [*]	0.006 [*]	–

Upper triangle indicates ANOSIM statistic based on Bray-Curtis dissimilarity between two different zinc concentrations applied to the soil. Lower triangle represents P -values of the test.


*

$P < 0.05$.

3.3. Variation in functional groups of microbial communities to zinc concentration in soil

Abundance-based [cluster analysis](#) of functional subcategories revealed four different bunches (I, II, III and IV) ([Fig. 2](#)). The highest variation in gene family abundances across the samples was noted for group III, encompassing virulence, carbon- and phosphorus-cycling and metabolic-related gene families. Virulence and stress-related

gene families that clustered as group II also exhibited increased variation across all samples (Fig. 2). Zinc transporter gene families exhibited increased abundance in control soil and soil amended with 5 kg Zn ha⁻¹ compared with soils supplemented with higher zinc concentrations in both field experiments (Fig. 2).



1. [Download high-res image \(589KB\)](#)
2. [Download full-size image](#)

Fig. 2. [Heatmap](#) and [cluster analysis](#) of samples based on the dissimilarity in gene family abundance correlated with [zinc](#) concentration. Gene families (rows) and samples (columns) were clustered based on microarray abundance data of gene families significantly correlated with zinc concentration in soil. The heatmap colors correspond to the normalized signal intensity of the gene families. The color gradient represents the signal values across all genes/functional subcategories and samples, varying from cold colors (lower signal) to warm colors (higher signal). On the x-axis, samples were reordered and clustered according to their color intensity profile. On the y-axis, genes were grouped by functional subcategory, and their abundance was summed.

Individual responses of gene families to increases in zinc concentration in soil are presented in [Table 5](#). The majority of the gene families that were significantly correlated with total content of zinc and its fractions exhibited a decrease in abundance with increasing zinc concentration in soil. A similar pattern was observed based on the differences in abundance for gene families between two zinc rate categories (0 and 5 kg Zn ha⁻¹ and 10 and 20 kg Zn ha⁻¹), indicating a threshold of zinc effect on gene family abundance between 5 and 10 kg Zn ha⁻¹ ([Fig. 3](#)). Microbial functional gene families associated with zinc transport (e.g., cation-transporting membrane facilitators), secondary metabolism (e.g., indole-3-pyruvic [acid](#) [metabolic](#) routes, [magnesium](#) chelatase, and [antioxidant enzymes](#) peroxidase), [oxidative stress](#) (e.g., [superoxide](#) dismutase), oxygen, [glucose](#), and phosphate limitation (e.g., transcriptional regulators), stress (e.g., regulator of the sigma factor, drought tolerance, envelope stress), infection (e.g., adherence, virulence, and dissolution of the extracellular matrix), secretion system (e.g., secretion protein, immune evasion), survival against host response (e.g., extracellular zinc metalloprotease, [biofilm](#) formation) and virulence (virulence protein, virulence [regulatory gene](#), [patulin](#) biosynthesis) exhibited reduced abundance with increasing zinc concentrations in soil ([Table 5](#)). Additionally, microbial functional gene families associated with carbon degradation (e.g., glyoxylate cycle/isocitrate [lyase](#), [pectin](#), [lignin](#), and phospholipids), [carbon fixation](#) (e.g., dicarboxylate/4-hydroxybutyrate cycle, reductive tricarboxylic acid cycle, 3-hydroxypropionate bicycle, Calvin cycle), [denitrification](#), dissimilatory nitrogen reduction, and polyphosphate degradation also exhibited reduced abundance with increasing zinc concentrations in soil ([Table 5](#)). Increased abundance was demonstrated for gene families associated with [methanogenesis](#), and secondary metabolism (e.g., [halogenation](#) and carotenoid) with increasing zinc concentrations in soil ([Table 5](#)). The microbial gene families 'siaC' and 'lip', which are associated with antiphagocytosis, exhibited contrasting responses to

increasing zinc concentrations in soil, with increases in abundance for the 'siaC' gene family. [Gene probe](#) origins for the gene families significantly correlated with zinc status in soil are presented in Supplementary Table S2.

Table 5. Gene families significantly correlated with [zinc](#) concentration in soil (total content of zinc and its fractions) in both [field experiments](#).


Gene families	Estimate	Gene families	Estimate
Zinc transport		Survival against host response	
MSC2	-0.4287	<i>siaC</i>	0.4626
YiiP (FieF)	-0.5333	lip	-0.4083
<i>ZntA</i>	-0.6826	<i>zmpA</i>	-0.4698
<i>ZnuC</i>	-0.4544	<i>amsG</i>	-0.4410
<i>Zrt2</i>	-0.4758	Virulence (others)	
Secondary metabolism		Vip	-0.4165
per arc	-0.5249	ler	-0.4629
per fun	-0.4274	patulin (6MSAS)	-0.4087
V BPO	0.4286	Carbon degradation	
mg chelatase	-0.5011	<i>AceA</i>	-0.4178
methoxyneurosporene desaturase	0.4494	Endopolygalacturonase	-0.4555
IPyA	-0.4198	Exopolygalacturonase	-0.5530
Oxidative stress		GLX	-0.4301
<i>sodA</i>	-0.4940	phospholipase C	-0.4310
Oxygen limitation		Carbon fixation	
Arca	-0.4698	C CoA hydratase DiC4HB	-0.4566
Glucose limitation		FBPase	-0.4211
<i>ccpA</i>	-0.4121	<i>frdA</i> (rTCA)	-0.4796
Phosphate limitation		mdh	-0.4344
<i>phoB</i>	-0.4110	MMCE	-0.4616
Stress (others)		Rubisco	-0.5825
<i>RseA</i>	-0.4166	<i>sdhA</i>	-0.4496
tre fun	-0.6035	<i>sucA</i> (DiC4HB)	-0.4583
CpxR	-0.4929	Methanogenesis	
<i>PspB</i>	-0.4079	<i>mrtH</i>	0.4240
Infection		MtaB	0.4528
pilin	-0.4746	Denitrification	

Gene families	Estimate	Gene families	Estimate
<i>pap</i>	-0.4730	<i>nirS</i>	-0.4356
<i>bfpB</i>	-0.4670	Dissimilatory N reduction	
<i>srt</i>	-0.4404	<i>Napa</i>	-0.4311
<i>hysA</i>	-0.5013	Phosphorus utilization	
Secretion system		5f1 (<i>ptxD</i>)	-0.4100
<i>hrpQ</i>	-0.5414	5f1 (ppn)	-0.5730
YopM	-0.5026		
<i>lcrD</i>	-0.4525		

Gene families displayed in the table with Spearman's rank coefficient at the $P < 0.05$ level. Gene categories are indicated in bold.

Zinc rate categories

□ 0 and 5 kg ha⁻¹
 ■ 10 and 20 kg ha⁻¹



1. [Download high-res image \(249KB\)](#)
2. [Download full-size image](#)

Fig. 3. Differences in abundance for gene families between two [zinc](#) rate categories revealed by using Statistical Analysis of [Metagenomic](#) Profiles (STAMP) software based on abundance of gene families correlated with zinc concentration in soil from both [field experiments](#) (Assis and Serra Azul).

4. Discussion

In agriculture, [zinc](#) is used as a nutritional supplement to promote plant growth. However, the incorporation of zinc into the soil may have complex effects on [soil microorganisms](#), and the functional consequences of zinc supplementation are still not

fully understood. In the present study, we used microarray-based high throughput [genomic](#) technologies to assess [functional response](#) at the community-level to zinc supplementation in [sugarcane](#) experimental fields. Our findings revealed significant differences in the functional gene structure of [microbial communities](#) and abundance of specific gene families across a gradient of zinc concentrations in these soils. However, the richness, diversity and evenness of functional genes did not differ across the zinc gradients in soil. This finding suggests that although the total number of functional genes within the microbial communities does not change, the identity of the present genes and the abundance of specific gene families are altered along the zinc gradient, with a clearer threshold in gene family abundance between 5 and 10 kg Zn ha⁻¹.

4.1. Impact of zinc inputs to the soil on richness, diversity, evenness and structure similarity of microbial functional genes

Classical observations on toxic effects of heavy metals on field organisms, such as population size or ecological aspects at the community level (richness, diversity and evenness), are often difficult to interpret due to unexplained variation (or absence of variation) and unclear causality. It is often hypothesized that due to functional redundancy, the diversity of species in a community is more sensitive to toxicant stress than community functioning itself ([Bååth, 1989](#), [Moffett et al., 2003](#)). However, [Lock and Janssen \(2005\)](#) showed that functional diversity of microbial communities decreased with increasing zinc concentrations in 11 [soils contaminated](#) with zinc runoff from galvanized electricity pylons. In the present study, analysis of similarity between the samples suggested that there is a threshold in gene family abundance between 5 and 10 kg Zn ha⁻¹, which did not caused a change in richness, diversity and evenness of microbial functional genes in soil. This finding could, however, be expected as functional richness, diversity and evenness are not only influenced by the zinc concentration but also by numerous other soil factors.

4.2. Impact of zinc inputs to the soil on microbial functional groups

4.2.1. Zinc transport

Cells require zinc transport mechanisms to allow efficient accumulation and distribution of zinc ions within the cell ([Eide, 2006](#)). In many bacteria, two well-characterized regulatory systems are used to ensure zinc [homeostasis](#), and these genetic pathways act by cooperatively controlling the expression of membrane-bound [transport systems](#) that either import or export zinc [cations](#) ([Hantke, 2001](#)). The zinc uptake

system, *Znu*, is employed by numerous Gram-negative bacteria to import zinc cations, and it is composed of an ABC-type [transporter](#), where *ZnuC* is the associated [ATPase](#) protein ([Chandra et al., 2007](#)). Additionally, the expression of the *ZnuABC* gene families is often controlled by a zinc-responsive transcriptional regulator of the *Fur* system called *Zur* ([Hantke, 2005](#)). The export of zinc from bacterial cells is often accomplished using the *ZntA* protein, which is an ATP-dependent transporter employed when cellular zinc concentrations are at toxic levels ([Hantke, 2001](#)). Previous studies have suggested that the efficient uptake of zinc could also play a critical role in virulence and during infection ([Hantke, 2005](#), [Ammendola et al., 2007](#), [Sheehan et al., 2015](#)). Our results are consistent with these previous observations given that zinc transport related gene families were adversely affected by increases in zinc concentrations, notably the cation-transporting membrane facilitators. The zinc effect on both microbial infection and virulence are discussed below.

4.3. Secondary metabolism

The [secondary metabolism](#) pathways potentially affected by zinc inputs were associated with the synthesis of [auxin indole-3-acetic acid](#) (IAA), [chlorophyll](#) and [heme](#) and the control of [oxygen-reactive species](#). The associated gene families exhibited decreased abundance under high zinc conditions. In microorganisms, the indole-3-pyruvic acid (IPyA) pathway is the main route for IAA synthesis ([Spaepen et al., 2007](#)). Interactions between IAA-producing bacteria and plants lead to diverse outcomes on the plant side, varying from [pathogenesis](#) to [phytostimulation](#) ([Spaepen et al., 2007](#)).

In [photosynthetic](#) organisms, the Mg-chelatase catalyzes the insertion of Mg^{2+} into protoporphyrin IX, which is the first step unique to the synthesis of (bacterio)chlorophyll. Magnesium-protoporphyrin IX chelatase (Mg-chelatase) is located at the branch point of tetrapyrrole [biosynthesis](#), at which point protoporphyrin IX is distributed for the synthesis of chlorophyll and heme. Regarding the control of oxygen-reactive species, microorganisms have developed efficient enzymatic and non-enzymatic mechanisms to eliminate [reactive oxygen](#) intermediates, e.g., [superoxide radicals](#), [hydrogen peroxide](#) and [hydroxyl radicals](#). Superoxide is eliminated by dismutation to H_2O_2 as catalyzed by superoxide dismutase, and the accumulation of H_2O_2 is prevented by the action of [catalases](#) and peroxidases ([Fridovich, 1978](#), [Zima et al., 1996](#)).

In contrast, [halogen](#) metabolism (vanadium-dependent bromoperoxidases 'V-BPO') and [carotenoid](#) biosynthesis ('Methoxyneurosporene desaturase')-related gene families exhibited an increase in abundance with zinc input. The 'V BPO' gene family is related to [antioxidative](#) capacities ([Ye et al., 2015](#)). Carotenoids play important biological roles

as accessory light-harvesting components of photosynthetic systems, photoprotecting antioxidants, and regulators of [membrane fluidity](#).

4.4. Stress (oxidative stress, and oxygen, glucose, and phosphate limitation)

Decreases in abundance for superoxide dismutase 'sodA' with increases in zinc concentration in our soils may be understood considering the key role played by SOD in the protection against harmful oxidative reaction resulting from metal stress ([Monk et al., 1989](#)). The 'sodA' is a gene family related to [oxidative stress](#). In regards to oxygen- and nutrient-responsive gene families ('arcA', 'ccpA' and 'phoB'), decreases in the expression of these proteins can alter transcriptional regulation, i.e., the means by which cells regulate the conversion of [DNA](#) to [RNA](#), thereby influencing gene activity. Additionally, the stress-related gene family ('RseA') also exhibited reduced abundance with zinc inputs to our soils. 'RseA' codes for a regulator of a sigma factor, a bacterial transcription [initiation factor](#) that enables specific binding of RNA polymerase to gene promoters. Taken together, these findings suggest adverse effects of zinc on the transcriptional process in microbial cells, which may be related to the fact that zinc is a component or activator of DNA-dependent RNA polymerase ([Coleman, 1974](#)), RNA-dependent DNA polymerase ([Auld et al., 1974](#)) and participates in the maintenance of [ribosome](#) integrity ([Prask and Plocke, 1971](#)) and the double helix of DNA ([Chvapil, 1973](#)). Finally, our results suggest a decrease in abundance for 'CpxR', a gene family related with envelope stress. The Cpx system responds to conditions associated with envelope stress, such as alkaline pH and overproduction of secreted proteins ([Langen et al., 2001](#), [Snyder et al., 1995](#)), and to attachment of cells to surfaces ([Otto and Silhavy, 2002](#)).

4.5. Infection

Adherence is the main microbial function related to infection that was adversely affected by zinc inputs to our soils. The zinc effect on microbial adherence appears to be inconsistent in the literature and could be organism dependent. Based on assays with Gram-negative and Gram-positive bacteria, [Sugarman et al. \(1982\)](#) suggested that zinc can bind to bacterial pili and increase bacterial adherence. [Crane et al. \(2007\)](#) showed that zinc reduced the abundance of the RNAs encoded by bundle-forming pilus (*bfp*), a protein virulence factor in *Escherichia coli*. A potential adverse effect of zinc input to the soil was also highlighted in the exoenzyme hyaluronidase (also called hyaluronate lyase) encoded by the 'hysA' gene. Hyaluronidases are important for virulence in a number of Gram-negative and Gram-positive bacteria ([Hart et al., 2009](#)). These findings

add evidence to the critical importance of tight regulation of zinc to soil microbial infection.

4.6. Secretion system

Microorganisms have evolved a wide variety of highly specialized macromolecular nanomachines that secrete a wide range of substrates, including [small molecules](#), proteins and DNA, which may be necessary for zinc homeostasis ([Serafini et al., 2014](#)). These substrates have key roles in the response of bacteria to its environment and in several physiological processes ([Costa et al., 2015](#)). In our soils, increases in zinc concentrations adversely affected secretion systems related to microbial functions encoded by '*hrpQ*', '*YopM*' and '*lcrD*' genes. In phytopathogenic bacteria, the '*hrp*' system (for [hypersensitive reaction](#) and pathogenicity) encodes components of type III secretion system (TTSS) ([Lindgren et al., 1986](#)). TTSS translocates effector proteins into [target cells](#) to disrupt or modulate host cell signaling pathways and establish replicative niches. *YopM*, a leucine-rich repeat TTSS protein, is a critical virulence factor in infection ([McCoy et al., 2010](#)). The *Yop* secretion system also involves *lcrD*, a gene of the *virA* locus, which encodes a 77-kDA inner [membrane protein](#). Homologues of *lcrD* are involved in [flagellar biogenesis](#) in bacteria ([Ramakrishnan et al., 1991](#)). Taken together, these effects can potentially result in differences in physiological processes in microorganisms inhabiting our soils, such as adhesion, pathogenicity, adaptation and survival.

4.7. Survival against host response

Both specific and nonspecific host responses are critical for host survival. Without them, microorganisms would multiply and invade vital tissues and organs, resulting in severe damage to the host. Some nonspecific responses are biochemical, whereas others are cellular. Our findings revealed a negative effect of zinc on gene families related to cellular microbial responses associated with antiphagocytosis ('*lip*') and [biofilm](#) formation ('*amsG*') along with biochemical microbial responses such as extracellular zinc metalloprotease ('*zmpA*'). Zinc metalloproteases have been described in *Burkholderia cepacia*, a soil [saprophyte](#) Gram-negative bacteria ([McKevitt et al., 1989](#), [Corbett et al., 2003](#)). The capsular gene families '*siaC*' and '*lip*' exhibited positive and negative responses to zinc inputs to our soils, respectively. Capsular [gene expression](#) is regulated both by phase variation via slipped-strand mispairing or reversible insertion of mobile elements ([Hammerschmidt et al., 1996a](#), [Hammerschmidt et al., 1996b](#)) and at the transcriptional level. Indeed, evidence suggests that capsule

biosynthesis and assembly are down-regulated during the early stages of the infectious cycle to facilitate adhesion to and invasion of the host cells ([Deghmane et al., 2002](#)).

4.8. Carbon degradation and fixation

The average composition of structural [carbohydrates](#) of sugarcane straw is 45–48% [cellulose](#), 26–31% [hemicellulose](#) and 7–20% [lignin](#) ([Singh et al., 2008](#)). [Endopolygalacturonase](#) and exopolygalacturonase are [enzymes](#) that act on [pectin](#), hydrolyzing its internal and external glycosidic bonds and producing shorter pectin [molecular structures](#) ([Souza et al., 2003](#)). Glyoxal [oxidase](#) (GLX) is presumed to generate radical species that participate in further reactions to degrade the lignin polymer ([Kirk and Farrell, 1987](#)). These enzymes involved in pectin and lignin degradation encoded by microbial genes were adversely affected by zinc inputs to our soils. Whereas microorganisms have metabolic requirements for [trace metals](#), the same metals are often toxic at concentrations only a few times greater than those required ([Hughes and Poole, 1991](#)). For example, [Davies and Sleep \(1979\)](#) demonstrated inhibition of [carbon fixation](#) as a function of zinc uptake in natural [phytoplankton](#) assemblages. C-fixation is a highly important source of C in soils. In our soils, gene families associated with C-fixation pathways were also adversely affected by zinc inputs to the soil.

4.9. Methanogenesis

Although the inhibition of zinc on [methanogens](#) has been widely studied ([Hayes and Theis, 1978](#), [DeWalle et al., 1979](#), [Bhattacharya et al., 1996](#)), our results revealed an increase in the gene families involved with both hydrogenotrophic ('*mtrH*') and acetoclastic [methanogenesis](#) ('*MtaB*'). Some studies have shown that, up to a certain dosage, metals can work as stimulating agents in [anaerobic digestion](#) processes with a consequent increase in [methane](#) production ([Demirel and Scherer, 2011](#), [Mudhoo and Kumar, 2013](#)).

4.10. Denitrification and dissimilatory nitrogen reduction

Microbial [denitrification](#) is a major pathway responsible for [nitrate](#) removal and a dominant source of [nitrous oxide](#) (N₂O) emissions from soil. [Zheng et al. \(2014\)](#) demonstrated that zinc substantially inhibited the gene expression and [catalytic activity](#) of key denitrifying enzymes in the same manner as noted in our findings. These negative effects of zinc on microbial denitrification may ultimately cause reduced nitrate removal and increased N₂O emissions, which is likely to exacerbate [global warming](#). An

inhibitory effect of zinc on nitrogen reduction has also been reported by [Stone et al. \(2006\)](#).

4.11. Phosphorus utilization

Utilization of organic P by microorganisms and plants requires [mineralization](#) (hydrolysis) of substrates by [phosphatase](#) enzymes that may be of either microbial or plant origin, with some evidence that microbial enzymes exhibit increased efficiency for P release ([Tarafdar et al., 2001](#), [Richardson and Simpson, 2011](#)). Zinc inputs to our soils reduced the abundance of gene families related to phosphorus oxidation and polyphosphate degradation. [Herbert \(1961\)](#) demonstrated that the distribution of P forms in microbial cells changes with environmental conditions, in particular with carbon and [nutrient availability](#). However, zinc effects on microbial phosphorus utilization were not reported previously in the literature.

5. Conclusion

Taken together, our findings demonstrate that increases in [zinc](#) concentration in sugarcane-cultivated soil can alter the functional gene structure of [microbial communities](#) and abundance of specific gene families, with a threshold in gene family abundance between 5 and 10 kg Zn ha⁻¹ across the analyzed zinc concentrations. Responses of functional gene family abundance in microbial communities to zinc status in our soils were at least in part associated with zinc transport, [antioxidative](#) capacity, virulence, [carbon-fixation](#), [methanogenesis](#), phosphorus utilization, metabolism of [halogen](#) and [carotenoid](#), and antiphagocytosis, which may reveal implications on soil microbial processes not only essential to [ecosystem function](#) but also for the sustainable management of [agricultural ecosystems](#). Finally, these findings could assist in the identification of gene families with potential use as management-indicators of [agricultural practices](#) based on specialized functioning of the soil conducted by microbial groups in [sugarcane](#) production fields.

Acknowledgments

This study was supported by a grant from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP – 2012/13321-7, 2015/09436-1 and 2011/51749-6). LN Lemos was supported by Brazilian National Council for Scientific and [Technological Development](#) (CNPq – 161931/2015-4). We wish to thank Wagner Picinini for his technical support during the [soil sampling](#); Joy D Van Nostrand from the Institute for [Environmental Genomics](#) (IEG), University of Oklahoma (OU) for her analytical suggestions; and Tong Yuan from IEG-OU for her technical laboratory support.

Appendix A. Supplementary data

The following is Supplementary data to this article:

[Download Word document \(43KB\)](#)[Help with docx files](#)

References

[Ammendola et al., 2007](#)

S. Ammendola, P. Pasquali, C. Pistoia, P. Petrucci, P. Petrarca, G. Rotilio, A. Battistoni **High-affinity Zn²⁺ uptake system *ZnuABC* is required for bacterial zinc homeostasis in intracellular environments and contributes to the virulence of *Salmonella enterica***
Infect. Immun., 75 (2007), pp. 5867-5876

[CrossRefView Record in Scopus](#)

[Andreini et al., 2006](#)

C. Andreini, L. Banci, I. Bertini, A. Rosato **Zinc through the three domains of life**
J. Proteome Res., 5 (2006), pp. 3173-3178

[CrossRefView Record in Scopus](#)

[Auld et al., 1974](#)

D.S. Auld, H. Kawaguchi, D.M. Livingston, B.L. Vallee **Reverse transcriptase from avian myeloblastosis virus: a zinc metalloenzyme**
Biochem. Biophys. Res. Commun., 57 (1974), pp. 967-972

[ArticleDownload PDFView Record in Scopus](#)

[Bååth, 1989](#)

E. Bååth **Effects of heavy metals in soil microbial processes and populations**
Water Air and Soil Poll., 47 (1989), pp. 335-379
(a review)

[View Record in Scopus](#)

[Bannert et al., 2011](#)

A. Bannert, K. Kleineidam, L. Wissing, C. Mueller-Niggemann, V. Vogelsang, G. Welzl, Z. Cao, M. Schlöter **Changes in diversity and functional gene abundances of microbial communities involved in nitrogen fixation nitrification, and denitrification in a tidal wetland versus paddy soils cultivated for different time periods**
Appl. Environ. Microbiol., 77 (2011), pp. 6109-6116

[CrossRefView Record in Scopus](#)

[Bardgett et al., 2008](#)

R.D. Bardgett, C. Freeman, N.J. Ostle **Microbial contributions to climate change through carbon cycle feedbacks**
ISME J., 2 (2008), pp. 805-814

[CrossRefView Record in Scopus](#)

[Benjamini and Hochberg, 1995](#)

Y. Benjamini, Y. Hochberg **Controlling the false discovery rate – a practical and powerful approach to multiple testing**

J. Roy. Stat. Soc., 57 (1995), pp. 289-300

[View Record in Scopus](#)

[Bhattacharya et al., 1996](#)

K.S. Bhattacharya, M. Qu, L.R. Madura **Effects of nitrobenzene and zinc on acetate utilizing methanogens**

Water Res., 30 (1996), pp. 3099-3100

[CrossRefView Record in Scopus](#)

[Brookes and McGrath, 1984](#)

P.C. Brookes, S.P. McGrath **Effects of metal toxicity on the size of the soil microbial biomass**

J. Soil Sci., 35 (1984), pp. 341-346

[CrossRefView Record in Scopus](#)

[Cambria et al., 1989](#)

S. Cambria, P.S. Boni, J. Strabelli **Estudos preliminares com micronutrientes: zinco**

Boletim Técnico Copersucar, 46 (1989), pp. 12-17

[Campoy et al., 2002](#)

S. Campoy, M. Jara, N. Busquets, A.M.P. de Rozas, I. Badiola, J. Barbé **Role of the high-affinity zinc uptake *znuABC* system in *Salmonella enterica* serovar Typhimurium virulence**

Infect. Immun., 70 (2002), pp. 4721-4725

[CrossRefView Record in Scopus](#)

[Chander et al.,](#)

[1995](#)

K. Chander, P.C. Brooks, S.A. Harding **Microbial biomass dynamics following addition of metal-enriched sewage sludges to a sandy loam**

Soil Biol. Biochem., 27 (1995), pp. 1409-1421

[ArticleDownload PDFView Record in Scopus](#)

[Chandra](#)

[a et al.,](#)

[2007](#)

B.R. Chandra, M. Yogavel, A. Sharma **Structural analysis of ABCfamily periplasmic zinc binding protein provides new insights into mechanism of ligand uptake and release**

J. Mol. Biol., 367 (2007), pp. 970-982

[ArticleDownload PDFView Record in Scopus](#)

M. Chvapil **New aspects in the biological role of zinc: a stabilizer of macromolecules and biological membranes**

Life Sci., 13 (1973), pp. 1041-1049

[ArticleDownload PDFView Record in Scopus](#)

[Clarke,
1993](#)

K.R. Clarke **Non-parametric multivariate analyses of changes in community structure**

Aust. J. Ecol., 18 (1993), pp. 117-143

[CrossRefView Record in Scopus](#)

[Coleman, 1974](#)

J.E. Coleman **The role of Zn (II) in transcription by T7 RNA polymerase**

Biochem. Biophys. Res. Commun., 60 (1974), pp. 641-648

[ArticleDownload PDFView Record in Scopus](#)

[Cong et al., 2015](#)

J. Cong, Y. Yang, X. Liu, H. Lu, X. Liu, J. Zhou, D. Li, H. Yin, J. Ding, Y. Zhang **Analyses of soil microbial community compositions and functional genes reveal potential consequences of natural forest succession**

Sci. Rep., 5 (2015), p. 10007

[Corbett et al., 2003](#)

C.R. Corbett, M.N. Burtnick, C. Kooi, D.E. Woods, P.A. Sokol **An extracellular zinc metalloprotease gene of *Burkholderia cepacia***

Microbiology, 149 (2003), pp. 2263-2271

[CrossRefView Record in Scopus](#)

[Costa et al., 2015](#)

T.R.D. Costa, C. Felisberto-Rodrigues, A. Meir, M.S. Prevost, A. Redzej, M. Trokter, G. Waksman **Secretion systems in Gram-negative bacteria: structural and mechanistic insights**

Nat. Rev. Microbiol., 13 (2015), pp. 343-359

[CrossRefView Record in Scopus](#)

[Costello et al., 2015](#)

L.C. Costello, Y. Liu, R.B. Franklin, M.C. Kennedy **Zinc inhibition of mitochondrial aconitase and its importance in citrate metabolism of prostate epithelial cells**

J. Biol. Chem., 272 (1997), pp. 28875-28881

[CrossRefView Record in Scopus](#)

[Crane et al., 2007](#)

J.K. Crane, T.M. Naeher, I. Shulgina, C. Zhu, E.C. Boedeker **Effect of zinc in enteropathogenic *Escherichia coli* infection**

Infect. Immun., 75 (2007), pp. 5974-5984

[CrossRefView Record in Scopus](#)

[Davies and Sleep, 1979](#)

A.G. Davies, J.A. Sleep **Inhibition of carbon fixation as a function of zinc uptake in natural phytoplankton assemblages**

J. Mar. Biol. Assoc. U. K., 59 (1979), pp. 937-949

[CrossRefView Record in Scopus](#)

[Davis et al., 2009](#)

L.M. Davis, T. Kakuda, V.J. DiRita **A *Campylobacter jejuni* *znuA* orthologue is essential for growth in low-zinc environments and chick colonization**

J. Bacteriol., 191 (2009), pp. 1631-1640

[CrossRefView Record in Scopus](#)

[DeWalle et al., 1979](#)

F.B. DeWalle, E.S.K. Chian, J. Brush **Heavy metal removal with a completely mixed anaerobic filter**

J. Wat. Pollut. Control Fed., 51 (1979), p. 1

[Deghmane et al., 2002](#)

A.E. Deghmane, D. Giorgini, M. Larribe, J.M. Alonso, M.K. Taha **Down-regulation of pili and capsule of *Neisseria meningitidis* upon contact with epithelial cells is mediated by CrgA regulatory protein**

Mol. Microbiol., 43 (2002), pp. 1555-1564

[CrossRefView Record in Scopus](#)

[Demirel and Scherer, 2011](#)

B. Demirel, P. Scherer **Trace element requirements of agricultural biogas digesters during biological conversion of renewable biomass to methane**

Biomass Bioenerg., 35 (2011), pp. 992-998

[ArticleDownload PDFView Record in Scopus](#)

[EMBRAPA, 2011](#)

EMBRAPA **Manual De Métodos De Análise De Solos**

Embrapa Solos, Rio de Janeiro, Brazil (2011), p. 230

Available online: <https://www.embrapa.br/busca-de-publicacoes/-/publicacao/990374/manual-de-metodos-de-analise-de-solo> (Accessed on 14 October 2016)

[Eide, 2006](#)

D.J. Eide **Zinc transporters and the cellular trafficking of zinc**

Biochim. Biophys. Acta, 1763 (2006), pp. 711-722

[ArticleDownload PDFView Record in Scopus](#)

[Fridovich, 1978](#)

I. Fridovich**The biology of oxygen radicals**

Science, 201 (1978), pp. 875-880

[View Record in Scopus](#)

[Hammerschmidt](#)

S. Hammerschmidt, R. Hilse, J.P. van Putten, R. Gerardy-

Schahn, A. Unkmeir, M.Frosch**Modulation of cell surface sialic acid expression in *Neisseria meningitidis* via a transposable genetic element**

EMBO J., 15 (1996), pp. 192-198

[View Record in Scopus](#)

[Hammerschmidt](#)

S. Hammerschmidt, A. Müller, H. Sillmann, M. Mühlenhoff, R. Borrow, A. Fox, J.van

Putten, W.D. Zollinger, R. Gerardy-Schahn, M. Frosch**Capsule phase variation in *Neisseria meningitidis* serogroup B by slipped-strand mispairing in the polysialyltransferase gene (*siaD*): correlation with bacterial invasion and the outbreak of meningococcal disease**

Mol. Microbiol., 20 (1996), pp. 1211-1220

[CrossRefView Record in Scopus](#)

[Hantke, 2001](#)

K. Hantke**Bacterial zinc transporters and regulators**

Biol. Met., 14 (2001), pp. 239-249

[CrossRefView Record in Scopus](#)

[Hantke, 2005](#)

K. Hantke**Bacterial zinc uptake and regulators**

Curr. Opin. Microbiol., 8 (2005), pp. 196-202

[ArticleDownload PDFView Record in Scopus](#)

[Hart et al., 2009](#)

M.E. Hart, M.J. Hart, A.J. Roop**Genotypic and phenotypic assessment of hyaluronidase among type strains of a select group of Staphylococcal species**

Int. J. Microbiol., 7 (2009), p. 1, [10.1155/2009/614371](#)

[CrossRefView Record in Scopus](#)

[Hayes and Theis](#)

T.D. Hayes, T.L. Theis**The distribution of heavy metals in anaerobic digestion**

J. Wat. Pollut. Control Fed., 50 (1978), p. 61

[View Record in Scopus](#)

[He et al., 2007](#)

Z. He, T.J. Gentry, C.W. Schadt, L. Wu, J. Liebich, S.C. Chong, Z. Huang, W. Wu, B. Gu, P.Jardin e, C. Criddle, J. Zhou **GeoChip: a comprehensive microarray for investigating biogeochemical, ecological and environmental processes**

ISME J., 1 (2007), pp. 67-77

[CrossRefView Record in Scopus](#)

[Herbert, 1961](#)

D. Herbert **The chemical composition of microorganisms as a function of their environment**

G.G. Meynell, H. Gooder (Eds.), Microbial Reaction to Environment: Eleventh Symposium of the Society for General Microbiology, Cambridge University Press, Cambridge (1961), pp. 391-416

[View Record in Scopus](#)

[Hughes and Poole](#)

M.N. Hughes, R.K. Poole **Metal speciation and microbial growth-the hard (and soft) facts**

J. Gen. Microbiol., 137 (1991), pp. 725-734

[CrossRefView Record in Scopus](#)

[Kandeler et al., 1996](#)

E. Kandeler, C. Kampichler, O. Horak **Influence of heavy metals on the functional diversity of soil microbial communities**

Biol. Fertil. Soils, 23 (1996), pp. 299-306

[CrossRefView Record in Scopus](#)

[Kelly et al., 1999](#)

J.J. Kelly, M. Häggblom, R.L. Tate III **Changes in soil microbial communities over time resulting from one time application of zinc: a laboratory microcosm study**

Soil Biol. Biochem., 31 (1999), pp. 1455-1465

[ArticleDownload PDFView Record in Scopus](#)

[Kirk and Farrell, 1987](#)

T.K. Kirk, R.L. Farrell **Enzymatic combustion: the microbial degradation of lignin**

Annu. Rev. Microbiol., 41 (1987), pp. 465-505

[View Record in Scopus](#)

[Koh et al., 1996](#)

J.Y. Koh, S.W. Suh, B.J.M. Gwag, Y.Y. He, C.Y. Hsu, D.W. Choi **The role of zinc in selective neuronal death after transient global cerebral ischemia**

Science, 272 (1996), pp. 1013-1016

[View Record in Scopus](#)

[Korndörfer et al., 1995](#)

G.H. Korndörfer, M.S. Beneditini, A.C. Rocha, D.A. Ferreira Neto **Avaliação de três variedades de cana (*Saccharum officinarum*) submetidas a adubação com micronutrientes. STAB Açúcar**

Álcool e Subprodutos, 14 (1995), pp. 23-26

[View Record in Scopus](#)

[Kunito et al., 2001](#)

T. Kunito, K. Saeki, S. Goto, H. Hayashi, H. Oyaizu, S. Matsumoto **Copper and zinc fractions affecting microorganisms in long-term sludge-amended soils**

Bioresour. Technol., 79 (2001), pp. 135-146

[ArticleDownload PDFView Record in Scopus](#)

[Lammel et al., 2015](#)

D.R. Lammel, B.J. Feigl, C.C. Cerri, N. Klaus **Specific microbial gene abundances and soil parameters contribute to C, N, and greenhouse gas process rates after land use change in Southern Amazonian Soils**

Front. Microbiol., 6 (2015), p. 1057

[Langen et al., 2001](#)

G.R. Langen, J.R. Harper, T.J. Silhavy, S.P. Howard **Absence of the outer membrane phospholipase A suppresses the temperature-sensitive phenotype of *Escherichia coli* degP mutants and induces the Cpx and E extracytoplasmic stress responses**

J. Bacteriol., 183 (2001), pp. 5230-5238

[CrossRefView Record in Scopus](#)

[Li et al., 2014](#)

X. Li, J. Rui, J. Xiong, J. Li, Z. He, J. Zhou, A.C. Yannarell, R.I. Mackie **Functional potential of soil microbial communities in the maize rhizosphere**

PLoS One, 9 (2014), p. e112609

[CrossRef](#)

[Liang et al., 2010](#)

Y.T. Liang, Z. He, L.Y. Wu, Y. Deng, G.H. Li, J.-Z. Zhou **Development of a common oligo reference standard (CORS) for microarray data normalization and comparison across different microbial communities**

Appl. Environ. Microbiol., 76 (2010), pp. 1088-1094

[CrossRefView Record in Scopus](#)

[Lindgren et al., 1986](#)

P.B. Lindgren, R.C. Peet, N.J.H. Panopoulos **Gene cluster of *Pseudomonas syringae* pv: phaseolicola controls pathogenicity on bean plants and hypersensitive response on nonhost plants**

J. Bacteriol., 168 (1986), pp. 512-522

[CrossRefView Record in Scopus](#)

[Lock and Janssen, 2005](#)

K. Lock, C.R. Janssen **Influence of soil zinc concentrations on zinc sensitivity and functional diversity of microbial communities**

Environ. Pollut., 136 (2005), pp. 275-281

[ArticleDownload PDFView Record in Scopus](#)

[Marinho and Albuquerque, 1981](#)

M.L. Marinho, G.A.C. Albuquerque **Efeito do cobre e do zinco na produção de cana-de-açúcar em solos de tabuleiros de Alagoas**

Brasil Açucareiro, 98 (1981), pp. 41-50

[View Record in Scopus](#)

[McCoy et al., 2010](#)

M.W. McCoy, M.L. Marré, C.F. Lesser, J. Mecsas **The C-terminal tail of *Yersinia pseudotuberculosis* YopM is critical for interacting with RSK1 and for virulence**

Infect. Immun., 78 (2010), pp. 2584-2598

[CrossRefView Record in Scopus](#)

[McKevitt et al., 1989](#)

A.I. McKevitt, S. Bajaksouzian, J.D. Klinger, D.E. Woods **Purification and characterization of an extracellular protease from *Pseudomonas cepacia***

Infect. Immun., 57 (1989), pp. 771-778

[View Record in Scopus](#)

[Mellis et al., 2016](#)

E.V. Mellis, J.A. Quaggio, G.R.G. Becari, L.A.J. Teixeira, H. Cantarella, F.L.F. Dias **Effect of micronutrients soil supplementation on sugarcane in different production environments**

Cane Plant Cycle. Agron. J., 108 (2016), pp. 2060-2070

[CrossRefView Record in Scopus](#)

[Moffett et al., 2003](#)

B.F. Moffett, F.A. Nicholson, N.C. Uwakwe, B.J. Chambers, J. Harris, T.C.J. Hill **Zinc contamination decreases the bacterial diversity of agricultural soil**

FEMS Microbiol. Ecol., 43 (2003), pp. 13-19

[ArticleDownload PDFCrossRefView Record in Scopus](#)

[Monk et al., 1989](#)

L.S. Monk, K.Y. Fagerstedt, R.M.M. Crawford **Oxygen toxicity and superoxide dismutase as an antioxidant in physiological stress**

Physiol. Plant., 76 (1989), pp. 456-459

[CrossRefView Record in Scopus](#)

[Mudhoo and Kumar, 2013](#)

A. Mudhoo, S. Kumar **Effects of heavy metals as stress factors on anaerobic digestion processes and biogas production from biomass**

Int. J. Environ. Sci. Technol., 10 (2013), pp. 1383-1398

[CrossRefView Record in Scopus](#)

[Otto and Silhavy, 2002](#)

K. Otto, T.J. Silhavy **Surface sensing and adhesion of *Escherichia coli* controlled by the Cpx-signaling pathway**

Proc. Natl. Acad. Sci. U. S. A., 99 (2002), pp. 2287-2292

[CrossRefView Record in Scopus](#)

[Outten and O'Halloran, 2001](#)

C. Outten, T. O'Halloran **Femtomolar sensitivity of metalloregulatory proteins controlling zinc homeostasis**

Science, 292 (2001), pp. 2488-2492

[CrossRefView Record in Scopus](#)

[Pająk et al., 2016](#)

M. Pająk, E. Błońska, M. Frać, K. Oszust **Functional diversity and microbial activity of forest soils that are heavily contaminated by lead and zinc**

Water Air Soil Pollut., 227 (2016), pp. 348-362

[Palmgren et al., 2008](#)

M.G. Palmgren, S. Clemens, L.E. Williams, U. Krämer, S. Borg, J.K. Schjørring, D. Sanders **Zinc biofortification of cereals: problems and solutions**

Trends Plant Sci., 13 (2008), pp. 464-473

[ArticleDownload PDFView Record in Scopus](#)

[Parks and Beiko, 2010](#)

D.H. Parks, R.G. Beiko **Identifying biologically relevant differences between metagenomic communities**

Bioinformatics, 26 (2010), pp. 715-721

[CrossRefView Record in Scopus](#)

[Prask and Plocke, 1971](#)

J.A. Prask, D.J. Plocke **A role for zinc in the structural integrity of the cytoplasmic ribosomes of Euglenagracilis**

Plant Physiol., 48 (1971), pp. 150-155

[CrossRefView Record in Scopus](#)

[R Core Team, 2015](#)

R Core Team **R: A Language and Environment for Statistical Computing**

R Foundation for Statistical Computing, Vienna Austria (2015)

(URL <http://www.R-project.org/>)

[Ramakrishnan et al., 1991](#)

G. Ramakrishnan, J.-L. Zhao, A. Newton **The cell cycle-regulated agellar gene bF of *Caulobacter crescentus* is homologous to a virulence locus (*IcrD*) of *Yersinia pestis***

J. Bacteriol., 173 (1991), pp. 7283-7292

[CrossRefView Record in Scopus](#)

[Richardson and Simpson, 2011](#)

A.E. Richardson, R.J. Simpson **Soil microorganisms mediating phosphorus availability update on microbial phosphorus**

Plant Physiol., 156 (2011), pp. 989-996

[CrossRefView Record in Scopus](#)

[Serafini et al., 2014](#)

A. Serafini, D. Pisu, G. Palù, G.M. Rodriguez, R. Manganelli **The ESX-3 secretion system is necessary for iron and zinc homeostasis in *Mycobacterium tuberculosis***
PLoS One, 8 (10) (2014), p. e78351

[Sheehan et al., 2015](#)

L.M. Sheehan, J.A. Budnick, R.M. Roop II, C.C. Caswella **Coordinated zinc homeostasis is essential for the wild-type virulence of *Brucella abortus***
J. Bacteriol., 197 (2015), pp. 1582-1591

[CrossRefView Record in Scopus](#)

[Sillanpää, 1990](#)

M. Sillanpää **Micronutrient Assessment at the Country Level: An International Study**
FAO Soils Bulletin No. 63, FAO, Rome (1990)

[Singh et al., 2008](#)

P. Singh, A. Suman, P. Tiwari, N. Arya, A. Gaur, A.K. Shrivastava **Biological pretreatment of sugarcane trash for its conversion to fermentable sugars**
World J. Microbiol. Biotechnol., 24 (2008), pp. 667-673

[CrossRefView Record in Scopus](#)

[Snyder et al., 1995](#)

W.B. Snyder, L.J. Davis, P.N. Danese, C.L. Cosma, T.J. Silhavy **Overproduction of NlpE, a new outer membrane lipoprotein, suppresses the toxicity of periplasmic LacZ by activation of the Cpx signal transduction pathway**
J. Bacteriol., 177 (1995), pp. 4216-4223

[CrossRefView Record in Scopus](#)

[Souza et al., 2003](#)

J.V.B. Souza, E.S. Silva, M.L.S. Maia, M.F.S. Teixeira **Screening of fungal strains for pectinolytic activity: endopolygalacturonase production by *Peacilomyces clavisporus* 2A. UMIDA .1**
Process Biochem, 39 (2003), pp. 455-458

[ArticleDownload PDFView Record in Scopus](#)

[Spaepen et al., 2007](#)

A. Spaepen, J. Vanderleyden, R. Remans **Indole-3-acetic acid in microbial and microorganism-plant signaling**
FEMS Microbiol. Rev., 31 (2007), pp. 425-448

[CrossRefView Record in Scopus](#)

[Stone et al., 2006](#)

J.J. Stone, W.R. Burgos, R.A. Royer, B.A. Dempsey **Impact of zinc on biological Fe(III) and nitrate reduction by *Shewanella putrefaciens* CN32**
Environ. Eng. Sci., 23 (2006), pp. 691-704

[CrossRefView Record in Scopus](#)

[Sugarman et al., 1982](#)

B. Sugarman, L.R. Epps, W.A. Stenback **Zinc and bacterial adherence**
Infect. Immun., 37 (1982), pp. 1191-1199

[View Record in Scopus](#)

[Tarafdar et al., 2001](#)

J.C. Tarafdar, R.S. Yadav, S.C. Meena **Comparative efficiency of acid phosphatase originated from plant and fungal sources**

J. Plant Nutr. Soil Sci., 164 (2001), pp. 279-282

[CrossRefView Record in Scopus](#)

[Tosi et al., 2016](#)

M. Tosi, O.S. Correa, M.A. Soria, J.A. Vogrig, O. Sydorenko, M.S. Montecchia **Land-use change affects the functionality of soil microbial communities: a chronosequence approach in the Argentinian Yungas**

Appl. Soil Ecol., 108 (2016), pp. 118-127

[ArticleDownload PDFView Record in Scopus](#)

[U.S. Environmental Protection Agency \(USEPA\), 1998](#)

U.S. Environmental Protection Agency (USEPA) **Method 3051A: Microwave Assisted Acid Digestion of Sediments, Sludges, Soils, and Oils; Test Methods for Evaluating Solid Waste**

Physical/Chemical Methods; Office of Solid Waste, U.S. Environmental Protection

Agency Washington, Washington (1998), pp. 1-20

[View Record in Scopus](#)

[Wang et al., 2005](#)

J.J. Wang, C.W. Kennedy, H.P. Viajator, A.E. Arceneaux, A.J. Guidry **Zinc fertilization of sugarcane in acid and calcareous soils**

J. ASSCT, 25 (2005), pp. 49-61

[View Record in Scopus](#)

[Weinberg, 1990](#)

E.D. Weinberg **Roles of trace metals in transcriptional control of microbial secondary metabolism**

Biol. Metals, 2 (1990), pp. 191-196

[CrossRefView Record in Scopus](#)

[Welch, 1947](#)

B.L. Welch **The generalization of Student's problem when several different population variances are involved**

Biometrika, 34 (1947), pp. 28-35

[CrossRefView Record in Scopus](#)

[Wu et al., 2006](#)

L. Wu, X. Liu, C.W. Schadt, J. Zhou **Microarray-based analysis of subnanogram quantities of microbial community DNAs by using whole-community genome amplification**

Appl. Environ. Microbiol., 72 (2006), pp. 4931-4941

[CrossRefView Record in Scopus](#)

[Ye et al., 2015](#)

N. Ye, X. Zhang, M. Miao **Saccharina genomes provide novel insight into kelp biology**
Nat. Commun., 6 (2015), p. 6986

[Zak et al., 1994](#)

J.C. Zak, M.R. Willig, D.L. Moorhead, H.G. Wildman **Functional diversity of microbial communities: a quantitative approach**
Soil Biol. Biochem., 26 (1994), pp. 1101-1108

[ArticleDownload PDFView Record in Scopus](#)

[Zheng et al., 2014](#)

X. Zheng, Y. Su, Y. Chen, R. Wan, K. Liu, M. Li, D. Yin **2014: Zinc oxide nanoparticles cause inhibition of microbial denitrification by affecting transcriptional regulation and enzyme activity**
Environ. Sci. Technol. 48, 1380 (2014), pp. 0-1

[Zima et al., 1996](#)

T. Zima, S. Stípek, J. Crkovská, K. Němecek, J. Pláteník, V. Bártová, V. Tesar **Antioxidant enzymes-superoxide dismutase and glutathione peroxidase-in haemodialyzed patients**
Blood Purif., 14 (1996), pp. 257-261

[CrossRefView Record in Scopus](#)