

UC Office of the President

Recent Work

Title

Characterizing the cellular architecture of dynamically remodeling vascular tissue using 3-D image analysis and virtual reconstruction

Permalink

<https://escholarship.org/uc/item/2t57042h>

Journal

Molecular Biology of the Cell, 31(16)

Author

Valentine, Megan T

Publication Date

2020-08-20

Peer reviewed

Characterizing the cellular architecture of dynamically remodeling vascular tissue using 3-D image analysis and virtual reconstruction.

Roopa Madhu*¹, Delany Rodriguez*², Claudia Guzik², Shambhavi Singh², Anthony W. De Tomaso², Megan T. Valentine³, and Dinah Loerke¹.

¹Department of Physics and Astronomy

University of Denver, CO 80208

²Department of Molecular, Cellular and Developmental Biology

³Department of Mechanical Engineering

University of California Santa Barbara, CA 93106

*These authors contributed equally to this work.

Correspondence: Dinah Loerke and Megan Valentine

Key words: Epithelial mechanics, Botryllus, morphology, blood vessels, stress fibers, cell shape, image analysis

Abstract

Epithelial tubules form critical structures in lung, kidney and vascular tissues. However, the processes that control their morphogenesis and physiological expansion and contraction are not well understood. Here we examine the dynamic remodeling of epithelial tubes *in vivo* using a novel model system: the extracorporeal vasculature of *Botryllus schlosseri*, in which the disruption of the basement membrane triggers rapid, massive vascular retraction without loss of barrier function. We developed and implemented 3-D image analysis and virtual reconstruction tools to characterize the cellular morphology of the vascular wall in unmanipulated vessels and during retraction. In both control and regressed conditions, cells within the vascular wall were planar polarized, with an integrin- and curvature-dependent axial elongation of cells and a robust circumferential alignment of actin bundles. Surprisingly, we found no measurable differences in morphology between normal and retracting vessels under ECM disruption. However, inhibition of integrin signaling through FAK inhibition caused disruption of cellular actin organization. Our results demonstrate that epithelial tubes can maintain tissue organization even during extreme remodeling events, but that the robust response to mechanical signals – such as the response to loss of vascular tension after ECM disruption - requires functional force sensing machinery via integrin signaling.

Introduction

Epithelial sheets in living organisms are highly dynamic, and must be able to remodel sheet geometry – through addition, removal or migration of cells within the sheet – while simultaneously maintaining barrier function through stable cell-cell adhesion. This balance places high demands on epithelial cells, requiring them to both generate and respond to forces through integration of mechanical and biochemical pathways. Thus, understanding epithelial organization and dynamics provides fundamental insights into biomechanical signaling pathways that are applicable to a wide range of biological processes, including embryonic development, angiogenesis, wound healing, and metastasis.

Some of the most tractable experimental model systems for epithelial dynamics are variations of cultured epithelial cells grown on glass coverslips; these can be imaged with high spatial and temporal resolution, and have the advantage of a well-controlled chemical environment through use of perfusion systems. For example, cultured monolayer models have enabled investigations of how vascular cells respond to various chemical and physical stimuli (Dewey, Bussolari et al. 1981, Galbraith, Skalak et al. 1998, Kaunas, Nguyen et al. 2005, Gu, Forostyan et al. 2011, Kutys and Chen 2016). One major drawback of these assays, however, is that glass or plastic substrates are much stiffer than the natural tissue environment of the cells, and in the case of vascular cells, the planar geometry is not physiological. Thus, it is unclear how relevant the insights gained in these artificial conditions ultimately are *in vivo*. Conversely, high-resolution imaging of epithelial sheets in their natural mechanical environment *in vivo* is not always feasible, since many epithelial and vascular tissues of interest are found deep within organisms – this constraint limits the available imaging approaches, and leads to significant signal extinction and scattering, fundamentally limiting any optical analysis.

Here we address this challenge using a unique model system: the extracorporeal vasculature of the ascidian (sea squirt) *Botryllus schlosseri*. *Botryllus* is an invertebrate chordate closely related to vertebrates (Gasparini, Manni et al. 2015, Holland 2016, Kassmer, Rodriguez et al. 2016), and has a large, transparent extracorporeal vascular network that allows unparalleled accessibility to study vascular biology *in vivo* (Tiozzo, Voskoboynik et al. 2008, Rodriguez, Braden et al. 2017). These blood vessels are large (with an average diameter of $\sim 50\mu\text{m}$) and consist of a single tubular monolayer of myoepithelial cells that create a massive branching vascular network encompassing areas over 10cm^2 . These vessels are embedded in a transparent, extracellular matrix called the tunic, can be directly manipulated, and are uniquely amenable to high-resolution imaging (Figure 1 and Figure S1). At the periphery of the vascular bed, the vessels terminate in structures called ampullae, which are bulbous, blind-ended extensions of the vessels themselves. The vascular cells are contractile, and the *Botryllus* vasculature consists of a single tube, without layers of smooth muscle or pericytes.

Finally, the vessel architecture is inverted in comparison to mammals: the vascular cells are not endothelial, rather the basal side of the vessel epithelium faces the lumen, with the basement membrane forming the inner layer of the vessel.

Under normal conditions, the *Botryllus* vasculature is highly dynamic, with individual vessels undergoing expansion and retraction on the order of millimeters on a daily basis, and during high growth periods the entire vascular bed can double in size every week. Angiogenesis can also be triggered experimentally: the extracorporeal vasculature can be surgically ablated; the vessels will clot, remodel and begin proliferating, resulting in regeneration the entire bed within 72 hours (Tiozzo, Voskoboynik et al. 2008, Braden, Taketa et al. 2014). Finally, we have recently demonstrated that we can also induce regression of the vasculature via direct manipulation of the stiffness of the basement membrane: Specifically, exposing the vasculature to β -aminopropionitrile (BAPN), which blocks collagen crosslinking, and which within a few hours of treatment completely changes the structure of collagen fibers in the basement membrane (Videos 1 and 2). We have shown that the corresponding change of ECM stiffness is sensed through integrin-dependent pathways and causes anoikis of a subset of cells, which are then extruded basally and removed by phagocytic cells in circulation (Rodriguez, Braden et al. 2017). This loss and removal of individual cells induces a rapid macroscopic regression of blood vessels on the order of millimeters in a time period of 16 hours.

Importantly, while this regression constitutes a massive macroscopic morphological change, it produces no pathological damage to the organism - regression progresses without bleeding or plasma leakage, indicating that the vessels maintain barrier function. Regression is also fully reversible: following removal of BAPN, the vessels begin proliferating and regenerate the entire vascular bed. As vessels are dynamic and regress even under normal conditions, we hypothesize that BAPN treatment induces normal physiological responses to changes in the mechanical environment. One interpretation is that the collapse of ECM tonus is interpreted mechanically by the epithelial sheet as cellular overcrowding, to which it responds with a homeostatic cell density reduction (Eisenhoffer, Loftus et al. 2012, Marinari, Mehonic et al. 2012). This suggests that under normal conditions, changes in the structure of the basement membrane and other mechanosensation events precede and trigger vascular regression or branching, as has been shown for epithelial remodeling events in other model systems (Ingber 1988). Finally, a number of small molecule inhibitors of the integrin pathway (e.g., inhibition of focal adhesion kinase (FAK)) can also induce vascular regression in manner similar to the disruption of the basement membrane (Rodriguez, Braden et al. 2017) (Videos 3 and 4). This suggests that cells sense loss of anchorage to the extracellular matrix due to collagen network disruption via an integrin-signaling pathway, and that loss of this tonic signal induces cell death (Ives, Eskin et al. 1986).

The ability to experimentally induce growth and regression of the *Botryllus* vasculature provides a unique model system in which to study morphogenesis and maintenance of epithelial tubes in an *in vivo*

vascular network. In this study, we focus on the induced regression event, which allows us to examine large-scale remodeling of epithelial sheets under physiological conditions. To capitalize fully on the strengths of this system, we aim to develop quantitative approaches to examine tissue morphology and multiscale organization to provide insight into the mechanical and molecular mechanisms that underlie dynamic remodeling of the vasculature at both tissue and cellular resolution. To achieve this, we developed and implemented a procedure for a detailed reconstruction of the cellular architecture and morphology using high-resolution 3-D confocal image stacks of immunostained samples of *Botryllus* vasculature. Using this approach, we examined changes in cellular organization and cytoskeletal structure under conditions of normal growth and chemically induced vascular regression. We found no stark differences in morphology between normal and retracting vessels under ECM disruption; however, inhibition of integrin signaling caused significant disruption of cellular actin organization. Our results provide insight into the organization and properties of cells within a dynamic remodeling tissue, while demonstrating the remarkable ability of organisms to maintain barrier function and structure in extreme conditions.

Results

Three-dimensional reconstruction of the epithelial sheet

Our reconstruction of the cellular architecture and morphology is based on high-resolution 3-D confocal image stacks of immunostained samples of *Botryllus* vasculature (Figure 2A). The surface of the cylindrical epithelial sheet was defined via staining with a cadherin antibody, followed by segmenting the 2-dimensional cross-sectional profile, while stepping this cross-section along the long axis of the blood vessel (the ‘vascular axis’) (Figure 2B), and attaching cylindrical coordinates to each surface point in the process (axial distance s , radial distance r from the center, and angle θ). Using that segmented surface, raw fluorescence intensities in the available color channels were interpolated into 2-dimensional intensity maps using regular-spaced bins of the axial distance s and of the circumferential angle θ . The resulting intensity maps thus conceptually represent the ‘unwrapped’ epithelial sheets in the different color channels (Figure 2C left panel, showing the example of the ‘unwrapped’ map for rhodamine-labeled phalloidin, scale bar 10 μ m; also see Figure S2A-C). We subsequently performed standard 2-dimensional image analysis on these 2-dimensional maps of the unwrapped sheets, using watershed segmentation to segment the cell-cell interfaces and then calculate the position of vertices and cell centroids (Figure 2C right panel). Importantly, we retain the inherently 3-dimensional structure of the sheet by using the surface points’ radial distance information from the original surface segmentation, such that every point on the 2-dimensional intensity map can be mapped back to its true position on the 3-dimensional curved surface. We used this approach to create a triangulated reconstruction of the complete vascular architecture, using the 3-dimensional positions of the cell vertices and cell centroids that were identified in the 2-dimensional

intensity map (Figure 2D; see Materials and Methods for additional details).

Actin bundles are preferentially oriented into circumferential direction

We first used this quantitative analysis approach to characterize the organization and properties of cells within the normal *Botryllus* vasculature, and then to compare the wild-type features to those of cells in regressing vessels in animals treated with BAPN to disrupt ECM organization, or the FAK inhibitor (FAKI) to disrupt integrin signaling. Figure S3 shows partial maximum projections of the blood vessels in homeostasis in wild-type, BAPN-treated and FAKI-treated animals. In all cases, fixation of BAPN-treated animals was performed 8 hours after the drug was applied, and 6 hours after FAKI application, which for both drugs constitutes the time point where vessels are regressing at maximum velocity (Rodriguez, Braden et al. 2017). Our sample size of analyzed images consisted of $n=30$ blood vessels in control conditions (from 10 unique animals), with a range of 26-219 cells per blood vessel for a total of $k=2,990$ cells; in BAPN-treated conditions, $n=20$ blood vessels (from 9 unique animals) with a range of 28-185 cells per blood vessel for a total of $k=1,660$ cells; in FAKI treated conditions, $n=13$ blood vessels (from 5 unique animals) with a range of 23-257 cells per blood vessel for a total of $k=1,391$ cells. This same data sample was used for all the analyses shown in Figures 3-5.

We found that most blood vessels in our sample ranged in diameter from approximately 25 to 55 μm (Figure 3A), where vessel diameter was calculated as the average diameter along each vessel's axis. The cell area (weighted mean \pm weighted SD) was $211 \pm 40 \mu\text{m}^2$ in control, $227 \pm 35 \mu\text{m}^2$ in BAPN-treated animals (difference from control not significant, 2-sample t -test), and $190 \pm 26 \mu\text{m}^2$ in FAKI-treated animals (difference from control not significant, difference from BAPN significant at $p=0.005$, 2-sample t -test) (Figure 3B). In all three cases, the tissue had a primarily hexagonal morphology (Figure 2C, Figure 3C-D). Thus, somewhat surprisingly, for these specific metrics (cell area and number of nearest neighbors) we found no drastic differences in basic tissue morphology, despite the presence of significant macroscopic tissue remodeling of blood vessels (Videos 1 and 3).

To test whether the cells responded to the anisotropies in their mechanical environment – since they experience lowest curvature within the tubular sheet into axial direction, and highest curvature into circumferential direction - we next probed for morphological anisotropies within the plane of the sheet.

The myoepithelial cells that form the lining of *Botryllus* blood vessels contain actin-rich bundles that are brightly labeled with the actin-binding dye phalloidin, and resemble the crossed-linked, contractile actin bundles called stress fibers in mammalian cells (Chazotte 2010, Rodriguez, Braden et al. 2017). We examined the preferential orientation of these actin bundles in the 2D intensity maps of phalloidin intensity by performing line

filtering with a variable filter orientation (*i.e.* variable spatial angle ϕ within the sheet) (see Material & Methods, Figure S4A-H). The filter response – which we calculate as the mean of the absolute image intensity after line filtering – is maximized when the orientation of the line filter element is aligned with the predominant orientation of any fiber-like features in the image. We observed that the line-filtered intensity was significantly increased for a filter orientation into the circumferential direction (representative sample plots in Figure 4A), indicating that the phalloidin-stained actin bundles have a predominantly circumferential orientation; this directional preference was extremely robust across all vessels and conditions (Figure S4I-L). This quantitative finding is consistent with visual inspection of the phalloidin-stained images, which often show a distinct visible horizontal orientation of fiber-like structures (Figure 4B, Figure S4C). Based on these results, we calculated directional preference mathematically as the contrast c between the maximum and minimum filter responses (*i.e.* the filtered intensities into those directions that either maximize or minimize signal), $c=(I_{max}-I_{min})/(I_{max}+I_{min})$. This directional preference is conceptually similar to the eccentricity of the polar plot in Figure 4A; thus, this metric designates relative directional preference without regard to absolute filter response magnitude. Numerically, this contrast metric will have a value of 0.2 if the maximum is 50% larger than the minimum, and ~ 0.1 if the maximum is 25% larger than the minimum. This directional preference value was not normally distributed (Figure 4C), with median (IQR = interquartile range) being 0.12 (IQR=0.10) in control, 0.12 (IQR=0.14) for BAPN, and 0.14 (IQR=0.12) for FAKI-inhibited animals (with the differences between control vs. BAPN and control vs. FAKI not statistically significant, two-sided Kolmogorov Smirnov-test). This suggests that the directional preference of the actin bundle architecture is mostly conserved in BAPN and in FAKI.

In addition, we examined the absolute magnitude of the maximum line filter response, which reflects the degree to which the image structures are fiber-like in appearance (as opposed to punctate, or homogeneous). This maximum line filter intensity was also not normally distributed (Figure 4D), with the median being 0.21 (IQR=0.09) in control, 0.20 (IQR=0.05) for BAPN, and 0.14 (IQR=0.05) for FAKI animals. In this case, the difference between control vs. BAPN-treatment was not statistically significant, but the difference between control vs. FAKI-treated animals was statistically significant at $p=0.016$ (both two-sided Kolmogorov Smirnov-test). This indicates that while the FAKI-treated animals do preserve the overall directional preference for circumferential orientation in the existing fibers, there is a substantial disruption of actin bundle organization overall, reflected in a reduction of the density and contrast of fiber-like features in the image (Figure S4I-K). This quantitative result is consistent with the visual impression of actin fiber organization being disrupted or even entirely absent in some FAKI-treated animals (Figure S4C).

Vascular epithelial cells are net axially elongated in homeostasis and induced regression

In examining the cells' major and minor axis lengths and orientations, we found that the major and minor axis lengths were (weighted mean \pm weighted SD) $19.7 \pm 1.7 \mu\text{m}$ vs $14.0 \pm 0.5 \mu\text{m}$ for control, $20.8 \pm 1.7 \mu\text{m}$ vs $14.4 \pm 1.3 \mu\text{m}$ for BAPN-treatment, and $19.2 \pm 1.4 \mu\text{m}$ vs $13.1 \pm 0.9 \mu\text{m}$ for FAKI-treatment (Figure 4E). The resulting aspect ratio was (weighted mean \pm weighted SD) 1.41 ± 0.08 for control, 1.44 ± 0.10 for BAPN (difference from control not significant, 2-sample *t*-test), and 1.46 ± 0.04 for FAKI (difference from control significant at $p=0.02$, 2-sample *t*-test).

Importantly, the orientation of the cells' major axis significantly favored the axial direction over the circumferential direction (Figure 4F) by roughly a factor of two (calculated by dividing axial frequency by circumferential frequency). This axial preference was increased relative to control in BAPN-treated animals, and decreased relative to control in FAKI animals. Specifically, the fractions of cells oriented into a 30° axial angle window vs. a 30° circumferential angle window were (weighted mean \pm weighted SD) $25.7 \pm 11.8 \%$ vs. $14.0 \pm 7.3 \%$ in control conditions (significance of axial-circumferential difference $p=7.65\text{e-}05$, 2-sample *t*-test), $31.8 \pm 18.7 \%$ vs. $12.9 \pm 10.6 \%$ in BAPN-treated animals (significance of axial-circumferential difference $p=9.0\text{e-}04$, 2-sample *t*-test), and $22.3 \pm 13.5\%$ vs. $13.2 \pm 6.6 \%$ in FAKI-treated animals (significance of axial-circumferential difference $p=0.049$, 2-sample *t*-test), with the differences between conditions being not statistically significant. Thus, the vascular morphology is planar polarized (anisotropic within the plane of the sheet) in all measured conditions, with a net preference of elongation of cells into the axial direction, and the strength of this preference varies slightly between conditions. In addition, we also found an axial preference in the orientation of cell-cell interfaces (Figure S5A), which is consistent with a 'ladder-like' morphology of the tissue, with almost rectangular cells in many cases (Figures S5B, C).

Planar polarity depends on vascular curvature

We have now established that the epithelial architecture within the blood vessels is typically planar polarized, where actin bundles within cells are net preferentially aligned into the circumferential direction, while the cells themselves are net preferentially elongated into the axial direction. To gain further insight into the relationship of these observations to curvature, we asked the following: (1) Does this observed polarity depend on epithelial sheet curvature, *i.e.* does polarity increase in highly curved vessels? (2) Does a cell's orientation depend on its relative elongation?

To answer these questions, we performed further directionally resolved analyses on cell properties. First, to address whether the degree of cellular polarization depends of vascular curvature, we measured the

distribution of cells' orientation angles as a function of vascular diameter. We designated the angular range of $90\pm 15^\circ$ as representing cells with 'axial' orientation, and the range of $0\pm 15^\circ$ as 'circumferential' orientation (Figure 5A and Figure S5D). Thus, the total range for axial and circumferential categories was 30° each; this means that for a perfectly isotropic case - in which cells have no directional preference whatsoever - we would expect to see $(30^\circ/180^\circ)=16.7\%$ of cells to fall into either the 'axial' or the 'circumferential' categories.

For blood vessels with diameters of $>40\mu\text{m}$, we observed that the proportions of 'axial' and 'circumferential' cell orientations were close to the reference values for an isotropic distribution in all conditions (Figure 5B,C); while all conditions had a net systematic axial directional preference, the differences were not statistically significant (see Table 1). However, as the vascular diameter decreased below $40\mu\text{m}$, both in control conditions and in BAPN-treated animals, cells showed a growing preference for axial orientation at the expense of the 'circumferential' one, reaching a fraction of $>50\%$ axially oriented cells in some individual blood vessels (Figure 5B,C), which constitutes a three-fold increase over an isotropic distribution; these differences were significant at $p < 10^{-3}$ (see Table 1). In FAKI cells, this increase of the axial preference for small vessel diameters was absent, and the observed small axial preference was similar to the baseline of large vessel diameters (Figure 5C).

Interestingly, in all observed blood vessels – both in control and BAPN-treated animals, and even in the FAKI animals that experience a strong loss of overall fiber content and organization - the existing actin fibers retained their preference for circumferential orientation (Figure 5D), regardless of whether cell orientation was isotropic or axially polarized, and across a wide range of vascular diameters. Thus, we conclude that the observation of planar polarity has two independent components: (1) the preferentially circumferential orientation of actin bundles, which is extremely robust; and (2) the preferentially axial orientation of cell axes, which is fairly weak for large vascular diameters (*i.e.* $>40\mu\text{m}$), but which becomes more pronounced for vascular diameters below $40\mu\text{m}$ in a FAK-dependent manner.

In addition, we observed visually that in some blood vessels with a strong axial orientation preference, the cells also appeared to be more elongated overall (e.g. Figure S5B). We speculated that this correlation might arise from a tendency of cells to change shape in order to reduce their net circumferential 'exposure' in the blood vessel. If this were the case, we would expect that, within the population of cells from a given blood vessel, the more elongated cells should have a stronger preference for being axially oriented than the round cells. We tested this hypothesis in those blood vessels that showed a polarized phenotype - defined as those in which the fraction of axially oriented cells exceeded 30%. For each blood vessel of this phenotype, we divided the cell population into quintiles according to their elongation (as measured by cells' major-to-minor axis aspect

ratio), and determined the relative probability density as a function of orientation angle in bins of 30° (Figure 5E). We found that cells indeed show a progressive axial orientation preference with increasing elongation: there was almost no observable preference for axial orientation in the least elongated quintile of cells, but the preference increased with increasing aspect ratio and reached its maximum in the most elongated quintile, for all conditions. Specifically, in control cells, the probability for axial orientation (mean \pm SD) in the least elongated group of cells was only $24.0 \pm 4.4\%$ (which is only a mild increase over the isotropic probability of 16.7%), compared to a $57.9 \pm 4.7\%$ probability for axial orientation among the most elongated group of cells (two-sided Kolmogorov-Smirnov test of the probabilities in these two quintiles against each other $p < 10^{-3}$). This result is consistent with the notion that within this polarized phenotype, cell morphology minimizes the cells' circumferential width. This effect - where the cells' probability of axial orientation increased from the least to the most elongated quintile - was also present for BAPN treatment, where the probability (mean \pm SD) increased from $25.5 \pm 11.3\%$ to $68.7 \pm 13.4\%$ ($p < 10^{-5}$), but somewhat less pronounced in FAKI, where the probability increased from $24.6 \pm 13.5\%$ to $53.2 \pm 25.2\%$ ($p = 0.08$). In contrast, in blood vessels of the isotropic phenotype (where the fraction of axially oriented cells was $< 30\%$), we found no strong relationship between cell elongation and orientation, suggesting that these cells show no strong systematic avoidance of circumferential exposure (Figure S5E).

Discussion

In this paper, we perform quantitative and high-resolution reconstruction of the epithelial architecture of the *Botryllus* vasculature by 'unwrapping' the 3D vascular tube into a 2-dimensional epithelial sheet; our method complements a recent demonstration of 'unrolled' tubular epithelia to enable analysis of developmental and other tissue systems (Heemskerk and Streichan 2015, Yang, Li et al. 2019). We observed that cells within the vascular wall were planar polarized, with robust preference for actin stress fiber orientation into the circumferential direction, and a preference for cell elongation into the axial direction that appears to minimize cells' circumferential dimension, which is both FAK-dependent and curvature-dependent.

A key motivator for this study was to understand the dynamic and dramatic tissue remodeling that could be triggered by collagen disruption in the *Botryllus* vasculature. Yet, strikingly, none of our measured morphological features (number of nearest neighbors, cell size and elongation) were significantly disrupted in BAPN-treatment relative to control conditions, even though we know that the vasculature is undergoing rapid cellular remodeling and macroscopic regression at this timepoint. There could be several possible explanations for this phenomenon: On the one hand, it is theoretically possible that not all regions of the vasculature are equally dynamic. If the bulk of BAPN-induced tissue remodeling were to be localized to regions outside of those

'linear' stretches of the vasculature that we typically select for imaging (e.g. instead to vascular junctions, or ampullae, or vessels close to the bodies), then our measurements could represent a population of less dynamic cells. However, we consider this scenario to be unlikely, since live-imaging data (Rodriguez, Braden et al. 2017) shows that apoptosis (followed by cell extrusion) is distributed fairly randomly across the entire vasculature, and no specialized extrusion zones have been observed thus far. Thus, we propose that apoptosis/extrusion most likely does occur in these linear stretches, and that our findings – i.e. the lack of obvious morphological defects - suggest that there must be a nearly immediate re-sealing of the epithelium after an individual cell is removed. We believe that these extrusion and recovery events happen so quickly that the fixed specimens examined here do not capture a sufficient number of active extrusion events to cause measurable defects in the morphology. Such fast re-sealing would have tremendous benefit for the organism since the vascular epithelium would be capable of maintaining homeostasis and barrier function even in the presence of large-scale dynamic rearrangement. This is consistent with the observation that the animals do not bleed during BAPN-induced regression and are capable of full regeneration after the drug is removed, and is in line with the idea that the dramatic regression in BAPN falls on a continuum of homeostatic (as opposed to catastrophic) responses to mechanical signaling. We hypothesize that the extrusion and resealing events that define the dynamic remodeling contribute to the considerable spread of the measured nearest neighbor number distribution (Figure 3C), which indicates some degree of topological disorder. While the precise topology of epithelial tissue of course depends on many factors – including tension and cell-cell adhesion – and is difficult to compare between different model organisms, we have observed similar features in the early germ band epithelium in *Drosophila melanogaster*. In particular, there is a significant increase in the number of cells with 5 or 7 nearest neighbors and corresponding decrease in the number of cells with 6 nearest neighbors after the onset of germ band extension (Figure S6), in which the tissue begins to remodel and transitions from a static to a highly dynamic state.

In addition, we found a system of planar polarity within the epithelial sheet, which was robust not just against regression but also partially against FAKI; one component of this polarity is cell orientation, the other is actin fiber orientation. With respect to cell orientation, larger-diameter blood vessels with less wall curvature (curvature radius $>40\mu\text{m}$) are generally more isotropic in terms of cell shape and orientation; while there is still a small degree of axial polarity, in our sample sizes it is not statistically significant. Strikingly, in smaller-diameter blood vessels with more wall curvature (curvature radius $<40\mu\text{m}$) this axially oriented phenotype becomes more pronounced in both control and BAPN-treated tissues, with axial orientation of cells preferred over circumferential orientation at a ratio of up to 3:1. Since within a given cell population, the preference for axial orientation increases with cell aspect ratio, we interpret this strongly polarized phenotype as an attempt to minimize the individual cell's circumferential 'exposure', by sensing cell bending or other force asymmetries. The

features of this phenotype, however, are weakened or absent in FAKI-treated animals, suggesting that the mechanical sensing directing the morphological response requires integrin signaling. We note that our approach to the quantitative measurement of cell morphometrics in fact likely underestimates the disruption under FAKI, since our watershed-based segmentation method requires somewhat regular cell boundaries in order to produce meaningful results, and we occasionally have to exclude an image stack when the fluorescence signal of the cell boundaries is too irregular (e.g. due to incomplete or irregular staining, see Figure S7). We observed that the fraction of excluded image stacks was significantly higher under FAK disruption conditions; in addition to a stronger degradation of intensities, we also observed the appearance of ‘rippling’ in the shape of cell-cell boundaries, which suggests loss of tension across the cell-cell junctions. While our current analysis tools do not yet fully capture these differences, they will be a focus of future work.

Previous studies of *in vitro* cultures of endothelial cells, which have an isotropic orientation at rest, have shown that cells become elongated and reorient in the direction of the fluid force when shear stress is applied (Dewey, Bussolari et al. 1981, Franke, Grafe et al. 1984, Wechezak, Viggers et al. 1985, Ives, Eskin et al. 1986, Iba and Sumpio 1991). *In vivo*, local hemodynamic forces dictate the shape of endothelial cells; the greater the force, the flatter and more elongated the cells become, and the more they align with the direction of blood flow (Flaherty, Pierce et al. 1972, Gotlieb, May et al. 1981, White and Fujiwara 1986). Thus, the preferential elongation and orientation of cells into the axial direction that we observed in *Botryllus* could be related to the mechanical shear effects of blood flow (Hellbach, Tiozzo et al. 2011). Unlike mammalian models, in *Botryllus*, individuals have a tubular heart that beats bidirectionally, where the direction of blood flow reverts every 2-3 minutes; thus, the shear force vectors should almost always point axially, along the vessel length, while the magnitude of the shear forces likely depends on the wall curvature of the vessel, producing stronger effects in smaller blood vessels.

In the majority of prior *in vitro* studies, however, the observed cytoskeletal structures (e.g. actin-rich stress fibers) were also aligned along the elongated cell axis, into the direction of fluid flow, which has been used as evidence to suggest that the stress fibers play a role in protection against fluid shear stress (Wong, Pollard et al. 1983, Sato and Ohashi 2005). This is a notable difference to our study – which is the first to measure actin fiber orientation in intact vasculature in *Botryllus*. We found a strong and robust preference for circumferential alignment of actin bundles, which are thus oriented perpendicular to the direction of fluid flow, and perpendicular to the direction of net cell elongation, both in control and BAPN-treated animals and regardless of cell size and shape. Not surprisingly, this polarized architecture was also integrin-dependent, as it was massively disrupted in FAKI-treated animals. Circumferential orientation of both actin fibers and cells was previously noted in the smooth muscle cells of rat blood vessel vein grafts (Liu 1998) , and was linked to the

presence of significant circumferential stress (*i.e.* 'hoop' stress) within the vessel wall arising from the applied arterial blood pressure. It is possible that a similar circumferential stress develops within the *Botryllus* vessel wall, but that the substantially different anatomy of the invertebrate system allows the decoupling of the actin orientation (which may be more sensitive to vessel contraction and blood pressure) and the cell elongation axis (which may be more sensitive to fluid flow).

Cell and actin orientation on curved surfaces has also been investigated in the past using *in vitro* model systems in the absence of fluid flow using microfabricated tubes or rods made from a variety of substrates including polydimethylsiloxane (PDMS), hydrogels and collagen (Shin, Matsuda et al. 2004, Fidkowski, Kaazempur-Mofrad et al. 2005, Chrobak, Potter et al. 2006, Fernandez, Bourget et al. 2007, Song, Cavnar et al. 2009). Cells on such rigid cylindrical substrates have shown a wide range of responses. On rod diameters ranging from 10-500 μm (a range that spans the diameters of capillaries to large blood vessels in the brain), Human Brain Microvasculature Endothelial Cells (HBMEC) presented circumferentially-oriented stress fibers, whereas Human Umbilical Vein Endothelial Cells (HUVEC) presented axially-aligned stress fibers (Ye, Sanchez et al. 2014). Epithelial cells (both Madin Darby Canine Kidney (MDCK) and retinal pigmented epithelial (RPE1) cells) showed strong circumferential alignment of both the cells and actin stress fibers on highly curved rods (radius < 40 μm), whereas 3T3 fibroblasts tended to align axially (Yevick, Duclos et al. 2015). In a separate study, it was shown that both mouse embryonic fibroblasts and human vascular smooth muscle cells aligned axially on highly curved rods (radius of \sim 40 μm), and that their actin stress fibers were polarized, with apical fibers aligned axially, and basal fibers aligned circumferentially (Bade, Kamien et al. 2017). Activation of Rho enhanced the circumferential alignment, particularly in confluent monolayers.

The results of these studies, in which cells are adhered to a rigid, solid cylinder, are difficult to compare to our results due to their relatively artificial environment, and our limited understanding of the role of geometry and boundary conditions on cellular structure, organization and remodeling dynamics. However, these prior *in vitro* results have generally been interpreted by considering an energy balance between stress fiber bending, which would bias toward axial orientation, and contraction, which would favor a circumferential orientation (Biton and Safran 2009). It is not clear that fiber bending would be penalized within a naturally curved surface, as found within the vessel wall, or in the absence of strong cell-substrate adhesions. Furthermore, a recent study of cells attached to regions of positive and negative Gaussian curvature demonstrated that at the unbounded apical surface, cells could minimize stress fiber bending by lifting away from the substrate and forming a straight chord between two distant focal adhesion sites within the three-dimensional volume (Bade, Xu et al. 2018). This suggests that when able to occupy a three-dimensional volume, cells can adapt their overall shape and cytoskeletal organization to minimize fiber bending. It is possible that

similar mechanisms are at play in *Botryllus* blood vessels, which are naturally suspended in the tunic. Finally, it is not established that the phalloidin-stained actin bundles that we observed in the *Botryllus* vasculature are identical in form and function to contractile actin stress fibers (or actomyosin bundles) in mammalian cells. If their orientation is not motivated by minimizing or resisting curvature, then they may fulfill an entirely different mechanical function, *e.g.* to resist variable hoop stress caused by changes in blood pressure during pulsatile beating.

Indeed, there is evidence that the actin-rich bundles we observe in *Botryllus* are not typical stress fibers. In mammalian cells, stress fibers are composed of complexes of actin filaments and the motor protein myosin II; these fibers are usually thick and very stable (Livne and Geiger 2016), and typically are anchored to focal adhesions that in turn connect to the extracellular matrix (ECM) (Tojkander, Gateva et al. 2012). In *Botryllus* we found no significant disruption or loss of planar polarity of actin bundles between control and BAPN-treated animals, despite the fact that BAPN treatment leads to collagen disruption and disassembly on the time scales of hours (Rodriguez, Braden et al. 2017). In fact, this massive disruption to the ECM is what we believe triggers the vessel retraction in the first place, via an integrin-dependent cell death pathway (*i.e.* anoikis). Such systemic disruption of collagen and the ECM would be expected to disrupt stress fiber formation as well, but we see no disruption of the actin bundles under BAPN treatment. It is possible that the focal adhesion complex in *Botryllus* has distinct mechanisms for regulating actin binding and ECM anchorage. Alternatively, it is also possible that in addition to ECM adhesion, the *Botryllus* vascular epithelium is mechanically supported by the external cellulose-rich tunic, and that this support provides additional pathways for cytoskeletal regulation, and perhaps even promotes the rapid resealing of the sheet. The latter case raises interesting possibilities for mechanically-supported vascular grafts using biocompatible hydrogel materials to create an artificial scaffold that stimulates cell dynamics and promotes organization into stable tubular structures.

Materials and Methods

***Botryllus* mariculture and induced vascular regression:**

Botryllus schlosseri colonies were retrieved from the harbor in Santa Barbara, California, and allowed to spawn on 3 x 2 in. glass slides. The progeny were subsequently grown in a mariculture system according to established protocols (Boyd, Brown et al. 1986), with circulating 0.5 mm-filtered seawater (FSW) at 18–21 °C and daily feedings with live algae for a period of 1-2 months, until they reached 5-8 zooids in size, as described (Rodriguez, Braden et al. 2017). Pharmacological inhibition of LOX activity with the specific small molecule inhibitor 3-aminopropionitrile fumarate (BAPN) disrupts the basement membrane, which in turn induces vascular regression of the entire extracorporeal vasculature within 16 hours (Rodriguez, Braden et al. 2017)

(Figure S1A and B). Induced vascular regression was obtained by incubating colonies of a single system 4-5 zooids 2-3 months old, in 400 μ M BAPN for 8 hours (full regression is obtained in 16 hours) or in 50 μ M FAK Inhibitor 14 (FAKI) 6 hours (Figure S1C and D) (full regression is obtained in 10 hours); colonies were then fixed and processed for immunostaining. Note that both BAPN and FAKI were resuspended in molecular biology grade water and therefore this water is used as a carrier control. Control colonies were treated with molecular biology grade water using the exact same volume used in BAPN (512.8 μ l in 500 ml of FSW) and they were incubated for 8 hours; colonies were then fixed and processed for immunostaining.

Live imaging:

Live imaging of colonies treated with either BAPN or FAK inhibitor was carried out using a motorized fluorescence stereomicroscope MZ16FA (Leica, Germany) as previously described (Braden, Taketa et al. 2014, Rodriguez, Braden et al. 2017). Live imaging of fluorescently labeled vasculature of *Botryllus* colonies labeled with BSA-Alexa 594 (A13101; Thermo Fisher Scientific, Waltham, MA), 2 μ l of the dye was microinjected directly into the vasculature of each colony and incubated for 24 hours before imaging using a motorized fluorescence stereomicroscope M205FCA (Leica, Germany).

Staining:

For each round of staining, colonies were transferred to glass coverslip-bottom dishes (0.17mm Delta TPG Dishes) (Bioptechs; 0420041500C) to enable high-resolution imaging while preserving the structure of the blood vessels throughout the staining process. After plating, the colonies were left undisturbed for 1 week to allow them to strongly adhere to the glass portion of the dish. 20 colonies were incubated in 400 μ M 3-aminopropionitrile fumarate (MP Biomedicals, Santa Ana, CA; 150105) (BAPN) in seawater for 8 hours, 23 colonies were incubated in 50 μ M FAK Inhibitor 14 (Santa Cruz Biotechnology, sc-203950) in seawater for 6 hours, and 25 colonies remained in seawater for control conditions. For the whole-mount immunofluorescence staining, the animals were first anesthetized using 790 μ M Trichane (TCI, Tokyo; T0941) in seawater for 10 minutes, then fixed for 3 hours using 4% paraformaldehyde (Fisher Scientific; 50-00-0) in 0.1 M MOPS buffer and 0.5 M NaCl. After three 10-minute washes in phosphate-buffered saline with 0.05% Tween (Fisher Scientific; BP337) (PBST), the specimens were blocked β 33342 (Invitrogen; H3570) (DAPI) and 1:100 rhodamine-phalloidin (Invitrogen, Oregon; R415) (Phalloidin) in PBST overnight and then washed with three 10-minute washes of PBST. We observed no obvious macroscopic differences in morphologies between live and fixed and stained cells.

Imaging:

Imaging of the vasculature of *Botryllus schlosseri* triple labeled with anti Pan-Cadherin, Hoechst 33342, and rhodamine-phalloidin was obtained using an SP8-TCS resonant scanning confocal microscope (Leica, Germany). 3-dimensional (3D) images were collected as z-stacks using a 40× water immersion objective with motorized correction collar and a numerical aperture (N.A.) of 1.1, and a digital 0.75× zoom. Laser excitation wavelength ranges were 350 - 461 nm, 496 - 519 nm, and 540 - 565 nm for visualization of Hoechst-labeled cell nuclei, immuno-labeled cadherin, and phalloidin-labeled actin, respectively. Each x-y image within the z-stack had size of 6704 x 2302 pixels, and all images were taken at a speed of 600 Hz. Of the images taken, 16 images of the β -aminopropionitrile fumarate (BAPN) treated animals, and 23 images of the control animals were selected for analysis due to their high signal-to-noise ratio.

Image analysis:

All the image and data analysis was performed using MATLAB (Mathworks); our custom code written for the image analysis procedures described in sections 1-4 below was also written in MATLAB, and will be available from the Loerke lab on request. Figures were edited using Adobe Illustrator and Adobe Photoshop.

(1) Mapping the surface of the epithelial sheet

From projections of the raw confocal z-stacks (with separate color channels for Hoechst dye, immunolabeled cadherin, and phalloidin-labeled actin), we selected 'straight' regions of the blood vessels, defined as stretches without branching points or overlap with other vessels (Figure 2A). First, raw 3D stacks (x-y-z pixel sizes 75 x 75 x 600 nm) of each channel (Figure 2A) were resized through interpolation to create image voxels of equal size into x-y-z direction. Subsequently, the vascular surface was identified through a watershed transformation on the individual axial sections of the vessel (*i.e.* sections into y-z dimension) (Figure 2B), using the images collected in the phalloidin/actin channel. The seeds for the watershed segmentation were propagated to subsequent axial sections after manually initializing them for the first axial section. After watershed segmentation, the resultant single pixel width segmentation lines were smoothed using a Savitzky-Golay filter with a polynomial of order 3 and a frame length of 23 μm . For every axial cross-section, each point on the segmentation line of the vascular surface is assigned to an angle θ and a distance r with respect to the center-of-mass of the cross-sectional profile (Figure 2B); these values are retained for subsequent analysis.

(2) Construction of 'unwrapped' 2-dimensional intensity maps

After this segmentation step, each pixel on the vascular surface was uniquely identifiable by its circumferential angle θ (and distance r) with respect to the center of the cross-sectional profile, as well as its

distance s along the length of the vascular axis. Using these parameters, the intensities (in each color channel) on the vascular surface were interpolated as 2-dimensional maps as a function of θ and s (Figure 2C, Figure S2A-C). These 2D intensity maps correspond to the mantle of the cylinder, *i.e.* what the epithelial sheets would look like if they were cut along the length of the cylinder, and then ‘unwrapped’ and flattened from their cylindrical shape. The bin size of the circumferential angle for the interpolation was chosen as $\Delta\theta=360^\circ/(\text{mean circumference of the blood vessel})$ so that the pixels in the resulting 2D intensity maps have approximately the same size in the (circumferential) x -direction as in the (axial) y -direction, with some size compressions and rarefactions into the circumferential direction depending on variations in radius (see further below). The resultant 2D intensity map (Figure 2C) was then segmented using a standard 2D seeded watershed segmentation algorithm. The initial segmentation seeds were extracted from cell nucleus positions in the 2D map of the DAPI channel (Figure S2A). Post-processing of the watershed segmentation (Vanderleest, Smits et al. 2018) yielded cell and vertex positions, as well as cell/vertex/interface connectivity.

(3) 3-dimensional (3D) reconstruction

While this unwrapping technique of the tubular sheet is a powerful means of data reduction for 2-dimensional (as opposed to 3-dimensional) data analysis, it creates some challenges for quantitative analysis. The pixels in the unwrapped intensity maps (Figure 2C) have the physical dimension of microns into the axial direction (representing the axial distance s), but the physical dimension of radians into the circumferential direction (representing the circumferential angle θ). As a result, quantitative distance measurements within the sheet must first dimensionally convert the circumferential distances from angles into the lengths of the corresponding circumferential arc segments. In perfectly cylindrical tubes – with constant tube radius - the circumferential arc distance and the radial angle are conveniently related by a constant linear factor, but in practice, the vasculature shows variations of the radial distance both in axial direction (as vessels gradually widen along their length) and in circumferential direction (as some vessels have slightly elliptical cross-sectional profiles).

We retained these radial variations for all experimental samples in corresponding maps of the true circumferential arc length (Figure S2E), and used them for pixel-by-pixel length correction for all 2-dimensional morphological measurements. Thus, the position of any feature that is detected in the 2D intensity map – *e.g.* the vertex between 3 specific cells – can be directly mapped back to a specific location on the 3D surface of the blood vessel, since any location on the 2D map is associated with axial length and the circumferential angle information; in addition, the original segmentation of the vascular surface retains the information where the center point of each axial cross-section was located, and at what distance from the center point the surface is located at any angular bin. In this way, we can reconstruct the ‘true’ 3D locations of all cell centroids and cell

vertices from the 2D segmentation results, and visualize the meshwork of connected points with standard 3D plotting tools (in our case using the *trisurf* function in MATLAB) (Figure 2D). Triangulation is performed on each cell by connecting the centroid of the cell with two of its connected vertices at a time.

To test the effectiveness of this strategy, we compared cell areas calculated using this 2-dimensional approach and a ‘true’ 3-dimensional approach (in which we directly measured cell area in the 3-dimensional triangulated surface), intentionally choosing as a test sample a blood vessel with some deviation from a perfect cylindrical shape (Figure S2D,E). While the values produced using 2-dimensional morphometrics with arc length correction (Figure S2D right panel) aligned well with ‘true’ 3-dimensional measurements, the 2-dimensional morphometrics without arc length correction (Figure S2D left panel) introduced a significant level of scatter, representing measurement ‘compressions’ and ‘rarefactions’ of the true circumferential distances due to arc length variations. Thus, we conclude that when using the convenient and computationally straightforward morphometric analysis on the ‘unwrapped’ 2-dimensional epithelial sheet, it is necessary to correct for non-uniform circumferential arc distance variation in order to obtain accurate morphometric measurements, and this approach was used for all quantitative analyses.

(4) Line Filtering

To eliminate any confounding effects due to the potentially polarized orientation of cell-cell interfaces (which are also efficiently stained with phalloidin), the interfaces were removed from the image for this analysis by masking off a dilated mask of the interface segmentation map, thereby isolating the image of the phalloidin signal only from the cell interiors. Line filtering was then performed by convolving the “masked” raw image with a 2-dimensional Ricker wavelet kernel of variable spatial angle.

The basic 2-dimensional wavelet profile is given by the equation 1, where $\psi(x, y)$ is the probability:

$$\psi(x, y) = \frac{2}{\sqrt{3\sigma}(\pi)^{0.25}} \left(1 - \left(\frac{x}{\sigma}\right)^2\right) \frac{x^2}{2\sigma^2}$$

The length of the kernel matrix in y -direction was $5\mu\text{m}$. The value of σ was selected to match the thickness of the fibers ($0.32\ \mu\text{m}$) in the raw image, slightly above the width of the PSF (Figure S4A-B), and the total width of the kernel matrix in x -direction was chosen to be 6σ (ranging from -3σ to $+3\sigma$), so as to encompass the first zero crossing (which occurs at $\pm\sigma$) while also making sure the kernel is not too large (which would significantly increase the computation time). Through the equation above, the filter kernel is already normalized in x -direction; to also normalize it in y -direction, each element was divided by the y -length of the kernel (in units of pixels). The spatial angle of the kernel with respect to the image coordinates was then varied using the MATLAB function *imrotate*.

The raw images were first normalized by subtraction of image mean and dividing by image standard

deviation before performing line filtering; subsequently, convolution of the image with the normalized and rotated kernel was performed using the MATLAB function *imfilter*.

(5) Quantitative morphometric and data analysis

Major and minor axes of a cell are determined in MATLAB's *regionprops* function by fitting an ellipse to the cell perimeter.

Statistical Analysis:

In Figures 3-5, unless explicitly stated otherwise, the results plots show mean \pm standard deviation (SD), where the mean and SD are calculated across the values from all unique blood vessels. Our sample size of analyzed images consisted of $n=30$ blood vessels in control conditions (from 10 unique animals), with a range of 26-219 cells per blood vessel for a total of $k=2,990$ cells; in BAPN-treated conditions, $n=20$ blood vessels (from 9 unique animals) with a range of 28-185 cells per blood vessel for a total of $k=1,660$ cells; in FAKI treated conditions, $n=13$ blood vessels (from 5 unique animals) with a range of 23-257 cells per blood vessel for a total of $k=1,391$ cells. Thus, the error bars in the figures represent the typical variation of the parameter between individual blood vessels (not between individual cells).

If the parameter of choice is cell-based (such as cell major axis), the parameter is first averaged within each blood vessel across the available cells in that vessel, before we then calculate the mean across blood vessels. Since each blood vessel contains a different number of cells (ranging from 23-257 cells per blood vessel), a weight corresponding to the number of cells is assigned to each blood vessel. Unless specified otherwise, p -values were calculated using a two-sample t -test on these weighted means (e.g. Control vs. BAPN, or circumferential vs. axial), with the number of observations in the t -test being n =the number of vessels (and not k =total number of cells). In cases where the value distributions were clearly non-normal, Kolmogorov-Smirnov tests were used as significance tests.

Acknowledgments:

This work was initiated through Research Corporation for Science Advancement (RCSA) 2016 Scialog fellowships to MV and DL, and supported by a collaborative 2016 Scialog award from RCSA and the Gordon and Betty Moore Foundation for the collaborative project 'Deconstructing the cell's mechanical circuits' to DL and MTV, as well as Adriana Dawes (Ohio State University) and Matthew Ferguson (Boise State University), who we thank for important technical input and useful discussions. The microscopy work and mariculture was partially supported by a grant from the G. Harold and Leila Y. Mathers Foundation Grant SB170066 (to MTV and ADT) and a UC Multicampus Research Programs and Initiatives Award #MRI-17-454791 (to MTV). CG was partially supported by a UCSB Faculty Senate Research Award (awarded to MTV). We acknowledge the use of the UCSB Neuroscience Research Institute/Department of Molecular Cellular and Developmental Biology Microscopy Facility and in particular the Leica SP8-TCS Resonant Scanning Confocal Microscope acquired through NSF MRI Grant DBI-1625770. We also thank Prof. JT Blankenship (University of Denver) for confocal images of the *drosophila* germ band epithelium.

Author Contributions: Conceived and designed experiments: DR, ADT, MTV and DL. Performed experiments: DR, CG, and SS. Conceived and designed image analysis: MTV and DL. Performed analysis: DL and RM. Manuscript was written by DR, RM, DL and MTV and edited by all authors.

References

- Bade, N. D., R. D. Kamien, R. K. Assoian and K. J. Stebe (2017). "Curvature and Rho activation differentially control the alignment of cells and stress fibers." *Sci Adv* **3**(9): e1700150.
- Bade, N. D., T. Xu, R. D. Kamien, R. K. Assoian and K. J. Stebe (2018). "Gaussian Curvature Directs Stress Fiber Orientation and Cell Migration." *Biophys J* **114**(6): 1467-1476.
- Biton, Y. Y. and S. A. Safran (2009). "The cellular response to curvature-induced stress." *Phys Biol* **6**(4): 046010.
- Boyd, H. C., S. K. Brown, J. A. Harp and I. L. Weissman (1986). "Growth and Sexual-Maturation of Laboratory-Cultured Monterey Botryllus-Schlosseri." *Biological Bulletin* **170**(1): 91-109.
- Braden, B. P., D. A. Taketa, J. D. Pierce, S. Kassmer, D. D. Lewis and A. W. De Tomaso (2014). "Vascular regeneration in a basal chordate is due to the presence of immobile, bi-functional cells." *PLoS One* **9**(4): e95460.
- Chazotte, B. (2010). "Labeling cytoskeletal F-actin with rhodamine phalloidin or fluorescein phalloidin for imaging." *Cold Spring Harb Protoc* **2010**(5): pdb prot4947.
- Chrobak, K. M., D. R. Potter and J. Tien (2006). "Formation of perfused, functional microvascular tubes in vitro." *Microvasc Res* **71**(3): 185-196.
- Dewey, C. F., Jr., S. R. Bussolari, M. A. Gimbrone, Jr. and P. F. Davies (1981). "The dynamic response of vascular endothelial cells to fluid shear stress." *J Biomech Eng* **103**(3): 177-185.
- Eisenhoffer, G. T., P. D. Loftus, M. Yoshigi, H. Otsuna, C. B. Chien, P. A. Morcos and J. Rosenblatt (2012). "Crowding induces live cell extrusion to maintain homeostatic cell numbers in epithelia." *Nature* **484**(7395): 546-549.
- Fernandez, P., C. Bourget, R. Bareille, R. Daculsi and L. Bordenave (2007). "Gene response in endothelial cells cultured on engineered surfaces is regulated by shear stress." *Tissue Eng* **13**(7): 1607-1614.
- Fidkowski, C., M. R. Kaazempur-Mofrad, J. Borenstein, J. P. Vacanti, R. Langer and Y. Wang (2005). "Endothelialized microvasculature based on a biodegradable elastomer." *Tissue Eng* **11**(1-2): 302-309.
- Flaherty, J. T., J. E. Pierce, V. J. Ferrans, D. J. Patel, W. K. Tucker and D. L. Fry (1972). "Endothelial nuclear patterns in the canine arterial tree with particular reference to hemodynamic events." *Circ Res* **30**(1): 23-33.
- Franke, R. P., M. Grafe, H. Schnittler, D. Seiffge, C. Mittermayer and D. Drenckhahn (1984). "Induction of human vascular endothelial stress fibres by fluid shear stress." *Nature* **307**(5952): 648-649.
- Galbraith, C. G., R. Skalak and S. Chien (1998). "Shear stress induces spatial reorganization of the endothelial cell cytoskeleton." *Cell Motil Cytoskeleton* **40**(4): 317-330.
- Gasparini, F., L. Manni, F. Cima, G. Zaniolo, P. Burighel, F. Caicci, N. Franchi, F. Schiavon, F. Rigon, D. Campagna and L. Ballarin (2015). "Sexual and asexual reproduction in the colonial ascidian *Botryllus schlosseri*." *Genesis* **53**(1): 105-120.
- Gotlieb, A. I., L. M. May, L. Subrahmanyam and V. I. Kalnins (1981). "Distribution of microtubule organizing centers in migrating sheets of endothelial cells." *J Cell Biol* **91**(2 Pt 1): 589-594.
- Gu, Y., T. Forostyan, R. Sabbadini and J. Rosenblatt (2011). "Epithelial cell extrusion requires the sphingosine-1-phosphate receptor 2 pathway." *J Cell Biol* **193**(4): 667-676.
- Heemskerk, I. and S. J. Streichan (2015). "Tissue cartography: compressing bio-image data by dimensional reduction." *Nat Methods* **12**(12): 1139-1142.
- Hellbach, A., S. Tiozzo, J. Ohn, M. Liebling and A. W. De Tomaso (2011). "Characterization of HCN and cardiac function in a colonial ascidian." *J Exp Zool A Ecol Genet Physiol* **315**(8): 476-486.

Holland, L. Z. (2016). "Tunicates." *Curr Biol* **26**(4): R146-152.

Iba, T. and B. E. Sumpio (1991). "Morphological response of human endothelial cells subjected to cyclic strain in vitro." *Microvasc Res* **42**(3): 245-254.

Ingber, D., Folkman, J (1988). "Inhibition of angiogenesis through modulation of collagen metabolism." *Laboratory Investigation; a Journal of Technical Methods and Pathology* **59**(1): 44-51.

Ives, C. L., S. G. Eskin and L. V. McIntire (1986). "Mechanical effects on endothelial cell morphology: in vitro assessment." *In Vitro Cell Dev Biol* **22**(9): 500-507.

Kassmer, S. H., D. Rodriguez and A. W. De Tomaso (2016). "Colonial ascidians as model organisms for the study of germ cells, fertility, whole body regeneration, vascular biology and aging." *Curr Opin Genet Dev* **39**: 101-106.

Kaunas, R., P. Nguyen, S. Usami and S. Chien (2005). "Cooperative effects of Rho and mechanical stretch on stress fiber organization." *Proc Natl Acad Sci U S A* **102**(44): 15895-15900.

Kutys, M. L. and C. S. Chen (2016). "Forces and mechanotransduction in 3D vascular biology." *Curr Opin Cell Biol* **42**: 73-79.

Liu, S. Q. (1998). "Influence of tensile strain on smooth muscle cell orientation in rat blood vessels." *J Biomech Eng* **120**(3): 313-320.

Livne, A. and B. Geiger (2016). "The inner workings of stress fibers - from contractile machinery to focal adhesions and back." *J Cell Sci* **129**(7): 1293-1304.

Marinari, E., A. Mehonic, S. Curran, J. Gale, T. Duke and B. Baum (2012). "Live-cell delamination counterbalances epithelial growth to limit tissue overcrowding." *Nature* **484**(7395): 542-545.

Rodriguez, D., B. P. Braden, S. W. Boyer, D. A. Taketa, L. Setar, C. Calhoun, A. D. Maio, A. Langenbacher, M. T. Valentine and A. W. De Tomaso (2017). "In vivo manipulation of the extracellular matrix induces vascular regression in a basal chordate." *Mol Biol Cell* **28**(14): 1883-1893.

Sato, M. and T. Ohashi (2005). "Biorheological views of endothelial cell responses to mechanical stimuli." *Biorheology* **42**(6): 421-441.

Shin, M., K. Matsuda, O. Ishii, H. Terai, M. Kaazempur-Mofrad, J. Borenstein, M. Detmar and J. P. Vacanti (2004). "Endothelialized networks with a vascular geometry in microfabricated poly(dimethyl siloxane)." *Biomed Microdevices* **6**(4): 269-278.

Song, J. W., S. P. Cavnar, A. C. Walker, K. E. Luker, M. Gupta, Y. C. Tung, G. D. Luker and S. Takayama (2009). "Microfluidic endothelium for studying the intravascular adhesion of metastatic breast cancer cells." *PLoS One* **4**(6): e5756.

Tiozzo, S., A. Voskoboynik, F. D. Brown and A. W. De Tomaso (2008). "A conserved role of the VEGF pathway in angiogenesis of an ectodermally-derived vasculature." *Dev Biol* **315**(1): 243-255.

Tojkander, S., G. Gateva and P. Lappalainen (2012). "Actin stress fibers--assembly, dynamics and biological roles." *J Cell Sci* **125**(Pt 8): 1855-1864.

Vanderleest, T. E., C. M. Smits, Y. Xie, C. E. Jewett, J. T. Blankenship and D. Loerke (2018). "Vertex sliding drives intercalation by radial coupling of adhesion and actomyosin networks during *Drosophila* germband extension." *Elife* **7**.

Wechezak, A. R., R. F. Viggers and L. R. Sauvage (1985). "Fibronectin and F-actin redistribution in cultured endothelial cells exposed to shear stress." *Lab Invest* **53**(6): 639-647.

White, G. E. and K. Fujiwara (1986). "Expression and intracellular distribution of stress fibers in aortic endothelium." *J Cell Biol* **103**(1): 63-70.

Wong, A. J., T. D. Pollard and I. M. Herman (1983). "Actin filament stress fibers in vascular endothelial cells in vivo." *Science* **219**(4586): 867-869.

Yang, R., E. Li, Y. J. Kwon, M. Mani and G. J. Beitel (2019). "QuBiT: a quantitative tool for analyzing epithelial tubes reveals unexpected patterns of organization in the *Drosophila* trachea." Development **146**(12).

Ye, M., H. M. Sanchez, M. Hultz, Z. Yang, M. Bogorad, A. D. Wong and P. C. Searson (2014). "Brain microvascular endothelial cells resist elongation due to curvature and shear stress." Sci Rep **4**: 4681.

Yevick, H. G., G. Duclos, I. Bonnet and P. Silberzan (2015). "Architecture and migration of an epithelium on a cylindrical wire." Proc Natl Acad Sci U S A **112**(19): 5944-5949.




Figure 1. The Extracorporeal Vasculature of *Botryllus schlosseri*. (A) Ventral view of a single colony of *Botryllus*, showing all individuals (zooid = z) at the center that are interconnected by an extracorporeal vasculature (ev), that extends out to blind end protrusions (ampulla = a). (Scale bar = 2mm). (B) Bright field micrograph showing blood vessels and ampullae. (Scale bar = 1mm). (C-F) Maximal projection of a confocal micrograph of a single blood vessel stained for DNA in blue (Scale bar = 25 μ m). (C), Pan-Cadherin in green (D), F-Actin in red (E). (F) Merged maximal projections.




Figure 2. Overview of Image Analysis Workflow. (A) Mean projection of the raw 3D image stack along the z-axis (Scale bar = 20 μ m). (B) Raw resized axial section of the vessel with the detected surface overlaid in red (Scale bar = 10 μ m). (C) Left panel: 2D projection of the vessel obtained by unwrapping the intensities along each axial section. Right panel: Intensity projection with cell outlines detected by watershed segmentation overlaid in red. Rectangular boxes correspond to non-repeat regions (Scale bar = 10 μ m). (D) 3D reconstruction of the blood vessel with each cell represented by a different color.




Figure 3. Cell Morphometrics Show No Significant Difference Between Control and BAPN Treated Vessels. (A) Distribution of vascular curvature, as assessed by the distribution of mean blood vessel diameters, analyzed under control (CTRL, upper, blue), BAPN-treatment (middle, red) and FAKI-treatment (lower, grey) conditions. (B) Distribution of cell areas for control (upper, blue), BAPN-treated (middle, red) and FAKI-treated (lower, grey) animals. (C) Distribution of the number of nearest cellular neighbors of control (blue), BAPN-treated (red) and FAKI-treated (grey) animals. Box plots show the central median, where edges of the boxes represent 25th and 75th percentiles (q_1 and q_3 , respectively); whiskers correspond to $q_1 - 1.5(\text{IQR})$ and $q_3 + 1.5(\text{IQR})$ with IQR=Inter Quartile Range. (D) 2D projection of the blood vessel with cells colored based on their neighbor number (Scale bar = 20 μm). For all the panels, the number of vessels (n) and number of cells (k) used were $n=30$, $k=2,990$ (ctrl); $n=20$, $k=1,660$ (BAPN-treated) and $n=13$, $k=1,391$ (FAKI-treated).




Figure 4. Both Cell Morphology and Actin Fiber Orientation Are Planar Polarized in Axial Vs Circumferential Direction. (A) Quantification of preferred actin fiber orientation, as determined by the response of a rotating line filter. The radial amplitude in the polar plot in each panel represents the line filter response in that direction (averaged over the entire 2D map) in one representative blood vessel in each condition (specifically, the vessel with the median response intensity for that condition). Control, CTRL = blue (upper plot), BAPN-treated = red (middle plot) and FAKI-treated = gray (lower plot) (B) Image of one representative control cell (with background masked out) with actin filaments stained with phalloidin (scale bar = 10 μ m). (C) Directional preference, defined as the contrast ratio of maximum vs minimum line filter responses like those shown in panel A, with $c=(\max-\min)/(\max+\min)$ of all vessels in Control, BAPN-treatment and FAKI-treatment conditions. (D) Absolute value of the maximum line filter response (like those shown in panel A) for each vessel for Control, BAPN- and FAKI-treated vessels. (E) Mean cellular major and minor axes lengths (averaged over all cells in each individual vessel) in Control, BAPN-treatment and FAKI-treatment conditions. (* $p<0.05$ and ** $p<10^{-8}$, 2-sample t -test). (F) Preferred major axis orientation angle of the cells. The radial amplitude in the polar plot represents the mean normalized probability – averaged over all vessels in that condition – where the radial lines are error bars representing standard deviation. For all the panels in this figure, number of vessels (n) and number of cells (k) were $n=30, k=2,990$ (CTRL); $n=20, k=1,660$ (BAPN-treated); and $n=13, k=1,391$ (FAKI-treated). Box plots show the central median, and edges of the boxes represent 25th and 75th percentiles (q_1 and q_3 , respectively); whiskers correspond to $q_1-1.5(\text{IQR})$ and $q_3+1.5(\text{IQR})$ with $\text{IQR}=\text{Inter Quartile Range}$.




Figure 5. Planar polarity is enhanced in smaller diameters. (A) Schematic showing categorization of cells into axial (75°-105°), circumferential (0°-15° and 165°-180°) and other (15°-75° and 105°-165°) categories based on their major axis orientation angle. (B) Fraction of cells with major axis oriented along axial (red) or circumferential (green) direction as a function of blood vessel diameter. (C) Polar histogram of the major axis orientation angle of the cells in small (Diameter $\leq 40 \mu\text{m}$) and large (Diameter $> 40 \mu\text{m}$) vessels; the radial amplitude represents the mean probability distribution and the thick black lines pointing radially along different polar angles represent the standard deviation. (D) Mean contrast ratio of filtered phalloidin intensity of cells as a function of blood vessel diameter. Contrast ratio = (circumferential response – axial response)/(circumferential response + axial response). (E) Heat maps of the normalized probability density as a function of major axis orientation angle (in angle bins of 30°) and of cell aspect ratio quintile for predominantly axially oriented blood vessels. Control = blue, BAPN-treatment = red and FAKI-treatment = gray. For all the panels number of vessels (n) and number of cells (k) used were $n=30$, $k=2,990$ (ctrl); $n=20$, $k=1,660$ (BAPN treated) and $n=13$, $k=1,391$ (FAK-inhibitor treated).

Table 1: Fraction of cells oriented into different planar directions (n =number of blood vessels, p = p -value of 2-sample t -test performed on weighted mean of axial and circumferential fractions). Significant p -values ($p<0.05$) are shown in bold font. Fractions are shown as (weighted mean \pm weighted SD).

	Control				BAPN				FAK			
	n	Axial fraction	Circumf fraction	p	n	Axial fraction	Circumf fraction	p	n	Axial fraction	Circumf fraction	p
d<40	14	0.34 \pm 0.14	0.10 \pm 0.06	4.8e-05	12	0.42 \pm 0.20	0.11 \pm 0.12	8.9e-04	3	0.24 \pm 0.03	0.18 \pm 0.02	0.11
d>40	16	0.21 \pm 0.08	0.16 \pm 0.07	0.07	8	0.23 \pm 0.12	0.14 \pm 0.09	0.13	10	0.22 \pm 0.14	0.13 \pm 0.06	0.08