

# UC Berkeley

## UC Berkeley Previously Published Works

### Title

Modeling the development of coarse-to-fine processing in the central visual pathway

### Permalink

<https://escholarship.org/uc/item/2ts8k55x>

### Journal

BMC Neuroscience, 14(Suppl 1)

### ISSN

1471-2202

### Author

Nirody, Jasmine A

### Publication Date

2013-07-08

### DOI

<http://dx.doi.org/10.1186/1471-2202-14-S1-P294>

Peer reviewed

POSTER PRESENTATION

Open Access


# Modeling the development of coarse-to-fine processing in the central visual pathway

Jasmine A Nirody

From Twenty Second Annual Computational Neuroscience Meeting: CNS\*2013  
Paris, France. 13-18 July 2013

The mode of information processing in neural sensory systems has been the subject of many experimental and computational studies. In the visual system, the existence of a sequential analysis of information, where coarse features are processed before fine details, is intuitive—when

looking quickly at a scene, we process general structures before focusing on individual objects. Recent studies have shown that cortical spatial coarse-to-fine processing is largely accounted for by feedforward input from the lateral geniculate nucleus (LGN) in adults [1]. However, the



Correspondence: jnirody@berkeley.edu  
Biophysics Graduate Group, University of California, Berkeley, Berkeley, CA,  
94720, USA

developmental aspect of this process has not yet been considered. We use experimental data from LGN neurons from three age groups: adult cats and kittens at 4 and 8 weeks postnatal [2] to construct a firing-rate based thalamocortical model. We consider how developmental changes in spatiotemporal structure affect feedforward thalamic contribution to cortical spatial frequency (SF) tuning, as well as the efficacy of cortical feedback in facilitating this dynamic. Our model predicts a non-monotone relationship between relative surround strength and shift in tuning peak, suggesting the existence of an optimal center-surround balance which maximizes SF shift. Taken together with previous data, our results suggest that the ratios for mature animals are distributed close to or around their “optimum”, while the distribution for kittens is centered at a ratio significantly lower than their peak value. As cortical feedback has been suggested to strengthen the antagonistic effects of the surround response, this implies that kittens early in the developmental process are more strongly affected by corticothalamic feedback than adults. Our results point to these recurrent connections as an important mechanism in facilitating the development of spatial coarse-to-fine processing.

#### Acknowledgements

The author thanks Ralph Freeman for the suggestion of studying the developmental process, and Bartlett Moore for many helpful discussions.

Published: 8 July 2013

#### References

1. Allen EA, Freeman RD: **Dynamic spatial processing originates in early visual pathways.** *J Neurosci* 2006, **26**(45):11763-11774.
2. Cai D, DeAngelis GC, Freeman RD: **Spatiotemporal receptive field organization in the lateral geniculate nucleus of cats and kittens.** *J Neurophysiol* 1997, **78**(2):1045-1061.

doi:10.1186/1471-2202-14-S1-P294

**Cite this article as:** Nirody: Modeling the development of coarse-to-fine processing in the central visual pathway. *BMC Neuroscience* 2013 **14** (Suppl 1):P294.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

