

# UCSF

## UC San Francisco Previously Published Works

### Title

Nicotine and cannabis vaping among adolescents in treatment for substance use disorders.

### Permalink

<https://escholarship.org/uc/item/2tv6736d>

### Authors

Young-Wolff, Kelly C  
Adams, Sara R  
Sterling, Stacy A  
[et al.](#)

### Publication Date

2021-06-01

### DOI

10.1016/j.jsat.2021.108304

Peer reviewed



Published in final edited form as:

*J Subst Abuse Treat.* 2021 June ; 125: 108304. doi:10.1016/j.jsat.2021.108304.

## Nicotine and cannabis vaping among adolescents in treatment for substance use disorders

Kelly C. Young-Wolff, PhD, MPH<sup>1,2</sup>, Sara R. Adams, MPH<sup>1</sup>, Stacy Sterling, DrPH, MSW<sup>1</sup>, Andy S.L. Tan, PhD, MPH, MBA, MBBS<sup>2</sup>, Ramzi G. Salloum, PhD<sup>3</sup>, Kira Torre, MS, MFT<sup>4</sup>, Lisa Carter-Harris, PhD, APRN, ANP-C, FAAN<sup>5</sup>, Judith J. Prochaska, PhD, MPH<sup>6</sup>

<sup>1</sup>Division of Research, Kaiser Permanente Northern California

<sup>2</sup>Annenberg School for Communication, University of Pennsylvania, Philadelphia, PA

<sup>3</sup>Department of Health Outcomes and Biomedical Informatics, and Institute for Child Health Policy, University of Florida College of Medicine, Gainesville, FL

<sup>4</sup>Addiction Medicine and Recovery Services, Kaiser Permanente Northern California

<sup>5</sup>Department of Psychiatry and Behavioral Sciences, Memorial Sloan Kettering Cancer Center, New York, NY

<sup>6</sup>Stanford Prevention Research Center, Department of Medicine, Stanford University

### Abstract

This study examined nicotine and cannabis vaping among adolescents in treatment for substance use disorders. Participants were 363 adolescents aged 12–17 (66% male, mean age=15.5 [SD=1.3], 46% non-Hispanic white) seen for a specialty addiction intake evaluation between 2017 and 2019 at one of six medical offices of a large, integrated health care system in Northern California. Multivariable logistic regression models tested for associations of sociodemographics, cigarette smoking, and substance use disorders with vaping behaviors. A majority of adolescents reported ever (68%) or current vaping (60%) of nicotine and/or cannabis; current vaping was similar for nicotine (50%) and cannabis (51%); 40% reported current vaping of both. Current smokers (6% of the sample) had higher odds of ever vaping (aOR=3.95, 95%CI:1.04–14.95). Black (versus non-Hispanic white) adolescents had lower odds of current nicotine vaping (aOR=0.08, 95%CI:0.02–0.37) and current vaping of both nicotine and cannabis (aOR=0.12, 95%CI:0.03–0.60). Having an alcohol use disorder was associated with current vaping (aOR=2.14, 95%CI:1.06–4.33). Those who endorsed that most friends get drunk/high (aOR=1.87,

---

**Send correspondence to:** Kelly C. Young-Wolff, PhD, MPH, Research Scientist, Division of Research, Kaiser Permanente Northern California, 2000 Broadway, Oakland, CA 94612, USA.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

#### Conflict of Interest

JPP has provided consultation to pharmaceutical and technology companies that make medications and other treatments for quitting smoking and has served as an expert witness in lawsuits against the tobacco companies. The other authors declare that they have no financial relationships with any organizations that might have an interest in the submitted work. Other authors report no conflicts of interest.

95% CI: 1.02–3.42) or that cannabis was their substance of choice (aOR=2.36, 95% CI: 1.16–4.81) had higher odds of current cannabis vaping. Higher neighborhood household income (\$80,000–\$120,000 and >\$120,000 vs. <\$80,000, aORs=2.05–9.48), never versus ever blunt use (aORs=2.47–8.68), and intakes in 2018 and 2019 versus 2017 (aORs=2.18–5.38) were associated with higher odds of all vaping outcomes. Vaping was common among adolescents in addiction treatment and varied with sociodemographics and substance-related factors. Research should assess how vaping impacts the development of substance use disorders and whether it interferes with addiction treatment.

## Keywords

E-cigarette; Adolescents; Vaping; Nicotine; Cannabis; Addiction treatment

## 1. Introduction

Vaping is an increasingly common mode of nicotine and cannabis administration among adolescents (Cullen et al., 2019; Dai, 2019; Miech, Patrick, O'Malley, Johnston, & Bachman, 2019; Wang et al., 2019). National data indicate that past-month nicotine vaping increased among U.S. high school students from 11% in 2016 to 20% in 2020, with 23% of users reporting daily use (Cullen et al., 2019; Wang et al., 2020). Similarly, from 2017 to 2019, past-month cannabis vaping increased from 4% to 13% among 10th grade students and from 5% to 14% among 12th grade students (Miech et al., 2019). Nicotine and cannabis vaping during adolescence is associated with significant harms (Singh et al., 2020; U.S. Department of Health and Human Services, 2016, 2019), and the U.S. Surgeon General has declared youth vaping an epidemic (U.S. Department of Health and Human Services, 2019).

A number of factors are associated with adolescent nicotine vaping, including use among peers, e-cigarette marketing, psychological problems, illicit drug use, cigarette smoking and harmful alcohol consumption, household exposure to smoking, and Hispanic or non-Hispanic white race/ethnicity (Barrington-Trimis et al., 2015; Barrington-Trimis et al., 2016; Fielding-Singh, Epperson, & Prochaska, 2020; Kwon, Seo, Lin, & Chen, 2018; Pike et al., 2019). Less is known about correlates of adolescent cannabis vaping, but initial studies indicate that it is more prevalent among current cigarette smokers, older adolescents, and Hispanic or non-Hispanic white adolescents (Dai, 2019).

Despite increased attention to adolescent vaping, few studies have examined vaping among adolescents in addiction treatment. Data from adults in addiction treatment suggest that one-quarter to one-third report current e-cigarette use, compared to 3% of the U.S. population in 2018 (Villarroel, Cha, & Vahratian, 2020), with quitting or reducing combustible cigarette smoking the most commonly endorsed reason for use (Campbell, Le, Gubner, & Guydish, 2019; Gubner, Andrews, Mohammad-Zadeh, Lisha, & Guydish, 2016; Peters et al., 2015; Stein et al., 2015). Studies of vaping among adolescents in addiction treatment and identifying covariates of use are critical to our understanding of at-risk subgroups.

This study capitalizes on the pilot implementation of a new intake template for adolescent specialty addiction treatment that included questions about nicotine and cannabis vaping in

six Kaiser Permanente Northern California (KPNC) medical offices as part of standard care. Study goals are to examine: 1) the prevalence of ever and current nicotine and cannabis vaping in this population, and 2) whether vaping behaviors differ by sociodemographics and behavioral health characteristics.

## 2. Materials and methods

### 2.1. Study setting and participants

KPNC is a nonprofit, integrated health care delivery system providing comprehensive health services to 4.5 million members. Adolescents for whom nicotine use is their only substance of abuse are typically treated within primary care. The KPNC Addiction Medicine and Recovery Services (AMRS) Department provides treatment to adolescents for other substance use disorders, including cannabis. The AMRS Department rolled out a new intake form with questions about ever and current vaping of nicotine and cannabis in six medical offices between January 1, 2017, and July 1, 2019. A total of 485 adolescents aged 12–17 received an intake for addiction treatment between the roll-out of the form at each office and December 31, 2019; of those, 363 patients (75%) completed the new intake form and are included in this study. The 122 patients who received an older version of the form without the vaping question did not differ significantly from those who were included on age, sex, or race/ethnicity; however, they were less likely to be diagnosed with a cannabis use disorder ( $p < .001$ ), perhaps in part because the form did not assess cannabis vaping specifically.

### 2.3. Measures

This study extracted data on age, race/ethnicity, sex, median neighborhood household income (geocoded from census data using patients' addresses), smoking status, and year of intake from the electronic health record (EHR). The study used *International Classification of Diseases, Tenth Revision, Clinical Modification* diagnosis codes to define anxiety disorders (F41, F42, F43, F930, excluding F43.21, F43.23); depressive disorders (F32, F33, F34.1, F43.21, F43.23, excluding F32.5, F33.42); and substance use disorders (excluding “in remission” codes), including alcohol use disorder (F10), cannabis use disorder (F12), other substance use disorders (for opioids; sedatives, hypnotics, or anxiolytics; cocaine; other stimulants; hallucinogens; inhalants; or other psychoactive substances; F11, F13–F16, F18–F19). The study derived ever and current vaping of nicotine and cannabis, substance of choice (cannabis versus other), and whether most of one's friends get drunk/high from standardized response options within the intake template. The study did not assess vaping of other substances, such as nicotine-free e-liquids, or cannabis use as an edible or joint. However, since smoking cannabis within a cigar casing is a common method among African American and Latinx adolescents and distinct enough to detect if reported in the EHR (Fairman, 2015; Montgomery & Mantey, 2017; Ream, Benoit, Johnson, & Dunlap, 2008; Schauer, Rosenberry, & Peters, 2017; Timberlake, 2009), a text search of the intake notes determined history of ever smoking a blunt. Outcomes included: 1) ever vaping nicotine and/or cannabis, 2) current nicotine vaping, 3) current cannabis vaping, 4) any current vaping (nicotine and/or cannabis), and 5) current vaping of both nicotine and cannabis.

## 2.4. Statistical analyses

The study reported prevalence of vaping outcomes specified here overall and by patients' sociodemographics and comorbidity status; chi-square tests assessed differences in vaping outcomes by patient characteristics. Multivariable logistic regression analyses evaluated whether outcomes were associated with age, sex, race/ethnicity, geocoded household income, alcohol use disorder, other substance use disorder, current smoking status, blunt use, whether most friends get drunk/high, whether cannabis was listed as a substance of choice, and intake year. Cannabis use disorder, depressive disorder, and anxiety disorder were not associated with vaping outcomes in bivariate analyses, and this study excluded them from multivariable models.

## 3. Results

The sample (N=363) had a mean age of 15.5 years (SD=1.3) and was 66% male; 13% Asian/Pacific Islander, 9% Black, 26% Hispanic, 46% non-Hispanic white, and 6% other/unknown race/ethnicity. Approximately two-thirds of the adolescents reported ever vaping (68%, Table 1); half reported current nicotine vaping (50%) or current cannabis vaping (51%), 60% reported any current vaping of nicotine and/or cannabis, and 40% reported current vaping of both. Six percent reported current cigarette smoking and 19% reported ever using blunts. Two-thirds (66%) had a cannabis use disorder, 16% an alcohol use disorder, and 14% another substance use disorder; 75% said cannabis was their only substance of choice and another 11% named cannabis as one of multiple substances of choice; 35% and 40% had an anxiety or depressive disorder, respectively.

Ever and current vaping of nicotine and/or cannabis was significantly more likely among those residing in neighborhoods of higher household income; who endorsed that most of their friends get drunk or high; who had an intake in 2018 or 2019 versus 2017; and among Asian/Pacific Islander and non-Hispanic white adolescents (Table 1). Ever and current vaping of nicotine and/or cannabis was less likely among Black adolescents and those of other/unknown race/ethnicity, and more likely among adolescents who had never used a blunt. Current nicotine vaping was significantly more likely among adolescents with alcohol use disorders.

In multivariable models (Table 2), those with higher incomes had greater odds of all vaping outcomes (\$80K–120K, aORs=2.05–3.34; >\$120K, aORs=3.68–9.48) relative to those with neighborhood household incomes <\$80K. Having an alcohol use disorder was associated with higher odds of current nicotine vaping (aOR=2.14, 95%CI:1.06–5.05), and current smokers had higher odds of ever vaping (aOR=3.95, 95%CI:1.04–15.95). Never blunt use was associated with higher odds of all vaping outcomes (vs. ever blunt use, aORs=2.47–8.68). Reporting most friends get drunk/high (aOR=1.87, 95%CI:1.02–3.42) or cannabis as a substance of choice (aOR 2.36, 95%CI:1.16–4.81) were associated with higher odds of current cannabis vaping. Having an intake in the last two years of the study (2018 and 2019) was associated with higher odds of all vaping outcomes (versus 2017, aORs=2.18–5.38). Finally, Black adolescents had lower odds of current nicotine vaping (aOR=0.08, 95%CI:0.02–0.37) and current vaping of both nicotine and cannabis (aOR=0.12, 95%CI:0.03–0.60) relative to non-Hispanic white adolescents. In post hoc analyses, we found that Black

adolescents were more likely to report ever blunt use (50%) compared to adolescents with non-Hispanic white (12%), Asian (24%), Hispanic (18%), and other/unknown (23%) race/ethnicity (chi-square  $p < .001$ ). Current smoking was very low for Black adolescents (3%) and adolescents with other/unknown race/ethnicity (0%) and slightly higher for adolescents with non-Hispanic white (7%), Asian (7%), and Hispanic (7%) race/ethnicity, although the difference was not statistically significant (Fisher's Exact Test  $p = .81$ ).

#### 4. Discussion

To our knowledge, this is the first study of vaping behaviors among adolescents entering addiction treatment, and findings indicate higher prevalence than surveillance estimates for adolescents in the U.S. (Cullen, et al., 2019, Miech et al., 2019). Most (85%) reported cannabis as one of their substances of choice, 60% reported current vaping of nicotine and/or cannabis, and about two-thirds had a cannabis use disorder. Of concern, cannabis vaping is associated with vaping use-associated lung injury (EVALI) (American Lung Association, 2020). The high prevalence of cannabis-related problems is consistent with national data on adolescent admissions to addiction treatment (Substance Abuse and Mental Health Services Administration & Center for Behavioral Health Statistics and Quality, 2017).

Notably, adolescents who reported ever using blunts had significantly lower odds of all vaping outcomes, suggesting that those who smoke cannabis via blunts may be less inclined to vape. Study findings also suggested racial/ethnic differences in mode of cannabis administration, with vaping of both nicotine and cannabis less likely, and ever blunt use more likely, among Black versus non-Hispanic white adolescents. Nicotine and cannabis vaping were more common among adolescents who lived in neighborhoods with higher versus lower median household incomes, consistent with prior research in California among young adults (Meng & Ponce, 2020). These findings may be mediated by greater vaping advertisements in higher income neighborhoods (Dai, 2019; Simon et al., 2018). However, other studies of adolescent vaping and socioeconomic status have reported no association (Barrington-Trimis et al., 2015; Moore et al., 2015) or increased risk with lower socioeconomic status (Simon et al., 2017). Research is needed to better understand the mechanisms that contribute to racial/ethnic and socioeconomic differences in vaping among adolescents in addiction treatment.

Interestingly, current smoking was rare (6%) and only associated with increased odds of ever vaping. The lack of association with current vaping behaviors may be due to the low prevalence of smoking in our sample and a potential lack of power to detect associations with vaping outcomes. Adolescents with an alcohol use disorder or who identified cannabis as their substance of choice had higher odds of current cannabis vaping. Other substance use disorders were not significantly associated with vaping outcomes.

This study took place over the period that cannabis was being legalized for adult use in California in January 2018, and we found that all adolescent vaping behaviors increased significantly from 2017 to 2019. Legalization could increase youth use of cannabis via greater availability and desirability, reduced prices, and increased access to cannabis vaping,

the latter of which is considered a more discreet, healthier and cost-effective mode of cannabis delivery than smoking (Budney, Sargent, & Lee, 2015; Malouff, Rooke, & Copeland, 2014). Initial data from Colorado indicate that adolescent addiction medicine providers perceived that adult-use cannabis legalization contributed to normalization of cannabis; validation of its consumption; greater access to new, potent cannabis products; and increased treatment need (Sobesky & Gorgens, 2016).

#### 4.1. Limitations

Participants were treated within a large health care delivery system and results may not generalize to adolescents outside of KPNC or to those without health care access. We cannot determine whether vaping started before or after the onset of other substance use. Outcomes are based on self-report during standard care and may underestimate the prevalence of vaping behaviors. The intake template did not systematically capture other modes of cannabis administration (e.g., edibles, joint) or ever smoking status. Blunt use was based on a text search of the intake notes and is likely under-detected. Finally, the lack of association between cannabis use disorders and vaping may be due to the high prevalence of cannabis use disorders in our sample. Future studies with larger samples should replicate this study's findings. Nevertheless, this study provides useful data on vaping among adolescents receiving addiction treatment that can inform future efforts to improve vaping screening and better assess the impact of vaping on addiction treatment in this population.

## 5. Conclusion

Given the high prevalence of vaping, results suggest that health care systems and clinicians who treat adolescent substance use may benefit from education and training in how to screen for and address vaping behaviors as part of addiction treatment. Further, the low prevalence of cigarette smoking indicates that exposure to nicotine may be overlooked if adolescent addiction treatment intake screenings for tobacco are limited to cigarettes and do not include nicotine vaping. Future large, longitudinal studies should look at the temporal relation between adolescent vaping and the development of substance use disorders and identify how vaping influences addiction treatment and recovery.

## Acknowledgments

### Sources of Support

Funding for this study was provided by a NIH NIDA K01 Award (DA043604) and a Kaiser Permanente Division of Research Behavioral Health, Aging, and Infectious Diseases Section Grant. NIDA and the Kaiser Permanente Division of Research had no role in the study design, collection, analysis or interpretation of the data, writing the manuscript, or the decision to submit the paper for publication.

## References

- American Lung Association. (2020). E-cigarette or vaping use-associated lung injury (EVALI). <https://www.lung.org/lung-health-diseases/lung-disease-lookup/evali>. Accessed September 30, 2020.
- Barrington-Trimis JL, Berhane K, Unger JB, Cruz TB, Huh J, Leventhal AM, et al. (2015). Psychosocial factors associated with adolescent electronic cigarette and cigarette use. *Pediatrics*, 136(2), 308–317. 10.1542/peds.2015-0639. [PubMed: 26216326]

- Barrington-Trimis JL, Berhane K, Unger JB, Cruz TB, Urman R, Chou CP, et al. (2016). The e-cigarette social environment, e-cigarette use, and susceptibility to cigarette smoking. *Journal of Adolescent Health*, 59(1), 75–80. 10.1016/j.jadohealth.2016.03.019.
- Budney AJ, Sargent JD, & Lee DC. (2015). Vaping cannabis (marijuana): parallel concerns to ecigs? *Addiction*, 110(11), 1699–1704. 10.1111/add.13036. [PubMed: 26264448]
- Campbell BK, Le T, Gubner NR, & Guydish J (2019). Health risk perceptions and reasons for use of tobacco products among clients in addictions treatment. *Addictive Behaviors*, 91, 149–155. 10.1016/j.addbeh.2018.08.037. [PubMed: 30206006]
- Cullen KA, Gentzke AS, Sawdey MD, Chang JT, Anic GM, Wang TW, et al. (2019). E-cigarette use among youth in the United States, 2019. *JAMA*. 10.1001/jama.2019.18387.
- Dai H (2019). Self-reported marijuana use in electronic cigarettes among US youth, 2017 to 2018. *JAMA*. 10.1001/jama.2019.19571.
- Fairman BJ. (2015). Cannabis problem experiences among users of the tobacco-cannabis combination known as blunts. *Drug and Alcohol Dependence*, 150, 77–84. 10.1016/j.drugalcdep.2015.02.014. [PubMed: 25746234]
- Fielding-Singh P, Epperson AE, & Prochaska JJ. (2020). Tobacco product promotions remain ubiquitous and are associated with use and susceptibility to use among adolescents. *Nicotine and Tobacco Research*. 10.1093/ntr/ntaa136.
- Gubner NR, Andrews KB, Mohammad-Zadeh A, Lisha NE, & Guydish J (2016). Electronic-cigarette use by individuals in treatment for substance abuse: A survey of 24 treatment centers in the United States. *Addictive Behaviors*, 63, 45–50. 10.1016/j.addbeh.2016.06.025. [PubMed: 27423097]
- Kwon E, Seo DC, Lin HC, & Chen Z (2018). Predictors of youth e-cigarette use susceptibility in a U.S. nationally representative sample. *Addictive Behaviors*, 82, 79–85. 10.1016/j.addbeh.2018.02.026. [PubMed: 29501011]
- Malouff JM, Rooke SE, & Copeland J (2014). Experiences of marijuana-vaporizer users. *Substance Abuse*, 35(2), 127–128. 10.1080/08897077.2013.823902. [PubMed: 24821347]
- Meng Y, & Ponce NA. (2020). The changing landscape: Tobacco and marijuana use among young adults in California. Los Angeles: UCLA Center for Health Policy Research. [https://healthpolicy.ucla.edu/publications/Documents/PDF/2020/Smoking\\_Landscape\\_June2020.pdf](https://healthpolicy.ucla.edu/publications/Documents/PDF/2020/Smoking_Landscape_June2020.pdf). Accessed July 16, 2020.
- Miech RA, Patrick ME, O'Malley PM, Johnston LD, & Bachman JG. (2019). Trends in reported marijuana vaping among US adolescents, 2017–2019. *JAMA*. 10.1001/jama.2019.20185.
- Montgomery L, & Mantey DS. (2017). Correlates of blunt smoking among African American, Hispanic/Latino, and White Adults: Results from the 2014 National Survey on Drug Use and Health. *Substance Use and Misuse*, 52(11), 1449–1459. 10.1080/10826084.2017.1284238. [PubMed: 28467153]
- Moore G, Hewitt G, Evans J, Littlecott HJ, Holliday J, Ahmed N, et al. (2015). Electronic-cigarette use among young people in Wales: Evidence from two cross-sectional surveys. *BMJ Open*, 5(4), e007072. 10.1136/bmjopen-2014-007072.
- Peters EN, Harrell PT, Hendricks PS, O'Grady KE, Pickworth WB, & Voci FJ. (2015). Electronic cigarettes in adults in outpatient substance use treatment: Awareness, perceptions, use, and reasons for use. *American Journal on Addictions*, 24(3), 233–239. 10.1111/ajad.12206.
- Pike JR, Tan N, Miller S, Cappelli C, Xie B, & Stacy AW. (2019). The effect of e-cigarette commercials on youth smoking: A prospective study. *American Journal of Health Behavior*, 43(6), 1103–1118. 10.5993/AJHB.43.6.8. [PubMed: 31662169]
- Ream GL, Benoit E, Johnson BD, & Dunlap E (2008). Smoking tobacco along with marijuana increases symptoms of cannabis dependence. *Drug and Alcohol Dependence*, 95(3), 199–208. 10.1016/j.drugalcdep.2008.01.011. [PubMed: 18339491]
- Schauer GL, Rosenberry ZR, & Peters EN. (2017). Marijuana and tobacco co-administration in blunts, spliffs, and mulled cigarettes: A systematic literature review. *Addictive Behaviors*, 64, 200–211. 10.1016/j.addbeh.2016.09.001. [PubMed: 27654966]
- Simon P, Camenga DR, Kong G, Connell CM, Morean ME, Cavallo DA, et al. (2017). Youth e-cigarette, blunt, and other tobacco use profiles: Does SES matter? *Tobacco Regulatory Science*, 3(1), 115–127. 10.18001/TRS.3.1.12. [PubMed: 29082301]



- Simon P, Camenga DR, Morean ME, Kong G, Bold KW, Cavallo DA, et al. (2018). Socioeconomic status and adolescent e-cigarette use: The mediating role of e-cigarette advertisement exposure. *Preventive Medicine*, 112, 193–198. 10.1016/j.ypmed.2018.04.019. [PubMed: 29673887]
- Singh S, Windle SB, Fillion KB, Thombs BD, O'Loughlin JL, Grad R, et al. (2020). E-cigarettes and youth: Patterns of use, potential harms, and recommendations. *Preventive Medicine*, 133, 106009. 10.1016/j.ypmed.2020.106009. [PubMed: 32027913]
- Sobesky M, & Gorgens K (2016). Cannabis and adolescents: Exploring the substance misuse treatment provider experience in a climate of legalization. *International Journal of Drug Policy*, 33, 66–74. 10.1016/j.drugpo.2016.02.008.
- Stein MD, Caviness CM, Grimone K, Audet D, Borges A, & Anderson BJ. (2015). E-cigarette knowledge, attitudes, and use in opioid dependent smokers. *Journal of Substance Abuse Treatment*, 52, 73–77. 10.1016/j.jsat.2014.11.002. [PubMed: 25483740]
- Substance Abuse and Mental Health Services Administration, & Center for Behavioral Health Statistics and Quality. (2017). Treatment Episode Data Set (TEDS): 2005–2015. National Admissions to Substance Abuse Treatment Services. Rockville, MD: Substance Abuse and Mental Health Services Administration. <https://www.samhsa.gov/data/sites/default/files/2015%20TEDS%20National%20Admissions.pdf>. Accessed September 30, 2020.
- Timberlake DS. (2009). A comparison of drug use and dependence between blunt smokers and other cannabis users. *Substance Use and Misuse*, 44(3), 401–415. 10.1080/10826080802347651. [PubMed: 19212929]
- U.S. Department of Health and Human Services. (2016). E-cigarette use among youth and young adults. A report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. [https://e-cigarettes.surgeongeneral.gov/documents/2016\\_SGR\\_Full\\_Report\\_non-508.pdf](https://e-cigarettes.surgeongeneral.gov/documents/2016_SGR_Full_Report_non-508.pdf). Accessed July 16, 2020.
- U.S. Department of Health and Human Services. (2019). Surgeon General's advisory on e-cigarette use among youth. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. [https://www.cdc.gov/tobacco/basic\\_information/e-cigarettes/surgeon-general-advisory/index.html](https://www.cdc.gov/tobacco/basic_information/e-cigarettes/surgeon-general-advisory/index.html). Accessed April 22, 2020.
- Villarroel MA, Cha AE, & Vahratian A (2020). Electronic cigarette use among U.S. adults, 2018. Hyattsville, MD.: National Center for Health Statistics. <https://www.cdc.gov/nchs/products/databriefs/db365.htm>. Accessed October 6, 2020.
- Wang TW, Gentzke AS, Creamer MR, Cullen KA, Holder-Hayes E, Sawdey MD, et al. (2019). Tobacco product use and associated factors among middle and high school students - United States, 2019. *MMWR: Surveillance Summaries*, 68(12), 1–22. 10.15585/mmwr.ss6812a1.
- Wang TW, Neff LJ, Park-Lee E, Ren C, Cullen KA, & King BA. (2020). E-cigarette use among middle and high school students - United States, 2020. *Morbidity and Mortality Weekly Report (MMWR)*, 69(37), 1310–1312. 10.15585/mmwr.mm6937e1. [PubMed: 32941408]

### Highlights

- This study examined nicotine and cannabis vaping in adolescents in addiction treatment.
- Ever and current vaping of nicotine and cannabis was common.
- Vaping was more common among adolescents residing in higher income neighborhoods.
- Vaping prevalence varied by race/ethnicity and alcohol use disorder status.
- Addiction medicine clinicians should screen for and address nicotine and cannabis vaping.

**Table 1**

Patient characteristics and the prevalence of vaping among adolescents in treatment for substance use disorders, overall and by patient characteristics.

Patient characteristics	Study cohort (N=363)		Ever vaped nicotine and/or cannabis (N=246)		Current nicotine vaping (N=180)		Current cannabis vaping (N=184)		Any current vaping (nicotine and/or cannabis) (N=217)		Current vaping of both nicotine and cannabis (N=147)	
	N (column %)	N (row %)	P value <sup>a</sup>	N (row %)	P value <sup>a</sup>	N (row %)	P value <sup>a</sup>	N (row %)	P value <sup>a</sup>	N (row %)	P value <sup>a</sup>	
Overall	363 (100.0%)	246 (67.8%)		180 (49.6%)		184 (50.7%)		217 (59.8%)		147 (40.5%)		
Sex			.24		.91		.17		.08		.78	
Female	124 (34.2%)	89 (71.8%)		62 (50.0%)		69 (55.6%)		82 (66.1%)		49 (39.5%)		
Male	239 (65.8%)	157 (65.7%)		118 (49.4%)		115 (48.1%)		135 (56.5%)		98 (41.0%)		
Age			.63		.48		.65		.39		.76	
12–15	164 (45.2%)	109 (66.5%)		78 (47.6%)		81 (49.4%)		94 (57.3%)		65 (39.6%)		
16–17	199 (54.8%)	137 (68.8%)		102 (51.3%)		103 (51.8%)		123 (61.8%)		82 (41.2%)		
Race/ethnicity			<.001		<.001		<.001		<.001		<.001	
Asian/Pacific Islander	46 (12.7%)	36 (78.3%)		30 (65.2%)		30 (65.2%)		32 (69.6%)		28 (60.9%)		
Black	34 (9.4%)	14 (41.2%)		2 (5.9%)		10 (29.4%)		10 (29.4%)		2 (5.9%)		
Hispanic	95 (26.2%)	58 (61.1%)		45 (47.4%)		42 (44.2%)		53 (55.8%)		34 (35.8%)		
Non-Hispanic White	166 (45.7%)	125 (75.3%)		94 (56.6%)		96 (57.8%)		111 (66.9%)		79 (47.6%)		
Other/unknown	22 (6.1%)	13 (59.1%)		9 (40.9%)		6 (27.3%)		11 (50.0%)		4 (18.2%)		
Geocoded median household income <sup>b</sup>			<.001		<.001		<.001		<.001		<.001	
<\$80K	122 (33.6%)	64 (52.5%)		33 (27.0%)		37 (30.3%)		49 (40.2%)		21 (17.2%)		
\$80K–120K	125 (34.4%)	86 (68.8%)		63 (50.4%)		63 (50.4%)		76 (60.8%)		50 (40.0%)		
>\$120K+	115 (31.7%)	96 (83.5%)		84 (73.0%)		84 (73.0%)		92 (80.0%)		76 (66.1%)		
Alcohol use disorder			.13		.03		.55		.10		.23	
Yes	59 (16.3%)	45 (76.3%)		37 (62.7%)		32 (54.2%)		41 (69.5%)		28 (47.5%)		
No	304 (83.7%)	201 (66.1%)		143 (47.0%)		152 (50.0%)		176 (57.9%)		119 (39.1%)		
Cannabis use disorder			.07		.08		.21		.09		.16	
Yes	240 (66.1%)	155 (64.6%)		111 (46.3%)		116 (48.3%)		136 (56.7%)		91 (37.9%)		
No	123 (33.9%)	91 (74.0%)		69 (56.1%)		68 (55.3%)		81 (65.9%)		56 (45.5%)		

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Patient characteristics	Study cohort (N=363)		Ever vaped nicotine and/or cannabis (N=246)		Current nicotine vaping (N=180)		Current cannabis vaping (N=184)		Any current vaping (nicotine and/or cannabis) (N=217)		Current vaping of both nicotine and cannabis (N=147)	
	N (column %)	N (row %)	P value <sup>a</sup>	N (row %)	P value <sup>a</sup>	N (row %)	P value <sup>a</sup>	N (row %)	P value <sup>a</sup>	N (row %)	P value <sup>a</sup>	P value <sup>a</sup>
Other substance use disorder <sup>c</sup>			.27		.04		.21		.16			.05
Yes	51 (14.0%)	38 (74.5%)		32 (62.7%)		30 (58.8%)		35 (68.6%)		27 (52.9%)		
No	312 (86.0%)	208 (66.7%)		148 (47.4%)		154 (49.4%)		182 (58.3%)		120 (38.5%)		
Depressive disorder			.32		.51		.59		.37			.75
Yes	147 (40.5%)	104 (70.7%)		76 (51.7%)		77 (52.4%)		92 (62.6%)		61 (41.5%)		
No	216 (59.5%)	142 (65.7%)		104 (48.1%)		107 (49.5%)		125 (57.9%)		86 (39.8%)		
Anxiety disorder			.65		.66		.89		.49			.92
Yes	127 (35.0%)	88 (69.3%)		65 (51.2%)		65 (51.2%)		79 (62.2%)		51 (40.2%)		
No	236 (65.0%)	158 (66.9%)		115 (48.7%)		119 (50.4%)		138 (58.5%)		96 (40.7%)		
Current smoker			.05		.36		.71		.20			.97
Yes	22 (6.1%)	19 (86.4%)		13 (59.1%)		12 (54.5%)		16 (72.7%)		9 (40.9%)		
No	341 (93.9%)	227 (66.6%)		167 (49.0%)		172 (50.4%)		201 (58.9%)		138 (40.5%)		
Ever blunt use			<.001		<.001		<.001		<.001			<.001
Yes	70 (19.3%)	27 (38.6%)		16 (22.9%)		9 (12.9%)		18 (25.7%)		7 (10.0%)		
No	293 (80.7%)	219 (74.7%)		164 (56.0%)		175 (59.7%)		199 (67.9%)		140 (47.8%)		
Most friends get drunk/high <sup>b</sup>			.005		.01		<.001		.01			<.001
Yes	270 (74.4%)	194 (71.9%)		145 (53.7%)		152 (56.3%)		173 (64.1%)		124 (45.9%)		
No	88 (24.2%)	49 (55.7%)		34 (38.6%)		31 (35.2%)		43 (48.9%)		22 (25.0%)		
Substance of choice = cannabis			.98		.16		.15		.69			.66
Yes <sup>d</sup>	310 (85.4%)	210 (67.7%)		149 (48.1%)		162 (52.3%)		184 (59.4%)		127 (41.0%)		
No	53 (14.6%)	36 (67.9%)		31 (58.5%)		22 (41.5%)		33 (62.3%)		20 (37.7%)		
Intake year			.005		<.001		<.001		<.001			<.001
2017	60 (16.5%)	30 (50.0%)		17 (28.3%)		16 (26.7%)		23 (38.3%)		10 (16.7%)		
2018	105 (28.9%)	74 (70.5%)		61 (58.1%)		61 (58.1%)		70 (66.7%)		52 (49.5%)		

Patient characteristics	Study cohort (N=363)		Ever vaped nicotine and/or cannabis (N=246)		Current nicotine vaping (N=180)		Current cannabis vaping (N=184)		Any current vaping (nicotine and/or cannabis) (N=217)		Current vaping of both nicotine and cannabis (N=147)	
	N (column %)	N (row %)	N (row %)	P value <sup>a</sup>	N (row %)	P value <sup>a</sup>	N (row %)	P value <sup>a</sup>	N (row %)	P value <sup>a</sup>	N (row %)	P value <sup>a</sup>
2019	198 (54.5%)	142 (71.7%)	102 (51.5%)		107 (54.0%)		124 (62.6%)		85 (42.9%)			

<sup>a</sup>  $\chi^2$  test P values; bold indicates  $p < .05$

<sup>b</sup> One patient was missing income information, and five patients were missing information for friends' substance use.

<sup>c</sup> Other substance use disorders included disorders related to opioids; sedatives, hypnotics, or anxiolytics; cocaine; other stimulants; hallucinogens; inhalants; or other psychoactive substances; ICD-10 codes F11, F13–F16, and F18–F19 were used, excluding “in remission” codes.

<sup>d</sup> Some patients listed substances in addition to cannabis.

**Table 2.**

Adjusted odds ratios with 95% confidence intervals of ever and current vaping among adolescents in treatment for substance use disorders.<sup>a</sup>

	Ever vaped nicotine and/or cannabis	Current nicotine vaping	Current cannabis vaping	Any current vaping (nicotine and/or cannabis)	Current vaping of both nicotine and cannabis
Age (per year)	1.02 (0.82, 1.26)	1.05 (0.85, 1.29)	0.98 (0.79, 1.21)	1.08 (0.87, 1.33)	0.95 (0.76, 1.19)
Female (Ref. = male)	1.49 (0.85, 2.64)	1.00 (0.58, 1.71)	1.61 (0.92, 2.82)	1.72 (0.99, 3.00)	0.94 (0.53, 1.68)
Race/ethnicity (Ref. = non-Hispanic White)					
Asian	1.43 (0.56, 3.66)	1.89 (0.80, 4.44)	2.24 (0.90, 5.56)	1.66 (0.67, 4.12)	2.67 (1.10, 6.49)
Black	0.52 (0.21, 1.26)	<b>0.08 (0.02, 0.37)</b>	0.84 (0.31, 2.29)	0.47 (0.18, 1.20)	<b>0.12 (0.03, 0.60)</b>
Hispanic	0.69 (0.37, 1.28)	1.11 (0.61, 2.01)	0.86 (0.47, 1.58)	0.93 (0.51, 1.69)	1.03 (0.55, 1.93)
Other/Unknown	0.68 (0.25, 1.90)	0.75 (0.27, 2.09)	0.33 (0.10, 1.02)	0.72 (0.26, 2.00)	0.30 (0.08, 1.09)
Geocoded household income (Ref. = <80K)					
\$80K–120K	<b>2.05 (1.12, 3.74)</b>	<b>2.91 (1.57, 5.39)</b>	<b>2.34 (1.26, 4.32)</b>	<b>2.44 (1.35, 4.42)</b>	<b>3.34 (1.69, 6.62)</b>
>\$120K	<b>3.68 (1.79, 7.55)</b>	<b>7.42 (3.68, 14.95)</b>	<b>6.08 (2.99, 12.37)</b>	<b>5.79 (2.84, 11.77)</b>	<b>9.48 (4.49, 20.00)</b>
Alcohol use disorder	1.68 (0.79, 3.57)	<b>2.14 (1.06, 4.33)</b>	1.31 (0.64, 2.65)	1.71 (0.83, 3.50)	1.75 (0.85, 3.62)
Other substance use disorder <sup>b</sup>	1.14 (0.52, 2.49)	1.41 (0.67, 2.96)	1.28 (0.60, 2.71)	1.26 (0.59, 2.67)	1.49 (0.69, 3.21)
Current smoking	<b>3.95 (1.04, 14.95)</b>	1.67 (0.61, 4.58)	1.69 (0.60, 4.75)	2.39 (0.80, 7.11)	1.25 (0.44, 3.56)
Never blunt use	<b>3.89 (2.05, 7.37)</b>	<b>2.47 (1.21, 5.05)</b>	<b>8.68 (3.76, 20.03)</b>	<b>4.33 (2.21, 8.49)</b>	<b>5.78 (2.24, 14.91)</b>
Most friends get drunk/high	1.68 (0.94, 3.00)	1.39 (0.77, 2.53)	<b>1.87 (1.02, 3.42)</b>	1.40 (0.79, 2.50)	<b>2.01 (1.04, 3.90)</b>
Substance of choice = cannabis <sup>c</sup>	1.35 (0.65, 2.79)	0.79 (0.39, 1.61)	<b>2.36 (1.16, 4.81)</b>	1.25 (0.61, 2.54)	1.57 (0.75, 3.31)
Intake year (Ref. = 2017)					
2018	<b>2.18 (1.01, 4.69)</b>	<b>3.88 (1.73, 8.73)</b>	<b>3.53 (1.54, 8.07)</b>	<b>3.30 (1.51, 7.23)</b>	<b>4.73 (1.86, 12.00)</b>
2019	<b>2.71 (1.36, 5.40)</b>	<b>3.60 (1.72, 7.55)</b>	<b>4.21 (1.97, 9.02)</b>	<b>3.37 (1.67, 6.81)</b>	<b>5.38 (2.23, 12.95)</b>

<sup>a</sup>N=357; 6 patients were excluded due to missing geocoded household income (n=1) or friends' substance use (n=5). Bold indicates p < .05.

<sup>b</sup>Other substance use disorders included disorders related to opioids; sedatives, hypnotics, or anxiolytics; cocaine; other stimulants; hallucinogens; inhalants; or other psychoactive substances; ICD-10 codes F11, F13–F16, and F18–F19 were used, excluding “in remission” codes.

<sup>c</sup>Some patients listed substances in addition to cannabis.